

## **Title page**

Voice acoustics allow classifying autism spectrum disorder with high accuracy.

## **Authors**

Frédéric Briend, PhD<sup>1,2\*</sup> ; Céline David, PhD<sup>2</sup> ; Silvia Silleresi, PhD<sup>3</sup> ; Joëlle Malvy, MD, PhD<sup>1,2</sup> ; Sandrine Ferré, PhD<sup>1,2</sup> ; Marianne Latinus, PhD<sup>1,2,4\*</sup>

## **Affiliations**

<sup>1</sup> EXAC·T, Centre Universitaire de Pédopsychiatrie, CHRU de Tours, Tours, France

<sup>2</sup> UMR 1253, iBrain, Université de Tours, INSERM, 37000 Tours, France

<sup>3</sup> University of Milano-Bicocca, Department of Psychology, Milan, Italy

<sup>4</sup> Centro de Estudios en Neurociencia Humana y Neuropsicología. Facultad de Psicología. Universidad Diego Portales, Santiago, Chile

\* These authors contributed equally to this work

Corresponding Author: Marianne Latinus: [marianne.latinus@univ-tours.fr](mailto:marianne.latinus@univ-tours.fr)

Word count: 3881

## **Running title**

Voice acoustics as a potential feature classification for autism

**Keywords**

Voice; acoustic; vocal biomarker; autism spectrum disorder; diagnostic methods; clustering analyses.

## **Abstract**

Early identification of children on the autism spectrum is crucial for early intervention with long-term positive effects on symptoms and skills. The need for improved objective autism detection tools is emphasized by the poor diagnostic power in current tools. Here, we aim to evaluate the classification performance of acoustic features of the voice in children with autism spectrum disorder (ASD) with respect to a heterogeneous control group (composed of neurotypical children, children with Developmental Language Disorder [DLD] and children with sensorineural hearing loss with Cochlear Implant [CI]). This retrospective diagnostic study was conducted at the Child Psychiatry Unit of Tours University Hospital (France). A total of 108 children, including 38 diagnosed with ASD ( $8.5 \pm 0.25$  years), 24 typically developing (TD;  $8.2 \pm 0.32$  years) and 46 children with atypical development (DLD and CI;  $7.9 \pm 0.36$  years) were enrolled in our studies. The acoustic properties of speech samples produced by children in the context of a nonword repetition task were measured. We applied a ROC (Receiving Operator Characteristic) supervised k-mean clustering algorithm combined with a Monte Carlo cross-validation to develop a classification model that can differentially classify a child with an unknown disorder.

We showed that voice acoustics classified autism diagnosis with an overall accuracy of 91% [CI95%, 90.40%-91.65%] against TD children, and of 85% [CI95%, 84.5%-86.6%] against an heterogeneous group of non-autistic children. Accuracy reported here with multivariate analysis combined with Monte Carlo cross-validation is higher than in previous studies. Our findings demonstrate that easy-to-measure voice acoustic parameters could be used as a diagnostic aid tool, specific to ASD.

## **Introduction**

Autism spectrum disorder (ASD) is a class of prenatal neurodevelopmental disorders<sup>1</sup> defined by the co-occurrence of two main diagnostic criteria: a socio-emotional impairment and a behavioral deficit manifested by repetitive behaviors and interests<sup>2</sup>. Socio-emotional impairments affect both the production and perception of social signal. To this day, there is no reliable biomarker of ASD, and diagnosis is based on a pluri-disciplinary clinical assessment of the child<sup>3</sup>. Finding a more objective and automated marker of ASD could help in the diagnosis of ASD making it simpler and more reliable<sup>4</sup>. Atypical voice prosody is one of the earliest markers of ASD<sup>5-8</sup>, evaluated in diagnostic tools such as ADOS<sup>9</sup>; here, we asked whether easy-to-measure vocal acoustic features could be used as an objective ASD-specific marker to help diagnosis.

The human voice carries a wealth of information regarding a speaker, its physical characteristics, state of mind and health. From birth, the voice is used to signal information on well-being to surrounding adults, and infant cries are part of the preliminary assessment of neonates' health. Atypical acoustic cry features are associated with central nervous system dysfunction in human neonates<sup>10</sup> and rodent pups<sup>11</sup>. Voice production involves the entire brain and is under the influence of both autonomic and somatic nervous systems<sup>12</sup>. Voice production starts with breathing. The air coming from the lungs is sent towards the larynx, where it induces the vibration of the vocal folds. The vibration of the vocal folds produces a buzzing sound with a particular fundamental frequency, and associated harmonics. This sound is then modulated by its passage through the vocal tract airways. Breathing is normally considered an automatic process, but during speech it can be controlled voluntarily yet unconsciously<sup>12,13</sup>. Muscles in the larynx are controlled by two different branches of the vagal nerve: the recurrent laryngeal nerve (RLN) and the superior laryngeal nerve (SLN). The RLN controls muscles of the larynx

that allow opening, closing, and adjusting the tension of the vocal folds; the SLN allows changing the tension of the vocal folds, therefore increasing fundamental frequency. Articulation of the sound depends on the position of the different elements forming the vocal tract airways and are under voluntary control by the primary motor cortex [e.g.,<sup>12</sup>]. Autism spectrum disorder is characterized by impaired functioning of both somatic and autonomic nervous systems, and these impairments have consequences in their vocal production. Consistently, previous studies have reported differences in the acoustic properties of autistic individuals. Nonetheless results are often contradictory and inconsistent (see<sup>5</sup>). Pitch, measured as the fundamental frequency (F0) of speech sounds, has been reported to be higher in autism<sup>14-18</sup>, although many studies do not show this result<sup>19-22</sup>. Jitter, a measure of cycle-to-cycle regularity in F0, and shimmer, a measure of cycle-to-cycle variation of F0 amplitude, are reported to be smaller in autism with regards to neurotypical individuals<sup>16,21</sup>; these observations suggest a greater stability of voicing during speech productions in ASD. It should be noted however that other studies report, using different measures, increased pitch variation in autism<sup>14,15,22</sup>, or lack of differences in pitch variability<sup>23</sup>. Studies of vocal tract features analysis also report discrepant results with higher formant frequencies<sup>18</sup> or a smaller formantic dispersion<sup>16</sup>. These data converge into demonstrating that there is something special in the voice of autistic individuals, that could help in diagnosis. Yet, univariate analysis of specific acoustic measures may not be powerful enough. Here, we describe a multivariate analysis of vocal acoustic parameters combined with machine learning techniques to develop potential tools to aid autism diagnosis.

Machine learning techniques are increasingly used for medical diagnosis, especially clustering which is a powerful tool for detecting patterns in datasets. Several studies have used clustering methods in order to develop diagnostic biomarkers of various pathologies in animal models<sup>4,24,25</sup> and in human trials<sup>26-28</sup>. A classical, non-supervised, and robust clustering

algorithm, the k-Means clustering algorithm<sup>29</sup>, also known as nearest centroid classifier when used in machine learning, yields high discriminating power to diagnose a single unknown subject in a given disorder state<sup>25,27</sup>. Here, we used voice acoustics (Fig. 1) as the selected features included in this common classifier to evaluate their classification performance of ASD relative to typical development and other pathologies. We evaluated the classification performance of voice acoustics in comparison not only to TD children (study 1) but also to children with other disorders sharing common deficits with ASD: sensorineural hearing loss and Developmental Language Disorder (DLD, study 2). These two pathologies were chosen due to observed commonalities in the language domain between children with DLD and autism<sup>30</sup> and between children with sensorineural hearing loss and with cochlear implants (CI) and autism<sup>31</sup>. The acoustic properties of speech samples produced by autistic children in the context of a nonword repetition task (NRT)<sup>32</sup> were examined. Data were analyzed with unsupervised and ROC (Receiving Operator Characteristic) supervised clustering algorithm.

## **Methods and Materials**

### **Participants**

One hundred and eight children were enrolled in our retrospective studies. Study 1 (Fig. 2a) is composed of 38 children on the autism spectrum (1 girl;  $8.5 \pm 0.25$  years) and 24 TD children (12 girls;  $8.2 \pm 0.32$  years), and Study 2 (Fig. 3a) additionally includes 21 children with DLD (9 girls;  $7.9 \pm 0.51$  years) and 25 children displaying severe-to-profound sensorineural hearing loss fitted with CIs (8 girls;  $8 \pm 0.22$  years). Data of 24 children were excluded from the analysis (see Experimental protocol and data acquisition). Therefore, the final sample comprised 84 children distributed as follows: 29 ASD (0 girl;  $8.4 \pm 0.29$  years; age range [6.3 12]; ADOS severity score:  $6.19 \pm 0.45$ ; CARS:  $27.7 \pm 0.7$ ), 20 TD (10 girls;  $7.99 \pm 0.33$  years; age range [6 10.5]), 20 CI (6 girls;  $8.2 \pm 0.19$  years; age range [6.5 9.9]); 12 with bilateral CI; 6 with right

CI; 2 with left CI; age at first implantation  $1.86 \pm 0.15$ ) and 15 DLD (7 girls;  $8.2 \pm 0.37$  years; age range [6.5 10.8]). Demographic and clinical information regarding the final sample is presented in Table 1. Youth with ASD received an expert clinical diagnosis based on Diagnostic and Statistical Manual of Mental Disorders – fifth Edition – (DSM-V)<sup>33</sup>; the Autism Diagnostic Interview-Revised<sup>34</sup>, and/or the Autism Diagnostic Observation Schedule<sup>35</sup> were used by experienced clinicians of the Excellence Center of Autism (Exac-t), Tours, France to inform diagnostic decisions. Children with DLD also received an expert clinical diagnosis based on the DSM-V<sup>33</sup> Nonverbal cognitive abilities were assessed either by Raven Progressive Matrices or Block Design and Matrix Reasoning of the WISC-IV (data of 5 TD children are missing). Only children with a minimum Mean Length of Utterances of 2.5 were included in the study<sup>36</sup> to ensure that language tests could be administered.

This study was carried out in accordance with the recommendations of the local ethics committee (Comité de Protection des Personnes [CPP] Tours Ouest 1, n°2006-RS), with written informed consent from all parents of the children and assent from the children, in accordance with the Declaration of Helsinki.

### **Experimental protocol and data acquisition**

Acoustic data were extracted from 20 speech samples recorded in the context of a nonword repetition task<sup>32</sup>, therefore reducing the influence of social interaction in voice production. The task focuses on complex phonological structures that have been identified as the source of errors in children with impaired phonology. Briefly, children had to repeat 50 or 70 nonwords of varying phonological complexity, presented with a computer either with only auditory or with both audio and visual information. Nonwords were created using 1, 2 or 3 of the three most common vowels among the languages of the world, namely [a], [i], [u], and from a concise list of consonants which included two stops ([k], [p]), two fricatives ([f], [s]), one liquid [l].

Nonwords had a maximum of 3 syllables to limit the influence of working memory on the repetition task. Nonwords had different syllable complexity: either a simple consonant vowel syllables, syllables with a final consonant or syllables with initial and median consonant clusters<sup>32</sup>. Phonological analysis of the data presented in the current manuscript are published elsewhere<sup>37,38</sup>. Among the 50 or 70 nonwords, the 20 ones with less phonological errors were chosen for acoustical analysis. The NWR took place in a quiet room and audio were digitally recorded using Zoom H4 microphones put on a table in front of the child. Overall performance in the NWR tasks is presented in the results section and was analyzed with a 1 factor Welch ANOVA. Acoustic parameters were analyzed using the open-source software Praat<sup>39</sup>. For each nonword, we extracted 9 acoustics parameters (Fig. 1): mean fundamental frequency (f0), mean formant frequencies (F1 to F4), mean formant dispersion (FD), mean harmonic-to-noise ratio (HNR), mean jitter (cycle-to-cycle variation in frequency of f0) and mean shimmer (cycle-to-cycle variation in intensity of f0); they were then averaged across the 20 nonwords. In addition, because ASD is characterized with increased intra-individual variability<sup>22,40,41</sup>, shape parameters (e.g., skewness and kurtosis) of f0, FD, HNR, jitter and shimmer were computed using Matlab2018b functions, leading to 19 variables (Fig. 1). Note that in the Matlab kurtosis function, the normal distribution has a kurtosis value of 3 (Fig. 1e).

Acoustic data are excluded according to two categories of rejection criteria: the nonword repetition task performance and acoustic rejection. For the first criteria, children whose performance in the repetition of vowels was considered outlier ( $[Q1-1.5 \times IQR]$  with Q1: lower quartile and IQR: interquartile range) were removed from the analysis (N = 9: 2 ASD, 2 DLD, 5 CI), to avoid bias due to the mispronunciation of certain vowels which can influence acoustics. For the second criteria, based on acoustical analysis (recording quality or outlier value of acoustic parameters with respect to the population), another 15 children (7 ASD; 4 DLD; 4 TD) were excluded from the analysis (ROC-supervised k-means classification results



including all participants but those with poor recordings quality are shown in Supplementary table 2).

### **Development of clustering diagnostic model**

Our goal was to determine if acoustic features of the voice could be used as a feature classification specific for autism by k-means classifying ASD against typical and other atypical development, thus we randomly dichotomized data in a diagnostic model group (train set) and an unknown data group (test set). In order to validate our model performance, we used Monte Carlo cross-validation. Note that this method is robust to imbalance gender across groups; indeed, with a clustering approach, if gender was an important factor the two identified clusters would reflect gender separation rather than diagnostic group.

To develop the diagnostic model, we randomly selected  $n$  ASD and  $n$  control as train data (70% of data), and then we applied k-means (50 iterations, Hartigan & Wong algorithm); this was repeated 500 times with random subsampling of the data from the entire population, e.g., Monte Carlo cross-validation. Because there is no general rule regarding the number of repetitions to use, we choose the value at which our main criteria (selectivity and sensitivity) appear stable beyond reasonable doubt, through multiple testing with different numbers of resampling (Supplementary Fig. 1). The number of clusters was set to two, since we aimed to determine ASD diagnostic against a control population (TD children only, or control children). We performed k-means clustering analysis (KCA) in an unsupervised way with the nine acoustic and derived acoustic variables ( $N = 19$ ) and assessed its performance. Then, in order to enhance our KCA, we performed receiver operating characteristic (ROC) as proposed by Nikas and colleagues<sup>25</sup> and used an AUC (Area Under the Curve) ROC curve, as measure of separability to evaluate the most discriminative acoustic parameters. This latter probability is an assessment of the discriminative power of a given variable with respect to two measures, here the two

groups involved. For example, with a given variable, an AUC of 1 is synonym of a separation between groups with 100% accuracy, and the given variable is considered as an excellent classifier. On the contrary, the worst discrimination between the two groups has an AUC = 0.50 (i.e., no discrimination capacity). In this way, the ROC curve allows us to optimize our KCA by supervising it using acoustic variables with the best discriminative performance. We used a threshold of AUC > 0.80 (80%) corresponding to a good discrimination <sup>42</sup>.

This model was then tested to identify the diagnostic group of the test data (30% of the entire data, corresponding to the data not used in model building) according to their KCA classification, always for each of the 500 bootstrap replications. To realize this, test data were added one-by-one for each participant and classified by supervised and unsupervised KCA. Hence, diagnostic of the participant was classified based on its data; accuracy was measured as the number of times a test participant was classified with the correct diagnostic, referred to as classification performance.

To assess the performance of our KCA, we measured selectivity, sensitivity and the classification performance of our model. Moreover, by the goodness of fit as quality criteria of KCA, we assess clustering effectiveness by a percent of variation (PV) corresponding to the total within-cluster sum of squares by the total of within and between-cluster sum of squares.

## **Results**

### **Classification of autistic children with respect to TD children**

Overall performance in the NWR task, measured on the entire set of items, differed between TD children (95.6%; performance range [84 100]) and children with ASD (80.4%; performance range [28 100]; Welch ANOVA:  $F(1, 31.7) = 22.3$ ;  $p < 0.001$ ).

To investigate the discriminative power of voice acoustics between autistic children and TD children, we performed K-means clustering analysis (KCA). Data were randomly split into a train group (N = 33; 20 ASD, 13 TD) and a test group (unknown data; N = 16; 9 ASD; 7 TD); this was done 500 times. Unsupervised KCA with the 19 acoustics variables was conducted on the train group, and the model was cross validated using repeated random sub-sampling using unknown data from the test group. We observed a percent of variation (PV, the dispersion between the two clusters; see Methods) of  $83.20\% \pm 1.33$ , a sensitivity of  $0.74 \pm 0.11$  and a specificity of  $0.92 \pm 0.12$  in the training group. The unsupervised KCA correctly classified 73.1% [71.8% 74.4%] of ASD and 92.3% [91.4% 93.1%] of TD children (Fig. 2b).

Our goal was to find optimum KCA settings, which best separate the ASD and the TD group to develop classification or diagnostic model. Therefore, we performed ROC curve analysis, to conduct ROC-supervised KCA<sup>25,43</sup> on training and test data with 500 bootstrap replications. The four most discriminant (Area Under the Curve [AUC] > 80%) acoustics parameters according to ROC analysis on the training group were mean F1, mean HNR, mean shimmer and jitter skewness. The ROC-supervised KCA setting yielded a considerable improvement over unsupervised KCA: as shown in Fig. 2, the ROC-supervised KCA had a PV of  $63.85\% \pm 3.00$ , a sensitivity of  $0.89 \pm 0.06$  and a specificity of  $0.94 \pm 0.10$ ; it classified correctly 89% [88.1% 90.0%] of ASD and 93% [92.7% 94.3%] of the TD group. ROC supervision allows decreasing false negatives.

### **Classification of autistic children with respect to a control population**

Overall performance in the NWR task, measured on the entire set of items, was affected by diagnostic group (Welch ANOVA:  $F(3, 33.3) = 62.1$ ;  $p < .001$ ). It was higher for TD children than all other groups (all pairwise comparisons  $p < 0.001$ ). Autistic children performed better

than SLI (49.3% [6 76]) and IC children (42.3% [10 80]); all pairwise comparisons  $p < 0.001$ ), who did not differ.

Next, we evaluated the classification performance of voice acoustics in comparison not only to TD children but also to children with other disorders sharing common deficits with ASD: sensorineural hearing loss and Developmental Language Delay. As previously described, we conducted a ROC-supervised KCA on the data of all participants, considering children with DLD, children with CI and TD children in the same group of heterogeneous control group (CTRL). See Supplementary table 1 for details about the unsupervised KCA.

Two acoustic parameters discriminated ASD from CTRL children according to ROC analysis: mean shimmer and F1, with an AUC, respectively of 85.23% and 82.36% (the separation is depicted in the Fig. 3c). The ROC-supervised KCA had a PV of  $57.42\% \pm 3.04$ , a sensitivity of  $0.86 \pm 0.05$  and a specificity of  $0.84 \pm 0.08$ ; it classified correctly 85.56% [84.5%-86.6%] of ASD and 84.2% [83.5%-85.0%] of the CTRL group. More specifically, in this latter heterogeneous population, 68.3% [66.8%-70.5%] of DLD, 85.6% [84.4%-86.8%] of CI and 93.9% [93.1%-94.7%] of TD were correctly classified (Fig. 3b).

## **Discussion**

### **Voice as clustering diagnostic approach?**

As classifier, the ROC-supervised KCA analysis, with classification performance around 90%, had an extremely high classification performance when separating ASD from TD children above previously reported classification value (between 80 and 89%<sup>4,6,22</sup>). Moreover, our method proves robust and reliable in discriminating autistic children from children without ASD, including other disorders (84%).

Importantly, acoustic factors predictive of autism diagnosis are mainly ones related to control of the vocal folds' vibrations (e.g., jitter, shimmer) rather than the  $f_0$  *per se*<sup>5</sup>, consistent with clinical description of a peculiar voice quality in autism and previous observations<sup>23</sup>. The pattern characteristics of autism, with respect to TD children, was lower average F1, higher HNR, higher shimmer and lower jitter skewness. A lower jitter skewness reflected a more normal distribution of jitter across nonwords, consistent with the observation of a greater stability in voice production<sup>16</sup>; TD children presented positively skewed and less tailed distribution, highlighting that most vocal sounds had similar shimmer and jitter. Note that we found that children on the autism spectrum have a higher shimmer than NT children, contrarily to what was reported in adults<sup>16</sup>, highlighting differences in the maturation of the vocal apparatus. A higher HNR suggested that vocal sounds of children on the autism spectrum are overall less noisy than those produced by TD children. The lower F1 in the ASD group did not reflect gender balance differences across groups as in the TD group male and female children had similar F1 values (938 Hz and 937 Hz). Note that when including the other pathologies, only mean F1 and mean shimmer remained classification features. Average F1 values were at the minimum 100 Hz lower in autistic children than in the other children; this is unlikely explained by gender imbalance as the second lowest F1 was observed for female (815 Hz) of the CI group, and F1 was the largest in male of the same group. F1 frequency is related to the length of the vocal tract<sup>44</sup> and tend to decrease with age; a lower F1 could reflect either an accelerated maturation of the vocal tract or differences in cranio-facial anatomy<sup>45</sup> and the presence of increased minor physical anomalies in autistic children<sup>46</sup>. Shimmer, which is a measure of cycle-to-cycle variation in amplitude of the  $f_0$ , presented an increased value of almost 16% in autism; although discriminant as in Guo et al.,<sup>6</sup> the opposite result was found in children speaking mandarin. Shimmer differences could reflect morphological differences or differences in control of the vocal cords of autistic children and other children<sup>47</sup>. Therefore,

these voice features could be an external marker of atypical neurodevelopment occurring before birth <sup>1</sup>. Note that in the current study we aimed to test whether voice could be used as a classification tool, and therefore we tested older children with stable diagnostic. Future studies should aim at studying the classification performance of infant's cries at birth or within the first year of life to test the validity of voice acoustics as a true biomarker of autism.

Previous studies that compared ASD to other populations are sparse and rely on different grouping strategies: Oller et al., <sup>48</sup> reported 62% accuracy in the classification of ASD children with DLD in a non-TD group, Bone et al., <sup>49</sup> reported a 78% of correct classification between ASD and DLD. Here, classifying children with DLD in the typically developing group, we obtained high classification rates of ASD not only with respect to TD, but also to other pathologies. The approach developed here combining feature selection, through ROC-supervision, a clustering analysis and Monte Carlo cross-validation demonstrates that voice features have a strong, specific, diagnosis power for ASD: accuracy was well-above chance for children with DLD and children with CI. This provides new information on the classifying power of voice features in ASD, in relation to other neurodevelopmental disorders in particular (e.g., DLD).

Central nervous system dysfunction affects vocal folds and by domino individual's voice. This is why automated voice analysis using recordings of patient speech is increasingly being used in psychiatry <sup>50</sup> and neurology as digital biomarkers of disease (i.e., in Major depressive disorder <sup>51</sup>, schizophrenia <sup>52</sup>, Parkinson's disease <sup>53</sup>, Alzheimer's Disease <sup>54</sup>, ...). However, this computational method should not be delegated solely to machines <sup>55</sup>, even if it is based on formal reasoning, this method should be used in complementarity to clinical diagnosis of experts.

This clustering method is a first step towards the development of an early diagnostic biomarker specific to ASD. Indeed, in this study, all children had had at some minimum, strong verbal

capabilities and data were selected to have the most optimum dataset; future studies should assess the classification performance of vocal acoustic based on non-linguistic vocal samples acquired in less controlled environment. In addition, data presented here comes from children between 6 and 12 while, in high-income countries the average age of ASD diagnosis is around age 4 <sup>56</sup>, and around 5 worldwide <sup>57</sup>. To be truly a biomarker of autism and understand its potential diagnostic value, these results should be replicated in younger children and possibly using cry features of babies. Moreover, in order to develop a sensitive diagnosis test, future works should include typical cases met in clinical practice, with disorders more often seen as comorbidities of ASD such as attention deficit hyperactivity disorder (ADHD), motor problems without social impairment, severe anxiety, and other behavior disorders <sup>56</sup>.

## **Conclusion**

Overall, our work suggests that easy-to-measure voice features, potentially linked to abnormal early neurodevelopment, can help in the diagnosis of autism spectrum disorder. Voice features in supervised clustering methods can be used as a potential feature classification for autism and paves the way to a new objective tool to aid clinical and differential diagnosis of ASD. The method developed here is in part automated, and in the future, a hand-in tool should be developed to automatically output diagnostic information. Early detection of ASD is crucial because it is likely to lead to an improved outcome. Thus, based on our simple clustering algorithm method, future work should investigate the acoustic cry features of baby as a potential biomarker for autism.

## **Author contributions**

S.F., M.L. designed the study. S.S., C.D., J.M. acquired the data. M.L. managed and coordinated the research activity planning and execution. F.B., S.F., S.S., C.D., M.L. analyzed the data. F.B. and M.L. wrote the original draft. All authors reviewed and edited the manuscript.

## Acknowledgments

The authors would like to thank all of the participants and their families for the time they spent to participate in this study, the staff of the Child Psychiatry Department of Tours University and the Neurofunctional Psychiatric Team at The University of Tours.

## Disclosure

### Funding

This work was supported by a grant from the Region Centre (Doctoral Grant awarded to Silvia Silleresi and Céline David) and the Fondation pour l'Audition.

### Competing interests

The authors declare no competing interests.

## References

- 1 Bonnet-Brilhault F, Rajerison TA, Paillet C, Guimard-Brunault M, Saby A, Ponson L *et al.* Autism is a prenatal disorder: Evidence from late gestation brain overgrowth. *Autism Res Off J Int Soc Autism Res* 2018; **11**: 1635–1642.
- 2 Lord C, Elsabbagh M, Baird G, Veenstra-Vanderweele J. Autism spectrum disorder. *Lancet Lond Engl* 2018; **392**: 508–520.
- 3 Bonnet-Brilhault F, Tuller L, Prevost P, Malvy J, Zebib R, Ferré S *et al.* A strategic plan to identify key neurophysiological mechanisms and brain circuits in autism. *J Chem Neuroanat* 2018; **89**: 69–72.
- 4 Rybner A, Jessen ET, Mortensen MD, Larsen SN, Grossman R, Bilenberg N *et al.* Vocal markers of autism: Assessing the generalizability of machine learning models. *Autism Res* 2022; **15**: 1018–1030.
- 5 Fusaroli R, Lambrechts A, Bang D, Bowler DM, Gaigg SB. Is voice a marker for Autism spectrum disorder? A systematic review and meta-analysis. *Autism Res Off J Int Soc Autism Res* 2017; **10**: 384–407.
- 6 Guo C, Chen F, Chang Y, Yan J. Applying Random Forest classification to diagnose autism using acoustical voice-quality parameters during lexical tone production. *Biomed Signal Process Control* 2022; **77**: 103811.




- 7 Kanner L. Autistic disturbances of affective contact. *Nerv Child* 1943; **2**: 217–250.
- 8 Paul R, Augustyn A, Klin A, Volkmar FR. Perception and production of prosody by speakers with autism spectrum disorders. *J Autism Dev Disord* 2005; **35**: 205–220.
- 9 Lord C, Risi S, Lambrecht L, Cook EH, Leventhal BL, DiLavore PC *et al*. The autism diagnostic observation schedule-generic: a standard measure of social and communication deficits associated with the spectrum of autism. *J Autism Dev Disord* 2000; **30**: 205–223.
- 10 LaGasse LL, Neal AR, Lester BM. Assessment of infant cry: acoustic cry analysis and parental perception. *Ment Retard Dev Disabil Res Rev* 2005; **11**: 83–93.
- 11 Scattoni ML, Crawley J, Ricceri L. Ultrasonic vocalizations: a tool for behavioural phenotyping of mouse models of neurodevelopmental disorders. *Neurosci Biobehav Rev* 2009; **33**: 508–515.
- 12 Kreidman J, Sidtis D. The Brain Behind the Voice: Cerebral Models of Voice Production and Perception. In: *Foundations of Voice Studies*. John Wiley & Sons, Ltd, 2011, pp 189–236.
- 13 Maclarnon A, Hewitt G. Increased breathing control: Another factor in the evolution of human language. *Evol Anthropol Issues News Rev* 2004; **13**: 181–197.
- 14 Filipe MG, Frota S, Castro SL, Vicente SG. Atypical prosody in Asperger syndrome: perceptual and acoustic measurements. *J Autism Dev Disord* 2014; **44**: 1972–1981.
- 15 Sharda M, Subhadra TP, Sahay S, Nagaraja C, Singh L, Mishra R *et al*. Sounds of melody--pitch patterns of speech in autism. *Neurosci Lett* 2010; **478**: 42–45.
- 16 Kissine M, Geelhand P. Brief Report: Acoustic Evidence for Increased Articulatory Stability in the Speech of Adults with Autism Spectrum Disorder. *J Autism Dev Disord* 2019; **49**: 2572–2580.
- 17 Bornstein MH, Costlow K, Truzzi A, Esposito G. Categorizing the cries of infants with ASD versus typically developing infants: A study of adult accuracy and reaction time. *Res Autism Spectr Disord* 2016; **31**: 66–72.
- 18 Mohanta A, Mittal VK. Analysis and classification of speech sounds of children with autism spectrum disorder using acoustic features. *Comput Speech Lang* 2022; **72**: 101287.
- 19 McAlpine A, Plexico L, Plumb A, Cleary J. Prosody in Young Verbal Children With Autism Spectrum Disorder. *Contemp Issues Commun Sci Disord* 2014; **41**: 120–132.
- 20 Brisson J, Martel K, Serres J, Sirois S, Adrien J-L. Acoustic Analysis of Oral Productions of Infants Later Diagnosed with Autism and Their Mother. *Infant Ment Health J* 2014; **35**: 285–295.
- 21 Diehl JJ, Watson D, Bennetto L, Mcdonough J, Gunlogson C. An acoustic analysis of prosody in high-functioning autism. *Appl Psycholinguist* 2009; **30**: 385–404.

- 22 Bonnef YS, Levanon Y, Dean-Pardo O, Lossos L, Adini Y. Abnormal speech spectrum and increased pitch variability in young autistic children. *Front Hum Neurosci* 2011; **4**: 237.
- 23 Bone D, Lee C-C, Black MP, Williams ME, Lee S, Levitt P *et al*. The Psychologist as an Interlocutor in Autism Spectrum Disorder Assessment: Insights From a Study of Spontaneous Prosody. *J Speech Lang Hear Res JSLHR* 2014; **57**: 1162–1177.
- 24 Cohen H, Zohar J, Matar MA, Kaplan Z, Geva AB. Unsupervised Fuzzy Clustering Analysis Supports Behavioral Cutoff Criteria in an Animal Model of Posttraumatic Stress Disorder. *Biol Psychiatry* 2005; **58**: 640–650.
- 25 Nikas JB, Low WC. Application of clustering analyses to the diagnosis of Huntington disease in mice and other diseases with well-defined group boundaries. *Comput Methods Programs Biomed* 2011; **104**: e133–e147.
- 26 Tokuda T, Yamashita O, Sakai Y, Yoshimoto J. Clustering of Multiple Psychiatric Disorders Using Functional Connectivity in the Data-Driven Brain Subnetwork. *Front Psychiatry* 2021; **12**.
- 27 Alashwal H, El Halaby M, Crouse JJ, Abdalla A, Moustafa AA. The Application of Unsupervised Clustering Methods to Alzheimer’s Disease. *Front Comput Neurosci* 2019; **13**.
- 28 Trevithick L, Painter J, Keown P. Mental health clustering and diagnosis in psychiatric inpatients. *BJPsych Bull* 2015; **39**: 119–123.
- 29 Forgy E. Cluster analysis of multivariate data : efficiency versus interpretability of classifications. *Biometrics* 1965; **21**.
- 30 Georgiou N, Spanoudis G. Developmental Language Disorder and Autism: Commonalities and Differences on Language. *Brain Sci* 2021; **11**: 589.
- 31 Robertson J. Children with cochlear implants and autism – challenges and outcomes: The experience of the National Cochlear Implant Programme, Ireland. *Cochlear Implants Int* 2013; **14**: S11–S14.
- 32 Dos Santos C, Ferré S. A Nonword Repetition Task to Assess Bilingual Children’s Phonology. *Lang Acquis* 2018; **25**: 58–71.
- 33 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM5*. American Psychiatric Publication Incorporated. Washington DC, 2013.
- 34 Lord C, Storoschuk S, Rutter M, Pickles A. Using the ADI-R to diagnose autism in preschool children. *Infant Ment Health J* 1993; **14**: 234–252.
- 35 Lord C, Rutter M, Goode S, Heemsbergen J, Jordan H, Mawhood L *et al*. Autism diagnostic observation schedule: A standardized observation of communicative and social behavior. *J Autism Dev Disord* 1989; **19**: 185–212.


- 36 Silleresi S, Tuller L, Delage H, Durrleman S, Bonnet-Brilhault F, Malvy J *et al.* Sentence repetition and language impairment in French-speaking children with ASD. In: Gavarro A (ed). *On the acquisition of the syntax of Romance*. John Benjamins Publishing Company: Amsterdam, 2018, pp 235–258.
- 37 David C, Tuller L, Schweitzer E, Lescanne E, Bonnet-Brilhault F, Gomot M *et al.* Does Phonological Complexity Provide a Good Index of Language Disorder in Children With Cochlear Implants? *J Speech Lang Hear Res JSLHR* 2021; **64**: 4271–4286.
- 38 Silleresi S, Prevost P, Zebib R, Bonnet-Brilhault F, Conte D, Tuller L. Identifying Language and Cognitive Profiles in Children With ASD via a Cluster Analysis Exploration: Implications for the New ICD-11. *Autism Res* 2020. doi:10.1002/aur.2268.
- 39 Boersma P. Praat, a system for doing phonetics by computer. *Glott Int* 2002; **5**.<https://dare.uva.nl/search?arno.record.id=109185> (accessed 8 Dec2020).
- 40 Latinus M, Mofid Y, Kovarski K, Charpentier J, Batty M, Bonnet-Brilhault F. Atypical sound perception in ASD explained by inter-trial (in)consistency in EEG. *Front Psychol* 2019; **10**: 1177.
- 41 Milne E. Increased intra-participant variability in children with autistic spectrum disorders: evidence from single-trial analysis of evoked EEG. *Front Psychol* 2011; **2**: 51.
- 42 Hosmer DW, Lemeshow S, Sturdivant RX. Application of Logistic Regression with Different Sampling Models. In: *Applied Logistic Regression*. John Wiley & Sons, Ltd, 2013, pp 227–242.
- 43 Vivre avec un enfant autiste. Fr. 2 Soirée Débat. 2020.
- 44 Vorperian HK, Kent RD. Vowel Acoustic Space Development in Children: A Synthesis of Acoustic and Anatomic Data. *J Speech Lang Hear Res JSLHR* 2007; **50**: 1510–1545.
- 45 Tripi G, Roux S, Matranga D, Maniscalco L, Glorioso P, Bonnet-Brilhault F *et al.* Cranio-Facial Characteristics in Children with Autism Spectrum Disorders (ASD). *J Clin Med* 2019; **8**. doi:10.3390/jcm8050641.
- 46 Tripi G, Roux S, Carotenuto M, Bonnet-Brilhault F, Roccella M. Minor Neurological Dysfunctions (MNDs) in Autistic Children without Intellectual Disability. *J Clin Med* 2018; **7**. doi:10.3390/jcm7040079.
- 47 Teston B. L'évaluation instrumentale des dysphonies. Etat actuel et perspectives. In: *Le bilan d'une dysphonie*. 2014, pp 105–169.
- 48 Oller DK, Niyogi P, Gray S, Richards JA, Gilkerson J, Xu D *et al.* Automated vocal analysis of naturalistic recordings from children with autism, language delay, and typical development. *Proc Natl Acad Sci* 2010; **107**: 13354–13359.

- 49 Bone D, Chaspari T, Audhkhasi K, Gibson J, Tsiartas A, Segbroeck MV *et al.* Classifying language-related developmental disorders from speech cues: the promise and the potential confounds. In: *INTERSPEECH*. 2013.
- 50 Fagherazzi G, Fischer A, Ismael M, Despotovic V. Voice for Health: The Use of Vocal Biomarkers from Research to Clinical Practice. *Digit Biomark* 2021; **5**: 78–88.
- 51 Zhang L, Duvvuri R, Chandra KKL, Nguyen T, Ghomi RH. Automated voice biomarkers for depression symptoms using an online cross-sectional data collection initiative. *Depress Anxiety* 2020; **37**: 657–669.
- 52 Parola A, Simonsen A, Bliksted V, Fusaroli R. Voice patterns in schizophrenia: A systematic review and Bayesian meta-analysis. *Schizophr Res* 2020; **216**: 24–40.
- 53 Tracy JM, Özkanca Y, Atkins DC, Hosseini Ghomi R. Investigating voice as a biomarker: Deep phenotyping methods for early detection of Parkinson’s disease. *J Biomed Inform* 2020; **104**: 103362.
- 54 König A, Satt A, Sorin A, Hoory R, Toledo-Ronen O, Derreumaux A *et al.* Automatic speech analysis for the assessment of patients with pre dementia and Alzheimer’s disease. *Alzheimers Dement Diagn Assess Dis Monit* 2015; **1**: 112–124.
- 55 Gross F. The Impact of Formal Reasoning in Computational Biology. In: *A Critical Reflection on Automated Science - Will Science Remain Human?* Springer, 2020, p 157.
- 56 Sheldrick RC, Maye MP, Carter AS. Age at First Identification of Autism Spectrum Disorder: An Analysis of Two US Surveys. *J Am Acad Child Adolesc Psychiatry* 2017; **56**: 313–320.
- 57 van ’t Hof M, Tisseur C, van Berckeleer-Onnes I, van Nieuwenhuyzen A, Daniels AM, Deen M *et al.* Age at autism spectrum disorder diagnosis: A systematic review and meta-analysis from 2012 to 2019. *Autism Int J Res Pract* 2021; **25**: 862–873.


## Figure and their legends



**Fig. 1. Experimental design.** **a**, Spectrogram of one of the nonwords produced in the nonword repetition task. **b**, Average power spectrum. **c**, Amplitude waveform. **d**, Zoom on the amplitude waveform to illustrate shimmer and jitter. **e**, Distribution of mean  $f_0$  measured in the 20 selected nonwords for a skewness (left panel; skewness = 0), and kurtosis (middle panel; kurtosis normalize = 0) corresponding to a normal distribution and for altered (right panel) skewness (3.1) and kurtosis (9.5).



**Fig. 2. Classification of autistic children with respect to TD children (study 1).** **a**, Procedure of inclusion of the participants and random dichotomization of the data in diagnostic model group and unknown data group. NRT: Nonword Repetition Task; ASD: Autism Spectrum Disorder; TD: typically developing children. **b**, Extrapolation of the classification on 100 subjects. Participants with good diagnosis are surrounded by a green rectangle (93%), the misclassified by a red one. **c**, Illustration of the acoustic profile by radar chart according to the four most significant voice features, namely, harmonic-to-noise ratio (HNR), formant frequencies 1 (F1), skewness of Jitter [s(Jitter)] and Shimmer generated the best ROC-supervised KCA setting.



**Fig. 3. Classification of autistic children with respect to a diverse control population (study**

**2).** **a,** Procedure of inclusion of the participants and random dichotomization of the data in diagnostic model group and unknown data group. OD: Other Disorders; NRT: Nonword Repetition Task; ASD: Autism Spectrum Disorder; CTRL: heterogeneous control group (composed of children with developmental language disorder [DLD] and cochlear implant [CI]). **b,** Extrapolation of the classification on 100 subjects. Participants with good diagnosis are surrounded by a green rectangle (84%), the misclassified by a red one. **c,** The top two most significant voice features, namely, formant frequencies 1 (F1) and Shimmer generated the best ROC-supervised KCA setting. Those three parameters are plotted against each other.

## **Data availability**

All data generated and/or analyzed during this study are available from the corresponding author (ML) on reasonable request. For all clustering runs, we used R (<http://cran.r-project.org>).

Code used in this manuscript is available on GitHub (<https://github.com/FredericBr/VoiceMarker>).



**Table**

	N	Male / Female	Age (years)	FRI $\pm$ sem (Percentile)	RPM $\pm$ sem (Percentile)	NRT Score $\pm$ sem (%)
<i>ASD</i>	29	29 / 0	8.42 $\pm$ 0.29	88.6 $\pm$ 5.56 (22)	50.7 $\pm$ 13.8 (7)	80 $\pm$ 3.1
<i>TD</i>	20	10 / 10	7.99 $\pm$ 0.33	121.7 $\pm$ 4.43 (15)		96 $\pm$ 0.92
<i>DLD</i>	15	8 / 7	8.15 $\pm$ 0.27	93 $\pm$ 3.14 (11)	36.9 $\pm$ 10.7 (4)	49 $\pm$ 6.3
<i>CI</i>	20	14 / 6	8.16 $\pm$ 0.16	96.6 $\pm$ 4.26 (20)		42 $\pm$ 4.6

**Table 1: Demographic and clinical information of the sample.** Numbers in brackets represent the number of participants contributing data to the measure. FRI: Fluid Reasoning Index calculated with the prorated sum of Block Design and Matrix Reasoning scores in WISC-IV; RPM: Raven's Progressive Matrices; NRT: Nonword Repetition Task; ASD: Autism Spectrum Disorder; TD: Typically developing children; DLD: children with Developmental language disorder and CI: children with Cochlear Implant.

## **Supplemental information**

**Table S1: Summary of the results of N = 500 KCA.** Each clustering method was tested both in an unsupervised and ROC-supervised way. The percent of variation of a clustering method assesses the clustering fitting of that method in a given setting. ASD: Autism Spectrum Disorder participants; CTRL: heterogeneous control group (composed of children with developmental language disorder [DLD] and cochlear implant [CI]); CI95%: Confidence interval of 95%. Data are presented as mean (Standard Deviation).

**Table S2: Summary of the results of N = 500 KCA after acoustic rejection.** Each clustering method was tested both in an unsupervised and ROC-supervised way. The percent of variation of a clustering method assesses the clustering fitting of that method in a given setting. ASD: Autism Spectrum Disorder participants; CTRL: heterogeneous control group (composed of children with developmental language disorder [DLD] and cochlear implant [CI]); CI95%: Confidence interval of 95%. Data are presented as mean (Standard Deviation).

**Supplementary Fig. 1. Number of bootstraps for K-means clustering Analysis decided by convergence criteria.** The number of bootstrap samples that it takes to get a stable histogram is the optimal, increasing bootstraps will not change the sensitivity and specificity. Here for the study 2, 500 bootstrap iterations were used and materialized by red vertical dotted line.