

# Non Conservative Products in Fluid Dynamics

Rinaldo M. Colombo<sup>a</sup>, Graziano Guerra<sup>b</sup>, Yannick Holle<sup>c</sup>

<sup>a</sup>*INdAM Unit & Department of Information Engineering, University of Brescia, via  
Branze 38, Brescia, 25123, Italy*

<sup>b</sup>*Dipartimento di Matematica e Applicazioni, Università degli Studi di Milano Bicocca, via  
R. Cozzi 55, Milano, 20126, Italy*

<sup>c</sup>*Institut für Mathematik, RWTH Aachen University, Templergraben 55, Aachen, 52062, , Germany*

---

## Abstract

Fluid flow in pipes with discontinuous cross section or with kinks is described through balance laws with a non conservative product in the source. At jump discontinuities in the pipes' geometry, the physics of the problem suggests how to single out a solution. On this basis, we present a definition of solution for a general **BV** geometry and prove an existence result, consistent with a limiting procedure from piecewise constant geometries. In the case of a smoothly curved pipe we thus justify the appearance of the curvature in the source term of the linear momentum equation.

These results are obtained as consequences of a general existence result devoted to abstract balance laws with non conservative source terms **in the non resonant case**.

**AMS subject Classification:** 35L65; 76N10.

*Keywords:* Fluid flows in pipes; Non conservative products in balance laws.

---

## 1. Introduction

Conservation laws in one space dimension, i.e., systems of partial differential equations in conservative form of the type

$$\partial_t u + \partial_x f(u) = 0 \quad t \geq 0, x \in \mathbb{R}, \quad (1.1)$$

allow to describe, for instance, the movement of a fluid along a rectilinear pipe with constant section. Assume that at a point  $\bar{x}$  the pipe's direction or its section changes. Then, equation (1.1) can be used, separately, where  $x < \bar{x}$  and where  $x > \bar{x}$ . At the point  $\bar{x}$ , on the basis of physical considerations, a further condition is necessary to prescribe the possible defect in the conservation of the various variables. Typically, such a condition is written as

$$\Psi \left( z^+, u(t, \bar{x}+), z^-, u(t, \bar{x}-) \right) = 0 \quad \text{for a.e. } t > 0, \quad (1.2)$$

where  $z^+$  and  $z^-$ , identify the physical parameters that change across  $\bar{x}$ . Alternatively, **introducing the function  $\Xi$  to measure the defect in the conservation of the  $u$  variable, under the assumptions taken below, (1.2) can be rewritten as**

$$f \left( u(t, \bar{x}+) \right) - f \left( u(t, \bar{x}-) \right) = \Xi \left( z^+, z^-, u(t, \bar{x}-) \right) \quad \text{for a.e. } t > 0 \quad (1.3)$$

(See Lemma 4.1 and Lemma 4.2 for further details). It is then natural to tackle the resulting Riemann Problem, that is, the Cauchy Problem consisting of (1.1)–(1.3) with an initial datum attaining two values, one for  $x < 0$  and another one for  $x > 0$ , as was accomplished, for instance, in [1, § 2] or [2, § 2]. The finite propagation speed, intrinsic to (1.1), allows then to extend the whole construction to any finite number of points  $\bar{x}_0, \bar{x}_1, \dots, \bar{x}_k$ , essentially solving the Cauchy Problem for the balance law

$$\begin{cases} \partial_t u + \partial_x f(u) = \sum_{i=1}^{k-1} \Xi(\zeta_k(\bar{x}_i+), \zeta_k(\bar{x}_i-), u(t, \bar{x}_i-)) \delta_{\bar{x}_i} \\ u(0, x) = u_o(x), \end{cases} \quad (1.4)$$

where  $\delta_{\bar{x}_i}$  denotes the Dirac measure at  $\bar{x}_i$  and  $\zeta_k$  is the piecewise constant function attaining the  $k+1$  constant values  $z_0, z_1, \dots, z_k$  on the intervals  $]-\infty, \bar{x}_1[$ ,  $]\bar{x}_1, \bar{x}_2[$ ,  $\dots$ ,  $]\bar{x}_k, +\infty[$ .

This paper provides a detailed description of the rigorous limit  $k \rightarrow +\infty$  of (1.4), covering the extension of (1.4) to the case of a general **BV** function  $\zeta \in \mathbb{R}^p$ . A standard assumption playing a key role throughout is the non resonance condition, i.e., we require that all eigenvalues of (1.1) be separated from 0. The resonant, or characteristic, case requires different techniques and allows for different results, see for instance [3, 4, 5, 6, 7] and the references therein.

In the general setting established below, not limited to fluid dynamics, solutions to (1.4) with initial datum  $u_o$  are shown to converge as  $k \rightarrow +\infty$  to solutions to

$$\begin{cases} \partial_t u + \partial_x f(u) = \sum_{x \in \mathcal{I}} \Xi(\zeta(x+), \zeta(x-), u(t, x-)) \delta_x + D_{v(x)}^+ \Xi(\zeta(x), \zeta(x), u(t, x)) \|\mu\| \\ u(0, x) = u_o(x). \end{cases} \quad (1.5)$$

Since  $\zeta \in \mathbf{BV}(\mathbb{R}; \mathbb{R}^p)$ , the right and left limits  $\zeta(x+)$  and  $\zeta(x-)$  are well defined and the distributional derivative  $D\zeta$  can be split in a discrete part and a continuous one, which may contain a Cantor part:

$$D\zeta = \sum_{x \in \mathcal{I}} (\zeta(x+) - \zeta(x-)) \delta_x + v \|\mu\|, \quad (1.6)$$

where the function  $v$  is Borel measurable with norm 1,  $\mu$  is the non atomic part of  $D\zeta$  and  $\mathcal{I}$  is the set of jump points in  $\zeta$ . In (1.5) we also used the (one sided) directional derivative

$$D_v^+ \Xi(z, z, u) = \lim_{t \rightarrow 0^+} \frac{\Xi(z + tv, z, u) - \Xi(z, z, u)}{t}. \quad (1.7)$$

Indeed, one of our motivating examples, namely the case of a curved pipe, leads to a function  $\Xi$  that admits directional derivatives but is not differentiable.

The setting introduced so far is motivated by fluid dynamics in pipes. From a more general point of view, we are addressing the issue of solving a balance law of the type

$$\partial_t u + \partial_x f(u) = D_{v(x)} \Xi(\zeta(x), \zeta(x), u) \|D\zeta(x)\| \quad (1.8)$$

with the notation (1.6)–(1.7). Whenever  $\zeta$  is not a  $\mathbf{W}^{1,1}$  function, the source term above may contain non conservative products that need further information to be uniquely determined. Our choice is detailed in (1.5) and is consistent with the limit of (1.4) to (1.5) as  $\zeta$  is approximated by piecewise constant functions.

In particular, as soon as  $\Xi$  is differentiable with respect to its first argument, the right hand side in (1.5) and (1.8) can be slightly simplified, to

$$D_{v(x)}^+ \Xi(a, a, u) \|\mu\| = D_1 \Xi(a, a, u) v(x) \|\mu\| = D_1 \Xi(a, a, u) \mu. \quad (1.9)$$

Moreover, in the case  $\Xi(z^+, z^-, u) = G(z^+) - G(z^-)$  for a suitable  $G \in \mathbf{C}^2(\mathbb{R}^p; \mathbb{R}^n)$ , the right hand side above takes a simpler form. Indeed, by [8, Theorem 3.96], (1.5) reduces to the conservative problem

$$\partial_t u + \partial_x f(u) = \partial_x (G \circ \zeta). \quad (1.10)$$

The existence of solutions to (1.5) is achieved sequentially combining wave front tracking [9, § 7.1], a nowadays classical technique that approximates solutions to conservation laws, with the approximation of the equation, in particular of the map  $\zeta$ . A key role is played by a very careful choice of these approximations. As a byproduct, we obtain the solutions to (1.5) as limits of (suitable subsequences of) solutions to (1.4).

Remark that the above general procedure, when applied to the case of a curved pipe with constant section, amounts to justify the role of the pipe's curvature on the fluid flow inside the pipe. Indeed, if  $x$  is the abscissa along the pipe and  $\Gamma = \Gamma(x)$  describes the pipe's shape, then the pipe's local direction that enters the equation for fluid flow is  $\zeta(x) = \Gamma'(x)$ . Problem (1.4) then corresponds to a piecewise linear pipe and (the second component of) (1.3) describes the change in the fluid linear momentum at a kink sited at  $\bar{x}$ . Assuming that the lack in the conservation of linear momentum depends on the angle in the pipe at  $\bar{x}$ , i.e.,  $\Xi(z^+, z^-, u) = K(\|z^+ - z^-\|, u)$  as in [10, 11], automatically implies in the smooth pipe limit, by Theorem 2.2, that the variation in the fluid momentum depends on the pipe's curvature  $\Gamma''$ , see § 3.1 for more details.

The current literature offers a variety of different conditions quantifying the lack in the conservation of linear momentum at a junction where the pipe's section changes, see for instance [12, 13, 14, 2, 15, 16, 17, 18]. As a consequence of Theorem 2.2, we can select those conditions that are consistent with the equations for a pipe with smoothly varying section, both in the isentropic and in the full  $3 \times 3$  cases, see § 3.2 and § 3.3 below.

While motivated by the above fluid dynamics problems, the present construction also suggests a criterion to select solutions to general balance laws with a non conservative product as a source term, see Definition 3.3. These solutions, whose existence follows from Theorem 2.2, are obtained as limits of solutions to the piecewise constant case (1.4).

The next section is devoted to the main results: the definition of solution and to the existence theorem. Section 3 presents applications to fluid dynamics and to general balance laws with non conservative product in the source. All technical proofs are deferred to Section 4.1.

## 2. Assumptions and Main Result

Throughout,  $|x|$  is the absolute value of the real number  $x$  while, as usual,  $\|v\|$  is the Euclidean norm of the vector  $v$  and  $\|\mu\|$  is the total variation of a measure  $\mu$ . The open ball in  $\mathbb{R}^n$  centered at  $u$  with radius  $\delta$  is denoted by  $B(u; \delta)$ , its closure is  $\overline{B(u; \delta)}$ . We also use the following standard notation for right/left limits and for differences at a point:

$$F(x-) = \lim_{\xi \rightarrow x^-} F(\xi), \quad F(x+) = \lim_{\xi \rightarrow x^+} F(\xi) \quad \text{and} \quad \Delta F(x) = F(x+) - F(x-).$$

The problem we tackle is defined by the flow  $f$  and by the functions  $\Xi$  and  $\zeta$ . Here we detail the key assumptions.

(f.1)  $f \in \mathbf{C}^2(\Omega; \mathbb{R}^n)$ ,  $\Omega$  being an open subset of  $\mathbb{R}^n$ ;

(f.2) the system (1.1) is strictly hyperbolic;

(f.3) each characteristic field is either genuinely nonlinear or linearly degenerate.

In the latter assumption we refer to the usual definitions by Lax [19], see also [20, § 7.5].

By (f.1) and (f.2) we know that, possibly restricting  $\Omega$ , the eigenvalues  $\lambda_1(u), \dots, \lambda_n(u)$  of  $Df(u)$  can be numbered so that, for all  $u \in \Omega$ ,

$$\lambda_1(u) < \lambda_2(u) < \dots < \lambda_n(u).$$

We choose  $i_o \in \{1, \dots, n-1\}$  and define the  $i_o$ -th non-characteristic set

$$A_{i_o} = \{u \in \Omega \mid \lambda_{i_o}(u) < 0 < \lambda_{i_o+1}(u)\}, \quad (2.1)$$

both the cases of characteristic speeds being either all positive or all negative being simpler.

On the function  $\Xi$  in (1.3), used to rewrite the coupling condition induced by  $\Psi$ , we require, **for a suitable open subset  $\mathcal{Z}$  of  $\mathbb{R}^p$** :

(E.1)  $\Xi : \mathcal{Z} \times \mathcal{Z} \rightarrow \mathbf{C}^1(\Omega; \mathbb{R}^n)$ , is a Lipschitz continuous map;

(E.2)  $\sup_{z^+, z^- \in \mathcal{Z}} \|\Xi(z^+, z^-, \cdot)\|_{\mathbf{C}^2(\Omega; \mathbb{R})} < +\infty$ ;

(E.3)  $\Xi(z, z, u) = 0$  for every  $z \in \mathcal{Z}$  and  $u \in \Omega$ ;

(E.4) There exists a non decreasing  $\sigma : [0, \bar{t}] \rightarrow [0, +\infty[$  with  $\sigma(t) \rightarrow 0$  as  $t \rightarrow 0$  such that for all  $(z, v, u) \in \mathcal{Z} \times \overline{B(0; 1)} \times \Omega$

$$\left\| \Xi(z + tv, z, u) - D_v^+ \Xi(z, z, u) t \right\| \leq \sigma(t) t$$

and moreover the map  $(z, v, u) \rightarrow D_v^+ \Xi(z, z, u)$  is Lipschitz continuous.

In the latter condition, recall the definition (1.7) of the Dini derivative. Our requiring this low regularity, i.e. the mere existence of the Dini derivative rather than differentiability, is motivated by the example of a pipe with angles, where  $\Xi$  depends on  $\|z^+ - z^-\|$ , see § 3.1.

Problem (1.5) requires the introduction of a further function, say  $\zeta : \mathbb{R} \rightarrow \mathbb{R}^p$  describing, for instance, geometrical aspects of the pipeline. We require that  $\zeta \in \mathbf{BV}(\mathbb{R}; \mathcal{Z})$ . Throughout, the map  $\zeta$  is assumed to be left continuous and the set of jump discontinuities in  $\zeta$  is denoted by  $\mathcal{I}$ , with  $\mathcal{I} \subset \mathbb{R}$ .

We now precisely state what we mean by *solution* to (1.5).

**Definition 2.1.** Let  $u_o \in \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \mathbb{R}^n)$ . A map  $u \in \mathbf{C}^0([0, +\infty[; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \mathbb{R}^n))$  with  $u(t) \in \mathbf{BV}(\mathbb{R}; \mathbb{R}^n)$  and left continuous for all  $t \in \mathbb{R}_+$ , is a solution to (1.5) if for all test functions  $\varphi \in \mathbf{C}_c^1([0, +\infty[ \times \mathbb{R}; \mathbb{R})$ ,

$$\begin{aligned} & - \int_0^{+\infty} \int_{\mathbb{R}} \left( u(t, x) \partial_t \varphi(t, x) + f(u(t, x)) \partial_x \varphi(t, x) \right) dx dt \\ & = \sum_{x \in \mathcal{I}} \int_0^{+\infty} \Xi(\zeta(x+), \zeta(x), u(t, x)) \varphi(t, x) dt \\ & \quad + \int_0^{+\infty} \int_{\mathbb{R}} D_{v(x)}^+ \Xi(\zeta(x), \zeta(x), u(t, x)) \varphi(t, x) d\|\mu\|(x) dt \end{aligned} \quad (2.2)$$

where  $\mathcal{I}$  is the set of jump points of  $\zeta$  and  $v, \mu$  are as in (1.6), and moreover  $u(0) = u_o$ .

In the last integral, the integrand is Borel measurable in  $(t, x)$  since, for instance, by the above assumptions on  $u$ , we have at every  $(t, x) \in \mathbb{R}_+ \times \mathbb{R}$

$$u(t, x) = \lim_{h \rightarrow 0} \frac{1}{h} \int_{x-h}^x u(t, y) dy .$$

Moreover, Borel measurability on  $\mathbb{R}^2$  ensures measurability with respect to the product measure.

Note that the value of the integrand in the first line above is independent of changes of the integrand on sets of Lebesgue measure 0, while the latter integrand is integrated with respect to the product measure  $\|\mu\| \otimes dt$ . Nevertheless, the above equality is meaningful, since  $u$  is prescribed *pointwise*, at every point and not merely almost everywhere. The construction below complies with this strict requirement, see (4.24)–(4.25)–(4.26)–(4.27)–(4.28).

The above definition is known not to guarantee uniqueness. Indeed, it is the *construction* of the solutions that allows to prove their uniqueness, as will be shown in a forthcoming work. The main result of this paper is the following.

**Theorem 2.2.** *Let  $\Omega \subseteq \mathbb{R}^n$  be open,  $f$  satisfy (f.1)–(f.3),  $\Xi$  satisfy (Ξ.1)–(Ξ.4) and  $\zeta \in \mathbf{BV}(\mathbb{R}; \mathcal{Z})$ . Fix  $\bar{u} \in A_{i_o}$ ,  $\bar{z} \in \mathcal{Z}$  and an initial datum  $u_o$  in  $\mathbf{L}_{\text{loc}}^1(\mathbb{R}; A_{i_o})$ , with  $A_{i_o}$  as defined in (2.1). Then, there exists a positive  $\delta$  such that if*

$$u_o(\mathbb{R}) \subseteq B(\bar{u}; \delta), \quad \text{TV}(u_o) < \delta \quad \text{and} \quad \zeta(\mathbb{R}) \subseteq B(\bar{z}; \delta), \quad \text{TV}(\zeta) < \delta \quad (2.3)$$

the Cauchy Problem for (1.5) with initial datum  $u_o$  admits a solution  $u_*$  in the sense of Definition 2.1. Moreover, there exists a sequence of piecewise constant approximations  $\zeta^h$  of  $\zeta$ , with  $\text{TV}(\zeta^h) < \delta$ , such that the corresponding solutions  $u^h$  converge to  $u_*$  pointwise in time and in  $\mathbf{L}_{\text{loc}}^1$  in space. In particular, at each discontinuity point  $y$  of  $\zeta^h$ ,  $u^h$  satisfies the junction condition

$$f(u^h(t, y+)) - f(u^h(t, y-)) = \Xi(\zeta^h(y+), \zeta^h(y-), u^h(t, y-)).$$

The above result is of a *local* nature. This is necessary in two respects. First, we used the Implicit Function Theorem to construct approximate solutions to Riemann Problems by means of Lax curves. Secondly, the smallness of the total variation is used in the interaction estimates that allow to apply Helly Compactness Theorem.

In any particular system where approximate solutions  $u^h$  corresponding to approximations  $\zeta^h$  of  $\zeta$  are available and satisfy (1), (2), (3), (4) in Theorem 4.12, the above result applies.

### 3. Applications

#### 3.1. Isentropic Gas in a Curved Pipe

The well known system of one dimensional isentropic gas dynamics within a pipe with constant section in Eulerian coordinates [20, Formula (7.1.12)] is

$$\begin{cases} \partial_t \rho + \partial_x q = 0 \\ \partial_t q + \partial_x P(\rho, q) = 0 \end{cases} \quad \text{where } P(\rho, q) = \frac{q^2}{\rho} + p(\rho), \quad \text{for a.e. } t \geq 0, x \in \mathbb{R}. \quad (3.1)$$

Here,  $x$  is the abscissa along the pipe,  $\rho \in ]0, +\infty[$  denotes the gas density,  $q \in \mathbb{R}$  the momentum density,  $p = p(\rho)$  the pressure and  $P = P(\rho, q)$  the momentum flux. The pressure law  $p$  satisfies

(p)  $p \in \mathbf{C}^2(]0, +\infty[, ]0, +\infty[)$ ,  $p'(\rho) \geq 0$  and  $p''(\rho) \geq 0$  for all  $\rho > 0$ .

Under this assumption, system (3.1) is strictly hyperbolic, except at the vacuum  $\rho = 0$ .

We aim to establish the existence of solutions to (3.1) in a curved pipeline with constant section lying in a horizontal plane. Parameterize the pipe's support by means of the arc length  $\Gamma: \mathbb{R} \rightarrow \mathbb{R}^2$ , so that  $\|\Gamma'(x)\| = 1$  for a.e.  $x \in \mathbb{R}$ . We assume that  $\zeta = \Gamma'$  is in  $\mathbf{BV}(\mathbb{R}; \mathbb{R}^2)$ .

As a first step, consider the case of (3.1) at a kink sited at  $\bar{x}$ , so that  $\Gamma$  is the gluing of two half lines. Therefore, to solve (3.1), we adopt the usual weak entropy solutions to (3.1) along the straight parts of  $\Gamma$  and match at the kink  $\bar{x}$  a coupling condition of the type

$$\begin{cases} q(t, \bar{x}+) - q(t, \bar{x}-) &= \Xi_1(\Gamma'(\bar{x}+), \Gamma'(\bar{x}-), (\rho, q)(t, \bar{x}-)) \\ P(\rho, q)(t, \bar{x}+) - P(\rho, q)(t, \bar{x}-) &= \Xi_2(\Gamma'(\bar{x}+), \Gamma'(\bar{x}-), (\rho, q)(t, \bar{x}-)) \end{cases} \quad (3.2)$$

We set  $\Xi_1 \equiv 0$  as it is necessary to comply with mass conservation. Physical considerations suggest that the defect in the conservation of linear momentum is a function, say  $K$ , of the norm of the difference in the orientations of the pipes on the sides of the kink:

$$\Xi(z^+, z^-, (\rho, q)) = \begin{bmatrix} 0 \\ K(\|z^+ - z^-\|, (\rho, q)) \end{bmatrix}. \quad (3.3)$$

This holds true in various instances of  $K$  considered in the literature. For instance, [11] first introduced the condition

$$K(\|z^+ - z^-\|, (\rho, q)) = -\alpha \|z^+ - z^-\| q \quad (3.4)$$

for a suitable  $\alpha > 0$ , motivated by

$$\|z^+ - z^-\| = \sqrt{2(1 - \cos \bar{\vartheta})} = 2 |\sin(\bar{\vartheta}/2)|,$$

$\bar{\vartheta}$  being the angle between the two sides of the kink. It is immediate to see that (E.1)–(E.3) all hold. Concerning (E.4), we have

$$D_v^+ \Xi(z, z, (\rho, q)) = \begin{bmatrix} 0 \\ -\alpha \|v\| q \end{bmatrix} \quad \text{with} \quad \sigma \equiv 0.$$

We stress that  $\Xi_2$  is *not* of class  $\mathbf{C}^1$ .

**Theorem 3.1.** *Let  $p$  satisfy (p) and  $(\bar{\rho}, \bar{q})$  be a subsonic state. Let  $\Gamma$  be piecewise  $\mathbf{C}^2(\mathbb{R}; \mathbb{R}^2)$ , such that  $\Gamma' \in \mathbf{BV}(\mathbb{R}; \mathbb{R}^2)$  and  $\|\Gamma'(x)\| = 1$  for all  $x \in \mathbb{R}$ . Let  $K \in \mathbf{C}^2([0, r] \times \Omega; \mathbb{R})$  for a positive  $r$ , with  $K(0, (\rho, q)) \equiv 0$ . Call  $\mathcal{I}$  the set of kink points of  $\Gamma$ . Then, there exists a positive  $\delta$  such that for all initial data  $(\rho_o, q_o)$  with*

$$\|(\rho_o, q_o) - (\bar{\rho}, \bar{q})\|_{\mathbf{L}^\infty(\mathbb{R}; \mathbb{R}^2)} < \delta, \quad \text{TV}(\rho_o, q_o) < \delta, \quad \text{TV}(\Gamma') < \delta$$

the problem

$$\left\{ \begin{array}{l} \partial_t \rho + \partial_x q = 0 \\ \partial_t q + \partial_x P(\rho, q) = - \sum_{y \in \mathcal{I}} K \left( \|\Gamma'(y+) - \Gamma'(y-)\|, (\rho, q)(t, y-) \right) \delta_y \\ \quad - \|\Gamma''(x)\| \partial_1 K(0, q) \\ (\rho, q)(0, x) = (\rho_o, q_o)(x) \end{array} \right. \quad (3.5)$$

admits a solution  $(\rho_*, q_*)$  in the sense of Definition 2.1. Moreover, there exists a sequence of piecewise linear approximations  $\Gamma^h$  of  $\Gamma$ , with  $\text{TV}((\Gamma^h)') < \delta$ , such that the corresponding solutions  $(\rho^h, q^h)$  converge to  $(\rho_*, q_*)$  pointwise in time and in  $\mathbf{L}_{\text{loc}}^1$  in space. In particular, at each discontinuity point  $\bar{x}$  of  $(\Gamma^h)'$ ,  $(\rho^h, q^h)$  satisfies condition (3.2) with  $\Xi$  as in (3.3).

The proof is deferred to § 4.4.

Remark that the second derivative  $\Gamma''$  appearing in the right hand side above confirms the relevance of the pipe's curvature. Nevertheless, Theorem 2.2 applies also to less regular functions  $\Gamma$ , but the above simpler formulation then needs to be replaced by the formulation used in Definition 2.1.

### 3.2. Isentropic Gas in a Pipe with Varying Section

The isentropic flow of a fluid in a pipe with smoothly varying section  $a = a(x)$  is described by

$$\left\{ \begin{array}{l} \partial_t \rho + \partial_x q = -\frac{a'}{a} q \\ \partial_t q + \partial_x P(\rho, q) = -\frac{a'}{a} \frac{q^2}{\rho} \end{array} \right. \quad \text{where } P(\rho, q) = \frac{q^2}{\rho} + p(\rho), \quad \text{for a.e. } t \geq 0, x \in \mathbb{R}, \quad (3.6)$$

see [17, 18, 21]. The case of  $a$  piecewise constant, i.e., the section of the pipe changes from  $a^-$  to  $a^+$  at a junction sited at  $\bar{x}$ , is covered in the literature supplementing the  $p$ -system (3.1) with a junction condition of the form

$$\left\{ \begin{array}{l} a^+ q(t, \bar{x}+) = a^- q(t, \bar{x}-) \\ P(\rho, q)(t, \bar{x}+) - P(\rho, q)(t, \bar{x}-) = \Xi_2(a^+, a^-, (\rho, q)(t, \bar{x}-)) \end{array} \right. \quad (3.7)$$

The former relation in (3.7) ensures the conservation of mass and fits in the framework of Section 2 setting in the first component of (1.3)

$$\Xi_1(a^+, a^-, (\rho^-, q^-)) = \left( \frac{a^-}{a^+} - 1 \right) q^-. \quad (3.8)$$

The literature offers a wide range of justifications, often phenomenological, for specific choices of the function  $\Xi_2$  in (3.7), see for instance [15, 17, 18]. Note that, as soon as  $\Xi_2$  is of class

$\mathbf{C}^2$  in all variables, with  $\Xi_2(a, a, (\rho, q)) = 0$ , and  $a$  is in  $\mathbf{BV}(\mathbb{R}; \mathbb{R})$ , then Theorem 2.2 applies ensuring the existence of solutions to

$$\left\{ \begin{array}{l} \partial_t \rho + \partial_x q = \sum_{x \in \mathcal{I}} \Xi_1(a(x+), a(x-), (\rho, q)(t, x-)) \delta_x \\ \quad + \partial_1 \Xi_1(a(x), a(x), (\rho, q)(t, x)) \mu \\ \partial_t q + \partial_x P(\rho, q) = \sum_{x \in \mathcal{I}} \Xi_2(a(x+), a(x-), (\rho, q)(t, x-)) \delta_x \\ \quad + \partial_1 \Xi_2(a(x), a(x), (\rho, q)(t, x)) \mu \end{array} \right. \quad (3.9)$$

where  $\mathcal{I}$  is the set of points of discontinuity of  $a$ . In (3.9) we also used (1.9) with  $\zeta(x) = a(x)$ . Note that as soon as  $a$  is smooth,  $\mu$  has density  $\partial_x a(x)$  with respect to the Lebesgue measure.

As an application of Theorem 2.2, we give sufficient conditions on  $\Xi$  that yield in the limit the case of the smooth pipe, i.e., equation (3.6).

**Theorem 3.2.** *Let  $p$  satisfy  $(\mathbf{p})$ ,  $(\bar{\rho}, \bar{q})$  be a subsonic state and  $\bar{a}$  be positive. Fix  $\Xi_1$  as in (3.8). For any  $\Xi_2$  of class  $\mathbf{C}^2$  with  $\Xi_2(a, a, (\rho, q)) = 0$  there exists a positive  $\delta$  such that for all initial data  $(\rho_o, q_o)$  and for all  $a \in \mathbf{BV}(\mathbb{R}; \mathbb{R})$  and*

$$\|(\rho_o, q_o) - (\bar{\rho}, \bar{q})\|_{\mathbf{L}^\infty(\mathbb{R}; \mathbb{R}^2)} < \delta, \quad \mathbf{TV}(\rho_o, q_o) < \delta, \quad \|a - \bar{a}\|_{\mathbf{L}^\infty(\mathbb{R}; \mathbb{R})} < \delta, \quad \mathbf{TV}(a) < \delta$$

problem (3.9) admits a solution  $(\rho_*, q_*)$ . Moreover, there exists a sequence of piecewise constant approximations  $a^h$  of  $a$ , with  $\mathbf{TV}(a^h) < \delta$ , such that the corresponding solution  $(\rho^h, q^h)$  converges to  $(\rho_*, q_*)$  pointwise in time and in  $\mathbf{L}_{\text{loc}}^1$  in space. In particular, at each discontinuity point  $y$  of  $a^h$ ,  $(\rho^h, q^h)$  satisfies the junction condition (3.7).

If  $a \in \mathbf{W}^{1,1}(\mathbb{R}; \mathbb{R})$  and  $\Xi_2$  satisfies

$$\partial_1 \Xi_2(a, a, (\rho, q)) = -\frac{1}{a} \frac{q^2}{\rho} \quad (3.10)$$

then  $(\rho_*, q_*)$  also solves (3.6).

The proof is deferred to § 4.4. Here, we only remark that the choice  $[\mathbf{S}]$  of  $\Xi_2$  in Table 1 satisfies (3.10).

We now test the above condition against various junction condition found in the literature, we refer in particular to [15] for the motivations and further information of the conditions considered below. More precisely, with reference to the labeling in Table 1, we consider definition  $[\mathbf{L}]$  from [14], condition  $[\mathbf{p}]$  from [12, 13], condition  $[\mathbf{P}]$  from [14, 2] and condition  $[\mathbf{S}]$  from [17, 18]. All these conditions differ only in the second component  $\Xi_2$ , the first one being fixed as in (3.8) to comply with mass conservation.

Simple computations lead to the results in Table 1, where the map  $a \rightarrow R(a; a^-, \rho^-, q^-)$  is the first component of the solution to the stationary version of (3.6), parameterized by the section  $a$ , i.e.,

$$\left\{ \begin{array}{l} \frac{d}{da} q = -\frac{1}{a} q \\ \frac{d}{da} (P(\rho, q)) = -\frac{1}{a} \frac{q^2}{\rho} \end{array} \right. \quad \begin{array}{l} \rho(a^-) = \rho^- \\ q(a^-) = q^- \end{array} \quad (3.11)$$



	$\Psi_2(a^-, (\rho^-, q^-), a^+, (\rho^+, q^+))$	$\Xi_2(a^+, a^-, (\rho^-, q^-))$	$D_1 \Xi_2(a, a, (\rho, q))$
[L]	$a^+ P(\rho^+, q^+) - a^- P(\rho^-, q^-)$	$\left(\frac{a^-}{a^+} - 1\right) \left(\frac{(q^-)^2}{\rho^-} + p(\rho^-)\right)$	$-\frac{1}{a} \left(\frac{q^2}{\rho} + p(\rho)\right)$
[p]	$p(\rho^+) - p(\rho^-)$	$\left(\left(\frac{a^-}{a^+}\right)^2 - 1\right) \frac{(q^-)^2}{\rho^-}$	$-\frac{2}{a} \frac{q^2}{\rho}$
[P]	$P(\rho^+, q^+) - P(\rho^-, q^-)$	0	0
[S]	$a^+ P(\rho^+, q^+) - a^- P(\rho^-, q^-)$ $- \int_{a^-}^{a^+} p(R(\alpha; a^-, \rho^-, q^-)) d\alpha$	$\left(\frac{a^-}{a^+} - 1\right) \left(\frac{(q^-)^2}{\rho^-} + p(\rho^-)\right)$ $+ \frac{1}{a^+} \int_{a^-}^{a^+} p(R(\alpha; a^-, \rho^-, q^-)) d\alpha$	$-\frac{1}{a} \frac{q^2}{\rho}$

Table 1: Various definitions of junction conditions, with the corresponding functions  $\Psi_2$  from (1.2),  $\Xi_2$  from (1.3) and its partial derivative  $\partial_1 \Xi_2$ . **The function  $R$  is the  $\rho$  component of the solution to (3.11).**

On the basis of Theorem 3.2, we know that condition [S] is compatible with the smooth limit (3.6). Moreover, Theorem 2.2 and Table 1, in particular the comparison of the rightmost column with (3.10), ensure that all the other conditions do *not* converge to (3.6) in the smooth pipe limit.

Remark that substituting in [S] any other smooth function  $R = R(a; a^-, \rho^-, q^-)$  such that  $R(a^-; a^-, \rho^-, q^-) = \rho^-$  yields a new condition at the junction compatible with the smooth limit.

**Observe that modifying  $\Xi_2$  by a factor 2 in condition [p] in Table 1 provides a further example satisfying (3.10), i.e., a condition coherent in the limit with smooth solutions.**

### 3.3. Full Gas Dynamics in Pipes with Varying Section

The full Euler system in a pipeline with smoothly varying section  $a = a(x)$  is

$$\begin{cases} \partial_t \rho + \partial_x(\rho v) = -\frac{a'}{a} \rho v \\ \partial_t(\rho v) + \partial_x(\rho v^2 + p) = -\frac{a'}{a} \rho v^2 \\ \partial_t\left(\frac{1}{2} \rho v^2 + \rho e\right) + \partial_x\left(v\left(\frac{1}{2} \rho v^2 + \rho e + p\right)\right) = -\frac{a'}{a} v\left(\frac{1}{2} \rho v^2 + \rho e + p\right) \end{cases} \quad (3.12)$$

see, for instance [16, 18, 21, 22]. Here,  $x$  is the abscissa along the pipe,  $\rho > 0$  denotes gas density,  $q \in \mathbb{R}$  the momentum density,  $p = p(\rho, s)$  the pressure and  $e = e(\rho, s)$  the energy density and  $s$  the entropy density. These two latter functions satisfy

(E)  $e \in \mathbf{C}^2(]0, +\infty[ \times \mathbb{R}, ]0, +\infty[)$  and  $\partial_s e(\rho, s) > 0$  for all  $\rho > 0$  and  $s \in \mathbb{R}$ .

(P)  $p \in \mathbf{C}^2(]0, +\infty[ \times \mathbb{R}, ]0, +\infty[)$ ,  $p(\rho, s) = \rho^2 \partial_\rho e(\rho, s)$ ,  $\partial_\rho p(\rho, s) > 0$  and  $\partial_{\rho\rho}^2(\rho p(\rho, s)) > 0$  for all  $\rho > 0$  and  $s \in \mathbb{R}$ .

We restrict our attention to the subsonic region where  $v \in ]0, \sqrt{\partial_\rho p(\rho, s)}[$ .

The conditions found in the literature, see [16], imposed at a point  $\bar{x}$  where the section suffers a discontinuity fit into the form

$$\begin{cases} \Delta(a \rho v)(t, \bar{x}) = 0 \\ \Delta(\rho v^2 + p)(t, \bar{x}) = \Xi_2(a(x+), a(x-), (\rho, v, s)(t, x-)) \\ \Delta\left(a v \left(\frac{1}{2} \rho v^2 + \rho e + p\right)\right)(t, \bar{x}) = 0 \end{cases} \quad (3.13)$$

The conservation of mass imposed by the first equality and the conservation of energy imposed by the third equality in (3.13) amount to setting

$$\begin{aligned} \Xi_1(a^+, a^-, (\rho^-, v^-, s^-)) &= \left(\frac{a^-}{a^+} - 1\right) \rho^- v^- \\ \Xi_3(a^+, a^-, u^-) &= \left(\frac{a^-}{a^+} - 1\right) \left(v^- \left(\frac{1}{2} \rho^- (v^-)^2 + \rho^- e^- + p^-\right)\right) \end{aligned}$$

so that

$$\begin{aligned} \partial_1 \Xi_1(a, a, (\rho, v, s)) &= -\frac{1}{a} \rho v \\ \partial_1 \Xi_3(a, a, u) &= -\frac{1}{a} \left(v \left(\frac{1}{2} \rho v^2 + \rho e + p\right)\right) \end{aligned}$$

The second equality in (3.13) is treated in different ways in the literature, giving rise to conditions analogous to those considered in § 3.2. Indeed, Table 1 directly extends to the present full  $3 \times 3$  case, simply understanding the map  $R$  as the  $\rho$  component  $a \rightarrow R(a; a^-, \rho^-, v^-, s^-)$  in the solution to the stationary Cauchy Problem

$$\begin{cases} \frac{d}{da}(\rho v) = -\frac{1}{a} \rho v & \rho(a^-) = \rho^- \\ \frac{d}{da}(\rho v^2 + p) = -\frac{1}{a} \rho v^2 & v(a^-) = v^- \\ \frac{d}{da}\left(v \left(\frac{1}{2} \rho v^2 + \rho e + p\right)\right) = -\frac{1}{a} v \left(\frac{1}{2} \rho v^2 + \rho e + p\right) & s(a^-) = s^- \end{cases} \quad (3.14)$$

#### 3.4. Balance Laws with Measure Valued Source Term

The theory developed in Section 2 allows to give a meaning to the following balance law, where the source term is *non conservative*:

$$\partial_t u + \partial_x f(u) = \partial_\zeta G(\zeta, u) D\zeta \quad (3.15)$$

where  $G$  is smooth and  $\zeta$  has bounded variation. In the case  $G$  independent of  $u$ , we recover the *conservative* case (1.10). In the general, *non conservative* case, (3.15) can be given different meanings.

A choice consists in setting

$$\Xi(z^+, z^-, u^-) = G(z^+, u^-) - G(z^-, u^-), \quad (3.16)$$

corresponding to the following condition at each point of jump:

$$f(u^+) - f(u^-) = G(z^+, u^-) - G(z^-, u^-).$$

The framework developed in the preceding section in connection with the Cauchy Problem (1.5) comprises (3.15). Therefore, we can particularize Definition 2.1 to the general case of non conservative products of the type (3.15).

**Definition 3.3.** Fix an initial datum  $u_o \in \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \mathbb{R}^n)$ . Let  $\Xi \in \mathbf{C}^2(\mathcal{Z} \times \mathcal{Z} \times \Omega; \mathbb{R}^n)$  be such that

$$D_1 \Xi(z, z, u) = D_z G(z, u). \quad (3.17)$$

Then, a map  $u \in \mathbf{C}^0([0, T]; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \mathbb{R}^n))$  with  $u(t) \in \mathbf{BV}(\mathbb{R}; \mathbb{R}^n)$  and left continuous for all  $t \in [0, T]$ , is a  $\Xi$ -solution to (1.5) if for all test function  $\varphi \in \mathbf{C}_c^1([0, T[ \times \mathbb{R}; \mathbb{R})$ ,

$$\begin{aligned} & - \int_0^{+\infty} \int_{\Omega} \left( u(t, x) \partial_t \varphi(t, x) + f(u(t, x)) \partial_x \varphi(t, x) \right) dx dt \\ &= \sum_{x \in \mathcal{I}} \int_0^{+\infty} \Xi(\zeta(x+), \zeta(x), u(t, x)) \varphi(t, x) dt \\ & \quad + \int_0^{+\infty} \int_{\mathbb{R}} D_z G(\zeta(x), u(t, x)) \varphi(t, x) D\mu(x) dt \end{aligned} \quad (3.18)$$

where  $\mathcal{I}$  is the set of jump points of  $\zeta$  and  $\mu$  is as in (1.6), and moreover  $u(0) = u_o$ .

This definition clearly separates those part of the solution that depend exclusively on (3.15) from those part, in the middle term in (3.18), that depend on the arbitrary choice of  $\Xi$ .

In particular, the choice (3.16) yields

$$\Xi(\zeta(x+), \zeta(x), u(t, x)) = G(\zeta(x+), u(t, x)) - G(\zeta(x), u(t, x)) \quad (3.19)$$

where we keep using the left continuous representatives. For completeness, we remark that the alternative choice  $\Xi(z^+, z^-, u^-) = G(z^+, u^+) - G(z^-, u^+)$  also meets condition (3.17).

A straightforward application of Theorem 2.2 now ensures the existence of  $\Xi$ -solutions to (3.15), as soon as  $G \in \mathbf{C}^2(\mathcal{Z} \times \Omega; \mathbb{R}^{n \times m})$ ,  $\zeta \in \mathbf{BV}(\mathbb{R}; \mathcal{Z})$ ,  $\Xi \in \mathbf{C}^2(\mathcal{Z} \times \mathcal{Z} \times \Omega; \mathbb{R}^n)$  and satisfies (3.17). Moreover, these solutions are limits of “discretized” approximations where (1.3) is imposed to the points of jump in  $\zeta$ .

#### 4. Technical Details

Below, by  $\mathcal{O}(1)$  we denote a constant depending exclusively on  $f$  and  $\Xi$ .

**Lemma 4.1.** Let  $f$  satisfy (f.1)–(f.3). Assume that  $z \in \mathcal{Z}$  and  $u \in A_{i_o}$ .

( $\alpha$ )  $\Psi: \mathcal{Z} \times \mathcal{Z} \rightarrow \mathbf{C}^1(\Omega \times \Omega; \mathbb{R}^n)$  is Lipschitz continuous.

( $\beta$ )  $\Psi(z, u, z, u) = 0$  for all  $z \in \mathcal{Z}$  and  $u \in \Omega$ .

Then, the following conditions are equivalent:

$$\det \left[ -D_{u^+} \Psi(z, u, z, u) [r_1(u) \cdots r_{i_o}(u)] \mid D_{u^-} \Psi(z, u, z, u) [r_{i_o}(u) \cdots r_n(u)] \right] \neq 0 \quad (4.1)$$

$$\det D_{u^+} \Phi(z, u, z, u) \neq 0 \quad (4.2)$$

where  $r_i(u)$  is a right eigenvector corresponding to  $\lambda_i(u)$ .

**Proof.** Differentiating  $\Psi(z, u, z, u) = 0$  with respect to  $u$ , we get  $D_{u^+} \Psi(z, u, z, u) + D_{u^-} \Psi(z, u, z, u) = 0$ , so that

$$\begin{aligned} & \det \left[ -D_{u^+} \Psi(z, u, z, u) [r_1(u) \cdots r_{i_o}(u)] \mid D_{u^-} \Psi(z, u, z, u) [r_{i_o}(u) \cdots r_n(u)] \right] \\ &= \det \left[ -D_{u^+} \Psi(z, u, z, u) [r_1(u) \cdots r_{i_o}(u)] \mid -D_{u^+} \Psi(z, u, z, u) [r_{i_o}(u) \cdots r_n(u)] \right] \\ &= \det \left( -D_{u^+} \Psi(z, u, z, u) [r_1(u) \cdots r_n(u)] \right) \\ &= (-1)^n \det D_{u^+} \Psi(z, u, z, u) \det [r_1(u) \cdots r_n(u)] \end{aligned}$$

proving that (4.1) and (4.2) are equivalent.  $\square$

Recall that condition (4.1) above ensures the well posedness of the solution to Riemann Problems at junctions, see [23, Condition (b), Section 2], [16, Condition (2.10)].

**Lemma 4.2.** *Let  $f$  satisfy (f.1)–(f.3). Assume that  $\bar{z} \in \mathcal{Z}$  and  $\bar{u} \in A_{i_o}$ . Let  $\Psi$  be as in Lemma 4.1 also satisfy (4.1). Then, there exists a map  $\Xi$  satisfying (Ξ.1) and (Ξ.3) in neighborhoods of  $\bar{z}$  and  $\bar{u}$  such that, locally,*

$$\Psi(z^+, u^+, z^-, u^-) = 0 \iff f(u^+) - f(u^-) = \Xi(z^+, z^-, u^-). \quad (4.3)$$

*Viceversa, if there exists a map  $\Xi$  satisfying (Ξ.1) and (Ξ.3), then there exists a  $\Psi$  satisfying the assumptions of Lemma 4.1 and (4.1) such that (4.3) holds.*

**Proof.** Use Lemma 4.1, then (4.2) to apply the Implicit Function Theorem in neighborhoods of  $z^- = \bar{z}$ ,  $z^+ = \bar{z}$ ,  $u^- = \bar{u}$  and  $u^+ = \bar{u}$  to obtain a function  $T$  such that  $\Psi(z^+, u^+, z^-, u^-) = 0$  is locally equivalent to  $u^+ = T(z^+, z^-, u^-)$ . Then, define  $\Xi(z^+, z^-, u^-) = f(T(z^+, z^-, u^-)) - f(u^-)$ . Viceversa, simply define  $\Psi(z^+, u^+, z^-, u^-) = f(u^+) - f(u^-) - \Xi(z^+, z^-, u^-)$  and observe that  $D_{u^+} \Psi(\bar{z}, \bar{u}, \bar{z}, \bar{u}) = D_u f(\bar{u})$ , which is non singular since  $\bar{u} \in A_{i_o}$ .  $\square$

#### 4.1. Preliminary Results

First, we prove a Lipschitz-type estimate on the map  $\Xi$  which we use throughout this paper.

**Lemma 4.3.** *Assume that (Ξ.1), (Ξ.3) hold. Then,*

$$\left\| \Xi(z^+, z^-, u_2) - \Xi(z^+, z^-, u_1) \right\| = \mathcal{O}(1) \left\| z^+ - z^- \right\| \|u_2 - u_1\|. \quad (4.4)$$

**Proof.** Since the map  $u \mapsto \Xi(z^+, z^-, u)$  is smooth, we can compute

$$\begin{aligned} & \left\| \Xi(z^+, z^-, u_2) - \Xi(z^+, z^-, u_1) \right\| \\ & \leq \|u_2 - u_1\| \int_0^1 \left\| D_u \Xi(z^+, z^-, u_1 + s(u_2 - u_1)) - D_u \Xi(z^-, z^-, u_1 + s(u_2 - u_1)) \right\| ds \\ & \leq \mathcal{O}(1) \left\| z^+ - z^- \right\| \|u_2 - u_1\|, \end{aligned}$$

where we used the equality  $D_u \Xi(z^-, z^-, u_1 + s(u_2 - u_1)) = 0$ .  $\square$

Introduce a map  $T$  related to the generalized Riemann problem.

**Lemma 4.4.** *Let  $f$  satisfy (f.1)–(f.3),  $\Xi$  satisfy (Ξ.1), (Ξ.3) and  $A_{i_o}$  be as in (2.1). Then, for any  $\bar{z} \in \mathcal{Z}$  and  $\bar{u} \in A_{i_o}$ , there exists  $\bar{\delta} > 0$  and a Lipschitz map*

$$T: B(\bar{z}; \bar{\delta})^2 \times B(\bar{u}; \bar{\delta}) \rightarrow A_{i_o}$$

such that

$$\begin{cases} f(u^+) - f(u^-) = \Xi(z^+, z^-, u^-) \\ z^+, z^- \in B(\bar{z}; \bar{\delta}) \\ u^+, u^- \in B(\bar{u}; \bar{\delta}) \end{cases} \iff u^+ = T(z^+, z^-, u^-).$$

Furthermore,

$$\left\| T(z^+, z^-, u^-) - u^- \right\| = \mathcal{O}(1) \left\| z^+ - z^- \right\|, \quad (4.5)$$

$$\left\| T(z^+, z^-, u_2) - T(z^+, z^-, u_1) - (u_2 - u_1) \right\| = \mathcal{O}(1) \left\| z^+ - z^- \right\| \|u_2 - u_1\|. \quad (4.6)$$

**Proof.** Since  $\bar{u} \in A_{i_o}$ , (f.1) and (f.2) ensure that the function  $f$  is locally invertible at  $\bar{u}$ . We define

$$T(z^+, z^-, u^-) = f^{-1} \left( f(u^-) + \Xi(z^+, z^-, u^-) \right). \quad (4.7)$$

By (Ξ.1), (Ξ.3) we compute

$$\begin{aligned} \left\| T(z^+, z^-, u^-) - u^- \right\| &= \left\| T(z^+, z^-, u^-) - f^{-1} \left( f(u^-) \right) \right\| \\ &= \mathcal{O}(1) \left\| \Xi(z^+, z^-, u^-) - \Xi(z^-, z^-, u^-) \right\| \\ &= \mathcal{O}(1) \left\| z^+ - z^- \right\|, \end{aligned}$$

proving (4.5). Introduce the smooth map

$$b(\xi, \Delta, v) = f^{-1} \left( f(u_1 + v) + \xi + \Delta \right) - f^{-1} \left( f(u_1) + \xi \right) - v.$$

Since  $b(\xi, 0, 0) = b(0, 0, v) = 0$ , the estimate

$$b(\xi, \Delta, v) = \mathcal{O}(1) [\|\xi\| \cdot \|v\| + \|\Delta\|]$$

holds, see [9, § 2.9]. The left hand side of (4.6) can be written as

$$\begin{aligned}
& \left\| T(z^+, z^-, u_2) - T(z^+, z^-, u_1) - (u_2 - u_1) \right\| \\
&= \left\| b \left[ \Xi(z^+, z^-, u_1), \Xi(z^+, z^-, u_2) - \Xi(z^+, z^-, u_1), u_2 - u_1 \right] \right\| \\
&\leq \mathcal{O}(1) \left[ \left\| \Xi(z^+, z^-, u_1) \right\| \cdot \|u_2 - u_1\| + \left\| \Xi(z^+, z^-, u_2) - \Xi(z^+, z^-, u_1) \right\| \right] \\
&\leq \mathcal{O}(1) \left\| z^+ - z^- \right\| \|u_2 - u_1\|.
\end{aligned}$$

□

#### 4.2. The Case $\zeta$ Piecewise Constant

In this section, we consider the case of  $\mathcal{I}$  being finite, with  $\zeta$  being piecewise constant. We index the points  $x \in \mathcal{I}$  so that  $x_i < x_j$  if and only if  $i < j$ . In this case, the general Definition 2.1 reduces to the following one, often found in the literature, see for instance [10, 16, 17, 11].

**Definition 4.5.** *A weak solution to the Cauchy Problem (1.5) with a piecewise constant  $\zeta$  is a map  $u \in \mathbf{C}^0([0, +\infty[; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$  with  $u(t) \in \mathbf{BV}(\mathbb{R}; \Omega)$ , left continuous, for all  $t \geq 0$ , such that for all  $\varphi \in \mathbf{C}_c^1([0, +\infty[ \times \mathbb{R}; \mathbb{R})$  whose support does not intersect  $[0, +\infty[ \times \mathcal{I}$*

$$\int_0^{+\infty} \int_{\mathbb{R}} (u \partial_t \varphi + f(u) \partial_x \varphi) \, dx \, dt = 0, \quad (4.8)$$

$u(0) = u_o$  and for all  $x \in \mathcal{I}$

$$f(u(t, x+)) - f(u(t, x)) = \Xi(\zeta(x+), \zeta(x), u(t, x)) \quad \text{for a.e. } t \in [0, +\infty[.$$

##### 4.2.1. The Generalized Riemann Problem

By *Generalized Riemann Problem* we consider the Cauchy Problem (1.5) with  $\zeta$  and the initial datum  $u_o$  being as follows:

$$\zeta(x) = z^- \chi_{]-\infty, 0[}(x) + z^+ \chi_{]0, +\infty[}(x) \quad \text{and} \quad u_o(x) = u^\ell \chi_{]-\infty, 0[}(x) + u^r \chi_{]0, +\infty[}(x). \quad (4.9)$$

For  $u \in A_{i_o}$ , call  $\sigma_i \rightarrow H_i(\sigma_i)(u)$  the Lax curve of the  $i$ -th family exiting  $u$ , see [9, § 5.2] or [20, § 9.3]. Introduce recursively the states  $w_0, \dots, w_{n+1}$  with  $w_0 = u^\ell$ ,  $w_{n+1} = u^r$  and

$$\begin{cases} w_{i+1} = H_{i+1}(\sigma_{i+1})(w_i) & \text{if } i = 0, \dots, i_o - 1, \\ w_{i_o+1} = T(z^+, z^-, w_{i_o}) \\ w_{i+1} = H_i(\sigma_i)(w_i) & \text{if } i = i_o + 1, \dots, n. \end{cases}$$

If  $z^+ - z^-$  is sufficiently small, [1, Lemma 3] ensures that the waves' sizes  $(\sigma_1, \dots, \sigma_n)$  and the states  $(w_1, \dots, w_n)$  exist, are uniquely defined and are Lipschitz continuous functions of  $z^-, z^+, u^\ell, u^r$ , which ensures also the well posedness of the Generalized Riemann Problem (1.5), (4.9). The following notation is of use below:

$$(\sigma_1, \dots, \sigma_n) = E(z^+, z^-, u^r, u^\ell). \quad (4.10)$$

We thus write the solution  $u$  to the Generalized Riemann Problem (1.5) (4.9), in the sense of Definition 4.5, as the gluing along  $x = 0$  of the Lax solutions to the (standard) Riemann Problems

$$\begin{cases} \partial_t u + \partial_x f(u) = 0 \\ u(0, x) = u^\ell \chi_{]-\infty, 0[}(x) + w_{i_o} \chi_{]0, +\infty[}(x), \end{cases} \quad \begin{cases} \partial_t u + \partial_x f(u) = 0 \\ u(0, x) = w_{i_o+1} \chi_{]-\infty, 0[}(x) + u^r \chi_{]0, +\infty[}(x). \end{cases}$$

#### 4.2.2. Interaction Estimates

We separate the waves with negative or positive propagation speed as follows:

$$\begin{aligned} \boldsymbol{\sigma}' &= (\sigma_1, \dots, \sigma_{i_o}, 0, \dots, 0), & \boldsymbol{\sigma}'' &= (0, \dots, 0, \sigma_{i_o+1}, \dots, \sigma_n), \\ \boldsymbol{\sigma} &= \boldsymbol{\sigma}' + \boldsymbol{\sigma}'' \in \mathbb{R}^n. \end{aligned} \tag{4.11}$$

Given two  $n$ -tuples of waves  $\boldsymbol{\alpha}$  and  $\boldsymbol{\beta}$ , the waves  $i$  with size  $\alpha_i$  and  $j$  with size  $\beta_j$  are *approaching* whenever  $i > j$  or  $\min\{\alpha_i, \beta_j\} < 0$ . Call  $\mathcal{A}_{\boldsymbol{\alpha}, \boldsymbol{\beta}}$  the set of these pairs.

In the following we recall several lemmas which are straightforward generalizations of results in [1].

**Lemma 4.6.** *Let  $f$  satisfy (f.1)–(f.3),  $\Xi$  satisfy (Ξ.1), (Ξ.3) and  $A_{i_o}$  be as in (2.1). Fix  $\bar{u} \in A_{i_o}$  and  $\bar{z} \in \mathcal{Z}$ . Then, there exists a  $\delta > 0$  such that if  $z^+, z^- \in B(\bar{z}; \delta)$ ,  $u^\ell, u^r \in B(\bar{u}; \delta)$ , we have*

$$\begin{aligned} \|u^r - u^\ell\| &= \mathcal{O}(1) \left( \|\boldsymbol{\sigma}\| + \|z^+ - z^-\| \right), \\ \|\boldsymbol{\sigma}\| &= \mathcal{O}(1) \left( \|u^r - u^\ell\| + \|z^+ - z^-\| \right). \end{aligned}$$

**Proof.** By Lemma 4.4, we get

$$\|u^r - u^\ell\| \leq \sum_{i=1}^{n+1} \|w_i - w_{i-1}\| = \mathcal{O}(1) \|\boldsymbol{\sigma}\| + \|T(z^+, z^-, w_{i_o}) - w_{i_o}\| = \mathcal{O}(1) \left( \|\boldsymbol{\sigma}\| + \|z^+ - z^-\| \right).$$

Recall  $E$  as defined in (4.10) and observe that  $E(z^+, z^-, T(z^+, z^-, u^\ell), u^\ell) = 0$ . By the Lipschitz continuity of  $E$ , we get

$$\begin{aligned} \|\boldsymbol{\sigma}\| &= \left\| E(z^+, z^-, u^r, u^\ell) - E(z^+, z^-, T(z^+, z^-, u^\ell), u^\ell) \right\| \\ &= \mathcal{O}(1) \|u^r - T(z^+, z^-, u^\ell)\| \\ &= \mathcal{O}(1) \left( \|u^r - u^\ell\| + \|u^\ell - T(z^+, z^-, u^\ell)\| \right) \\ &= \mathcal{O}(1) \left( \|u^r - u^\ell\| + \|z^+ - z^-\| \right) \end{aligned}$$

completing the proof. □

**Lemma 4.7** ([1, Lemma 5]). *Let  $f$  satisfy (f.1)–(f.3) and  $A_{i_o}$  be as in (2.1). Fix  $\bar{u} \in A_{i_o}$ . For  $u \in A_{i_o}$  sufficiently close to  $\bar{u}$  and  $y_1, y_2, \alpha \in \mathbb{R}^n$  sufficiently small, we have*

$$\|y_2 + H(\alpha)(u) - H(\alpha)(u + y_1)\| = \mathcal{O}(1) (\|\alpha\| \|y_1\| + \|y_1 - y_2\|).$$

**Lemma 4.8.** *Let  $f$  satisfy (f.1)–(f.3),  $\Xi$  satisfy (Ξ.1), (Ξ.3) and  $A_{i_o}$  be as in (2.1). Fix  $\bar{u} \in A_{i_o}$  and  $\bar{z} \in \mathcal{Z}$ . For  $u \in A_{i_o}$  sufficiently close to  $\bar{u}$ ,  $z^+, z^- \in \mathcal{Z}$  sufficiently close to  $\bar{z}$  and  $\alpha \in \mathbb{R}^n$  sufficiently small, we have*

$$\left\| T(z^+, z^-, H(\alpha)(u)) - H(\alpha)(T(z^+, z^-, u)) \right\| = \mathcal{O}(1) \|\alpha\| \|z^+ - z^-\|.$$

**Proof.** Applying Lemma 4.7 with  $y_1 = T(z^+, z^-, u) - u$  and  $y_2 = T(z^+, z^-, H(\alpha)(u)) - H(\alpha)(u)$  gives

$$\begin{aligned} & \left\| T(z^+, z^-, H(\alpha)(u)) - H(\alpha)(T(z^+, z^-, u)) \right\| \\ & \leq \mathcal{O}(1) \left( \|\alpha\| \|T(z^+, z^-, u) - u\| + \left\| T(z^+, z^-, H(\alpha)(u)) - H(\alpha)(u) - T(z^+, z^-, u) + u \right\| \right). \end{aligned}$$

The result follows from Lemma 4.4.  $\square$

**Lemma 4.9.** *Let  $f$  satisfy (f.1)–(f.3),  $\Xi$  satisfy (Ξ.1), (Ξ.3) and  $A_{i_o}$  be as in (2.1). Fix  $\bar{u} \in A_{i_o}$  and  $\bar{z} \in \mathcal{Z}$ . Then, there exists a  $\delta > 0$  such that if  $u^r, u^\ell \in B(\bar{u}; \delta)$ ,  $z^+, z^- \in B(\bar{z}; \delta)$  and*

$$\begin{aligned} u^{m-} &= H(\alpha)(u^\ell), \\ u^r &= H(\beta'') \left( T(z^+, z^-, H(\beta')(u^{m-})) \right), \\ u^r &= H(\sigma'') \left( T(z^+, z^-, H(\sigma')(u^\ell)) \right), \end{aligned}$$

with  $\alpha, \beta, \sigma \in \mathbb{R}^n$  and  $u^{m-} \in B(\bar{u}; \delta)$ , we have

$$\sum_{i=1}^n |\sigma_i - \alpha_i - \beta_i| = \mathcal{O}(1) \left( \sum_{(i,j) \in \mathcal{A}_{\alpha,\beta}} |\alpha_i \beta_j| + \|z^+ - z^-\| \sum_{i > i_o} |\alpha_i| \right), \quad (4.12)$$

where  $\mathcal{A}_{\alpha,\beta}$ , as above, denotes the set of approaching waves. Analogously, if

$$\begin{aligned} u^{m+} &= H(\alpha'') \left( T(z^+, z^-, H(\alpha')(u^\ell)) \right), \\ u^r &= H(\beta)(u^{m+}), \\ u^r &= H(\sigma'') \left( T(z^+, z^-, H(\sigma')(u^\ell)) \right), \end{aligned}$$

then,

$$\sum_{i=1}^n |\sigma_i - \alpha_i - \beta_i| = \mathcal{O}(1) \left( \sum_{(i,j) \in \mathcal{A}_{\alpha,\beta}} |\alpha_i \beta_j| + \|z^+ - z^-\| \sum_{i < i_o} |\beta_i| \right). \quad (4.13)$$



**Proof.** It is sufficient to prove (4.12), since (4.13) is proved analogously. We set

$$\begin{aligned}\tilde{u} &= H(\boldsymbol{\alpha}'' + \boldsymbol{\beta}'') \left( T \left( z^+, z^-, H(\boldsymbol{\alpha}' + \boldsymbol{\beta}')(u^\ell) \right) \right), \\ u_1 &= H(\boldsymbol{\beta}'') \circ H(\boldsymbol{\alpha}'') \left( T \left( z^+, z^-, H(\boldsymbol{\alpha}')(u^\ell) \right) \right), \\ u_2 &= H(\boldsymbol{\beta}'') \left( T \left( z^+, z^-, H(\boldsymbol{\alpha})(u^\ell) \right) \right).\end{aligned}$$

By the Lipschitz continuity of  $E$ , we obtain

$$\begin{aligned}\|\boldsymbol{\sigma} - (\boldsymbol{\alpha} + \boldsymbol{\beta})\| &= \left\| E(z^+, z^-, u^r, u^\ell) - E(z^+, z^-, \tilde{u}, u^\ell) \right\| \\ &= \mathcal{O}(1) \|u^r - \tilde{u}\| \\ &= \mathcal{O}(1) (\|u^r - \tilde{u} + u_1 - u_2\| + \|u_1 - u_2\|).\end{aligned}$$

To estimate the first term we consider the function  $u^r - \tilde{u} + u_1 - u_2$  which is  $\mathbf{C}^2$  w.r.t.  $\boldsymbol{\alpha}, \boldsymbol{\beta}$ .

**First**, we assume that there are no approaching waves and obtain

$$\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_i, 0, \dots, 0), \quad \boldsymbol{\beta} = (0, \dots, 0, \beta_j, \dots, \beta_n), \quad i \leq j.$$

The case  $i = j$  **corresponds to** two rarefaction waves, **i.e.**,  $\alpha_i, \beta_i \geq 0$ . If  $i \leq i_o$ , then  $\boldsymbol{\alpha}'' = 0$ ,  $H(\boldsymbol{\alpha}' + \boldsymbol{\beta}')(u) = H(\boldsymbol{\beta}') \circ H(\boldsymbol{\alpha}')(u)$ , whence  $u^r = \tilde{u}$  and  $u_1 = u_2$ . If  $i > i_o$ , then  $\boldsymbol{\beta}' = 0$  and  $H(\boldsymbol{\alpha}'' + \boldsymbol{\beta}'')(u) = H(\boldsymbol{\beta}'') \circ H(\boldsymbol{\alpha}'')(u)$ , whence  $u^r = u_2$  and  $\tilde{u} = u_1$ . In all cases we get  $u_0^+ - \tilde{u}^+ + u_1 - u_2 = 0$ .

**Second**, standard considerations (see e.g. [9, § 7.3], [20, § 13.3] or [24]) and Lemma 4.4 lead to

$$\|u^r - \tilde{u} + u_1 - u_2\| = \mathcal{O}(1) \sum_{(i,j) \in \mathcal{A}_{\boldsymbol{\alpha}, \boldsymbol{\beta}}} |\alpha_i \beta_j|$$

in the general case.

Concerning  $\|u_1 - u_2\|$ , we get

$$\|u_1 - u_2\| \leq \mathcal{O}(1) \left\| H(\boldsymbol{\alpha}'') \left( T(z^+, z^-, H(\boldsymbol{\alpha}')(u^\ell)) \right) - T(z^+, z^-, H(\boldsymbol{\alpha})(u^\ell)) \right\|.$$

The equality  $H(\boldsymbol{\alpha})(u) = H(\boldsymbol{\alpha}'') \circ H(\boldsymbol{\alpha}')(u)$  and Lemma 4.8 with  $u = H(\boldsymbol{\alpha}')(u^\ell)$  lead to

$$\|u_1 - u_2\| = \mathcal{O}(1) \left\| z^+ - z^- \right\| \sum_{i > i_o} |\alpha_i|.$$

The result follows.  $\square$

Lemma 4.9 suggests that the quantity  $\|z^+ - z^-\|$  is a convenient way to measure the strength of the zero-waves associated to the coupling condition. More precisely, we define the strength of the zero-wave at a junction with parameters  $z^+, z^- \in \mathcal{Z}$  as  $\sigma = \|z^+ - z^-\|$ .

*Wave-front tracking approximate solutions*

We adapt the wave-front tracking techniques from [1, 9, 17, 18] to construct a sequence of approximate solutions to the Cauchy problem (1.5) and prove uniform **BV**-estimates in space. The approximate solutions converge towards a solution to the Cauchy problem with finitely many junctions. First, we define the approximations.

**Definition 4.10.** *Let  $\zeta \in \mathbf{BV}(\mathbb{R}; \mathcal{Z})$  be piecewise constant. For  $\varepsilon > 0$ , a continuous map*

$$u^\varepsilon : [0, +\infty[ \rightarrow \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \mathbb{R}^n)$$

*is an  $\varepsilon$ -approximate solution to (1.5) if the following conditions hold:*

- *$u^\varepsilon$  as a function of  $(t, x)$  is piecewise constant with discontinuities along finitely many straight lines in the  $(t, x)$ -plane. There are only finitely many wave-front interactions and at most two waves interact with each other. There are four types of discontinuities: shocks (or contact discontinuities), rarefaction waves, non-physical waves and zero-waves. We distinguish these waves' indexes in the sets  $\mathcal{J} = \mathcal{S} \cup \mathcal{R} \cup \mathcal{NP} \cup \mathcal{ZW}$ , the generic index in  $\mathcal{J}$  being  $\alpha$ .*

- *At a shock (or contact discontinuity)  $x_\alpha = x_\alpha(t)$ ,  $\alpha \in \mathcal{S}$ , the traces  $u^+ = u^\varepsilon(t, x_\alpha+)$  and  $u^- = u^\varepsilon(t, x_\alpha-)$  are related by  $u^+ = H_{i_\alpha}(\sigma_\alpha)(u^-)$  for  $1 \leq i_\alpha \leq n$  and wave-strength  $\sigma_\alpha$ . If the  $i_\alpha$ -th family is genuinely nonlinear, the Lax entropy condition  $\sigma_\alpha < 0$  holds and*

$$\left| \dot{x}_\alpha - \lambda_{i_\alpha}(u^+, u^-) \right| \leq \varepsilon,$$

*where  $\lambda_{i_\alpha}(u^+, u^-)$  is the wave speed described by the Rankine-Hugoniot conditions.*

- *For a rarefaction wave  $x_\alpha = x_\alpha(t)$ ,  $\alpha \in \mathcal{R}$  the traces are related by  $u^+ = H_{i_\alpha}(\sigma_\alpha)(u^-)$  for a genuinely nonlinear family  $1 \leq i_\alpha \leq n$  and wave-strength  $0 < \sigma_\alpha \leq \varepsilon$ . Moreover,*

$$\left| \dot{x}_\alpha - \lambda_{i_\alpha}(u^+) \right| \leq \varepsilon.$$

- *All non-physical fronts  $x = x_\alpha(t)$ ,  $\alpha \in \mathcal{NP}$  travel at the same speed  $\dot{x}_\alpha = \hat{\lambda}$  with  $\hat{\lambda} > \sup_{u,i} |\lambda_i(u)|$ . The total strength of all non-physical fronts is uniformly bounded by*

$$\sum_{\alpha \in \mathcal{NP}} \|u^\varepsilon(t, x_\alpha+) - u^\varepsilon(t, x_\alpha-)\| \leq \varepsilon \quad \text{for all } t > 0.$$

- *Zero-waves are located at the junctions  $x_\alpha \in \mathcal{I}$ . At a zero-wave  $x_\alpha$ ,  $\alpha \in \mathcal{ZW}$ , the traces are related by the coupling condition  $u^+ = T(\zeta(x_\alpha+), \zeta(x_\alpha-), u^-)$  for all  $t > 0$  except at the interaction times.*

- *The initial data satisfies  $\|u^\varepsilon(0, \cdot) - u_o\|_{\mathbf{L}^1(\mathbb{R}; \mathbb{R}^n)} \leq \varepsilon$ .*

Next, we prove the existence of  $\varepsilon$ -approximate solutions.

**Theorem 4.11.** *Let  $\Omega \subseteq \mathbb{R}^n$  be open,  $f$  satisfy **(f.1)**–**(f.3)** and  $\Xi$  satisfy **(Ξ.1)**–**(Ξ.3)**. Fix  $\bar{u} \in A_{i_o}$  and  $\bar{z} \in \mathcal{Z}$ . Then, there exist  $\delta > 0$  such that for all piecewise constant  $\zeta \in \mathbf{BV}(\mathbb{R}; \mathcal{Z})$  with*

$$\zeta(\mathbb{R}) \subseteq B(\bar{z}; \delta) \quad \text{and} \quad \text{TV}(\zeta) < \delta$$

and for all initial data  $u_o$  with

$$u_o(\mathbb{R}) \subseteq B(\bar{u}; \delta), \quad \text{TV}(u_o) < \delta,$$

for every  $\varepsilon$  sufficiently small there exists an  $\varepsilon$ -approximate solution to (1.5) in the sense of Definition 4.10. Moreover, the following quantities are bounded uniformly in  $\varepsilon$  sufficiently small,  $t \in \mathbb{R}_+$ ,  $x \in \mathbb{R}$ , for every piecewise constant  $\zeta$  with  $\text{TV}(\zeta) < \delta$  and for any open interval  $J$  with  $J \subseteq \mathbb{R} \setminus \{x_\alpha : \alpha \in \mathcal{ZW}\}$ :

- (a) The total variation in space  $\text{TV}(u^\varepsilon(t, \cdot))$ .
- (b) The total variation in time  $\text{TV}(u^\varepsilon(\cdot, x))$ .
- (c) The  $\mathbf{L}^1(\mathbb{R}; \mathbb{R})$ -Lipschitz constant of the map  $t \rightarrow u^\varepsilon(t, \cdot)$ .
- (d) The  $\mathbf{L}^1(\mathbb{R}_+; \mathbb{R}^n)$ -Lipschitz constant of the map  $x \rightarrow u^\varepsilon(\cdot, x)$  defined on  $J$ .

**Proof.** *Description of the wave front tracking algorithm.* Let  $\varepsilon$  and  $\text{TV}(\zeta)$  be sufficiently small, then we construct the approximate solution in the following way. (For notational convenience, we drop the  $\varepsilon$ .)

- To obtain piecewise constant approximate solutions, we discretize the rarefactions as in [9]. For a fixed small parameter  $\delta_R$ , each rarefaction of size  $\sigma$  is divided into  $m = \lceil \sigma/\delta_R \rceil + 1$  wave-fronts<sup>1</sup>, each one with size  $\sigma/m \leq \delta_R$ .
- Given initial data  $u_o$ , we can define a piecewise constant approximation  $u(0, \cdot)$  satisfying the requirements of Definition 4.10 and

$$\text{TV}(u(0, \cdot)) \leq \text{TV}(u_o).$$

For small  $t$ ,  $u(t, x)$  is constructed by solving the generalized Riemann problem at every point  $x_\alpha$  with  $\alpha \in \mathcal{ZW}$  and by solving the homogeneous Riemann Problem at every remaining discontinuity in  $u(0, \cdot)$ .

- At every interaction point, a new Riemann Problem arises. Notice that because of their fixed speed, two non-physical fronts cannot interact with each other, neither can the zero-waves. Moreover, by a slight modification of the speed of some waves (only among shocks, contact discontinuities and rarefactions), it is possible to achieve the property that not more than two wave-fronts interact at a point.

After each interaction time, the number of wave-fronts may increase. In order to prevent this number to become infinite in finite time, a specific treatment has been proposed for waves whose strength is below a threshold value  $\rho$  by means of a simplified Riemann solver [9, § 7.2].

Suppose that two wave-fronts of strengths  $\sigma, \sigma'$  interact at a given point  $(t, x)$ . If  $x \neq x_\alpha$ ,  $\alpha \in \mathcal{ZW}$ , we use the classical accurate or simplified homogeneous Riemann solver as in [9, § 7.2]. Assume now that  $x = x_\alpha$ ,  $\alpha \in \mathcal{ZW}$ . We briefly recall the different situations that can occur, see [1] for more details.

---

<sup>1</sup> $\lceil x \rceil$  stands for the integer part of  $x$ .

- If the wave approaching the zero wave is physical and  $|\sigma \sigma'| \geq \rho$  we use the (accurate) generalized Riemann solver.
- If the wave approaching the zero wave is physical and  $|\sigma \sigma'| < \rho$ , we use a simplified Riemann solver. Assume that the wave-front on the right is the zero-wave. Let  $u_l, u_m = H_i(\sigma)(u_l), u_r = T(\zeta(x_\alpha+), \zeta(x_\alpha-), u_m)$  be the states before the interaction. We define the auxiliary states

$$\tilde{u}_m = T(\zeta(x_\alpha+), \zeta(x_\alpha-), u_l), \quad \tilde{u}_r = H_i(\sigma)(\tilde{u}_m).$$

Then, three fronts propagate after the interaction: the zero-wave  $(u_l, \tilde{u}_m)$ , the physical front  $(\tilde{u}_m, \tilde{u}_r)$  and the non-physical one  $(\tilde{u}_r, u_r)$ . Due to the commutation defect, we use Lemma 4.8 to ensure that the introduced error, i.e. the size of the generated non-physical wave, is of second order.

- Suppose now that the wave on the left belongs to  $\mathcal{NP}$ . Again we use a simplified solver: let  $u_l, u_m, u_r = T(\zeta(x_\alpha+), \zeta(x_\alpha-), u_m)$  be the states before the interaction and define the new state  $\tilde{u}_l = T(\zeta(x_\alpha+), \zeta(x_\alpha-), u_l)$ . After the interaction time, only two fronts propagate: the zero-wave  $(u_l, \tilde{u}_l)$  and the non-physical wave  $(\tilde{u}_l, u_r)$ . Lemma 4.4 ensures that the error we made is quadratic.

*Stability of the algorithm.* We recall how junctions are taken care in [1], within the Glimm functionals [25]:

$$V(t) = \sum_{\alpha \in \text{SUR} \cup \mathcal{NP} \cup \mathcal{ZW}} |\sigma_\alpha|, \quad Q(t) = \sum_{\alpha, \beta \in \tilde{\mathcal{A}}} |\sigma_\alpha \sigma_\beta|, \quad (4.14)$$

measuring respectively the total wave strengths and the interaction potential in  $u(t, \cdot)$ . Remember that if  $\alpha \in \mathcal{ZW}$  then the strength of the wave located in  $x_\alpha$  is given by  $\sigma_\alpha = \|\zeta(x_\alpha+) - \zeta(x_\alpha-)\|$ . Notice that there exists a constant  $C > 1$  (see Lemma 4.6) such that

$$\frac{1}{C} (\text{TV}(u(t, \cdot)) + \text{TV}(\zeta)) \leq V(t) \leq C (\text{TV}(u(t, \cdot)) + \text{TV}(\zeta)).$$

Thus, according to the estimates in Lemma 4.4 and Lemma 4.9 and to the classical ones [9, Lemma 7.2], at every time  $\tau$  when two waves of strengths  $\sigma, \sigma'$  interact, we get:

$$V(\tau+) - V(\tau-) \leq C |\sigma \sigma'|, \quad (4.15)$$

$$Q(\tau+) - Q(\tau-) \leq (C V(\tau-) - 1) |\sigma \sigma'|. \quad (4.16)$$

Therefore, if  $V$  is sufficiently small, (4.16) implies

$$Q(\tau+) - Q(\tau-) \leq -\frac{1}{2} |\sigma \sigma'|. \quad (4.17)$$

By (4.15) and (4.17) we can choose a constant  $C$  large enough and  $\delta_* > 0$  so that (4.17) holds and the quantity

$$\Upsilon(t) = V(t) + C Q(t) \quad (4.18)$$

decreases at every interaction time  $\tau$  provided that  $V(\tau-)$  is sufficiently small. Thus, by standard arguments [1], choosing initial data  $u_o$  satisfying

$$\text{TV}(u_o) + \text{TV}(\zeta) \leq \delta, \quad (4.19)$$

ensures that the  $\varepsilon$ -approximate solution satisfies for any  $t \geq 0$ ,

$$\mathrm{TV}(u(t, \cdot)) + \mathrm{TV}(\zeta) \leq \delta_*. \quad (4.20)$$

The same arguments used in [1] allow to control the total number of wave fronts, that the maximal strength of a rarefaction wavelet is bounded by  $\mathcal{O}(1)\varepsilon$ , that the sum of the strengths of all  $\mathcal{NP}$  waves is also bounded by  $\mathcal{O}(1)\varepsilon$ .

Now, (a) follows from (4.20) which, together with the boundedness of wave speeds, proves (c). The same techniques in [20, § 14.5, in particular Formula (14.19)] allow to verify (b), the non resonance condition playing here an essential role. Again, this condition and (b) imply (d), see also [1, Theorem 2, Definition 2 and Remark 3].  $\square$

*Passing to the Limit  $\varepsilon \rightarrow 0$*

**Theorem 4.12.** *Let  $\Omega \subseteq \mathbb{R}^n$  be open,  $f$  satisfy (f.1)–(f.3) and  $\Xi$  satisfy (Ξ.1)–(Ξ.3). Fix  $\bar{u} \in A_{i_o}$  and  $\bar{z} \in \mathcal{Z}$ . Then, there exist  $\delta > 0$  such that for all piecewise constant  $\zeta \in \mathbf{BV}(\mathbb{R}; \mathcal{Z})$  with*

$$\zeta(\mathbb{R}) \subseteq B(\bar{z}; \delta) \quad \text{and} \quad \mathrm{TV}(\zeta) < \delta$$

and for all initial data  $u_o$  with

$$u_o(\mathbb{R}) \subseteq B(\bar{u}; \delta), \quad \mathrm{TV}(u_o) < \delta,$$

the Cauchy Problem (1.5) admits a solution  $u$  in the sense of Definition 2.1 enjoying the properties:

- (1) *The maps  $t \rightarrow \mathrm{TV}(u(t, \cdot))$  and  $t \rightarrow \|u(t, \cdot)\|_{\mathbf{L}^\infty(\mathbb{R}; \mathbb{R}^n)}$  are uniformly bounded and the map  $x \rightarrow u(t, x)$  is left continuous, for all  $t \geq 0$ .*
- (2) *For all  $x \in \mathbb{R}$ , the map  $t \rightarrow u(t, x)$  admits a representative  $\tilde{u}_x$  such that  $\mathrm{TV}(\tilde{u}_x)$  is uniformly bounded.*
- (3) *For all  $t \geq 0$ ,  $u(t, \cdot) \in \mathbf{L}_{\mathrm{loc}}^1(\mathbb{R}; \mathbb{R}^n)$  and the map  $t \rightarrow u(t, \cdot)$  is  $\mathbf{L}^1(\mathbb{R}; \mathbb{R}^n)$ -Lipschitz continuous.*
- (4) *For all  $T > 0$  and for all open interval  $J \subseteq \mathbb{R} \setminus \mathcal{I}$ , the map  $x \rightarrow u(\cdot, x)$  is  $\mathbf{L}^1([0, T]; \mathbb{R}^n)$ -Lipschitz continuous, with a Lipschitz constant independent of  $J$ ,  $\mathcal{I}$  being the set of points of jump of  $\zeta$ .*

**Proof.** For  $\varepsilon > 0$  sufficiently small, fix an  $\varepsilon$ -approximate solution  $u^\varepsilon$ . By Theorem 4.11,  $u^\varepsilon$  satisfies (1)–(2)–(3)–(4), all bounds being uniform in  $\varepsilon$ . By Helly Theorem as extended in [9, § 2.5], there exists a map  $u: [0, +\infty[ \times \mathbb{R} \rightarrow \mathbb{R}^n$  such that, up to a subsequence,  $u^\varepsilon(t, \cdot)$  converges to  $u(t, \cdot)$  in  $\mathbf{L}_{\mathrm{loc}}^1(\mathbb{R}; \mathbb{R}^n)$  for all  $t \in [0, +\infty[$  and  $u$  satisfies (1), (3).

We now prove that  $u$  satisfies (4). By possibly passing to subsequences, we may assume that  $u^\varepsilon(\cdot, x) \rightarrow u(\cdot, x)$  for a.e.  $x \in \mathbb{R}$  in  $\mathbf{L}^1([0, T]; \mathbb{R}^n)$ . Since  $u^\varepsilon$  satisfies (4), we may pass the Lipschitz continuity of  $x \rightarrow u^\varepsilon(\cdot, x)$  to the limit  $\varepsilon \rightarrow 0$  for a.e.  $x \in \mathbb{R}$ . The limit  $u$  is left continuous in the space variable  $x$  by (1), hence  $u$  satisfies (4).

We now prove that for all  $x \in \mathbb{R}$ ,  $u^\varepsilon(\cdot, x) \rightarrow u(\cdot, x)$  in  $\mathbf{L}^1([0, T]; \mathbb{R}^n)$ . To this aim, fix an arbitrary  $x \in \mathbb{R}$  and  $y < x$  such that  $]y, x[ \subset \mathbb{R} \setminus \mathcal{I}$  and  $u^\varepsilon(\cdot, y) \rightarrow u(\cdot, y)$  in  $\mathbf{L}^1([0, T]; \mathbb{R}^n)$ . Both  $x \rightarrow u^\varepsilon(\cdot, x)$  and  $x \rightarrow u(\cdot, x)$  are Lipschitz continuous, hence

$$\int_0^T \|u^\varepsilon(t, x) - u(t, x)\| dt \leq \mathcal{O}(1)|x - y| + \int_0^T \|u^\varepsilon(t, y) - u(t, y)\| dt ;$$

$$\limsup_{\varepsilon \rightarrow 0} \int_0^T \|u^\varepsilon(t, x) - u(t, x)\| dt \leq \mathcal{O}(1) |x - y|.$$

Letting now  $y \rightarrow x$  we obtained the desired convergence.

Note that  $\text{TV}(u^\varepsilon(\cdot, x))$  is bounded uniformly **in  $\varepsilon$  and in  $x$** , so that  $u(\cdot, x)$  admits a **BV** representative, proving **(2)**.

Finally, we prove that  $u$  solves (1.5). Choose  $\varphi \in \mathbf{C}_c^1(]0, T[ \times \mathbb{R}; \mathbb{R})$  and  $K$  so that  $\text{spt } \varphi \subseteq ]0, T[ \times ]-K, K[$ . Then,

$$\int_0^T \int_{-K}^K (u^\varepsilon \partial_t \varphi + f(u^\varepsilon) \partial_x \varphi) dx dt = \int_0^T \sum_{\alpha \in \mathcal{J}} e_{\varepsilon, \alpha}(t) \varphi(t, x_\alpha(t)) dt,$$

where  $e_{\varepsilon, \alpha}(t)$  measures the error in the Rankine–Hugoniot conditions along the discontinuity supported on  $x = x_\alpha(t)$ ,  $\alpha \in \mathcal{J}$ , i.e.,

$$e_{\varepsilon, \alpha}(t) = \dot{x}_\alpha (u^\varepsilon(t, x_\alpha(t)+) - u^\varepsilon(t, x_\alpha(t))) - (f(u^\varepsilon(t, x_\alpha(t)+)) - f(u^\varepsilon(t, x_\alpha(t)))).$$

By Definition 4.10 and standard estimates,

$$\sum_{\alpha \in \mathcal{J} \setminus \mathcal{ZW}} |e_{\varepsilon, \alpha}(t)| \leq \mathcal{O}(1) \varepsilon.$$

Since the coupling condition (1.3) holds along the zero–waves  $\alpha \in \mathcal{ZW}$ , we obtain

$$\left\| \int_0^T \int_{-K}^K (u^\varepsilon \partial_t \varphi + f(u^\varepsilon) \partial_x \varphi) dx dt + \int_0^T \sum_{\alpha \in \mathcal{ZW}} \varphi(t, x_\alpha) \Xi(\zeta(x_\alpha+), \zeta(x_\alpha), u^\varepsilon(t, x_\alpha)) dt \right\| \leq C \varepsilon. \quad (4.21)$$

As  $\varepsilon \rightarrow 0$  the first integrand above converges to the integrand on the left hand side of (4.8). Using **(Ξ.1)** and the convergence  $u^\varepsilon(\cdot, x_\alpha) \rightarrow u(\cdot, x_\alpha)$  in  $\mathbf{L}^1([0, T]; \mathbb{R}^n)$ , we prove the convergence of the second integrand in the left hand side of (4.21), obtaining

$$\int_0^{+\infty} \int_{\mathbb{R}} (u \partial_t \varphi + f(u) \partial_x \varphi) dx dt + \int_0^{+\infty} \sum_{\alpha \in \mathcal{ZW}} \varphi(t, x_\alpha) \Xi(\zeta(x_\alpha+), \zeta(x_\alpha), u(t, x_\alpha)) dt = 0,$$

completing the proof.  $\square$

### 4.3. Convergence Towards a General $\zeta$

**Proof of Theorem 2.2.** The proof consists of different steps.

*Step 1: Approximation of  $\zeta$ .* Let  $\zeta \in \mathbf{BV}(\mathbb{R}; \mathcal{Z})$ . Call  $\mathcal{I}$  the, possibly infinite, set of points of jump in  $\zeta$ . Recall that  $D\zeta$  is a finite measure. By Lusin Theorem [26, Theorem 2.24], for any  $h > 0$ , there exists a  $g^h \in \mathbf{C}_c^0(\mathbb{R}; \mathbb{R}^p)$  such that  $\|g^h(x)\| \leq 1$  and

$$\|D\zeta\| \left( \left\{ x \in \mathbb{R} : g^h(x) \neq v(x) \right\} \right) < h. \quad (4.22)$$

Introduce points  $\{x_1, \dots, x_{N_h-1}\} \in \mathbb{R}$  such that<sup>2</sup>:

<sup>2</sup>Everywhere,  $\sharp A$  stands for the (finite) cardinality of the set  $A$ .

- (i)  $x_0 = -\infty$ ,  $x_1 < -1/h$ ,  $x_{i-1} < x_i$  for  $i = 2, \dots, N_h - 1$ ,  $x_{N_h-1} > 1/h$  and  $x_{N_h} = +\infty$ .
- (ii)  $\sum_{x \in \mathcal{I}^h} \|\Delta \zeta(x)\| < h$  for a suitable set of points  $\mathcal{I}^h$  contained in  $\{x_1, x_2, \dots, x_{N_h-1}\}$ .
- (iii) Whenever  $x_i \in \mathcal{I}^h$ ,  $\text{TV}(\zeta, [x_{i-1}, x_i]) < h/(1 + \#\mathcal{I}^h)$ .
- (iv)  $\text{TV}(\zeta, ]x_{i-1}, x_i[) < h$  for all  $i = 1, \dots, N_h$ .
- (v)  $\|g^h(x') - g^h(x'')\| < h$  for  $x', x'' \in ]x_{i-1}, x_i[$ ,  $i = 1, \dots, N_h$ .
- (vi)  $x_i - x_{i-1} \in ]0, h[$  for all  $i = 2, \dots, N_h - 1$ .

Points satisfying (i) are easily constructed. Then, adding more points, one fulfills also (ii). Iteratively continuing to add points, we satisfy also (iii), (iv), (v) and (vi), in this order.

We approximate  $\zeta$  by means of the piecewise constant left continuous function

$$\zeta^h(x) = \zeta(-\infty) \chi_{]-\infty, x_1]}(x) + \sum_{i=2}^{N_h-1} \zeta(x_{i-1}+) \chi_{]x_{i-1}, x_i]}(x) + \zeta(x_{N_h-1}+) \chi_{]x_{N_h-1}, +\infty[}(x). \quad (4.23)$$

By Theorem 4.12, there exists a solution  $u^h$  to the Cauchy Problem (1.5) with  $\zeta$  substituted by  $\zeta^h$  as in (4.23). We prove that as  $h \rightarrow 0$  the solution  $u^h$  converges in  $\mathbf{L}_{\text{loc}}^1$  to a solution to (1.5), possibly up to a subsequence.

*Step 2: Select a Convergent Subsequence..* We claim that there exists a map  $u$  and a convergent subsequence, which we keep denoting  $u^h$ , such that

$$u^h(t, \cdot) \rightarrow u(t, \cdot) \text{ in } \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \mathbb{R}^n) \text{ for all } t; \quad (4.24)$$

$$u^h(\cdot, x) \rightarrow u(\cdot, x) \text{ in } \mathbf{L}_{\text{loc}}^1([0, +\infty[; \mathbb{R}^n) \text{ for all } x; \quad (4.25)$$

$$t \rightarrow u(t, \cdot) \text{ is Lipschitz continuous in } \mathbf{L}^1(\mathbb{R}; \mathbb{R}^n); \quad (4.26)$$

$$\text{TV}(u(t, \cdot)) \text{ is bounded uniformly in } t; \quad (4.27)$$

$$x \rightarrow u(t, x) \text{ is left continuous for all } t \geq 0. \quad (4.28)$$

Indeed, by (1) and (3) in Theorem 4.12, we can apply Helly Theorem as presented in [9, § 2.5] obtaining the existence of a map  $u$  satisfying (4.24), (4.26), (4.27) and (4.28).

We are left with the convergence (4.25). Introduce a point  $y < x$  and all the points of jump  $\bar{x}_0, \dots, \bar{x}_{M+1}$  (for a suitable  $M \geq 0$ ) in  $\zeta^h$  such that

$$-\infty \leq \bar{x}_0 < y \leq \bar{x}_1 < \bar{x}_2 < \dots < \bar{x}_M < x \leq \bar{x}_{M+1} \leq +\infty.$$

We now estimate

$$\begin{aligned} \text{TV}(\zeta^h; [y, x]) &= \sum_{i=1}^M \|\Delta \zeta^h(\bar{x}_i)\| = \sum_{i=1}^M \|\zeta(\bar{x}_i+) - \zeta(\bar{x}_{i-1}+)\| \\ &\leq \|\zeta(\bar{x}_1-) - \zeta(\bar{x}_0+)\| + \|\zeta(\bar{x}_1+) - \zeta(\bar{x}_1-)\| + \sum_{i=2}^M \|\zeta(\bar{x}_i+) - \zeta(\bar{x}_{i-1}+)\| \\ &\leq \text{TV}(\zeta, ]\bar{x}_0, \bar{x}_1]) + \text{TV}(\zeta, ]y, x]) \\ &\leq h + \text{TV}(\zeta, ]y, x]). \end{aligned} \quad (4.29)$$

where to get to the last line above we used **(iv)**.

Fix a positive  $T$ . By the triangle inequality, **(4)** in Theorem 4.12, inequality (4.29) and Lemma 4.4, since  $u^h(t, \bar{x}_{i+}) = T \left( \zeta^h(\bar{x}_{i+}), \zeta^h(\bar{x}_i), u^h(t, x_i) \right)$ ,

$$\begin{aligned}
& \int_0^T \left\| u^h(t, x) - u^h(t, y) \right\| dt \\
& \leq \int_0^T \left\| u^h(t, x) - u^h(t, \bar{x}_M+) \right\| dt + \int_0^T \left\| u^h(t, \bar{x}_M+) - u^h(t, \bar{x}_M) \right\| dt \\
& \quad + \sum_{i=1}^{M-1} \left( \int_0^T \left\| u^h(t, \bar{x}_{i+1}) - u^h(t, \bar{x}_i+) \right\| dt + \int_0^T \left\| u^h(t, \bar{x}_i+) - u^h(t, \bar{x}_i) \right\| dt \right) \\
& \quad + \int_0^T \left\| u^h(t, \bar{x}_1) - u^h(t, y+) \right\| dt + \int_0^T \left\| u^h(t, y+) - u^h(t, y) \right\| dt \\
& \leq \mathcal{O}(1) |x - \bar{x}_M| + \mathcal{O}(1) \left\| \Delta \zeta^h(\bar{x}_M) \right\| \\
& \quad + \mathcal{O}(1) \sum_{i=1}^{M-1} \left( |\bar{x}_{i+1} - \bar{x}_i| + \left\| \Delta \zeta^h(\bar{x}_i) \right\| \right) + \mathcal{O}(1) |\bar{x}_1 - y| + \mathcal{O}(1) \left\| \Delta \zeta^h(y) \right\| \\
& \leq \mathcal{O}(1) \left( |x - y| + \text{TV} \left( \zeta^h, [y, x] \right) \right) \\
& \leq \mathcal{O}(1) \left( |x - y| + h + \text{TV} \left( \zeta, [y, x] \right) \right). \tag{4.30}
\end{aligned}$$

Since  $u^h$  converges to  $u$  in  $\mathbf{L}_{\text{loc}}^1([0, +\infty[ \times \mathbb{R}, \mathbb{R}^n)$  too, possibly passing to a subsequence, we may assume that for a.e.  $x \in \mathbb{R}$  we have  $u^h(\cdot, x) \rightarrow u(\cdot, x)$  in  $\mathbf{L}_{\text{loc}}^1([0, +\infty[; \mathbb{R}^n)$ . Pass to the limit  $h \rightarrow 0$  in (4.30) and obtain that for a.e.  $x, y \in \mathbb{R}$  with  $y < x$ ,

$$\int_0^T \left\| u(t, x) - u(t, y) \right\| dt \leq \mathcal{O}(1) \left( |x - y| + \text{TV} \left( \zeta, [y, x] \right) \right). \tag{4.31}$$

By the left continuity of  $x \rightarrow u(t, x)$  and of the right hand side of (4.31) (with respect to both  $x$  and  $y$ ), the inequality (4.31) holds for *all*  $x, y \in \mathbb{R}$  with  $y < x$ .

Fix now an arbitrary  $x \in \mathbb{R}$  and choose  $y \in \mathbb{R}$  with  $y < x$  and such that  $u^h(\cdot, y) \rightarrow u(\cdot, y)$ . By the triangle inequality, (4.30) and (4.31), we have

$$\int_0^T \left\| u^h(t, x) - u(t, x) \right\| dt \leq \mathcal{O}(1) \left( |x - y| + h + \text{TV} \left( \zeta, [y, x] \right) \right) + \int_0^T \left\| u^h(t, y) - u(t, y) \right\| dt.$$

Hence, for almost every  $y < x$ ,

$$\limsup_{h \rightarrow 0} \int_0^T \left\| u^h(t, x) - u(t, x) \right\| dt \leq \mathcal{O}(1) \left( |x - y| + \text{TV} \left( \zeta, [y, x] \right) \right)$$

which proves the convergence for every  $x \in \mathbb{R}$ , since the latter right hand side vanishes as  $y \rightarrow x^-$ .

*Step 3: The Limit is a Solution..* Fix  $\varphi \in \mathbf{C}_c^1([0, +\infty[ \times \mathbb{R}; \mathbb{R})$  such that  $\text{spt } \varphi \subseteq [0, T] \times [-K, K]$  for suitable  $T, K > 0$ . Showing that the left hand side below vanishes in the limit



$h \rightarrow 0$  completes the proof.

$$\begin{aligned} & \left\| - \int_0^T \int_{-K}^K (u \partial_t \varphi + f(u) \partial_x \varphi) dt dx - \sum_{x \in \mathcal{I}, |x| \leq K} \int_0^T \Xi(\zeta(x+), \zeta(x), u(t, x)) \varphi(t, x) dt \right. \\ & \quad \left. - \int_0^T \int_{-K}^K D_{v(x)}^+ \Xi(\zeta(x), \zeta(x), u(t, x)) \varphi(t, x) d\|\mu\|(x) dt \right\| \\ & \leq \mathcal{E}_1^h + \mathcal{E}_2^h + \mathcal{E}_3^h + \mathcal{E}_4^h + \mathcal{E}_5^h + \mathcal{E}_6^h + \mathcal{E}_7^h + \mathcal{E}_8^h + \mathcal{E}_9^h + \mathcal{E}_{10}^h, \end{aligned}$$

the terms  $\mathcal{E}_1^h, \dots, \mathcal{E}_{10}^h$  are defined below. To this aim, consider the terms on the right hand side separately:

**Term  $\mathcal{E}_1^h$ :** By the  $L_{\text{loc}}^1$  convergence proved in Step 2.

$$\begin{aligned} \mathcal{E}_1^h & \stackrel{\text{def}}{=} \left\| - \int_0^T \int_{-K}^K (u \partial_t \varphi + f(u) \partial_x \varphi) dx dt + \int_0^T \int_{-K}^K (u^h \partial_t \varphi + f(u^h) \partial_x \varphi) dx dt \right\| \\ & \rightarrow 0 \quad \text{as } h \rightarrow 0. \end{aligned}$$

**Term  $\mathcal{E}_2^h$ :** Each  $u^h$  is a solution associated with  $\zeta^h$ , hence

$$- \int_0^T \int_{-K}^K (u^h \partial_t \varphi + f(u^h) \partial_x \varphi) dt dx = \sum_{i: |x_i| \leq K} \int_0^T \Xi(\zeta^h(x_{i+}), \zeta^h(x_i), u^h(t, x_i)) \varphi(t, x_i) dt$$

so that

$$\begin{aligned} \mathcal{E}_2^h & \stackrel{\text{def}}{=} \left\| - \int_0^T \int_{-K}^K (u^h \partial_t \varphi + f(u^h) \partial_x \varphi) dt dx \right. \\ & \quad \left. - \sum_{i: |x_i| \leq K} \int_0^T \Xi(\zeta^h(x_{i+}), \zeta^h(x_i), u^h(t, x_i)) \varphi(t, x_i) dt \right\| \\ & = 0. \end{aligned}$$

**Term  $\mathcal{E}_3^h$ :** Recall that by (4.23),  $\zeta^h(x_i) = \zeta(x_{i-1+})$  and  $\zeta^h(x_{i+}) = \zeta(x_{i+})$ . By the Lipschitz continuity of  $\Xi$  and (iii)

$$\begin{aligned} \mathcal{E}_3^h & \stackrel{\text{def}}{=} \left\| \sum_{i: |x_i| \leq K, x_i \in \mathcal{I}^h} \int_0^T \Xi(\zeta^h(x_{i+}), \zeta^h(x_i), u^h(t, x_i)) \varphi(t, x_i) dt \right. \\ & \quad \left. - \sum_{i: |x_i| \leq K, x_i \in \mathcal{I}^h} \int_0^T \Xi(\zeta(x_{i+}), \zeta(x_i), u^h(t, x_i)) \varphi(t, x_i) dt \right\| \\ & \leq \mathcal{O}(1) \sum_{i: |x_i| \leq K, x_i \in \mathcal{I}^h} \|\zeta(x_{i-1+}) - \zeta(x_i)\| \end{aligned}$$

$$\begin{aligned}
&\leq \mathcal{O}(1) \#\mathcal{I}^h \frac{h}{1 + \#\mathcal{I}^h} \\
&\leq \mathcal{O}(1) h \\
&\rightarrow 0 \quad \text{as } h \rightarrow 0.
\end{aligned}$$

**Term  $\mathcal{E}_4^h$ :** Recall that if  $x_i \notin \mathcal{I}$ , then  $\zeta(x_{i+}) = \zeta(x_i)$ , which implies the equality  $\Xi(\zeta(x_{i+}), \zeta(x_i), u^h(t, x_i)) = 0$ . Hence, by **(Ξ.1)**, **(Ξ.3)** and **(ii)** we compute

$$\begin{aligned}
\mathcal{E}_4^h &\stackrel{\text{def}}{=} \left\| \sum_{i: |x_i| \leq K, x_i \in \mathcal{I}^h} \int_0^T \Xi(\zeta(x_{i+}), \zeta(x_i), u^h(t, x_i)) \varphi(t, x_i) dt \right. \\
&\quad \left. - \sum_{x \in \mathcal{I}, |x| \leq K} \int_0^T \Xi(\zeta(x+), \zeta(x), u^h(t, x)) \varphi(t, x) dt \right\| \\
&\leq \left\| \sum_{x \in \mathcal{I} \setminus \mathcal{I}^h, |x| \leq K} \int_0^T \Xi(\zeta(x+), \zeta(x), u^h(t, x)) \varphi(t, x) dt \right\| \\
&\leq \mathcal{O}(1) \sum_{x \in \mathcal{I} \setminus \mathcal{I}^h, |x| \leq K} \|\Delta\zeta(x)\| \\
&\leq \mathcal{O}(1) h \\
&\rightarrow 0 \quad \text{as } h \rightarrow 0.
\end{aligned}$$

**Term  $\mathcal{E}_5^h$ :** Using Lemma 4.3

$$\begin{aligned}
\mathcal{E}_5^h &\stackrel{\text{def}}{=} \left\| \sum_{x \in \mathcal{I}, |x| \leq K} \int_0^T \Xi(\zeta(x+), \zeta(x), u^h(t, x)) \varphi(t, x) dt \right. \\
&\quad \left. - \sum_{x \in \mathcal{I}, |x| \leq K} \int_0^T \Xi(\zeta(x+), \zeta(x), u(t, x)) \varphi(t, x) dt \right\| \\
&\leq \mathcal{O}(1) \sum_{x \in \mathcal{I}} \left( \|\Delta\zeta(x)\| \int_0^T \|u^h(t, x) - u(t, x)\| dt \right) \\
&\rightarrow 0 \quad \text{as } h \rightarrow 0.
\end{aligned}$$

The last limit is due to (4.25) and the convergence of the series  $\sum_{x \in \mathcal{I}} \|\Delta\zeta(x)\|$ . This concludes the convergence to the discrete part of the measure.

**Term  $\mathcal{E}_6^h$ .** Recall that by (4.23) we have  $\zeta^h(x_i) = \zeta(x_{i-1+})$  and  $\zeta^h(x_{i+}) = \zeta(x_{i+})$ . We use below also **(ii)**:

$$\mathcal{E}_6^h \stackrel{\text{def}}{=} \left\| \sum_{i: |x_i| \leq K, x_i \notin \mathcal{I}^h} \int_0^T \Xi(\zeta^h(x_{i+}), \zeta^h(x_i), u^h(t, x_i)) \varphi(t, x_i) dt \right\|$$

$$\begin{aligned}
& - \sum_{i: |x_i| \leq K, x_i \notin \mathcal{I}^h} \int_0^T \Xi \left( \zeta(x_{i-1}+) + \mu(\cdot]x_{i-1}, x_i], \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) dt \Big\| \\
& \leq \mathcal{O}(1) \sum_{i: |x_i| \leq K, x_i \notin \mathcal{I}^h} \left\| \zeta(x_i+) - \zeta(x_{i-1}+) - \mu(\cdot]x_{i-1}, x_i] \right\| \\
& = \mathcal{O}(1) \sum_{i: |x_i| \leq K, x_i \notin \mathcal{I}^h} \left\| D\zeta(\cdot]x_{i-1}, x_i] - \mu(\cdot]x_{i-1}, x_i] \right\| \\
& \leq \mathcal{O}(1) \sum_{x \in \mathcal{I} \setminus \mathcal{I}^h} \left\| \Delta\zeta(x) \right\| \\
& = \mathcal{O}(1) h \\
& \rightarrow 0 \quad \text{as } h \rightarrow 0.
\end{aligned}$$

**Term  $\mathcal{E}_7^h$ .** Using (iii),

$$\begin{aligned}
\mathcal{E}_7^h & \stackrel{\text{def}}{=} \left\| \sum_{i: |x_i| \leq K, x_i \notin \mathcal{I}^h} \int_0^T \Xi \left( \zeta(x_{i-1}+) + \mu(\cdot]x_{i-1}, x_i], \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) dt \right. \\
& \quad \left. - \sum_{i: |x_i| \leq K} \int_0^T \Xi \left( \zeta(x_{i-1}+) + \mu(\cdot]x_{i-1}, x_i], \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) dt \right\| \\
& \leq \mathcal{O}(1) \sum_{i: |x_i| \leq K, x_i \in \mathcal{I}^h} \text{TV}(\zeta; \cdot]x_{i-1}, x_i]) \\
& \leq \mathcal{O}(1) \#\mathcal{I}^h \frac{h}{1 + \#\mathcal{I}^h} \\
& \leq \mathcal{O}(1) h \\
& \rightarrow 0 \quad \text{as } h \rightarrow 0.
\end{aligned}$$

**Term  $\mathcal{E}_8^h$ .** Introduce now  $\delta_i = \|\mu\|(\cdot]x_{i-1}, x_i])$ ,  $\mathcal{J} = \{i \in \{1, \dots, N_h\} : \delta_i \neq 0\}$  and for  $i \in \mathcal{J}$ , let  $v_i = \mu(\cdot]x_{i-1}, x_i])/\delta_i$ . Below, we use (E.4) with  $\delta_i$  for  $t$  and  $v_i$  for  $v$ , and (iv):

$$\begin{aligned}
\mathcal{E}_8^h & \stackrel{\text{def}}{=} \left\| \sum_{i: |x_i| \leq K} \int_0^T \Xi \left( \zeta(x_{i-1}+) + \mu(\cdot]x_{i-1}, x_i], \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) dt \right. \\
& \quad \left. - \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_0^T \delta_i D_{v_i}^+ \Xi \left( \zeta(x_{i-1}+), \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) dt \right\| \\
& \leq \mathcal{O}(1) \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_0^T \left\| \Xi \left( \zeta(x_{i-1}+) + \mu(\cdot]x_{i-1}, x_i], \zeta(x_{i-1}+), u^h(t, x_i) \right) \right. \\
& \quad \left. - \delta_i D_{v_i}^+ \Xi \left( \zeta(x_{i-1}+), \zeta(x_{i-1}+), u^h(t, x_i) \right) \right\| dt
\end{aligned}$$

$$\begin{aligned}
&\leq \mathcal{O}(1) \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \sigma(\delta_i) \delta_i \\
&\leq \mathcal{O}(1) \sigma(h) \text{TV}(\zeta) \\
&\rightarrow 0 \quad \text{as } h \rightarrow 0.
\end{aligned}$$

**Term  $\mathcal{E}_9^h$ .** Use **(Ξ.4)** and recall that by (1.6),  $v_i = (1/\delta_i) \int_{]x_{i-1}, x_i[} v(y) d\|\mu\|(y)$ , while clearly  $v(x) = (1/\delta_i) \int_{]x_{i-1}, x_i[} v(x) d\|\mu\|(y)$ . We also use  $g^h$ , that is defined in **Step 1** and satisfies (4.22).

$$\begin{aligned}
\mathcal{E}_9^h &\stackrel{\text{def}}{=} \left\| \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_0^T \delta_i D_{v_i}^+ \Xi \left( \zeta(x_{i-1}+), \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) dt \right. \\
&\quad \left. - \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_0^T \int_{]x_{i-1}, x_i[} D_{v(x)}^+ \Xi \left( \zeta(x_{i-1}+), \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) d\|\mu\|(x) dt \right\| \\
&\leq \mathcal{O}(1) \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_{]x_{i-1}, x_i[} \|v(x) - v_i\| d\|\mu\|(x) \\
&\leq \mathcal{O}(1) \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \frac{1}{\delta_i} \int_{]x_{i-1}, x_i[^2} \|v(x) - v(y)\| d(\|\mu\| \otimes \|\mu\|)(x, y) \\
&\leq \mathcal{O}(1) \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \frac{1}{\delta_i} \int_{]x_{i-1}, x_i[^2} \left[ \|v(x) - g^h(x)\| + \|g^h(y) - v(y)\| \right] d(\|\mu\| \otimes \|\mu\|)(x, y) \quad (4.32) \\
&\quad + \mathcal{O}(1) \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \frac{1}{\delta_i} \int_{]x_{i-1}, x_i[^2} \|g^h(x) - g^h(y)\| d(\|\mu\| \otimes \|\mu\|)(x, y). \quad (4.33)
\end{aligned}$$

The two terms in the integral in (4.32) are estimated in the same way, using (4.22), as

$$\begin{aligned}
&\sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_{]x_{i-1}, x_i[^2} \frac{1}{\delta_i} \|v(x) - g^h(x)\| d(\|\mu\| \otimes \|\mu\|)(x, y) \\
&\leq \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_{]x_{i-1}, x_i[} \|v(x) - g^h(x)\| d\|\mu\|(x) \\
&\leq \int_{\mathbb{R}} \|v(x) - g^h(x)\| d\|\mu\|(x) \\
&\leq \int_{\{x \in \mathbb{R}: v(x) \neq g^h(x)\}} \left( \|v(x)\| + \|g^h(x)\| \right) d\|\mu\|(x) \\
&\leq 2h \\
&\rightarrow 0 \quad \text{as } h \rightarrow 0.
\end{aligned}$$

We now estimate the term (4.33) by means of **(v)**:

$$\sum_{i: |x_i| \leq K, i \in \mathcal{J}} \frac{1}{\delta_i} \int_{]x_{i-1}, x_i[^2} \|g^h(x) - g^h(y)\| d(\|\mu\| \otimes \|\mu\|)(x, y)$$

$$\begin{aligned}
&\leq h \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \frac{1}{\delta_i} \int_{]x_{i-1}, x_i]^2} d(\|\mu\| \otimes \|\mu\|)(x, y) \\
&\leq h \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \delta_i \\
&\leq h \text{TV}(\zeta) \\
&\rightarrow 0 \quad \text{as } h \rightarrow 0.
\end{aligned}$$

**Term  $\mathcal{E}_{10}^h$ .** Using **(E.4)**

$$\begin{aligned}
\mathcal{E}_{10}^h &\stackrel{\text{def}}{=} \left\| \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_0^T \int_{]x_{i-1}, x_i[} D_{v(x)}^+ \Xi \left( \zeta(x_{i-1}+), \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) d\|\mu\|(x) dt \right. \\
&\quad \left. - \int_0^T \int_{-K}^K D_{v(x)}^+ \Xi \left( \zeta(x), \zeta(x), u(t, x) \right) \varphi(t, x) d\|\mu\|(x) dt \right\| \\
&= \left\| \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_0^T \int_{]x_{i-1}, x_i[} D_{v(x)}^+ \Xi \left( \zeta(x_{i-1}+), \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) d\|\mu\|(x) dt \right. \\
&\quad \left. - \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_0^T \int_{]x_{i-1}, x_i[} D_{v(x)}^+ \Xi \left( \zeta(x), \zeta(x), u(t, x) \right) \varphi(t, x) d\|\mu\|(x) dt \right\| \\
&\leq \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_0^T \int_{]x_{i-1}, x_i[} \left\| D_{v(x)}^+ \Xi \left( \zeta(x_{i-1}+), \zeta(x_{i-1}+), u^h(t, x_i) \right) \varphi(t, x_i) \right. \\
&\quad \left. - D_{v(x)}^+ \Xi \left( \zeta(x), \zeta(x), u(t, x) \right) \varphi(t, x) \right\| d\|\mu\|(x) dt \\
&\leq \mathcal{O}(1) \sum_{i: |x_i| \leq K, i \in \mathcal{J}} \int_0^T \int_{]x_{i-1}, x_i[} \left( \left\| \zeta(x_{i-1}+) - \zeta(x) \right\| + \left\| u^h(t, x_i) - u^h(t, x) \right\| \right. \\
&\quad \left. + \left\| u^h(t, x) - u(t, x) \right\| + |x_i - x| \right) d\|\mu\|(x) dt.
\end{aligned}$$

Observe that

$$\begin{aligned}
\left\| \zeta(x_{i-1}+) - \zeta(x) \right\| &\leq h && \text{by (iv)} \\
|x_i - x| &\leq h && \text{by (vi)} \\
\int_0^T \left\| u^h(t, x_i) - u^h(t, x) \right\| dt &\leq \mathcal{O}(1) h && \text{by (4.30), (iv) and (vi)}
\end{aligned}$$

while by (4.25), Fubini Theorem and the Dominated Convergence Theorem,

$$\int_{\mathbb{R}} \int_0^T \left\| u^h(t, x) - u(t, x) \right\| dt d\|\mu\|(x) \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

The proof is completed.  $\square$

#### 4.4. Proof Relative to Section 3

**Proof of Theorem 3.1.** It is immediate to check that (f.1)–(f.3) hold, thanks to (p). Define  $\Xi$  as in (3.3). Then, conditions (Ξ.1) and (Ξ.2) follow from the assumed  $\mathbf{C}^2$  regularity of  $K$  in all its variables. Condition (Ξ.3) follows from (3.3) and  $K(0, (\rho, q)) \equiv 0$ . Concerning (Ξ.4), we have

$$D_v^+ \Xi(z, z, v) = \begin{bmatrix} 0 \\ \partial_1 K(0, (\rho, q)) \|v\| \end{bmatrix}$$

indeed, for  $v$  such that  $\|v\| \leq 1$ , we can estimate

$$\begin{aligned} & \left\| K(t\|v\|, (\rho, q)) - \|v\| \partial_1 K(0, (\rho, q)) t \right\| \\ &= \left\| \int_0^1 \left( \partial_1 K(st\|v\|, (\rho, q)) - \partial_1 K(0, (\rho, q)) \right) t \|v\| ds \right\| \\ &\leq \|K\|_{\mathbf{C}^2([0,r] \times \Omega; \mathbb{R})} t^2 \end{aligned}$$

proving (Ξ.4) with  $\sigma(t) = \|K\|_{\mathbf{C}^2([0,r] \times \Omega; \mathbb{R})} t$ .

Theorem 2.2 can then be applied, exhibiting the existence of a solution in the sense of Definition 2.1.

To obtain the formulation (3.5) from (2.2), only the two terms in the right hand side of the second equations need to be considered. The first one is immediate: it only requires the substitution (3.3). Concerning the second one, recall that by (1.6) and by the  $\mathbf{C}^2$  regularity of  $\Gamma$ ,  $d\mu(x) = \Gamma''(x) dx$ , so that  $d\|\mu\|(x) = \|\Gamma''(x)\| dx$ , and that  $v(x) = \frac{\Gamma''(x)}{\|\Gamma''(x)\|}$  for a.e.  $x$  with respect to the measure  $\|\mu\|$ .

Hence, since  $\|v(x)\| = 1$  for a.e.  $x$  with respect to the measure  $\|\mu\|$ ,

$$D_{v(x)}^+ \Xi(\Gamma'(x), \Gamma'(x), (\rho, q)) d\|\mu\|(x) = \partial_1 K(0, (\rho, q)(x)) \|\Gamma''(x)\| dx$$

completing the proof.  $\square$

**Proof of Theorem 3.2.** Condition (p) ensures that (f.1)–(f.3) hold. The choice (3.8) and the assumptions on  $\Xi_2$  imply that (Ξ.1)–(Ξ.4) hold. Since the distributional derivative of  $a$  has neither Cantor part nor atomic part, due to (3.10) problem (3.9) reduces to (3.6).  $\square$

**Acknowledgment.** The first and second authors were partly supported by the GNAMPA 2020 project "From Wellposedness to Game Theory in Conservation Laws". The work of the third author has been funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Projektnummer 320021702/GRK2326 Energy, Entropy, and Dissipative Dynamics (EDDy).

The authors express their deep gratitude to the anonymous referee that dived into the deepest details of this paper providing non trivial corrections as well as hints that helped to improve the presentation.

## References

- [1] D. Amadori, L. Gosse, G. Guerra, Global BV entropy solutions and uniqueness for hyperbolic systems of balance laws, *Arch. Ration. Mech. Anal.* 162 (4) (2002) 327–366. doi:10.1007/s002050200198.  
URL <https://doi.org/10.1007/s002050200198>
- [2] R. M. Colombo, M. Garavello, On the Cauchy problem for the  $p$ -system at a junction, *SIAM J. Math. Anal.* 39 (5) (2008) 1456–1471. doi:10.1137/060665841.  
URL <https://doi.org/10.1137/060665841>
- [3] B. Boutin, F. Coquel, P. G. LeFloch, Coupling techniques for nonlinear hyperbolic equations. II. Resonant interfaces with internal structure, *Netw. Heterog. Media* 16 (2) (2021) 283–315. doi:10.3934/nhm.2021007.  
URL <https://doi.org/10.3934/nhm.2021007>
- [4] G.-Q. Chen, J. Glimm, Global solutions to the compressible Euler equations with geometrical structure, *Comm. Math. Phys.* 180 (1) (1996) 153–193.  
URL <http://projecteuclid.org/euclid.cmp/1104287238>
- [5] P. Goatin, P. G. LeFloch, The Riemann problem for a class of resonant hyperbolic systems of balance laws, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 21 (6) (2004) 881–902. doi:10.1016/j.anihpc.2004.02.002.  
URL <https://doi.org/10.1016/j.anihpc.2004.02.002>
- [6] E. Isaacson, B. Temple, Nonlinear resonance in systems of conservation laws, *SIAM J. Appl. Math.* 52 (5) (1992) 1260–1278. doi:10.1137/0152073.  
URL <https://doi.org/10.1137/0152073>
- [7] W.-C. Lien, Hyperbolic conservation laws with a moving source, *Comm. Pure Appl. Math.* 52 (9) (1999) 1075–1098. doi:10.1002/(SICI)1097-0312(199909)52:9<1075::AID-CPA2>3.3.CO;2-W.  
URL [https://doi.org/10.1002/\(SICI\)1097-0312\(199909\)52:9<1075::AID-CPA2>3.3.CO;2-W](https://doi.org/10.1002/(SICI)1097-0312(199909)52:9<1075::AID-CPA2>3.3.CO;2-W)
- [8] L. Ambrosio, N. Fusco, D. Pallara, Functions of bounded variation and free discontinuity problems, *Oxford Mathematical Monographs*, The Clarendon Press Oxford University Press, New York, 2000.
- [9] A. Bressan, *Hyperbolic systems of conservation laws*, Vol. 20 of Oxford Lecture Series in Mathematics and its Applications, Oxford University Press, Oxford, 2000, the one-dimensional Cauchy problem.
- [10] R. M. Colombo, H. Holden, Isentropic fluid dynamics in a curved pipe, *Z. Angew. Math. Phys.* 67 (5) (2016) Art. 131, 10. doi:10.1007/s00033-016-0725-0.  
URL <https://doi.org/10.1007/s00033-016-0725-0>
- [11] H. Holden, N. H. Risebro, Riemann problems with a kink, *SIAM J. Math. Anal.* 30 (3) (1999) 497–515. doi:10.1137/S0036141097327033.  
URL <https://doi.org/10.1137/S0036141097327033>
- [12] M. K. Banda, M. Herty, A. Klar, Coupling conditions for gas networks governed by the isothermal Euler equations, *Netw. Heterog. Media* 1 (2) (2006) 295–314. doi:10.3934/nhm.2006.1.295.
- [13] M. K. Banda, M. Herty, A. Klar, Gas flow in pipeline networks, *Netw. Heterog. Media* 1 (1) (2006) 41–56. doi:10.3934/nhm.2006.1.41.

- [14] R. M. Colombo, M. Garavello, On the  $p$ -system at a junction, in: Control methods in PDE-dynamical systems, Vol. 426 of Contemp. Math., Amer. Math. Soc., Providence, RI, 2007, pp. 193–217. doi:10.1090/conm/426/08189.
- [15] R. M. Colombo, M. Garavello, On the 1D modeling of fluid flowing through a junction, Discrete Contin. Dyn. Syst. Ser. B 25 (10) (2020) 3917–3929. doi:10.3934/dcdsb.2019149.  
URL <https://doi.org/10.3934/dcdsb.2019149>
- [16] R. M. Colombo, F. Marcellini, Coupling conditions for the  $3 \times 3$  Euler system, Netw. Heterog. Media 5 (4) (2010) 675–690. doi:10.3934/nhm.2010.5.675.  
URL <https://doi.org/10.3934/nhm.2010.5.675>
- [17] R. M. Colombo, F. Marcellini, Smooth and discontinuous junctions in the  $p$ -system, J. Math. Anal. Appl. 361 (2) (2010) 440–456. doi:10.1016/j.jmaa.2009.07.022.  
URL <https://doi.org/10.1016/j.jmaa.2009.07.022>
- [18] G. Guerra, F. Marcellini, V. Schleper, Balance laws with integrable unbounded sources, SIAM J. Math. Anal. 41 (3) (2009) 1164–1189. doi:10.1137/080735436.  
URL <https://doi.org/10.1137/080735436>
- [19] P. D. Lax, Hyperbolic systems of conservation laws. II, Comm. Pure Appl. Math. 10 (1957) 537–566. doi:10.1002/cpa.3160100406.  
URL <https://doi.org/10.1002/cpa.3160100406>
- [20] C. M. Dafermos, Hyperbolic conservation laws in continuum physics, 4th Edition, Vol. 325 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer-Verlag, Berlin, 2016. doi:10.1007/978-3-662-49451-6.  
URL <https://doi.org/10.1007/978-3-662-49451-6>
- [21] T. P. Liu, Quasilinear hyperbolic systems, Comm. Math. Phys. 68 (2) (1979) 141–172.  
URL <http://projecteuclid.org/euclid.cmp/1103905309>
- [22] T. P. Liu, Nonlinear stability and instability of transonic flows through a nozzle, Comm. Math. Phys. 83 (2) (1982) 243–260.  
URL <http://projecteuclid.org/euclid.cmp/1103920803>
- [23] R. M. Colombo, G. Guerra, On general balance laws with boundary, J. Differential Equations 248 (5) (2010) 1017–1043. doi:10.1016/j.jde.2009.12.002.  
URL <https://doi.org/10.1016/j.jde.2009.12.002>
- [24] W.-A. Yong, A simple approach to Glimm’s interaction estimates, Appl. Math. Lett. 12 (2) (1999) 29–34. doi:10.1016/S0893-9659(98)00144-X.  
URL [https://doi.org/10.1016/S0893-9659\(98\)00144-X](https://doi.org/10.1016/S0893-9659(98)00144-X)
- [25] J. Glimm, Solutions in the large for nonlinear hyperbolic systems of equations, Comm. Pure Appl. Math. 18 (1965) 697–715. doi:10.1002/cpa.3160180408.  
URL <https://doi.org/10.1002/cpa.3160180408>
- [26] W. Rudin, Real and complex analysis, 3rd Edition, McGraw-Hill Book Co., New York, 1987.