



OPEN

Severe neurological outcomes after very early bilateral nephrectomies in patients with autosomal recessive polycystic kidney disease (ARPKD)

Kathrin Burgmaier¹, Gema Ariceta², Martin Bald³, Anja Katrin Buescher⁴, Mathias Burgmaier⁵, Florian Erger^{6,7}, Michaela Gessner⁸, Ibrahim Gokce⁹, Jens König¹⁰, Claudia Kowalewska¹¹, Laura Massella¹², Antonio Mastrangelo¹³, Djalila Mekahli^{14,15}, Lars Pape^{4,16}, Ludwig Patzer¹⁷, Alexandra Potemkina¹⁸, Gesa Schalk¹⁹, Raphael Schild²⁰, Rukshana Shroff²¹, Maria Szczepanska²², Katarzyna Taranta-Janusz²³, Marcin Tkaczyk²⁴, Lutz Thorsten Weber¹, Elke Wühl²⁵, Donald Wurm²⁶, Simone Wygoda²⁷, Ilona Zagozdzon²⁸, Jörg Dötsch¹, Jun Oh^{20,80}, Franz Schaefer^{25,80}, Max Christoph Liebau^{1,7,80}✉ & The ARegPKD consortium*

¹Department of Pediatrics, Faculty of Medicine, University Hospital Cologne and University of Cologne, Kerpener Str. 62, 50937 Cologne, Germany. ²Department of Pediatric Nephrology, University Hospital Vall d'Hebron, Barcelona, Spain. ³Department of Pediatric Nephrology, Klinikum Stuttgart, Olga Children's Hospital, Stuttgart, Germany. ⁴Department of Pediatrics II, University Hospital Essen, Essen, Germany. ⁵Department of Internal Medicine I, University Hospital of the RWTH Aachen, Aachen, Germany. ⁶Institute of Human Genetics, University Hospital of Cologne, Cologne, Germany. ⁷Center for Molecular Medicine Cologne, University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne, Germany. ⁸Department of General Pediatrics and Hematology/Oncology, Children's University Hospital Tuebingen, Tuebingen, Germany. ⁹Division of Pediatric Nephrology, Research and Training Hospital, Marmara University, Istanbul, Turkey. ¹⁰Department of General Pediatrics, University Hospital Muenster, Muenster, Germany. ¹¹The Children's Memorial Health Institute, Warsaw, Poland. ¹²Division of Nephrology, Department of Pediatric Subspecialties, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy. ¹³Pediatric Nephrology, Dialysis and Transplant Unit, Fondazione IRCCS Cà Granda, Ospedale Maggiore Policlinico, Milan, Italy. ¹⁴Department of Development and Regeneration, PKD Research Group, KU Leuven, Leuven, Belgium. ¹⁵Department of Pediatric Nephrology, University Hospitals Leuven, Leuven, Belgium. ¹⁶Department of Pediatric Kidney, Liver and Metabolic Diseases, Hannover Medical School, Hannover, Germany. ¹⁷Children's Hospital St. Elisabeth and St. Barbara, Halle (Saale), Germany. ¹⁸Department of Paediatrics and Adolescent Medicine, Medical University Vienna, Vienna, Austria. ¹⁹Department of Pediatrics, University Hospital Bonn, Bonn, Germany. ²⁰University Children's Hospital, University Medical Center Hamburg Eppendorf, Hamburg, Germany. ²¹UCL Great Ormond Street Hospital for Children Institute of Child Health, UCL, London, UK. ²²Department of Pediatrics, Faculty of Medical Sciences in Zabrze, SUM in Katowice, Katowice, Poland. ²³Department of Paediatrics and Nephrology, Medical University of Bialystok, Bialystok, Poland. ²⁴Department of Pediatrics, Immunology and Nephrology, Polish Mother's Memorial Hospital Research Institute, Lodz, Poland. ²⁵Division of Pediatric Nephrology, Center for Pediatrics and Adolescent Medicine, University Hospital Heidelberg, Heidelberg, Germany. ²⁶Department of Pediatrics, Klinikum Saarbrücken, Saarbrücken, Germany. ²⁷Clinic for Children and Adolescents, Hospital St. Georg, Leipzig, Germany. ²⁸Department of Pediatrics, Nephrology and Hypertension, Medical University of Gdansk, Gdansk, Poland. ⁸⁰These authors jointly supervised this work: Jun Oh, Franz Schaefer and Max Christoph Liebau. *A list of authors and their affiliations appears at the end of the paper ✉email: max.liebau@uk-koeln.de

To test the association between bilateral nephrectomies in patients with autosomal recessive polycystic kidney disease (ARPKD) and long-term clinical outcome and to identify risk factors for severe outcomes, a dataset comprising 504 patients from the international registry study ARegPKD was analyzed for characteristics and complications of patients with very early (≤ 3 months; VEBNE) and early (4–15 months; EBNE) bilateral nephrectomies. Patients with very early dialysis (VED, onset ≤ 3 months) without bilateral nephrectomies and patients with total kidney volumes (TKV) comparable to VEBNE infants served as additional control groups. We identified 19 children with VEBNE, 9 with EBNE, 12 with VED and 11 in the TKV control group. VEBNE patients suffered more frequently from severe neurological complications in comparison to all control patients. Very early bilateral nephrectomies and documentation of severe hypotensive episodes were independent risk factors for severe neurological complications. Bilateral nephrectomies within the first 3 months of life are associated with a risk of severe neurological complications later in life. Our data support a very cautious indication of very early bilateral nephrectomies in ARPKD, especially in patients with residual kidney function, and emphasize the importance of avoiding severe hypotensive episodes in this at-risk cohort.

Abbreviations

ARPKD	Autosomal recessive polycystic kidney disease
CLKTx	Combined liver and kidney transplantation
CVVH	Continuous veno-venous hemofiltration
EBNE	Early bilateral nephrectomies
FTT	Failure to thrive
HR	Hazard ratio
HD	Hemodialysis
KRT	Kidney replacement therapy
NE	Nephrectomy/nephrectomies
P	Percentile
PD	Peritoneal dialysis
PTP	Pole-to-pole
SDS	Standard deviation scores
TKV	Total kidney volume
VEBNE	Very early bilateral nephrectomies
VED	Very early dialysis

Autosomal recessive polycystic kidney disease (ARPKD) is a severe and mostly early-onset hepatorenal disorder mainly caused by mutations in the *PKHD1* gene^{1–3}. Mutations in a second gene, *DZIP1L*, have recently been associated with an ARPKD phenotype^{3,4}. Affected newborns may suffer from massive kidney enlargement and may require early kidney replacement therapy (KRT). Peritoneal dialysis (PD) is then considered the dialysis modality of choice⁵. It has been suggested that large kidneys may contribute to respiratory problems and may hamper nutritional support^{6,7}. The respiratory situation in ARPKD neonates can be challenging due to pulmonary hypoplasia and additional abdominal pressure by PD. Furthermore, concerns about the feasibility of PD in cases of massive renal enlargement have been raised. In some infants with severe ARPKD, uni- or bilateral nephrectomies are therefore performed with the aim to improve the respiratory, nutritional and dialysis conditions^{6–8}. However, nephrectomies have not been shown to improve the respiratory situation and the rationale for nutritional improvement is based on small studies regarding unilateral nephrectomy^{9–11}. Even in severely affected infants, residual kidney function may be preserved in ARPKD and may even improve during the first months of life¹². Obviously, bilateral nephrectomies result in a life-long need for KRT. Additionally, arterial hypotension has been reported after bilateral nephrectomies and may contribute to neurological or ophthalmological complications in young PD patients^{13–16}. The association of early bilateral nephrectomies in general and of the timing of the second nephrectomy in particular with long-term outcome in ARPKD patients remains unclear. To address this topic, we analyzed the comprehensive pre-, peri- and postnatal information captured in the ARegPKD registry study^{17,18}.

Results

Characteristics of patient subgroups. At the timepoint of analysis, 504 patients with the clinical diagnosis of ARPKD and sufficient data quality were registered in the ARegPKD registry. Of these, 19 patients (3.8%) with very early bilateral nephrectomies (within first 3 months of life, VEBNE) and further 9 patients (1.8%) with early bilateral nephrectomies (first nephrectomy within first 15 months of life, second nephrectomy at age 4–15 months, EBNE) were identified. We included a control group of 11 patients, who had very large total kidney volumes (TKV) comparable to VEBNE patients (TKV > ~200 ml, measured by ultrasound within the first 3 months of life), but were not treated with early nephrectomy or dialysis (TKV control group). Pole-to-pole lengths, single kidney volumes and total kidney volumes did not differ significantly between VEBNE and TKV control patients but TKV control patients were slightly older at measurement (Table 1). To account for effects of very early dialysis (VED) onset, an additional control group was defined with 12 ARPKD patients with dialysis onset within the first 3 months of life without bilateral nephrectomies in their disease course (VED). Patients were grouped according to their clinical status at time of last documented observation in the registry or at time prior to their death. The patient characteristics regarding nephrectomies and KRT are depicted in Table 2.

	VEBNE n = 19	TKV Control group n = 11	P
Age at kidney sonography, n	16/19	11/11	
Mean (SD), mo	0.7 (0.4)	1.5 (0.9)	0.009
Median (IQR), mo	0.6 (0.3–1.0)	1.5 (0.6–2.4)	
PTP left kidney, n	7/19	8/11	
Mean (SD), cm	9.6 (1.5)	9.7 (1.2)	0.96
Median (IQR), cm	10.0 (9.2–10.5)	10.0 (8.5–10.7)	
SDS, n	4/19	8/11	
Mean (SD)	20.1 (2.1)	22.9 (3.9)	0.28
Median (IQR)	20.6 (17.8–21.8)	24.0 (18.9–26.6)	
PTP right kidney, n	6/19	8/11	
Mean (SD), cm	9.8 (1.6)	9.6 (0.8)	0.23
Median (IQR), cm	10.4 (9.1–10.6)	9.5 (9.2–10.1)	
SDS, n	4/19	8/11	
Mean (SD)	19.7 (4.7)	22.7 (2.8)	0.28
Median (IQR)	19.0 (15.7–24.4)	23.0 (20.0–24.6)	
Volume left kidney, n	11/19	11/11	
Mean (SD), ml	188.3 (84.7)	138.6 (64.8)	0.15
Median (IQR), ml	208.0 (105.0–260.0)	105.0 (96.0–162.0)	
SDS, n	10/19	9/11	
Mean (SD)	50.2 (25.3)	39.9 (17.9)	0.50
Median (IQR)	59.3 (22.6–66.7)	32.7 (24.6–49.0)	
Volume right kidney, n	11/19	11/11	
Mean (SD), ml	189.5 (79.3)	156.0 (53.6)	0.48
Median (IQR), ml	217.0 (103.0–250.0)	144.0 (119.0–173.0)	
SDS, n	9/19	9/11	
Mean (SD)	48.6 (24.1)	44.5 (14.4)	0.86
Median (IQR)	48.8 (23.4–68.0)	45.5 (32.9–52.9)	
Total kidney volume, n	9/19	11/11	
Mean (SD), ml	369.9 (176.9)	294.6 (116.2)	0.60
Median (IQR), ml	437.0 (205.0–535.0)	257.0 (217.0–335.0)	
Min–max, ml	130–580	194–580	

Table 1. Sonographic kidney measurements in patients with very early bilateral nephrectomies and patients from the TKV control group, kidney sonography performed within the first 3 months of life. Continuous variables are expressed as mean (\pm SD) and median (IQR). *P* values (right column) were derived using Mann–Whitney *U* tests. *PTP* pole-to-pole, *SDS* standard deviation score.

Indications for nephrectomies. In VEBNE patients, indications for nephrectomies encompassed respiratory failure and/or ventilation problems, abdominal distension and/or massive kidney enlargement and onset of or problems with PD (Table 2). Most patients with sequential nephrectomies started dialysis after the first nephrectomy (11/17, 65%). Two patients of the VED group received very early unilateral nephrectomy. In the TKV control group, two patients received unilateral nephrectomy during combined liver and kidney transplantation surgery later in life.

Pre- and perinatal characteristics. Detailed information on patient characteristics is provided in Table 3. VEBNE and TKV control patients more frequently showed poor postnatal respiratory adaptation and were more often admitted to neonatal intensive care units (NICU). Other characteristics—including gestational age, birth weight, length, head circumference as well as Apgar scores and requirement of respiratory support—did not show relevant differences between the groups.

Long-term complications with focus on neurological development and risk factors for severe neurological complications. In a first step, we addressed the occurrence of relevant symptoms or complications including blood pressure alterations, neurological symptoms, sepsis, cardiopulmonary resuscitation and death in a descriptive manner in the four different patient groups. Among all complications investigated, the most profound differences were observed in the neurological outcome: seizures and severe neurological complications encompassing ischemic defects, hypoxic brain damage, brain infarct, parenchymal defect, severe neurodevelopmental disorder and optic neuropathy with vision loss occurred more frequently in VEBNE patients. While 12/19 (63%) VEBNE patients were affected by severe neurological complications, this was true for only 2/9 (22%) EBNE, 2/12 (17%) VED and none of the 11 TKV control patients. Of the 12 affected VEBNE patients, three suffered from cerebral ischemia, three from infarction, bleeding or parenchymal defects, three from optic neuropathy (with vision loss), two from combined cerebral ischemia or hypoxemia and optic neuropathy, and one from cerebral ischemia with bleeding. The two EBNE patients developed sinus vein thrombosis and bilateral cerebral infarctions, the two VED patients developed posthypoxic ischemic lesions after a very difficult postnatal adaptation and hydrocephalus with brain atrophy. Neurodevelopmental delay occurred most frequently in VEBNE patients. In a re-survey with specific questions on the psychomotoric development, only one of 17 (6%) studied VEBNE patients was classified to have a normal development compared to 2/8 (25%) EBNE patients, 5/11 (46%) VED patients and 5/8 (63%) TKV control patients. Approximately one third of VEBNE patients suffered from severely or very severely affected psychomotoric development each. Younger ages at the timepoint of second nephrectomy were more frequently documented in those patients with severe neurological and devel-

	VEBNE n = 19		EBNE n = 9		VED n = 12	TKV control n = 11
Age at 1st NE, n	19		9		2	2
Mean (SD), mo	1.0 (0.8)		4.1 (2.0)		1.8 (0.3)	75.3 (44.8)
Median (IQR or min;max), mo	0.8 (0.4–1.3)		4.2 (2.7–6.0)		1.8 (1.6;2.0)	75.3 (43.6;107.0)
Age at 2nd NE, n	19		9			
Mean (SD), mo	1.5 (0.8)		7.4 (3.7)		–	–
Median (IQR), mo	1.6 (0.7–2.0)		5.9 (4.5–11.5)			
Age at simultaneous bilateral NE, n	2					
Mean (SD), mo	2.6 (0.6)		–		–	–
Median (IQR or min;max), mo	2.6 (2.2;3.0)					
Localisation 1st NE right, n (%)	9/17 (53%)		4/9 (44%)		1/2 (50%)	2/2 (100%)
Indications for NE (multiple answering possible)	1st	2nd	1st	2nd	1st	1st
Respiratory failure/ventilation (problems), n (%)	7/19 (37%)	4/19 (21%)	4/9 (44%)	2/9 (22%)	1/2 (50%)	–
Nutritional problems/FTT, n (%)	1/19 (5%)	1/19 (5%)	0/9 (0%)	2/9 (22%)	–	–
Arterial hypertension, n (%)	1/19 (5%)	1/19 (5%)	2/9 (22%)	2/9 (22%)	–	–
Onset of PD, n (%)	8/19 (42%)	3/19 (16%)	0/9 (0%)	0/9 (0%)	–	–
PD problems/inefficacy, n (%)	1/19 (5%)	5/19 (26%)	1/9 (11%)	1/9 (11%)	–	–
Abdominal distension/kidney enlargement, n (%)	7/19 (37%)	7/19 (37%)	1/9 (11%)	1/9 (11%)	–	–
Massive growth of kidney after 1st NE, n (%)	–	3/19 (16%)	–	0/9 (0%)	–	–
Indication unknown, n (%)	2/19 (11%)	4/19 (21%)	2/9 (22%)	3/9 (33%)	1/2 (50%)	–
Others, n (%)	–	–	–	–	–	2/2 (100%) CLKTx
Age at onset of dialysis/KRT, n	19		7		12	2
Mean (SD), mo	0.8 (0.8)		5.4 (4.1)		1.1 (1.0)	72.0 (49.6)
Median (IQR or min;max), mo	0.6 (0.2–1.2)		5.0 (1.8–8.1)		0.6 (0.4–1.9)	72.0 (37.0;107.0)
Type of dialysis/KRT	PD 16/19, HD 1/19, CVVH 2/19		PD 5/7, HD 2/7		PD 10/12, HD 1/12, CVVH 1/12	HD 1/2, CLKTx 1/2
Onset of dialysis prior to 1st NE, n (%)	7/19 (37%) (5 PD, 2 CVVH)		2/7 (29%) (1 PD, 1 HD)		No NE at all: 10/12 (83%) (9 PD, 1 CVVH)	1/2 (50%) (1 HD)
Onset of dialysis after 1st NE, n (%)	11/19 (58%) (10 PD, 1 HD)		3/7 (43%) (3 PD)		2/12 (17%) (1 PD, 1 CVVH)	–
Onset of dialysis after 2nd NE, n (%)	1/19 (5%) (1 PD)		2/7 (29%) (1 PD, 1 HD)		–	–

Table 2. Characteristics of patient subgroups regarding nephrectomies and kidney replacement therapy. Categorical values given as n/n total (percentage); continuous variables, as mean \pm SD and median (IQR or min;max) with min;max indicated in case of $n \leq 3$. *CLKTx* combined liver and kidney transplantation, *CVVH* continuous veno-venous hemofiltration, *EBNE* early bilateral nephrectomies, *FTT* failure to thrive, *HD* hemodialysis, *KRT* kidney replacement therapy, *NE* nephrectomy, *PD* peritoneal dialysis, *TKV* total kidney volume, *VEBNE* very early bilateral nephrectomies, *VED* very early dialysis.

ompmental complications, especially timepoints of second nephrectomy within the first 3 months of life (Suppl. Figure S1). The proportions of patients receiving some form of specific therapies for developmental complications were high in all four patient groups, ranging from 55% in VED to 94% in VEBNE patients. Microcephaly was hardly found at birth but was most frequently documented in VEBNE patients in the further course (11/19, 58%). Severe hyper- and hypotensive episodes as well as cardiopulmonary resuscitation (CPR) were reported for subcohorts of VEBNE and EBNE patients, without reports in VED and TKV control patients. Proportions of patients with a septic episode prior to the evaluated neurological complication did not differ significantly between the groups (Table 4).

Follow-up time was similar for VEBNE, EBNE and TKV control patients, but shorter for VED patients (Table 4). Severe neurological complications were further studied over time using Kaplan–Meier analyses and Cox regression analyses. Survival without severe neurological complications after 7 years was approximately 30% in VEBNE, 80% in EBNE, 70% in VED patients and 100% in TKV control patients according to Kaplan–Meier survival analysis (Fig. 1). Thus, VEBNE patients showed significantly worse outcome than EBNE or TKV control patients with a similar trend compared to VED patients. Severe neurological complications in the VEBNE group were evident within the 14 months of life in most cases (10/12). Only in two patients, atrophy of the optic nerve was diagnosed at the ages of 2.8 and 6.6 years.

	VEBNE n = 19	EBNE n = 9	VED n = 12	TKV control n = 11	P
Sex (male), n (%)	8/19 (42%)	5/9 (56%)	6/12 (50%)	5/11 (46%)	0.92
Prenatal anomalies, n (%)	19/19 (100%)	7/9 (78%)	11/12 (92%)	10/10 (100%)	0.10
Oligo- or anhydramnios, n (%)	17/18 (94%)	6/9 (67%)	10/12 (83%)	10/10 (100%)	0.11
Gw at diagnosis, n	15	6	9	9	0.12
Mean (SD)	27.2 (3.8)	31.8 (4.1)	30.9 (5.6)	29.1 (5.9)	
Median (IQR)	27.0 (24.0–30.0)	31.5 (28.0–34.5)	31.0 (27.0–36.0)	29.0 (25.0–34.0)	
Increased echogenicity, n (%)	10/13 (77%)	3/7 (43%)	5/9 (56%)	8/8 (100%)	0.07
Gw at diagnosis, n	9	3	5	7	0.70
Mean (SD)	28.9 (2.4)	27.7 (5.5)	30.6 (6.5)	28.7 (6.7)	
Median (IQR or min;max)	28.0 (27.5–30.5)	28.0 (22.0;33.0)	32.0 (25.0–35.5)	29.0 (24.0–36.0)	
Renal hyperplasia, n (%)	13/16 (81%)	4/8 (50%)	6/9 (67%)	7/7 (100%)	0.13
Renal cysts, n (%)	13/16 (81%)	5/9 (56%)	7/11 (64%)	7/7 (100%)	0.16
Perinatal information					
Normal vaginal delivery, n (%)	5/18 (28%)	4/9 (44%)	9/11 (82%)**§	3/10 (30%)	0.03
Perinatal problems, n (%)	17/19 (90%)	7/9 (78%)	7/12 (58%)	10/10 (100%)	0.06
NICU, n (%)	17/19 (90%)	6/9 (67%)	7/12 (58%)	10/10 (100%) [†]	0.04
NICU days, n	15	6	7	9	0.06
Mean (SD)	60 (35)	131 (207)	91 (134)	30 (52)	
Median (IQR)	50 (35–85)	39 (14–248)	45 (20–83)	7 (3–39)	
Poor postnatal adaptation, n (%)	16/19 (84%) ^{##}	3/9 (33%)	7/12 (58%)	9/10 (90%) [#]	0.02
Assisted breathing, n (%)	17/19 (90%)	7/9 (78%)	7/12 (58%)	9/10 (90%)	0.16
Pulmonary Hypertension, n (%)	8/17 (47%)	2/7 (29%)	3/11 (27%)	3/9 (33%)	0.69
Postnatal information					
Gestational age at birth, n	17	9	10	10	0.31
Mean (SD), weeks	35.0 (1.7)	35.9 (2.6)	36.2 (3.1)	34.8 (2.5)	
Median (IQR), weeks	35.0 (34.0–36.0)	36.0 (33.5–38.5)	37.0 (34.8–38.3)	35.0 (32.8–37.3)	
Birth weight, n	18	7	9	10	0.46
Mean (SD), kg	3.01 (0.67)	3.19 (0.62)	3.07 (0.84)	2.70 (0.57)	
Median (IQR), kg	3.00 (2.61–3.55)	3.18 (2.71–3.54)	3.41 (2.46–3.60)	2.70 (2.19–3.22)	
Birth length, n	14	6	7	9	0.52
Mean (SD), cm	47.8 (4.1)	48.5 (2.8)	49.1 (5.9)	46.3 (3.4)	
Median (IQR), cm	49.0 (45.3–51.0)	50.0 (45.5–50.3)	49.0 (45.0–52.0)	45.0 (43.3–49.5)	
Head circumference, n	14	4	8	8	0.98
Mean (SD), cm	33.0 (1.9)	33.0 (1.8)	33.4 (2.4)	32.6 (2.1)	
Median (IQR), cm	33.3 (32.4–34.5)	33.0 (31.3–34.8)	33.3 (32.0–34.6)	32.0 (31.0–35.0)	
Head circumference percentile, n	13	4	7	8	0.42
Mean (SD)	68 (26)	58 (38)	53 (30)	76 (14)	
Median (IQR)	73 (54–90)	74 (18–82)	55 (26–76)	82 (61–88)	
Microcephaly at birth (< P3), n (%)	0/14 (0%)	1/4 (25.0%)	1/8 (13%)	0/8 (0%)	0.20
Apgar 1, n	14	6	9	10	0.37
Mean (SD)	4.6 (2.3)	5.5 (3.0)	6.1 (2.4)	6.1 (2.4)	
Median	5.0 (3.0–6.3)	5.5 (3.3–8.3)	6.0 (5.0–8.0)	6.0 (5.0–8.3)	
Apgar 5, n	14	6	8	10	0.29
Mean (SD)	6.1 (2.0)	6.0 (2.2)	7.0 (2.3)	7.1 (2.1)	
Median	6.5 (5.0–8.0)	5.5 (4.5–8.3)	7.5 (6.3–8.8)	8.0 (6.0–8.3)	
Apgar 10, n	13	5	8	10	0.30
Mean (SD)	7.5 (1.3)	7.8 (1.3)	7.8 (1.7)	8.2 (2.0)	
Median	8.0 (7.5–8.0)	8.0 (6.5–9.0)	8.0 (7.3–9.0)	9.0 (7.8–9.3)	
Genetic information					
Variant detection in <i>PKHD1</i> n (%)	20/22 (91%) (n = 11)	5/10 (50%) (n = 5)	8/12 (67%) (n = 6)	14/14 (100%) (n = 7)	
Two truncating <i>PKHD1</i> variants, n (%)	3/11 (27%)	0/5 (0%)	1/6 (17%)	2/7 (29%)	

Table 3. Pre-, peri- and postnatal information by patient group. Binary or categorical values given as n/n total (percentage); continuous variables, as mean \pm SD and median (IQR or min;max) with min;max indicated in case of $n \leq 3$. *P* values (right column) were derived using Kruskal–Wallis or chi-squared tests. *EBNE* early bilateral nephrectomies, *Gw* gestational week, *NICU* neonatal intensive care unit, *P* Percentile, *TKV* total kidney volume, *VEBNE* very early bilateral nephrectomies, *VED* very early dialysis. ** $P < 0.01$ for comparison with *VEBNE*; # $P < 0.05$ for comparison with *EBNE*; ## $P < 0.01$ for comparison with *EBNE*; [†] $P < 0.05$ for comparison with *VED*; [§] $P < 0.05$ for comparison with *TKV* control.

	VEBNE n = 19	EBNE n = 9	VED n = 12	TKV control n = 11	P
Follow-up—complications					
Follow-up time, mo, median (IQR)	78.0 (27.0–106.0)	99.0 (84.0–156.0)	43.0 (13.0–56.0)	59.0 (32.0–106.0)	0.06
Neurodevelopmental delay, n (%)	17/19 (90%) ^{§§§}	7/9 (78%) [§]	6/12 (50%)	3/11 (27%)	0.003
Specific therapies, n (%)	15/16 (94%)	6/8 (75%)	6/11 (55%)	7/8 (88%)	0.09
Physiotherapy, n (%)	14/16 (88%)	6/8 (75%)	6/11 (55%)	4/8 (50%)	0.16
Speech therapy, n (%)	10/16 (63%) ^{††}	4/8 (50%) ^{††}	0/11 (0%)	2/8 (25%)	0.007
Ergotherapy, n (%)	10/16 (63%)	4/7 (57%)	2/11 (18%)	3/7 (43%)	0.14
Special teacher, n (%)	6/15 (40%)	2/8 (25%)	0/11 (0%)	1/7 (14%)	0.10
Regular school, n (%)	3/9 (33%)	2/5 (40%)	1/1 (100%)	4/4 (100%)	0.10
Psychomotoric development	n = 17	n = 8	n = 11	n = 8	
Normal, n (%)	1/17 (6%)	2/8 (25%)	5/11 (46%) [*]	5/8 (63%) ^{**}	0.02
Mild disorder, n (%)	4/17 (24%)	4/8 (50%)	4/11 (36%)	3/8 (38%)	0.61
Severe disorder, n (%)	6/17 (35%) [†]	1/8 (13%)	0/11 (0%)	0/8 (0%)	0.04
Very severe disorder, n (%)	6/17 (35%)	1/8 (13%)	2/11 (18%)	0/8 (0%)	0.20
Seizures, n (%)	14/19 (74%) ^{*†§}	2/9 (22%)	4/12 (33%)	3/11 (27%)	0.02
Severe neurological complication, n (%)	12/19 (63%) ^{*†§§}	2/9 (22%)	2/12 (17%)	0/11 (0%)	0.001
Documentation of microcephaly (< P3), n (%)	11/19 (58%) ^{§§}	2/9 (22%)	4/12 (33%) [§]	0/11 (0%)	0.01
Age at 1st report, n	10	2	3		
Mean (SD), mo	42.1 (51.0)	15.5 (13.4)	0.9 (1.0)	–	0.64
Median (IQR or min;max), mo	19.5 (2.8–86.4)	15.5 (6.0; 25.0)	0.4 (0.2;2.0)		
Severe hypertensive episodes, n (%)	6/19 (32%) [§]	1/9 (11%)	0/12 (0%)	0/11 (0%)	0.03
Severe hypotensive episodes, n (%)	5/19 (26%)	2/9 (22%)	0/12 (0%)	0/11 (0%)	0.08
s/p Septic episode, n (%)	5/19 (26%)	1/9 (11%)	3/12 (25%)	3/11 (27%)	0.81
s/p CPR, n (%)	5/19 (26%)	1/9 (11%)	0/12 (0%)	0/11 (0%)	0.07
Death, n (%)	3/19 (16%)	2/9 (22%)	6/12 (50%)	1/11 (9%)	0.09
Age, n	3	2	6	1	
Mean (SD), mo	28.3 (39.5)	57.7 (24.5)	4.8 (4.6)	0.10	0.08
Median (IQR or min;max), mo	7.8 (3.2;73.8)	57.7 (40.4;75.0)	3.8 (1.5–7.6)	0.10 (-)	

Table 4. Follow-up-complications by patient group. Binary or categorical values given as n/n total (percentage); continuous variables, as mean ± SD and median (IQR or min;max) with min;max indicated in case of n ≤ 3. P values (right column) were derived using log-rank, Kruskal–Wallis or chi-squared tests. CPR cardiopulmonary resuscitation, EBNE early bilateral nephrectomies, s/p state post, TKV total kidney volume, VEBNE very early bilateral nephrectomies, VED very early dialysis. *P < 0.05 for comparison with VEBNE; **P < 0.01 for comparison with VEBNE; †P < 0.05 for comparison with EBNE; ††P < 0.05 for comparison with VED; †††P < 0.01 for comparison with VED; §P < 0.05 for comparison with TKV control; §§P < 0.01 for comparison with TKV control; §§§P < 0.001 for comparison with control.

Overall, severe neurological complications occurred in 16 of 51 (31%) analyzed patients (Table 5). According to Cox regression analysis, the report of severe neurological complications was associated with multiple factors, such as documentation of microcephaly during the course of follow-up as well as weaker and more common neurological symptoms like seizures and reports of severe hypertensive and hypotensive episodes. Furthermore, patients with very early bilateral nephrectomies seemed to be at special risk.

To identify independent risk factors associated with severe neurological complications, multivariate Cox regression analysis with consecutive backward selection was performed (including seizures, documentation of microcephaly, severe hypertensive episodes, severe hypotensive episodes and VEBNE), which revealed the report of a severe hypotensive episode and very early bilateral nephrectomies to be independent risk factors for severe neurological complications (Table 5). In this multivariate analysis, patients with very early bilateral nephrectomies were at fivefold elevated risk for severe neurological complications.

Deaths. Three VEBNE patients died due to cerebral ischemia (age 3.2 months), respiratory failure (7.8 months) and massive postoperative bleeding after revision of dialysis catheter (age 73.8 months). Two EBNE patients died, one due to a septic episode in leukemia (age 40.4 months) and the other patient due to aspiration (75.0 months). Causes of death in six VED patients were indicated as sepsis (ages 5.4 months, 5.7 months, 13.3 months), lung hemorrhage (age 1.8 months) and parental wish to withdraw care in two cases (ages 0.8 months, 2.2 months). Importantly, four of the six deceased VED patients did not display major or severe neurological complications before death. Three out of the four VED patients that passed away without withdrawal of care died after the fifth month of life. One patient from the TKV control group died shortly post-natally due to respiratory failure.




Figure 1. Kaplan–Meier survival without severe neurological complications by patient group. Censored observations (last documented follow-up or death) are marked with a cross. *P* values were derived using log rank tests. *EBNE* early bilateral nephrectomies, *TKV* total kidney volume, *VEBNE* very early bilateral nephrectomies, *VED* very early dialysis.

	With severe neurological complication (n = 16)	Without severe neurological complication (n = 35)	HR (95% CI)	<i>P</i>
Cox regression analysis				
<i>Pre- and perinatal information</i>				
Oligo- or anhydramnios, n (%)	14/16 (88%)	29/33 (88%)	0.86 (0.20–3.78)	0.84
Prenatal increased echogenicity, n (%)	10/13 (77%)	16/24 (67%)	1.34 (0.37–4.89)	0.66
Prenatal renal hyperplasia, n (%)	13/16 (81%)	17/24 (71%)	1.30 (0.37–4.59)	0.68
Prenatal renal cysts, n (%)	10/14 (71%)	22/29 (76%)	0.76 (0.24–2.44)	0.65
Vaginal delivery, n (%)	8/16 (50%)	13/32 (41%)	1.40 (0.52–3.74)	0.50
Gestational age at birth, mean (SD), weeks	35.1 (2.5) (n = 16)	35.5 (2.4) (n = 30)	0.94 (0.77–1.14)	0.54
Microcephaly at birth, n (%)	1/12 (8%)	1/22 (5%)	1.18 (0.15–9.17)	0.87
Head circumference (Percentile Fenton), mean (SD)	68 (29) (n = 11)	64 (26) (n = 21)	1.01 (0.98–1.03)	0.70
Poor postnatal adaptation, n (%)	12/16 (75%)	23/34 (68%)	1.35 (0.44–4.20)	0.60
Assisted breathing and/or ventilation, n (%)	15/16 (94%)	25/34 (74%)	4.00 (0.53–30.30)	0.18
<i>Follow-up—complications</i>				
Seizures, n (%)	14/16 (88%)	9/35 (26%)	12.29 (2.76–54.65)	0.001
Documentation of microcephaly (<P3), n (%)	11/16 (69%)	6/35 (17%)	4.71 (1.63–13.61)	0.004
Severe hypertensive episodes, n (%)	5/16 (31%)	2/35 (6%)	3.12 (1.08–9.05)	0.04
Severe hypotensive episodes, n (%)	7/16 (44%)	0/35 (0%)	13.92 (4.22–45.87)	<0.001
s/p Septic episode, n (%)	4/16 (25%)	8/35 (23%)	0.89 (0.29–2.76)	0.84
s/p CPR, n (%)	3/16 (19%)	3/35 (9%)	1.60 (0.45–5.62)	0.47
<i>Patient groups</i>				
VEBNE, n (%)	12/16 (75%)	7/35 (20%)	6.28 (2.02–19.57)	0.002
EBNE, n (%)	2/16 (13%)	7/35 (20%)	0.46 (0.10–2.01)	0.30
VED, n (%)	2/16 (13%)	10/35 (29%)	0.76 (0.17–3.35)	0.72
TKV control, n (%)	0/16 (0%)	11/35 (31%)	0.03 (0.00–3.07)	0.14
Cox multiple regression analysis				
Severe hypotensive episodes, n (%)	7/16 (44%)	0/35 (0%)	11.42 (3.21–40.71)	<0.001
VEBNE, n (%)	12/16 (75%)	7/35 (20%)	5.12 (1.60–16.39)	0.006

Table 5. Cox regression analysis of pre- and postnatal factors associating with severe neurological complications. Cox multiple regression analysis with independent risk factors for severe neurological complications. *P* values (right column) were derived from Cox single-variable regression analysis or Cox multiple regression analysis. Binary or categorical values given as n/n total (percentage); continuous variables, as mean ± SD. *CPR* cardiopulmonary resuscitation, *EBNE* early bilateral nephrectomies, *HR* Hazard Ratio, *s/p* state post, *TKV* total kidney volume, *VEBNE* very early bilateral nephrectomies, *VED* very early dialysis.

Discussion

The management of severely affected neonates and infants with ARPKD remains a matter of intense debate. The aim of this study was to examine the association of the timing of bilateral nephrectomies with long-term complications in ARPKD patients.

In our analysis, ARPKD patients with very early bilateral nephrectomies (VEBNE) within the first 3 months of life showed significantly more neurological complications than patients who underwent bilateral nephrectomies receiving their second nephrectomy after the third month of life (EBNE). The severe neurological anomalies observed in VEBNE patients encompassed cerebral ischemia or infarcts, parenchymal brain defects, brain volume decrease, subdural bleeding and (anterior ischemic) optic neuropathy. These complications occurred predominantly during the first year of life.

The rationale of drawing a line for evaluation of consequences of bilateral nephrectomies at the age of 3 months was based on two considerations: firstly, major maturation processes of autonomic cardiovascular control seem to occur within the first 3 months of life^{19,20}. Animal models suggest that renal sympathetic nerve control matures postnatally²¹. Secondly, both plasma renin activities and aldosterone levels^{22,23} have been reported to peak within the first 3 months with a steep decline afterwards. Data from healthy infants demonstrated a remarkable increase of systolic blood pressure during the first 6 weeks and a physiologic dip of diastolic blood pressure after birth with a gradual increase during the first year of life from month two to three on²⁴. Summarizing these hypotheses, the removal of both kidneys in this early phase of life might result in the maximum impact by abolishing blood pressure autoregulation. Plotting the occurrence of severe neurological complications against the age at second nephrectomy supports the choice of drawing a line at the age of 3 months as critical time point (Suppl. Figure S1).

Interestingly, our analysis demonstrated that VEBNE and severe hypotensive episodes were independently associated with severe neurological complications. In addition to VEBNE additional factors leading to hypotensive episodes seem to be involved, e.g. acute illnesses²⁵. Previous smaller case series reported an anephric status and ARPKD as risk factors for development of anterior ischemic optic neuropathy in patients on PD²⁶. Hypovolemia has been reported in half of the affected infants with anterior ischemic optic neuropathy on PD²⁶. In our study, optic neuropathy was frequently observed in the VEBNE group. One might speculate about common pathophysiologic mechanisms underlying to all types of cerebral defects, such as reduced cerebral autoregulation and chronic hypotension with increased vulnerability in acute hypotensive episodes. Fluid management per se can pose major challenges in infants on PD, but might affect blood pressure more immediately in anephric infants. In our analysis, we did not have comprehensive data on the fluid status of patients. It is worth noting that cardiopulmonary resuscitation was documented most frequently in the VEBNE patient group. Despite the challenges of fluid management in anephric infants, PD remains the preferred modality for ARPKD infants¹¹. A recent analysis showed that technique survival in ARPKD is comparable to that observed in other early-onset kidney diseases, although PD in ARPKD may require smaller fill volumes and/or more cycles²⁷. Our analysis emphasizes in particular the association of documented pronounced hypotensive episodes with severe neurological complications in ARPKD patients receiving early bilateral nephrectomies and underlines the recommendation of very tight blood pressure control for all anephric children on PD. Furthermore, measures to avoid hypotensive episodes on PD need to be considered, e.g. by avoiding high glucose solutions or by sodium supplementation in dialysate and nutritionally^{25,28,29}. On the other hand, ARPKD specific therapeutic options and clinical studies are eagerly awaited for this severely affected patient population.

None of the TKV control patients with massively enlarged kidneys developed severe neurological complications. Caution is required not to anticipate postnatal severe courses in all cases of newborns with pre- and perinatal anomalies³⁰.

Severe neurological complications were observed in two VED patients: in one patient cerebral ischemia occurred during a very difficult postnatal adaptation requiring intense cardiorespiratory support and unilateral nephrectomy. This patient passed away at the age of 4 weeks after withdrawal of intensive care treatment upon parental request. The other patient died from sepsis at 13 months of age after documentation of hydrocephalus with brain atrophy 1 month earlier. All four of the six deceased VED infants who did not experience severe neurological complications deceased within the first 8 months of life. This is important for the interpretation of the Kaplan–Meier survival analysis for severe neurological complications (Fig. 1). Due to their early death, these patients were no longer at risk for severe neurological complications. Thus, death could be a competing risk for severe neurological complications in this specific subcohort. Survival without severe neurological complications tended to be better in VED compared to VEBNE patients, although statistical significance was missed due to small sample size. Importantly, the majority of the deceased VED patients died well beyond 3 months of age. EBNE patients were also severely affected with similar Apgar values as VEBNE and early dialysis dependency. Yet, only two of nine EBNE patients suffered from severe neurological sequelae, arguing against an intrinsic ARPKD-specific neurologic phenotype.

The current study faces limitations: as severely affected and/or early deceased patients are likely to be underrepresented in our registry study, the obtained sample sizes of the four groups are rather small. Furthermore, there might be a center bias in deciding for nephrectomies in infants. The risk factor severe hypotensive episodes needs to be interpreted carefully: the report and documentation might be biased by the personal perception of the attending physicians and severely affected children might spend more time in hospital, where blood pressure documentation could reveal anomalies more frequently. Hypotensive episodes are known to frequently occur within hypovolemia in infants on PD^{25,31,32}. Our study describes interesting associations of severe hypotensive episodes and the decision to perform VEBNE with severe neurological complications, but we cannot prove causality due to the retrospective registry data underlying the analysis. Importantly, in severely affected infants there may be situations where the decision for uni- or bilateral nephrectomies appears inevitable. Given our

data a thorough discussion of all relevant aspects both with the parents and the attending medical team seems important.

Conclusion

Neurological complications occur more frequently in patients with bilateral nephrectomies performed during the first 3 months of life compared to patients with the second nephrectomy performed between 4 and 15 months of age. Our data suggest utmost caution in the decision-making process for very early bilateral nephrectomies, especially in infants with partially conserved residual renal function, and support postponing the second nephrectomy when possible. Importantly, every effort should be made to avoid pronounced hypotensive episodes in this population.

Methods

Registry. The international cohort study ARegPKD follows patients with the clinical diagnosis of ARPKD according to the previously described protocol^{17,18}. In summary, basic data and regular follow-up data sets are obtained and are subject to regular data quality control. The study protocol was approved by the Ethics Committee of the Faculty of Medicine of Cologne University and the Institutional Review Boards of participating sites. Informed consent was obtained from all subjects or, if subjects are under 18, from a parent and/or legal guardian according to applicable local regulations. All methods were carried out in accordance with relevant guidelines and regulations.

For this specific analysis ischemic brain defects, hypoxic brain damage, brain infarct, parenchymal defect, severe neurodevelopmental disorder and optic neuropathy with vision loss were rated as severe neurological complication. For detailed classification of neurocognitive development, we re-surveyed data on head circumferences at birth and during further course, school education, specific therapies, developmental milestones and psychological/intelligence testing for the studied subcohort: psychomotoric development was classified as normal or mildly, severely or very severely disordered. Head circumferences at birth were graded in percentiles according to the revised Fenton growth chart³³.

All *PKHD1* variants were classified according to criteria of the American College of Medical Genetics (ACMG)³⁴. Sonography-based kidney volumes were calculated according to the ellipsoid formula (length × width × depth × $\pi/6$) and standardized to standard deviation scores (SDS) to pediatric normal values^{35,36}.

Statistics. Data analysis was performed on the dataset available in September 2018 using SPSS 25 (IBM Corp., Armonk, NY, USA) for statistical analyses. Data completeness varied by variable. Continuous variables were described using the number of non-missing values, mean and standard deviation (SD) as well as median and interquartile range (IQR). For binary or categorical variables, absolute and relative frequencies were provided. Event-free survival rates were estimated by the Kaplan–Meier method. Follow-up duration in the four groups were calculated using the reverse Kaplan–Meier method, displayed as median (IQR) and compared using log-rank test. Differences of (continuous) sonographic parameters between VEBNE and control group were compared by Mann–Whitney U test. Differences between the four patient groups were assessed by Kruskal–Wallis tests for continuous and by chi-squared tests for binary or categorical variables. The statistical tests did not adjust formally for multiplicity due to the exploratory nature of the analysis. No imputation was performed.

To investigate the predictive value of clinical risk factors for severe neurological outcome, Cox regression analysis was performed for each risk factor separately. The parameters with a *p* value below 0.10 were then included in a Cox multiple regression analysis with consecutive backward selection for variables with a *p* value below 0.10.

To assess the differences of survival without severe neurological complications in the four patient groups, Kaplan–Meier analysis and log rank tests were used. All analyses are exploratory and *p* values of less than 0.05 were considered significant in a descriptive manner in distinguishing between the groups.

Data availability

The data that support the findings of this study are available, on reasonable request, from the corresponding author.

Received: 11 April 2020; Accepted: 24 August 2020

Published online: 29 September 2020

References

1. Ward, C. J. *et al.* The gene mutated in autosomal recessive polycystic kidney disease encodes a large, receptor-like protein. *Nat Genet* **30**, 259–269 (2002).
2. Onuchic, L. F. *et al.* PKHD1, the polycystic kidney and hepatic disease 1 gene, encodes a novel large protein containing multiple immunoglobulin-like plexin-transcription-factor domains and parallel beta-helix 1 repeats. *Am J Hum Genet* **70**, 1305–1317 (2002).
3. Bergmann, C. *et al.* Polycystic kidney disease. *Nat Rev Dis Primer* **4**, 50 (2018).
4. Lu, H. *et al.* Mutations in DZIP1L, which encodes a ciliary-transition-zone protein, cause autosomal recessive polycystic kidney disease. *Nat Genet* **49**, 1025–1034 (2017).
5. Zurawska, A. M. *et al.* Clinical practice recommendations for the care of infants with stage 5 chronic kidney disease (CKD5). *Pediatr. Nephrol. Berl. Ger.* **28**, 1739–1748 (2013).
6. Luoto, T. T., Pakarinen, M. P., Jahnukainen, T. & Jalanko, H. Liver disease in autosomal recessive polycystic kidney disease: clinical characteristics and management in relation to renal failure. *J. Pediatr. Gastroenterol. Nutr.* **59**, 190–196 (2014).
7. Hoyer, P. F. Clinical manifestations of autosomal recessive polycystic kidney disease. *Curr. Opin. Pediatr.* **27**, 186–192 (2015).
8. Beaunoyer, M. *et al.* Optimizing outcomes for neonatal ARPKD. *Pediatr. Transplant.* **11**, 267–271 (2007).
9. Bean, S. A., Bednarek, F. J. & Primack, W. A. Aggressive respiratory support and unilateral nephrectomy for infants with severe perinatal autosomal recessive polycystic kidney disease. *J. Pediatr.* **127**, 311–313 (1995).

10. Shukla, A. R., Kiddoo, D. A. & Canning, D. A. Unilateral nephrectomy as palliative therapy in an infant with autosomal recessive polycystic kidney disease. *J. Urol.* **172**, 2000–2001 (2004).
11. Guay-Woodford, L. M. *et al.* Consensus expert recommendations for the diagnosis and management of autosomal recessive polycystic kidney disease: report of an international conference. *J. Pediatr.* **165**, 611–617 (2014).
12. Cole, B. R., Conley, S. B. & Stapleton, F. B. Polycystic kidney disease in the first year of life. *J. Pediatr.* **111**, 693–699 (1987).
13. Riechardt, S., Koch, M., Oh, J. & Fisch, M. Early bilateral nephrectomy in neonatal autosomal recessive polycystic kidney disease: Improved prognosis or unnecessary effort?. *Urol. Aug.* **56**, 882–886 (2017).
14. Hassinger, A. B. & Garimella, S. Refractory hypotension after bilateral nephrectomies in a Denys–Drash patient with phenylketonuria. *Pediatr. Nephrol. Berl. Ger.* **28**, 345–348 (2013).
15. Al-Kaabi, A. *et al.* Bilateral anterior ischaemic optic neuropathy in a child on continuous peritoneal dialysis: case report and literature review. *Sultan Qaboos Univ. Med. J.* **16**, e504–e507 (2016).
16. van Lieburg, A. F. & Monnens, L. A. Persistent arterial hypotension after bilateral nephrectomy in a 4-month-old infant. *Pediatr. Nephrol. Berl. Ger.* **16**, 604–605 (2001).
17. Ebner, K. *et al.* Rationale, design and objectives of ARegPKD, a European ARPKD registry study. *BMC Nephrol.* **16**, 22 (2015).
18. Ebner, K., Schaefer, E., Liebau, M. C. & ARegPKD Consortium. Recent Progress of the ARegPKD Registry Study on autosomal recessive polycystic kidney disease. *Front. Pediatr.* **5**, 18 (2017).
19. Yiallourou, S. R., Sands, S. A., Walker, A. M. & Horne, R. S. C. Maturation of heart rate and blood pressure variability during sleep in term-born infants. *Sleep* **35**, 177–186 (2012).
20. Yiallourou, S. R., Sands, S. A., Walker, A. M. & Horne, R. S. C. Postnatal development of baroreflex sensitivity in infancy. *J. Physiol.* **588**, 2193–2203 (2010).
21. Booth, L. C. *et al.* Maturation-related changes in the pattern of renal sympathetic nerve activity from fetal life to adulthood. *Exp. Physiol.* **96**, 85–93 (2011).
22. Bauer, J. H. Age-related changes in the renin-aldosterone system. Physiological effects and clinical implications. *Drugs Aging* **3**, 238–245 (1993).
23. Siegler, R. L., Crouch, R. H., Hackett, T. N., Walker, M. & Jubiz, W. Potassium-renin-aldosterone relationships during the first year of life. *J. Pediatr.* **91**, 52–55 (1977).
24. Report of the Second Task Force on Blood Pressure Control in Children. Task force on blood pressure control in children. National Heart, Lung, and Blood Institute, Bethesda, Maryland. *Pediatrics* **1987**(79), 1–25 (1987).
25. Schaefer, F. Peritoneal dialysis in infants: never lose sight of-and from-arterial hypotension!. *Perit. Dial. Int. J. Int. Soc. Perit. Dial.* **35**, 123–124 (2015).
26. Dufek, S. *et al.* Anterior ischemic optic neuropathy in pediatric peritoneal dialysis: risk factors and therapy. *Pediatr. Nephrol. Berl. Ger.* **29**, 1249–1257 (2014).
27. Akarkach, A. *et al.* Maintenance peritoneal dialysis in children with autosomal recessive polycystic kidney disease: a comparative cohort study of the international pediatric peritoneal dialysis network registry. *Am. J. Kidney Dis. Off. J. Natl. Kidney Found.* **75**, 460–464 (2020).
28. Di Zazzo, G. *et al.* Anterior ischemic optical neuropathy in children on chronic peritoneal dialysis: report of 7 cases. *Perit. Dial. Int. J. Int. Soc. Perit. Dial.* **35**, 135–139 (2015).
29. KDOQI Work Group. KDOQI Clinical Practice Guideline for Nutrition in Children with CKD: 2008 update Executive summary. *Am. J. Kidney Dis. Off. J. Natl. Kidney Found.* **53**, 11–104 (2009).
30. Burgmaier, K. *et al.* Risk factors for early dialysis dependency in autosomal recessive polycystic kidney disease. *J. Pediatr.* <https://doi.org/10.1016/j.jpeds.2018.03.052> (2018).
31. Vidal, E. & Schaefer, F. Hypotension in infants on chronic peritoneal dialysis: mechanisms, complications, and management. *Adv. Perit. Dial. Conf. Perit. Dial.* **31**, 54–58 (2015).
32. Rippe, B. & Venturoli, D. Optimum electrolyte composition of a dialysis solution. *Perit. Dial. Int. J. Int. Soc. Perit. Dial.* **28**(Suppl 3), S131–136 (2008).
33. Fenton, T. R. & Kim, J. H. A systematic review and meta-analysis to revise the Fenton growth chart for preterm infants. *BMC Pediatr.* **13**, 59 (2013).
34. Richards, S. *et al.* Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet. Med. Off. J. Am. Coll. Med. Genet.* **17**, 405–424 (2015).
35. Scholbach, T. & Weitzel, D. Body-surface-area related renal volume: a common normal range from birth to adulthood. *Scientifica* **2012**, 949164 (2012).
36. Weitzel, D. Nieren und ableitende Harnwege. In *Die Ultraschalluntersuchung des Kindes* (eds Deeg, K.-H. *et al.*) 289–324 (Springer, Berlin, Heidelberg, 1997).

Acknowledgements

We thank Ms Regina Brinster and Mr Samuel Kilian for support in conducting statistical analysis. Parts of this work have been presented as a poster presentation at ASN “Kidney Week 2019” in Washington, DC, USA on November 7th 2019 (please see <https://www.asn-online.org/education/kidneyweek/2019/program-abstract.aspx?controlId=3232194> as accessed on the 11th of April 2020).

Author contributions

K.B. and M.L. drafted the manuscript. K.B. and M.B. performed statistical analysis. K.B., G.A., M.B., A.K.B., F.E., M.G., I.G., J.K., C.K., L.M., A.M., D.M., L.P., L.P., A.P., G.S., R.S., R.S., M.S., K.T.J., M.T., L.T.W., E.W., D.W., S.W., I.Z., J.D., J.O., F.S. and M.L. coordinated clinical care of patients and gathered clinical or genetic information. K.B., J.O., F.S. and M.L. set up this substudy, F.S. and M.L. set up the ARegPKD database, M.L. had overall oversight of the project. All authors reviewed and approved the final manuscript.

Funding

We thank the German Society for Pediatric Nephrology (GPN) and the ESCAPE Network for their support. ML was supported by grants of the GPN, the European Society for Paediatric Nephrology (ESPN), the German PKD foundation, the Koeln Fortune program, the GEROK program of the Medical Faculty of University of Cologne, and the Marga and Walter Boll-Foundation. FS and ML are supported by the the German Federal Ministry of Research and Education (BMBF grant 01GM1515 and 01GM1903). KB was supported by the Koeln Fortune program of the Medical Faculty of University of Cologne and the Marga and Walter Boll-Foundation. This work was generated within the European Reference Network for Rare Kidney Disorders (ERKNet). Open Access funding provided by Projekt DEAL.

Competing interests

ML has received honoraria for scientific lectures from Pfizer. Representing the University Hospital of Cologne ML has been counselling Otsuka in an advisory board. DM, represented by KU Leuven University, received an educational grant from Otsuka and participated in an advisory board. The other authors declare no potential conflict of interest.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-020-71956-1>.

Correspondence and requests for materials should be addressed to M.C.L.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020

The ARegPKD consortium

Loai Akram Eid²⁹, Klaus Arbeiter¹⁸, Nadejda Ranguelov³⁰, Laure Collard³¹, Aurélie De Mul³², Markus Feldkoetter³³, Tomas Seeman³⁴, Julia Thumfart³⁵, Franziska Grundmann³⁶, Matthias Galiano³⁷, Björn Buchholz³⁸, Rainer Buescher⁴, Karsten Häffner³⁹, Oliver Gross⁴⁰, Wanja Bernhardt⁴¹, Anke Doyon²⁵, Michael Henn²⁷, Jan Halbritter⁴², Ute Derichs⁴³, Günter Klaus⁴⁴, Bärbel Lange-Sperandio⁴⁵, Barbara Uetz⁴⁶, Marcus Benz⁴⁷, Andrea Titieni¹⁰, Hagen Staude⁴⁸, Heinz E. Leichter⁴⁹, Neveen A. Soliman⁵⁰, Luis Enrique Lara², Francisco de la Cerda Ojeda⁵¹, Jerome Harambat⁵², Bruno Ranchin⁵³, Marc Fila⁵⁴, Claire Dossier⁵⁵, Olivia Boyer⁵⁶, Matko Marlais²¹, Stella Stabouli⁵⁷, Nakysa Hooman⁵⁸, Francesca Mencarelli⁵⁹, William Morello¹³, Germana Longo⁶⁰, Francesco Emma¹², Dovile Ruzgiene⁶¹, Anna Wasilewska²³, Irena Balasz-Chmielewska²⁸, Monika Miklaszewska⁶², Malgorzata Stanczyk²⁴, Przemyslaw Sikora⁶³, Mieczyslaw Litwin¹¹, Aurelia Morawiec-Knysak²², Ana Teixeira⁶⁴, Gordana Milosevski-Lomic⁶⁵, Larisa Prikhodina⁶⁶, Rina Rus⁶⁷, Houweyda Jilani⁶⁸, Engin Melek⁶⁹, Ali Duzova⁷⁰, Alper Soylu⁷¹, Cengiz Candan⁷², Lale Sever⁷³, Alev Yilmaz⁷⁴, Neslihan Cicek⁹, Nurver Akinci⁷⁵, Sevgi Mir⁷⁶, Ismail Dursun⁷⁷, Yilmaz Tabel⁷⁸, Hulya Nalcacioglu⁷⁹,

²⁹Pediatric Nephrology Department, Dubai Kidney Center of Excellence, Dubai Hospital, Dubai, UAE.

³⁰Department of Pediatrics, Université Catholique de Louvain Medical School, Saint-Luc Academic Hospital,

Brussels, Belgium. ³¹Centre de référence de Néphrologie Pédiatrique Sud, Clinique de l'Espérance, Montegnée,

Belgium. ³²Hôpitaux Universtaires de Genève, Unité Romande de Néphrologie Pédiatrique, Geneva, Switzerland.

³³Pediatric Nephrology Unit, University Children's Hospital, Zurich, Switzerland. ³⁴Department of Pediatrics,

University Hospital Motol, 2nd Faculty of Medicine, Charles University Prague, Prague, Czech Republic.

³⁵Department of Pediatric Nephrology, Charité–Universitätsmedizin Berlin, Corporate Member of Freie Universität

Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany. ³⁶Department II of Internal

Medicine, University Hospital of Cologne, Cologne, Germany. ³⁷Department of Pediatrics and Adolescent Medicine,

University of Erlangen-Nürnberg (FAU), Erlangen, Germany. ³⁸Department of Nephrology and Hypertension,

University of Erlangen-Nürnberg, Erlangen, Germany. ³⁹Department of General Pediatrics, Adolescent Medicine

and Neonatology, Freiburg University Hospital, Freiburg, Germany. ⁴⁰Clinic for Nephrology and Rheumatology,

University Medical Center Goettingen, Goettingen, Germany. ⁴¹Nephrology Clinic Hannover, Hannover, Germany.

⁴²Division of Nephrology, University Hospital Leipzig, Leipzig, Germany. ⁴³Pediatric Nephrology, Center for

Paediatric and Adolescent Medicine, University Medical Clinic, Mainz, Germany. ⁴⁴KfH Center of Paediatric

Nephrology, University Hospital of Marburg, Marburg, Germany. ⁴⁵Dr. von Haunersches Kinderspital, Ludwigs

Maximilian University, Munich, Germany. ⁴⁶KfH Center of Pediatric Nephrology, Children's Hospital Munich

Schwabing, Munich, Germany. ⁴⁷Klinik für Kinder-und Jugendmedizin, Klinikum Dritter Orden, München,

Germany. ⁴⁸Pediatric Nephrology, University Children's Hospital Rostock, Rostock, Germany. ⁴⁹Olga Children's

Hospital, Clinic Stuttgart, Stuttgart, Germany. ⁵⁰Department of Pediatrics, Center of Pediatric Nephrology and

Transplantation, Kasr Al Ainy School of Medicine, Cairo University, Cairo, Egypt. ⁵¹Hospital Virgen del Rocío, Sevilla,

Spain. ⁵²Department of Pediatrics, Bordeaux University Hospital, Bordeaux, France. ⁵³Pediatric Nephrology Unit,

Hôpital Femme Mere Enfant, Hospices Civils de Lyon, Lyon, France. ⁵⁴Pediatric Nephrology Unit, CHU Arnaud de Villeneuve-Université de Montpellier, Montpellier, France. ⁵⁵Service de Néphrologie Pédiatrique, Hôpital Robert-debré, Paris, France. ⁵⁶Department of Pediatric Nephrology and Kidney Transplantation, Necker Hospital, Paris, France. ⁵⁷First Department of Pediatrics, Hippokraton Hospital, Aristotle University of Thessaloniki, Thessaloniki, Greece. ⁵⁸Department of Pediatric Nephrology, Ali-Asghar Children Hospital, University of Medical Sciences, Tehran, Iran. ⁵⁹Nephrology and Dialysis Unit, Department of Pediatrics, Azienda Ospedaliero Universitaria Sant'Orsola-Malpighi, Bologna, Italy. ⁶⁰Nephrology, Dialysis and Transplant Unit, Department of Women's and Children's Health, University Hospital of Padova, Padova, Italy. ⁶¹Clinic of Children Diseases, Institute of Clinical Medicine, Vilnius University, Vilnius, Lithuania. ⁶²Department of Pediatric Nephrology and Hypertension, Faculty of Medicine, Jagiellonian University Medical College, Kraków, Poland. ⁶³Department of Pediatric Nephrology, Medical University of Lublin, Lublin, Poland. ⁶⁴Pediatric Nephrology, Centro Hospitalar São João, Porto, Portugal. ⁶⁵Department of Nephrology, University Children's Hospital, Belgrade, Serbia. ⁶⁶Department of Inherited and Acquired Kidney Diseases, Research Clinical Institute for Pediatrics, Pirogov Russian National Research Medical University, Moscow, Russia. ⁶⁷Department of Paediatric Nephrology, University Children's Hospital, University Medical Centre Ljubljana, Ljubljana, Slovenia. ⁶⁸Service des Maladies Congénitales et Hérititaires, CHU Mongi Slim La Marsa, Sidi Daoud La Marsa, Tunis, Tunisia. ⁶⁹Department of Pediatric Nephrology, Faculty of Medicine, Cukurova University, Adana, Turkey. ⁷⁰Department of Pediatrics, Division of Pediatric Nephrology, Faculty of Medicine, Hacettepe University, Ankara, Turkey. ⁷¹Department of Pediatric Nephrology, Dokuz Eylul University Medical Faculty, Balçova, Izmir, Turkey. ⁷²Division of Pediatric Nephrology, Istanbul Medeniyet University, Göztepe Hospital, Istanbul, Turkey. ⁷³Department of Pediatric Nephrology, Cerrahpaşa School of Medicine, Istanbul University, Istanbul, Turkey. ⁷⁴Pediatric Nephrology Department, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey. ⁷⁵Division of Pediatric Nephrology, Sisli Etfal Training and Research Hospital, Istanbul, Turkey. ⁷⁶Department of Pediatric Nephrology, Ege University Medical Faculty, Izmir, Turkey. ⁷⁷Department of Pediatric Nephrology, Faculty of Medicine, Erciyes University, Kayseri, Turkey. ⁷⁸Department of Pediatric Nephrology, Faculty of Medicine, İnönü University, Malatya, Turkey. ⁷⁹Division of Pediatric Nephrology, Faculty of Medicine, Ondokuz Mayıs University, Samsun, Turkey.