

Higher spins and Finsler geometry

Alessandro Tomasiello

Dipartimento di Matematica, Università degli Studi di Milano–Bicocca,
Via Cozzi 55, 20126 Milano, Italy

and

INFN, sezione di Milano–Bicocca

`alessandro.tomasiello@unimib.it`

Finsler geometry is a natural generalization of (pseudo-)Riemannian geometry, where the line element is not the square root of a quadratic form but a more general homogeneous function. Parameterizing this in terms of symmetric tensors suggests a possible interpretation in terms of higher-spin fields. We will see here that, at linear level in these fields, the Finsler version of the Ricci tensor leads to the curved-space Fronsdal equation for all spins, plus a Stueckelberg-like coupling. Nonlinear terms can also be systematically analyzed, suggesting a possible interacting structure. No particular choice of spacetime dimension is needed. The Stueckelberg mechanism breaks gauge transformations to a redundancy that does not change the geometry. This is however not enough to eliminate non-transverse modes, at least for some versions of Finsler dynamics.

Contents

1	Introduction	1
2	Fronsdal equation	4
2.1	Spin two	4
2.2	Symmetric products	5
2.3	Differential operators	6
2.4	Flat space	8
2.5	Curved space	9
3	Finsler geometry	10
3.1	Distance	11
3.2	Connection	12
3.3	Curvature	15
4	Linearized analysis	15
4.1	Deformations	16
4.2	Parametrization	17
4.3	Sum over spins	18
5	Nonlinearities	20
5.1	General expansion	20
5.2	Spin-six nonlinear equation	21
6	Challenges for Finsler dynamics	22
6.1	Equations of motion	23
6.2	Broken Fronsdal gauge transformations	24
6.3	Perturbations	27
6.4	Discussion	29
A	More details on the symmetrized derivative	30

1 Introduction

The dynamics of fields with spin higher than two has an intricate structure. Their free equations of motion are already quite complicated when they have a mass [1], but simplifies in the massless case, where their equation of motion mimics that of spin

two, using the so-called Fronsdal operator \mathcal{F} [2]. Their interactions have a highly constrained dynamics; for reviews see for example [3–7]. It has long been surmised that they are important for theories of quantum gravity. String theory does have a tower of such fields, with masses related to the string tension. On the other hand, theories of Vasiliev type describe massless higher spins in (A)dS [8–12]. Their status as quantum theories is bolstered by their holographic interpretation [13, 14]. Higher spin fields and symmetries are expected more generally in holography, although most often in a broken version [15]. Some versions of these theories can indeed be even obtained as limits of string theory [16, 17].

Finsler geometry is a natural generalization of Riemannian and pseudo-Riemannian geometry; for some recent introductions see [18–20]. In the former case, instead of writing the length of a curve as $\int d\tau \sqrt{2g_2}$, where $g_2 \equiv \frac{1}{2}g_{\mu\nu}\dot{x}^\mu\dot{x}^\nu$, one considers a more general $\int d\tau F(x, \dot{x})$. The homogeneity rule $F(x, \lambda\dot{x}) = \lambda F(x, \dot{x})$ is needed to ensure that such a length does not depend on how the curve is parameterized, but this still leaves many possibilities. A homogeneous analogue of a power series leads to a fairly general expression:¹

$$F^2 = 2g_2 + \frac{\phi_4}{g_2} + \frac{\phi_6}{g_2^2} \dots, \quad \phi_s \equiv \frac{1}{s!} \phi_{\mu_1 \dots \mu_s} \dot{x}^{\mu_1} \dots \dot{x}^{\mu_s}. \quad (1.1)$$

So the first term in this expansion leads to the usual notion of distance; the others are new. One can also think of this as a line element $ds^2 = ds_0^2 + (2/ds_0^2)\phi_{\mu\nu\rho\sigma}dx^\mu dx^\nu dx^\rho dx^\sigma + \dots$, where $ds_0^2 = g_{\mu\nu}dx^\mu dx^\nu$ is of the customary (pseudo-)Riemannian type.

Analogues of the usual notions of connection and curvature have been studied for Finsler geometry for a long time. These notions are most natural when F is considered as a function on the tangent bundle TM ,² with \dot{x}^μ now promoted to a coordinate y^μ along the fiber.

Mathematically it looks natural to build a Finsler modification of general relativity (GR); this has indeed been explored at length in the literature (see for example [21, 22] for reviews). But from a physical point of view, no model of gravity can be considered to be well-motivated unless it improves on Einstein gravity on the crucial issue of its quantum behavior. This important issue appears to have been relatively unexplored.

In any case, the appearance of the symmetric tensors $\phi_{\mu_1 \dots \mu_s}$ in (1.1) suggests a relation to higher-spin theories. Given the promising status of the latter as quantum

¹Odd s can also be introduced, as we will discuss later.

²More precisely, because of the homogeneity constraint on F , one has to work either with the *slit* tangent bundle TM –(zero section), or with the *sphere* bundle SM inside it, where each fiber is quotiented by overall rescalings.

theories, such a link would also make Finsler geometry a lot more interesting physically. The $\int d\tau F$ would then be interpreted as a natural coupling of the $\phi_{\mu_1 \dots \mu_s}$ to a particle.

A version of (1.1) with $s = 2$ and one $s > 2$ was considered in this context [23, (5.8)], and an approximate gauge transformation for the particle action was noticed. In [24] an analogue of Finsler geometry (without the homogeneity constraint) was used for W-gravity in low dimensions. Most relevant for us, (1.1) was found in [25] for $s = 4$ and $g_2 = \eta_2$ (flat space) to be related to the spin-4 Fronsdal equation [2].

In this paper I will explore (1.1) more systematically, in the spirit of taking the ϕ_s to be Finsler deformations around an ordinary (pseudo-)Riemannian geometry described by $g_2 = \frac{1}{2}g_{\mu\nu}y^\mu y^\nu$. The first intriguing result is that the Fronsdal kinetic operator $\mathcal{F}(\phi_s)$ appears for all spins $s > 2$ and around any g_2 . Among the curvature tensors, one that we will call ρ is the analogue of the ordinary Ricci tensor. We will see that

$$\rho \sim R_{\alpha\beta}^0 y^\alpha y^\beta + \sum_{s>2} g_2^{1-s/2} \left(-\frac{1}{2}\mathcal{F}(\phi_s) + \alpha_s d^2 \phi_{s-2} \right), \quad (1.2)$$

to linear order in the ϕ_s . The first term contains the ordinary Ricci tensor of g_2 ; the rest is a series similar to (1.1). In a condensed notation to be fully explain below, d represents the symmetrized derivative $\nabla_{(\mu_1} \phi_{\mu_2 \dots \mu_s}^{s-2}$. The coefficient $\alpha_s = \frac{1}{8}(s-4)(s+D-4)$, where D is the spacetime dimension, which we will leave unspecified throughout. A perhaps deeper expression is

$$\rho \sim R_{\alpha\beta}^0 y^\alpha y^\beta + \frac{1}{2}\mathcal{F}(\delta F^2) \quad (1.3)$$

with an appropriate understanding of the action of \mathcal{F} on arbitrary functions of y . This is valid for Finsler deformations δF^2 around any (pseudo-)Riemannian geometry, without even using the expansion (1.1).

A second interesting point is that the Finsler Ricci tensor is not linear in the ϕ_s ; going to higher orders gives rise to expressions that are more complicated but still manageable, especially for low s . It is natural to interpret these as higher spin interactions, similar to the graviton interactions appearing in the perturbative expansion of GR around Minkowski space.

Unfortunately there is also bad news. What makes the Fronsdal operator especially important is that it admits gauge transformations: under $\delta\phi_s = d\lambda_{s-1}$, $\delta\mathcal{F}(\phi_s) = \frac{1}{2}d^3\lambda'_{s-1}$, with λ'_{s-1} representing the trace $\lambda^\mu{}_{\mu\mu_2 \dots \mu_s}$. Thus the traceless λ_{s-1} give a large set of gauge transformations.³ On the other hand, the double symmetrized derivative d^2 in (1.2) is not invariant. The two terms combine to transform as $\delta(\mathcal{F}(\phi_s) - 2\alpha_s d^2 \phi_{s-2}) =$

³This traceless condition is often viewed with suspicion; there is a way to get rid of it [26]. In the context of Finsler geometry we will see that in a sense this condition is rather natural.

$d^3(-\frac{1}{4}\lambda'_{s-1} + \alpha_s \lambda_{s-3})$. The gauge transformation of each field is “eaten” by that of a higher spin, in a sort of Stückelberg mechanism. Unfortunately, the surviving transformations can be shown to be a trivial rearrangement of fields that leaves the geometry unchanged. (An equivalent, sleeker analysis is possible from (1.3).)

This essential lack of gauge transformations (beyond the usual diffeomorphisms) is worrisome. We will see that the equations of motion for a Finsler analogue of GR, the simplest of which just reads $\rho = 0$, have many perturbative solutions that are not transverse to the momentum, and thus would presumably create trouble upon quantization. These solutions also have many free parameters; this might give hope that a gauge transformation can indeed be defined for this system. Unfortunately I have found no such candidate, but I have not excluded it either.

There are of course many other possible problems with a theory of higher spins [3]. The Vasiliev theory manages to avoid them in a rather sophisticated fashion. In contrast, the idea of a Finsler higher spin theory should be regarded as moving here one of its first steps. Nevertheless, the present results make it worth exploring further in the future.

In Sec. 2 some aspects of higher spin field theory are reviewed, setting up notation. Sec. 3 is a lightning introduction to Finsler geometry. In Sec. 4 I connect the two, showing that a linearization of the Finsler analogue ρ of the Ricci tensor contains infinitely many Fronsdal operators, as promised above. In Sec. 5 a partial analysis is given beyond the linear order in the ϕ_s . Sec. 6 considers the physical issues of gauge transformations and of perturbative solutions, with some details in App. A.

2 Fronsdal equation

The Fronsdal equation [2] describes a massless free field in a totally symmetric representation. We will give here a very basic review of some its features. There are many deeper discussions in the literature [3–5].

2.1 Spin two

As a warm-up, let us recall the equation for a spin-two field. This can be obtained by applying the vacuum equations of general relativity, $R_{\mu\nu} = 0$, to an infinitesimal perturbation $g_{\mu\nu} + h_{\mu\nu}$ of a background metric $g_{\mu\nu}$. The connection is deformed by

$$\delta\Gamma_{\nu\rho}^{\mu} = g^{\mu\sigma} \left(\nabla_{(\nu} h_{\rho)\sigma} - \frac{1}{2} \nabla_{\sigma} h_{\nu\rho} \right) \quad (2.1)$$

and the Riemann tensor by

$$\delta R_{\nu\rho} = 2\nabla_{[\mu}\delta\Gamma_{\nu]\rho}^{\mu} = -\frac{1}{2}\nabla^2 h_{\nu\rho} + \nabla^{\mu}\nabla_{(\nu}h_{\rho)\mu} - \frac{1}{2}\nabla_{\nu}\nabla_{\rho}h', \quad (2.2)$$

where $h' = g^{\mu\nu}h_{\mu\nu}$. So setting the right-hand side to zero is the equation of motion for $h_{\mu\nu}$.

Sometimes it can be useful to rewrite (2.2) in terms of the *Lichnerowicz operator* Δ :

$$\delta R_{\nu\rho} = -\frac{1}{2}\Delta\delta g_{\nu\rho} + \nabla_{(\nu}\nabla^{\mu}\delta g_{\rho)\mu} - \frac{1}{2}\nabla_{\rho}\nabla_{\nu}\delta g', \quad (2.3a)$$

$$\Delta\delta g_{\nu\rho} \equiv \nabla^2\delta g_{\nu\rho} - 2R_{(\nu}{}^{\sigma}\delta g_{\rho)\sigma} + 2R_{\mu\nu\rho\sigma}h^{\mu\sigma}. \quad (2.3b)$$

The gauge transformations of general relativity are infinitesimal coordinate changes; if we take

$$\delta h_{\mu\nu} = 2\nabla_{(\mu}\xi_{\nu)}, \quad (2.4)$$

(2.2) gives

$$\delta_{\xi}R_{\mu\nu} = \xi^{\rho}\nabla_{\rho}R_{\mu\nu} + 2\nabla_{(\mu}\xi^{\rho}R_{\nu)\rho} = L_{\xi}R_{\mu\nu} \quad (2.5)$$

after using the familiar

$$[\nabla_{\mu}, \nabla_{\nu}]v^{\rho} = R^{\rho}{}_{\sigma\mu\nu}v^{\sigma} \quad (2.6)$$

and its generalization on tensors. If the original metric is a solution of the vacuum equation, $R_{\mu\nu} = 0$, and so (2.2) is invariant.

2.2 Symmetric products

To set the stage for a generalization to higher spins, let us consider now a condensed notation. A version of this formalism is used one way or another in much of the higher-spin literature; see for example [3, (2.10)]. My definitions are adapted to Finsler geometry, so some normalizations might be a bit unfamiliar.

We introduce a formal variable y^{μ} (which will later acquire an interpretation as a velocity vector). To a completely symmetric tensor $\phi_{\mu_1\dots\mu_s}$ we associate a polynomial

$$\phi_s \equiv \frac{1}{s!}\phi_{\mu_1\dots\mu_s}y^{\mu_1}\dots y^{\mu_s}; \quad (2.7)$$

this is of course similar to the notation in (1.1). It is convenient to also define

$$\begin{aligned} \phi_{\mu}^s &\equiv \partial_{y^{\mu}}\phi_s = \frac{1}{(s-1)!}\phi_{\mu_2\dots\mu_s}y^{\mu_2}\dots y^{\mu_s}, \\ \phi_{\mu\nu}^s &\equiv \partial_{y^{\mu}}\partial_{y^{\nu}}\phi_s = \frac{1}{(s-2)!}\phi_{\mu_3\dots\mu_s}y^{\mu_3}\dots y^{\mu_s}, \end{aligned} \quad (2.8)$$

and so on. The s is a label and can be written up or down as convenient, and will sometimes be omitted when clear from the context. Notice that

$$y^\mu \phi_\mu^s = s \phi_s. \quad (2.9)$$

The trace is defined as

$$\phi'_s \equiv \phi_{s\mu}^\mu = g^{\mu\nu} \partial_{y^\mu} \partial_{y^\nu} \phi_s. \quad (2.10)$$

A product of two polynomials $\phi_s \psi_{s'}$ represents in this language the symmetric product $\frac{(s+s')!}{s!s'!} \phi_{(\mu_1 \dots \mu_s \mu_{s+1} \dots \mu_{s+s'})}$. We will often encounter products $g_2 \phi_s$ or more generally $g_2^k \phi_s$, with $g_2 = \frac{1}{2} g_{\mu\nu} y^\mu y^\nu$ and $g_{\mu\nu}$ a metric. It is useful to compute the trace of such objects:

$$(g_2 \phi_s)' = g^{\mu\nu} \partial_{y^\mu} (y_\nu \phi_s + g_2 \phi_\nu^s) = D \phi_s + 2y^\mu \phi_\mu^s + g_2 \phi'_s = (D + 2s) \phi_s + g_2 \phi'_s. \quad (2.11)$$

One can rewrite this more abstractly by introducing the operators

$$t \equiv g^{\mu\nu} \partial_{y^\mu} \partial_{y^\nu} = \partial_{y^\mu} \partial_{y^\mu}, \quad \text{deg} \equiv y^\mu \partial_{y^\mu}, \quad (2.12)$$

so that $t \phi_s = \phi'_s$ and $\text{deg} \phi_s = s \phi_s$, respectively. With the usual commutator rules, it is easy to work out

$$[t, g_2 \cdot] = \partial_{y^\mu} y^\mu + y^\mu \partial_{y^\mu} = D + 2 \text{deg}, \quad (2.13)$$

where $g_2 \cdot$ is multiplication by g_2 . This reproduces (2.11).

Any ϕ_s can be written in terms of traceless tensors:

$$\phi_s = \phi_{s,0} + g_2 \phi_{s,1} + \dots + g_2^{s/2} \phi_{s,s}, \quad \phi'_{s,k} = 0. \quad (2.14)$$

The degree $(s - 2k)$ fields $\phi_{s,k}$ can also be written in terms of the iterated traces $\phi^{(j)}$ (in a notation where $\phi^{(1)} = \phi'$, $\phi^{(2)} = \phi''$ and so on):

$$\phi_{s,0} = \phi_s + t_{s1} g_2 \phi'_s + t_{s2} \frac{g_2^2}{2} \phi''_s + \dots, \quad \phi_{s,j} = (-)^j t_{s-j+1,j} (\phi^{(j)})_0, \quad (2.15)$$

where

$$t_{s1} \equiv -\frac{1}{D + 2s - 4}, \quad t_{s,j} \equiv -\frac{t_{s,j-1}}{D + 2s - 2j - 2}. \quad (2.16)$$

2.3 Differential operators

The placeholder variable y is inert under derivatives. So $\nabla_\mu \phi_s = \frac{1}{s!} \nabla_\mu \phi_{\mu_1 \dots \mu_s} y^{\mu_1} \dots y^{\mu_s}$. We will often encounter the operator

$$d \equiv y^\mu \nabla_\mu. \quad (2.17)$$

This name is inspired by $d = dx^\mu \wedge \partial_\mu = dx^\mu \wedge \nabla_\mu$ in exterior algebra, which represents an antisymmetrized derivative, similar to how (2.17) represents a *symmetrized* derivative:

$$d\phi_s = \frac{1}{s!} \nabla_{(\mu} \phi_{\mu_1 \dots \mu_s)} y^\mu y^{\mu_1} \dots y^{\mu_s}. \quad (2.18)$$

(Writing the symmetrizer on the indices is here of course optional, as the product of y 's enforces it anyway.) A crucial difference is that the d in exterior algebra squares to zero, while (2.17) does not.

In this language, the equation of motion $R_{\mu\nu} = 0$ from (2.2), namely $\frac{1}{2} \nabla^2 h_{\nu\rho} - \nabla^\mu \nabla_{(\nu} h_{\rho)\mu} + \frac{1}{2} \nabla_\nu \nabla_\rho h' = 0$, can be rewritten by multiplying by $y^\nu y^\rho$ as

$$\nabla^2 h_2 - \nabla_\mu d h_2^\mu + \frac{1}{2} d^2 h' = 0. \quad (2.19)$$

The gauge transformation (2.4) reads

$$\delta h_2 = d\xi_1. \quad (2.20)$$

We seem not to have gained much by this rewriting. However, the nice properties of (2.19) now suggest to replace h_2 in (2.19) by a polynomial ϕ in y , obtaining the equation of motion

$$\mathcal{F}(\phi) \equiv \nabla^2 \phi - \nabla_\mu d\phi^\mu + \frac{1}{2} d^2 \phi' = 0 \quad (2.21)$$

irrespectively of the degree of ϕ . This is the *Fronsdal equation* for massless totally symmetric fields. It is much simpler than its explicit expression: for $\phi = \phi_s$ as in (2.7),

$$\nabla^2 \phi_{\mu_1 \dots \mu_s} - s \nabla_\mu \nabla_{(\mu_1} \phi_{\mu_2 \dots \mu_s)}^\mu + \frac{s(s-1)}{2} \nabla_{(\mu_1} \nabla_{\mu_2} \phi_{\mu_3 \dots \mu_s)}^{\mu} = 0. \quad (2.22)$$

We can also rewrite

$$\mathcal{F} = \nabla^2 - \nabla^\mu d \partial_{y^\mu} + \frac{1}{2} d^2 t = \Delta - d d^\dagger + \frac{1}{2} d^2 t, \quad (2.23)$$

with $\Delta = \nabla^2 - R_{\mu\nu} y^\mu \partial_{y^\nu} + R_{\mu\rho\nu\sigma} y^\mu y^\nu \partial_{y^\rho} \partial_{y^\sigma}$ as in (2.3b), recalling (2.12), and with the definition

$$d^\dagger \equiv g^{\mu\nu} \partial_{y^\mu} \nabla_\nu = \partial_{y^\mu} \nabla_\mu. \quad (2.24)$$

We note for later use:

$$[t, d] = 2d^\dagger, \quad [t, d^\dagger], \quad [d^\dagger, g_2 \cdot] = d, \quad [d, g_2 \cdot] = 0, \quad [d, d^\dagger] = -2\nabla^2 + \Delta. \quad (2.25)$$

2.4 Flat space

We will now see that (2.21) inherits some of the nice properties of (2.19). In this subsection we work in flat space, $g_{\mu\nu} = \eta_{\mu\nu}$.

In flat space, a ϕ_s give a representation of the Lorentz group; it is irreducible if $\phi'_s = 0$. Otherwise, (2.14) can be viewed as a decomposition into s irreducible representations.

Generalizing the gauge transformation (2.20), we take that of a field $\phi = \phi_s$ to be

$$\delta\phi_s = d\lambda_{s-1}. \quad (2.26)$$

In components this reads $\delta\phi_{\mu_1\dots\mu_s} = s\partial_{(\mu_1}\lambda_{\mu_2\dots\mu_s)}$. We find

$$\delta\phi^\mu = \partial_{y_\mu}(y^\alpha\partial_\alpha\lambda) = \partial^\mu\lambda + d\lambda^\mu, \quad \delta\phi^{\mu\nu} = 2\partial^{(\mu}\lambda^{\nu)} + y^\alpha\partial_\alpha\lambda^{\mu\nu}. \quad (2.27)$$

(As in (2.21), when the equations don't depend on the degree/spin we sometimes omit it, to get more readable expressions.) So for the Fronsdal equation (2.21):

$$\begin{aligned} \delta\mathcal{F}(\phi) &= \mathcal{F}(d\lambda) = \partial^2 d\lambda - d\partial_\mu(\partial^\mu\lambda + d\lambda^\mu) + \frac{1}{2}d^2(2\partial_\mu\lambda^\mu + d\lambda') \\ &= \frac{1}{2}d^3\lambda'. \end{aligned} \quad (2.28)$$

In particular, $\mathcal{F}(\phi_s)$ is invariant under λ_{s-1} that are traceless:

$$\lambda' = \lambda^\alpha_\alpha = 0. \quad (2.29)$$

Recall that more explicitly this means $\lambda^\alpha_{\alpha\mu_3\dots\mu_{s-1}} = 0$.

Trying to write an action for the Fronsdal equation leads to a second constraint. Up to integration by parts,

$$S = \int d^Dx \phi \mathcal{G}, \quad \mathcal{G} \equiv \mathcal{F}(\phi) - \frac{1}{2}\eta_2\mathcal{F}'(\phi). \quad (2.30)$$

This action is in general not gauge invariant: $\delta S = \int d^Dx \lambda \partial_\mu\mathcal{G}^\mu \neq 0$. Fortunately, a few more cancellations yield

$$\partial_\mu\mathcal{G}^\mu = -\frac{1}{4}d^3\phi''. \quad (2.31)$$

We see that the action is invariant if we also impose that the double trace of the field vanishes:

$$\phi'' = \phi^{\alpha\beta}_{\alpha\beta} = 0. \quad (2.32)$$

(Recall that more explicitly this means $\phi^{\alpha\beta}_{\alpha\beta\mu_5\dots\mu_s} = 0$.) In this case ϕ consists of two irreducible representations, its trace ϕ' and its traceless part.

With this constraint, the variation $\delta S = \int d^D x \delta \phi \mathcal{G} = 0$ now only imposes that the double-traceless part of \mathcal{G} vanishes. Fortunately, it is simple to see (just by counting the number of available indices) that (2.32) in fact directly implies $\mathcal{F}'' = \mathcal{G}'' = 0$. So the action does imply $\mathcal{G} = 0$. This in turn sets to zero \mathcal{G}' , which is proportional to \mathcal{F}' . So the equation of motion is indeed $\mathcal{F} = 0$.

For spin two, the first constraint (2.29) was automatic because in that case λ_1 was in fact ξ_μ , which only has one index. (2.31) becomes the linearized flat-space version of the spin-two identity $\nabla^\mu (R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu}) = 0$, a consequence of the Bianchi identity. The double-trace constraint (2.32) is also automatic, since the field only has two indices.

We conclude with a quick review of gauge fixing. First we notice

$$\delta \left(\partial_\mu \phi^\mu - \frac{1}{2} d\phi' \right) = \partial^2 \lambda, \quad \left(\partial_\mu \phi^\mu - \frac{1}{2} d\phi' \right)' = -\frac{1}{2} d\phi'' = 0. \quad (2.33)$$

So we can always find a λ such that $\partial_\mu \phi^\mu - \frac{1}{2} d\phi' = 0$, by inverting the operator ∂^2 . This simplifies the Fronsdal equation to $\partial^2 \phi = 0$. There is still a residual gauge invariance, consisting of λ such that $\partial^2 \lambda = 0$ (and which are traceless).

We now go to momentum space, where we have $p^2 \phi(p) = 0$, $p_\mu \phi^\mu = \frac{1}{2} p_1 \phi'$, with $p_1 = y^\mu p_\mu$ as by now familiar. We pick a vector u^μ such that $u^\mu p_\mu = 1$. Using (2.27),

$$\delta(u^\mu \phi_\mu) = \lambda + u_\mu y^\alpha p_\alpha \lambda^\mu = M \lambda, \quad M \equiv 1 + p_1 u^\mu \partial_{y^\mu}. \quad (2.34)$$

It can be shown that M is invertible, with inverse $M^{-1} = \sum_l (-)^l (p_1^l / l!) \partial_{y^{\mu_1}} \dots \partial_{y^{\mu_l}}$. Moreover $M^{-1}(-u^\mu \phi_\mu) = \sum_l (-)^l (p_1^l / l!) u_{\mu_1} \dots u_{\mu_l} \phi^{\mu_1 \dots \mu_l}$ can be shown to be traceless. So by picking this λ we can set $u^\mu \phi_\mu$ to zero. Since M is invertible, it has no kernel, and so there is no residual gauge transformation. Finally, $p_\mu \phi^\mu = \frac{1}{2} p_1 \phi'$ implies $2p^\nu \phi_{\mu\nu} = p_\mu \phi' + p_1 \phi'_\mu$; contracting this with u_μ we find now that $\phi' = 0$, and so also that $p_\mu \phi^\mu = 0$. This is the analogue of the transverse traceless gauge for spin-two fields.

2.5 Curved space

In curved space, the gauge transformation of ϕ_s is still (2.26), but it now represents $\delta \phi_{\mu_1 \dots \mu_s} = s \nabla_{(\mu_1} \lambda_{\mu_2 \dots \mu_s)}$, with covariant derivatives. The transformation of (2.21) is more complicated: some terms that canceled in (2.28) no longer do so, because of the non-commutativity of covariant derivatives recalled in (2.6).

Besides the explicit indices, one also needs to take care of the tensor indices inside ϕ_s , made explicit in (2.7). In this respect, the following identity is useful:

$$[\nabla_\mu, \nabla_\nu] \phi = -R^\rho{}_{\sigma\mu\nu} y^\sigma \partial_{y^\rho} \phi. \quad (2.35)$$

Using also the second Bianchi identity we obtain

$$\delta\mathcal{F}(\phi) = \frac{1}{2}d^3\lambda' - y^\alpha y^\beta (\lambda^\nu \nabla_\nu R_{\alpha\beta} + 2\nabla_{(\alpha} \lambda^\nu R_{\beta)\nu}) + d\rho_{\mu\nu} \lambda^{\mu\nu} + 2\rho_{\mu\nu} d\lambda^{\mu\nu}. \quad (2.36)$$

We have introduced

$$\rho_{\mu\nu} \equiv R_{\mu\alpha\nu\beta} y^\alpha y^\beta \quad (2.37)$$

which will be useful later on. The parenthesis with the Ricci terms in (2.36) is reminiscent of the Lie derivative $L_\xi R_{\alpha\beta} = \xi^\mu \nabla_\mu R_{\alpha\beta} + 2\nabla_{(\alpha} \xi^\nu R_{\beta)\nu}$, except that $\lambda^\mu = \lambda_{s-1}^\mu = \frac{1}{(s-1)!} \lambda^{\mu_2 \dots \mu_{s-1}}$ has $(s-2)$ hidden indices. For $s=2$ this expression does reduce to the Lie derivative in (2.5).

The appearance of the Riemann tensor in (2.36) is famously problematic [27]. We can still take $\lambda' = 0$ to make the first term disappear. The Ricci tensor appears in the ordinary gravity equations, and in particular it simply vanishes for vacuum GR, so those terms can also be reasonably set to zero. However, setting the Riemann tensor to any particular value appears unjustified. So the gauge invariance of the Fronsdal equation is broken in general.

If one simply postulates that the background geometry is maximally symmetric, as in AdS_D or dS_D , then

$$R_{\mu\nu\rho\sigma} = \frac{\Lambda}{D-1} (g_{\mu\rho} g_{\nu\sigma} - g_{\mu\sigma} g_{\nu\rho}). \quad (2.38)$$

The constant is chosen so that $R_{\mu\nu} = \Lambda g_{\mu\nu}$. (2.36) now becomes the simpler

$$\delta\mathcal{F}(\phi_s) = \frac{1}{2}d^3\lambda'_{s-1} + \frac{4\Lambda}{D-1}g_2 d\lambda'_{s-1} - \Lambda\mu_s^2 d\lambda_{s-1}, \quad \mu_s^2 \equiv \frac{2(s-1)(D+s-3)}{D-1}. \quad (2.39)$$

(We reinstated the spin labels because now the equation does explicitly depend on s .) Recall that $g_2 = \frac{1}{2}g_{\mu\nu}y^\mu y^\nu$, per our notation (2.7). This is still non-zero even if $\lambda' = 0$. But the modified equation

$$\mathcal{F}(\phi_s) + \Lambda\mu_s^2\phi_s = 0 \quad (2.40)$$

does have a gauge invariance.

3 Finsler geometry

We will give here a relatively quick introduction to Finsler geometry. Some relatively recent accounts are [18–20], in increasing order of detail.

3.1 Distance

In Riemannian geometry, the length γ of a curve is given by the integral $\int_{\gamma} d\tau \sqrt{g_{\mu\nu}(x)\dot{x}^{\mu}\dot{x}^{\nu}}$, with τ a coordinate on γ and $\dot{x}^{\mu} \equiv \partial_{\tau}x^{\mu}$. The idea of Finsler geometry is to generalize this to

$$\int d\tau F(x, \dot{x}) \tag{3.1}$$

with $F(x, \dot{x})$ any function of x^{μ} , \dot{x}^{μ} that is homogeneous of degree one in velocity:

$$F(x, \lambda\dot{x}) = \lambda F(x, \dot{x}). \tag{3.2}$$

This requirement is needed so that the length (3.1) of γ is invariant under $\tau \rightarrow \tau'(\tau)$ and does not depend on how we parameterize it. Actually it is customary to call

$$\dot{x}^{\mu} \equiv y^{\mu} \tag{3.3}$$

the velocity variable, and we will do so. This can be viewed as a coordinate along the fiber of the tangent bundle TM . In this sense, F can be viewed as a function on TM , a fact that will be useful later.

Clearly there are many new options opened by the generalization (3.1), (3.2). Two simple possibilities that have been considered in the mathematical literature are the *Randers* choice $F = \phi_1 + \sqrt{2g_2}$ and the *s-th root* choice $F = (\phi_s)^{1/s}$, with $\phi_s = \frac{1}{s!}\phi_{\mu_1\dots\mu_s}y^{\mu_1}\dots y^{\mu_s}$ as in (2.7) and (1.1). That equation also shows infinitely many new possibilities, which will be explored in Sec.4 below.

Most of the mathematical literature has been devoted to the case where F is positive, thus defining a notion of distance that generalizes Riemannian geometry.⁴ In that case, F is also taken to be smooth, and the *fundamental tensor*

$$g_{\mu\nu} = \frac{1}{2}\partial_{y^{\mu}}\partial_{y^{\nu}}F^2 \tag{3.4}$$

is required to be positive-definite for all x and y . When there is danger of confusion between (3.4) and a Riemannian metric, I will call the latter $g_{\mu\nu}^0$. Notice that (3.4) depends on y , except if $F^2 = 2g_2^0 = g_{\mu\nu}^0(x)y^{\mu}y^{\nu}$, where $g_{\mu\nu} = g_{\mu\nu}^0(x)$ is an ordinary Riemannian metric. (3.4) is used to lower indices in Finsler geometry; given that it is positive-definite, its inverse $g^{\mu\nu}$ is used to raise them.

For physics, we are more interested in generalizing pseudo-Riemannian spacetimes; for this, we require instead (3.4) to have signature $(-+++)$. Now in spite of its name

⁴Not all metric spaces are of this form, as pointed out in [28]. There exist even more general notions of distance that cannot be obtained as integrals of a function $F(x, \dot{x})$.

F^2 is no longer assumed to be positive, similar to how the symbol ds^2 can have either sign in ordinary pseudo-Riemannian geometry. We take $\int d\tau\sqrt{F^2}$ to measure distances, and $\int d\tau\sqrt{-F^2}$ to measure proper time.

Even with the understanding that F^2 can have either sign, some choices are not defined for all y . An example is the pseudo-Randers $F^2 = \phi_1 + \sqrt{-2g_2}$, with $g_{\mu\nu}$ pseudo-Riemannian. Another example is encountered when generalizing (1.1) to include odd s , with terms $\phi_3/\sqrt{-g_2} + \phi_5/(-g_2)^{3/2} + \dots$. Physically this appears to be fine as long as, for every x , F^2 is real above a cone in the tangent space T_xM . Definitions in the mathematical literature vary as to whether to impose F^2 to be defined for all y or not [29]. The rest of our discussion in this Section will be basically valid for either signature.

Before we proceed further, we note that the homogeneity condition (3.2) has several interesting consequences. Recall the *Euler theorem* for a homogeneous function of degree k :

$$\phi(\lambda y) = \lambda^k \phi(y) \quad \Rightarrow \quad y^\mu \partial_{y^\mu} \phi = k\phi. \quad (3.5)$$

A monomial such as (2.7) is obviously homogeneous of degree s , as we saw in (2.9). If $\phi = \phi(x, y)$ is analytic around $y = 0$, it can be Taylor expanded in such a basis:

$$\phi = \phi_0 + \phi_\mu y^\mu + \frac{1}{2} \phi_{\mu\nu} y^\mu y^\nu + \dots. \quad (3.6)$$

In this sense, such a function can be seen as a collection of completely symmetric tensors $\phi_{\mu_1 \dots \mu_s}$.

F^2 is homogeneous of degree two; moreover, each ∂_{y^μ} lowers the homogeneity degree by one, so for example (3.4) has degree zero. This gives the following useful identities:

$$y^\mu \partial_{y^\mu} F^2 = 2F^2, \quad y^\mu g_{\mu\nu} = \frac{1}{2} \partial_{y^\nu} F^2, \quad y^\mu y^\nu g_{\mu\nu} = F^2, \quad y^\mu \partial_{y^\mu} g_{\nu\rho} = 0. \quad (3.7)$$

3.2 Connection

Just like in ordinary (pseudo-)Riemannian geometry, a connection is needed in order to write derivatives that transform well under coordinate changes. A first piece of information is obtained by varying (3.1): this gives the familiar-looking geodesic equation $\ddot{x}^\mu + \gamma_{\nu\rho}^\mu \dot{x}^\nu \dot{x}^\rho = 0$ with

$$\gamma_{\nu\rho}^\mu = \frac{1}{2} g^{\mu\sigma} (\partial_\nu g_{\rho\sigma} + \partial_\rho g_{\nu\sigma} - \partial_\sigma g_{\nu\rho}). \quad (3.8)$$

Recall however that g now also depends on $y^\mu = \dot{x}^\mu$. The combination appearing in the geodesic equation appears often enough that we give it a name, *spray coefficients*:

$$G_\mu \equiv \gamma_{\mu\nu\rho} y^\nu y^\rho = \frac{1}{2} (y^\nu \partial_\nu \partial_{y^\mu} - \partial_\mu) F^2. \quad (3.9)$$

The second equality uses (3.7).

We noticed earlier that F is a function on TM . It proves useful to also think of the derivatives ∂_μ and ∂_{y^μ} as vector fields on that bundle. Under a coordinate change $x^\mu \rightarrow x'^\mu = x'^\mu(x)$, the velocity variables should transform as $y^\mu \rightarrow y'^\mu = \frac{\partial x'^\mu}{\partial x^\nu} y^\nu$. If we now view both formulas together as a coordinate change on TM , it follows that

$$\partial_\mu \rightarrow \partial'_\mu = \frac{\partial x'^\nu}{\partial x^\mu} \frac{\partial}{\partial x'^\nu} + \frac{\partial y'^\nu}{\partial x^\mu} \frac{\partial}{\partial y'^\nu} = \frac{\partial x'^\nu}{\partial x^\mu} \partial'_\nu + y^\rho \frac{\partial^2 x'^\nu}{\partial x^\mu \partial x^\rho} \partial_{y'^\nu}, \quad (3.10a)$$

$$\partial_{y^\mu} \rightarrow \partial_{y'^\mu} = \frac{\partial y'^\nu}{\partial y^\mu} \frac{\partial}{\partial y'^\nu} = \frac{\partial x'^\nu}{\partial x^\mu} \partial_{y'^\nu}. \quad (3.10b)$$

The ∂_μ transform in a complicated fashion, but it turns out that the combination

$$\delta_\mu \equiv \partial_\mu - N^\nu{}_\mu \partial_{y^\nu}, \quad N^\nu{}_\mu \equiv \frac{1}{2} \partial_{y^\mu} G^\nu \quad (3.11)$$

transforms without the second term in (3.10a). A dual issue appears when considering forms: on TM the dy^μ transform in a complicated fashion, and this is solved by introducing

$$\delta y^\mu \equiv dy^\mu + N^\mu{}_\nu dx^\nu. \quad (3.12)$$

To see the importance of (3.11), consider the case of (pseudo-)Riemannian geometry, $F^2 = 2g_2^0 = g_{\mu\nu}^0 y^\mu y^\nu$. The fundamental tensor is $g_{\mu\nu} = g_{\mu\nu}^0$, and $N^\mu{}_\nu = \gamma_{\nu\rho}^\mu y^\rho$.⁵ On each of the monomials $T_s = \frac{1}{s!} T_{\mu_1 \dots \mu_s} y^{\mu_1} \dots y^{\mu_s}$ appearing in (1.1), we obtain $\delta_\mu T_s = \frac{1}{s!} \nabla_\mu T_{\mu_1 \dots \mu_s} y^{\mu_1} \dots y^{\mu_s}$, where ∇ is the covariant derivative relative to the (pseudo-)Riemannian metric $g_{\mu\nu}^0$. So for ordinary (pseudo-)Riemannian geometry δ_μ coincides with the usual covariant derivative ∇_μ , even when acting on objects that depend on y .

Returning to the general discussion, the importance of (3.11) suggests introducing the *Chern connection*

$$\Gamma_{\nu\rho}^\mu = \frac{1}{2} g^{\mu\sigma} (\delta_\nu g_{\rho\sigma} + \delta_\rho g_{\nu\sigma} - \delta_\sigma g_{\nu\rho}). \quad (3.13)$$

This has zero torsion, and satisfies

$$D_\mu g_{\nu\rho} \equiv \delta_\mu g_{\nu\rho} - \Gamma_{\mu\nu}^\sigma g_{\sigma\rho} - \Gamma_{\mu\rho}^\sigma g_{\nu\sigma} = 0. \quad (3.14)$$

We define the operator D_μ on any tensor, with Γ acting on the indices in the way that is familiar in (pseudo-)Riemannian geometry, and the partial derivative replaced by δ :⁶

$$\begin{aligned} D_\mu T_{\nu_1 \dots \nu_l}^{\mu_1 \dots \mu_k} &= \delta_\mu T_{\nu_1 \dots \nu_l}^{\mu_1 \dots \mu_k} + \Gamma_{\mu\rho}^{\mu_1} T_{\nu_1 \dots \nu_l}^{\rho \mu_2 \dots \mu_k} + \Gamma_{\mu\rho}^{\mu_2} T_{\nu_1 \dots \nu_l}^{\mu_1 \rho \mu_3 \dots \mu_k} + \dots + \Gamma_{\mu\rho}^{\mu_k} T_{\nu_1 \dots \nu_l}^{\mu_1 \dots \mu_{k-1} \rho} \\ &\quad - \Gamma_{\mu\nu_1}^\rho T_{\rho\nu_2 \dots \nu_l}^{\mu_1 \dots \mu_k} - \Gamma_{\mu\nu_2}^\rho T_{\nu_1 \rho \nu_3 \dots \nu_l}^{\mu_1 \dots \mu_k} - \dots - \Gamma_{\mu\nu_l}^\rho T_{\nu_1 \dots \nu_{l-1} \rho}^{\mu_1 \dots \mu_k}. \end{aligned} \quad (3.15)$$

⁵This is not true in general, because $\gamma_{\nu\rho}^\mu$ depends on y . By using (3.7) one can show that $\partial_{y^\nu} G_\mu = 2\gamma_{\mu\nu\rho} y^\rho$, but $N^\nu{}_\mu$ contains the derivative of G^μ with an *upper* index.

⁶In the literature this derivative is usually denoted by a “slash” at the end: $T^{\mu_1 \dots \mu_k}{}_{\nu_1 \dots \nu_l | \mu}$.

Once again, this reduces to ∇_μ in (pseudo-)Riemannian geometry, even if T depends on y . In this sense, (3.14) is more natural than the analogous equality containing $\gamma_{\nu\rho}^\mu$ and ordinary partial derivatives.

At a more conceptual level, on TM there is no connection that preserves the metric and has zero torsion; this is unlike in (pseudo-)Riemannian geometry, where (3.8) is the Levi-Civita connection and does have both properties. (3.8) and (3.13) both have zero torsion, but both fail to respect the metric in the y directions of TM . If on TM we write the exterior differential $d_{TM} \equiv dx^\mu \partial_\mu + dy^\mu \partial_{y^\mu} = dx^\mu \delta_\mu + \delta y^\mu \partial_{y^\mu}$, (3.14) implies

$$d_{TM} g_{\mu\nu} = \delta y^\rho \partial_{y^\rho} g_{\mu\nu} = \frac{1}{2} \delta y^\rho \partial_{y^\mu} \partial_{y^\nu} \partial_{y^\rho} F^2. \quad (3.16)$$

This property is called *almost compatibility* with the metric of the Chern connection (3.13). Other connections exist with nice properties, the most notable being the Cartan connection, which also has components along the y coordinates; it is metric-compatible but has torsion [30, Thm. 1.4.2].

The coefficients we introduced previously are related to the Chern connection:

$$\Gamma_{\nu\rho}^\mu y^\rho = N^\mu{}_\nu, \quad N^\mu{}_\nu y^\nu = G^\mu. \quad (3.17)$$

The second is a simple consequence of G^μ being homogeneous in y with degree two; the first is a little more involved.

It is also useful to notice that

$$[\partial_{y^\mu}, \delta_\nu] = -\partial_{y^\mu} N^\rho{}_\nu \partial_{y^\rho} = -(L_{\mu\nu}^\rho + \Gamma_{\mu\nu}^\rho) \partial_{y^\rho}, \quad L_{\mu\nu}^\rho \equiv y^\sigma \partial_{y^\mu} \Gamma_{\nu\sigma}^\rho. \quad (3.18)$$

We introduced the *Landsberg tensor* $L_{\mu\nu}^\rho$. As an application,

$$\partial_{y^\mu} \delta_\nu T = D_\nu \partial_{y^\mu} T - L_{\mu\nu}^\rho \partial_{y^\rho} T \quad (3.19)$$

on a function $T = T(x, y)$. One can show that

$$y^\nu L_{\mu\nu}^\rho = 0, \quad (3.20)$$

so in particular (3.19) implies

$$y^\nu \partial_{y^\mu} \delta_\nu T = y^\nu D_\nu \partial_{y^\mu} T = d \partial_{y^\mu} T. \quad (3.21)$$

In the last step we have defined

$$d \equiv y^\mu D_\mu \quad (3.22)$$

by analogy with the d in (2.17), to which it reduces in the (pseudo-)Riemannian case.

The contraction

$$L^\mu = g^{\nu\rho} L_{\nu\rho}^\mu \quad (3.23)$$

is called *mean Landsberg tensor*. On the other hand,

$$L_{\mu\nu}^\mu = y^\rho \partial_{y^\nu} \Gamma_{\mu\rho}^\mu = \frac{1}{2} y^\rho \partial_{y^\nu} \delta_\rho \log \det g = dI_\nu, \quad (3.24)$$

where

$$I_\mu = \frac{1}{2} \partial_{y^\mu} \log \det g, \quad (3.25)$$

is called the *mean Cartan tensor*.

3.3 Curvature

In view of (3.13), we now define

$$\frac{1}{2} R^\mu{}_{\nu\rho\sigma} \equiv \delta_{[\rho} \Gamma_{\sigma]\nu}^\mu + \Gamma_{[\rho|\lambda}^\mu \Gamma_{|\sigma]\nu}^\lambda. \quad (3.26)$$

It has many of the usual properties of the ordinary Riemann tensor: $R^\mu{}_{\nu(\rho\sigma)} = 0$, $R^\mu{}_{[\nu\rho\sigma]} = 0$. However, it is not antisymmetric in the first two indices: rather, it is possible to show

$$R_{(\mu\nu)\rho\sigma} = -\frac{1}{2} \partial_{y^\alpha} g_{\mu\nu} R^\alpha{}_{\beta\rho\sigma} y^\beta. \quad (3.27)$$

A particular contraction of (3.26) with the y can be reexpressed in terms of $N^\mu{}_\nu$ or of G^μ :

$$\rho^\mu{}_\rho \equiv R^\mu{}_{\alpha\rho\beta} y^\alpha y^\beta = 2y^\sigma \delta_{[\rho} N^\mu{}_{\sigma]} \quad (3.28a)$$

$$= \partial_\rho G^\mu - \frac{1}{4} \partial_{y^\rho} G^\nu \partial_{y^\nu} G^\mu - \frac{1}{2} y^\nu \partial_\nu \partial_{y^\rho} G^\mu + \frac{1}{2} G^\nu \partial_{y^\nu} \partial_{y^\rho} G^\mu. \quad (3.28b)$$

(3.27) implies that $\rho_{\mu\rho} = \rho_{\rho\mu}$, $\rho_{\mu\rho} \equiv g_{\mu\nu} \rho^\nu{}_\rho$. We already encountered $\rho_{\mu\nu}$ in (2.37) in the (pseudo-)Riemannian case.⁷ The trace contains the Finsler extension of the Ricci tensor:

$$\rho \equiv \rho^\mu{}_\mu = R_{\mu\nu}^0 y^\mu y^\nu \quad (\text{when } F^2 = g_{\mu\nu}^0 y^\mu y^\nu). \quad (3.29)$$

4 Linearized analysis

We would now like to obtain some formulas for infinitesimal deformations of Finsler structures, inspired by (2.1), (2.2) in the ordinary (pseudo-)Riemannian case. As anticipated, this will lead to the appearance of the Fronsdal operator.

⁷In the literature on Finsler geometry, $\rho^\mu{}_\nu$ is called $(1/F^2)R^\mu{}_\nu$. I have decided to change this notation in order to avoid confusion with the Ricci tensor in (pseudo-)Riemannian geometry. Even more confusingly, $\rho^\mu{}_\mu$ is often called ‘‘Ricci scalar’’; we will not follow this custom.

4.1 Deformations

We first take a small detour and show a formula for a *finite* deformation. Given a F^2 , consider a second Finsler structure \tilde{F}^2 . It determines a new fundamental tensor, connection and curvature; we will denote them all with a tilde. After recalling (3.11), one finds

$$\tilde{G}^\mu = G^\mu + \frac{1}{2}\tilde{g}^{\mu\rho}(y^\nu\partial_{y^\rho}\delta_\nu - \delta_\rho)\tilde{F}^2. \quad (4.1)$$

It can be shown [18, Thm. 3.3.1] that the geodesics of \tilde{F}^2 coincide with those of F^2 iff $\delta_\nu\tilde{F} = y^\mu\partial_{y^\nu}\delta_\mu\tilde{F}$; in this case $\tilde{G}^\mu = G^\mu + Py^\mu$, with $P = \tilde{F}^{-1}y^\nu\delta_\nu\tilde{F}$. The two Finsler \tilde{F} and F are said to be *projectively equivalent*.

Coming now to the case of an infinitesimal deformation, (4.1) gives

$$\delta G^\mu = \frac{1}{2}g^{\mu\rho}(y^\nu\partial_{y^\rho}\delta_\nu - \delta_\rho)\delta F^2 = \frac{1}{2}g^{\mu\rho}(\mathrm{d}\partial_{y^\rho} - \delta_\rho)\delta F^2. \quad (4.2)$$

In the second step we have used (3.21), (3.22). (δF^2 is to be understood as $\delta(F^2)$.)

Recall from (3.29) that $\rho = \rho^\mu{}_\mu$ is the analogue of the Ricci tensor. So to obtain the analogue of (2.2) we need $\delta\rho = \delta\rho^\mu{}_\mu$. We can use either (3.28a) or (3.28b) to obtain

$$\delta\rho = D_\mu\delta G^\mu + L^\mu{}_{\mu\nu}\delta G^\nu - \frac{1}{2}\mathrm{d}\partial_{y^\mu}\delta G^\mu. \quad (4.3)$$

Combining with (4.2):

$$\begin{aligned} \delta\rho &= \frac{1}{2}g^{\mu\nu} \left(D_\mu\mathrm{d}\partial_{y^\nu} - D_\mu\delta_\nu - \frac{1}{2}\mathrm{d}^2\partial_{y^\mu}\partial_{y^\nu} \right) \delta F^2 \\ &\quad - \frac{1}{2}\mathrm{d}(L^\rho\partial_{y^\rho}\delta F^2) + \frac{3}{2}\mathrm{d}I_\nu\delta G^\nu + \frac{1}{2}I_\mu\mathrm{d}\delta G^\mu. \end{aligned} \quad (4.4)$$

(4.3) and (4.4) are similar to the first and second step in (2.2).

These results hold for a general deformation of a Finsler structure F^2 . If the latter is in fact a (pseudo-)Riemannian geometry, (4.2) becomes

$$\delta G^\mu = \frac{1}{2}(\mathrm{d}\partial_{y^\mu} - \nabla^\mu)\delta F^2. \quad (4.5)$$

As for (4.4), since neither g nor $\Gamma^\rho{}_{\mu\nu}$ depend on y , we have $L^\rho{}_{\mu\nu} = 0 = I_\mu$, so the second line in (4.4) vanishes. The last term on the first line contains $g^{\mu\nu}\partial_{y^\mu}\partial_{y^\nu}$; we saw back in (2.10) that this gives the trace operator t on monomials such as (2.7), and now we can consider it as extending it on arbitrary (even non-polynomial) functions of y . This point will be crucial later. With this understanding, the first line of (4.4) is nothing but the Fronsda operator (2.21):

$$\delta\rho = -\frac{1}{2}\mathcal{F}(\delta F^2), \quad (4.6)$$

as anticipated in the introduction.

In the next subsection we will motivate the parameterization (1.1), and apply it to (4.6).

4.2 Parametrization

We will now justify the expansion (1.1). The idea is to start from a (pseudo-)Riemannian geometry, and to deform it into the more general Finsler type.

(1.1) is not quite a series expansion, since F^2 should be homogeneous. In Euclidean signature, one way to deal with this constraint is to think of it as a function on a sphere. For fixed x , $F^2(x, y)$ is a function on $S^{D-1} \cong \mathbb{R}^D/\mathbb{R}_+$, where the quotient acts by rescaling. A basis of functions on the round sphere is given by the spherical harmonics. These can be viewed as polynomials of the type ϕ_s in (2.7) restricted to $\{2\delta_2 \equiv \delta_{\mu\nu}y^\mu y^\nu = 1\}$, or alternatively as functions on \mathbb{R}^D made homogeneous of degree zero by dividing them by an appropriate power of δ_2 :

$$\frac{1}{s!} \frac{1}{\delta_2^{s/2}} \phi_{\mu_1 \dots \mu_s} \quad (4.7)$$

where s is any non-negative integer, and the tensors $\phi_{\mu_1 \dots \mu_s}$ are irreducible: $\phi'_s = 0$.

This suggests that a good parameterization for F^2 can be obtained generalizing (4.7) by replacing δ_2 with $g_2 = \frac{1}{2}g_{\mu\nu}^0 y^\mu y^\nu$. More precisely, F^2 is homogeneous of degree two, not zero; this can be repaired by multiplying by an overall g_2 . This leads us to

$$F^2 = 2g_2 + \frac{\phi_3}{\sqrt{g_2}} + \frac{\phi_4}{g_2} + \dots = 2g_2 + \sum_{s>2} g_2^{1-s/2} \phi_s. \quad (4.8)$$

This extends (1.1) by including odd s . Actually their presence creates a possible issue in the pseudo-Lorentzian case because of the roots, as we discussed at the end of Sec. 3.1. It becomes more sensible then to make F^2 defined above the light cone by changing $(g_2)^{s/2} \rightarrow (-g_2)^{s/2}$. Our discussion in the following is not influenced by this issue.

As mentioned earlier, the ϕ_s should be irreducible for an expansion that is complete and non-redundant. But no harm is done if the ϕ_s are not traceless and (4.8) is redundant. In that case, F^2 is trivially invariant (even at a finite level) under

$$\delta_\kappa \phi_{s_0} = \kappa_{s_0}, \quad \delta_\kappa \phi_{s_0+2} = -g_2 \kappa_{s_0}. \quad (4.9)$$

This redundancy will provide useful cross-checks of our results. It can be fixed to make the ϕ_s traceless again, or fixed only partially by enforcing a weaker constraint such as $\phi''_s = 0$. (This was imposed in (2.32), but only in order to obtain a gauge-invariant

action for (2.21); here we will not be using that action.) Alternatively one can proceed by expanding each ϕ_s in (4.8) in terms of its traceless components as in (2.14), collecting powers of g_2 . This gives an expansion like (4.8) whose degree s coefficient is now the traceless

$$\sum_{j=0}^{\infty} \phi_{s+2j,j}. \quad (4.10)$$

Similar to (4.9), there is an invariance where we deform

$$\delta g_2 = \kappa_2, \quad \delta \phi_s = (2 - s/2)\kappa_2 g_{s-2}. \quad (4.11)$$

One could even use this to connect higher spins around a curved space to higher spins around flat space, at the price of making the ϕ_s maximally non-irreducible.

Strictly speaking, a complete parameterization would have included lower spins in (4.8); we have not done so because of our focus on higher spins. A spin-one term $\phi_1 \sqrt{g_2}$ term can be generated by changing $\phi_3 \rightarrow \phi_3 + g_2 \phi_1$ in (4.8), or in other words (4.9) for $s_0 = 1$. A spin-zero term $\phi_0 g_2$ is equivalent to rescaling the first term in (4.8), or (after renaming $g_2 \rightarrow g_2(1 + \phi_0)^{-1}$) to rescaling $\phi_s \rightarrow (1 + \phi_0)^{s/2-1}$. In the future it might be interesting to investigate the effect of including this scalar; I have not done so in what follows. It would be especially intriguing to explore the limit in which $\phi_0 \rightarrow -1$, and the first term in (4.8) would disappear.

An alternative to (4.8) would be obtained by applying the same logic directly to F , leading to $F = \sqrt{2g_2} + f_3/g_2 + f_4(g_2)^{3/2} + \dots$. Adding the spin-one term to this parameterization leads to a generalization of the Randers choice mentioned in Sec. 3.1.

We will use an expansion similar to (4.8) also for other y -dependent quantities, such as the connection and curvature:

$$G^\mu = \gamma_{0\alpha\beta}^\mu y^\alpha y^\beta + \sum_{s>2} g_2^{1-s/2} G_s^\mu, \quad \rho^\mu{}_\nu = R_{0\alpha\nu\beta}^\mu y^\alpha y^\beta + \sum_{s>2} g_2^{1-s/2} \rho_{s\nu}^\mu, \quad (4.12)$$

where as usual a 0 denotes quantities associated to the (pseudo-)Riemannian metric $g_{\mu\nu}^0$. Notice that the coefficients in this expansion are typically not traceless even if the ϕ_s are.

4.3 Sum over spins

Let us take the ϕ_s to be small, and work at linear order in them. In this case we can use the formalism developed in Sec. 4.1. Namely, we take $F^2 = 2g_2 = g_{\mu\nu} y^\mu y^\nu$ to be of (pseudo-)Riemannian type, and \tilde{F}^2 to be (4.8).

Recall that both D_μ and δ_μ reduce to the usual covariant derivative ∇_μ in the (pseudo-)Riemannian case. In particular, these operators have trivial action on the powers of g_2 in (4.12). On the other hand, the y derivatives do act nontrivially.

For G^μ , (4.5) gives

$$G_{Ls}^\mu = \frac{1}{2}(\mathrm{d}\phi_s^\mu - \nabla^\mu\phi_s) + \frac{4-s}{4}y^\mu\mathrm{d}\phi_{s-2}. \quad (4.13)$$

Here and later, the label L stands for ‘‘at linear order in the ϕ_s ’’. Recall that $\phi_s = 0$ for $s < 2$, so the last term in (4.13) is actually absent for G_3^μ .

G^μ is related to the analogue of a connection, while (4.13) is a tensor.⁸ The first term in its expansion (4.12) does contain a connection; the following terms are to be thought of as deformations, and it is a familiar result that the difference of two connections is a tensor.

A similar application of (4.8) to (4.6) gives

$$\rho_s^L \equiv (\rho_s^L)^\mu{}_\mu = -\frac{1}{2}\mathcal{F}(\phi_s) + \alpha_s\mathrm{d}^2\phi_{s-2}, \quad \alpha_s \equiv \frac{1}{8}(s-4)(D+s-4), \quad (4.14)$$

again with L denoting linear order in ϕ_s . (4.14) are tensors, as manifest from (4.4) but not in (3.28). This result was anticipated in the introduction as (1.2).

It is instructive to check invariance under the trivial transformation (4.9). Since it does not change F^2 , it should not change the geometric quantities above; let us focus on (4.14). From (2.27) (with $\partial_\mu \rightarrow \nabla_\mu$) we obtain the identity

$$\mathcal{F}(-g_2\phi_{s-2}) = -g_2\mathcal{F}(\phi_{s-2}) + \left(3 - \frac{D}{2} - s\right)\mathrm{d}^2\phi_{s-2}. \quad (4.15)$$

Fixing a particular $s_0 > 4$, there are two affected ϕ_s , which appear in a total of four terms:

$$\begin{aligned} \delta_\kappa\rho^L &= g_2^{-s_0/2}\mathrm{d}^2\delta_\kappa\phi_{s_0} + g_2^{1-s_0/2}\left(-\frac{1}{2}\mathcal{F}(\delta_\kappa\phi_{s_0}) + \alpha_{s_0}\mathrm{d}^2\delta_\kappa\phi_{s_0-2}\right) - \frac{1}{2}g_2^{2-s_0/2}\mathcal{F}(\delta_\kappa\phi_{s_0-2}) \\ &= g_2^{1-s_0/2}\left(-\alpha_{s_0+2} + \alpha_{s_0} + \frac{1}{2}(s_0-3) + \frac{D}{4}\right)\mathrm{d}^2\kappa_{s_0-2}. \end{aligned} \quad (4.16)$$

In the second step we have used (4.15). Imposing $\delta_\kappa\rho^L = 0$ for all s_0 gives a recurrence relation for the α_s , which is indeed satisfied by the value in (4.14). This also shows that the $\mathrm{d}^2\phi_{s-2}$ term in (4.14) is unavoidable.⁹

⁸One of the advantages of working with (4.5) is that it makes this property more or less manifest. It is of course possible, but more laborious, to get the same results from (3.9).

⁹A curious possibility is to use (4.15) to express $\mathrm{d}\phi_{s-2}$ in terms of Fronsdal operators, and to substitute in (4.14). This results in $\rho_s^L = -\frac{1}{2}\mathcal{F}(\psi_s)$, where $\psi_s = \frac{\alpha_{s+4}}{\alpha_{s+4}-\alpha_{s+2}}\phi_s - \frac{\alpha_s}{\alpha_{s+2}-\alpha_s}g_2\phi_{s-2}$, thus getting rid of the second term in (4.14). Inverting this relation one would naively seem to arrive at $F^2 = \sum g_2^{1-s/2}\psi_s$. However, the last step only works by summing a series that does not really converge.

A more complicated computation also gives

$$(\rho_s^L)^\mu{}_\rho = -\frac{1}{2}\mathcal{F}^\mu{}_\rho(\phi_s) + \frac{4-s}{4}\left(y^\mu\nabla_\rho d\phi_{s-2} + g_{\nu\rho}y^{[\nu}d\nabla^{\mu]}\phi_{s-2} - \frac{1}{2}\delta_\rho^\mu d^2\phi_{s-2} - g_{\nu\rho}y^{(\mu}d^2\phi_{s-2}^{\nu)}\right) - \frac{1}{16}(4-s)(6-s)y^\mu y_\rho d^2\phi_{s-4}, \quad (4.17)$$

where we introduced a Fronsdal “precursor”

$$\mathcal{F}^\mu{}_\rho(\phi_s) = \nabla_\rho\nabla^\mu\phi_s - \nabla_\rho d\phi_s^\mu - g^{\mu\nu}d\nabla_{[\nu}\phi_{\rho]}^s + \frac{1}{2}d^2\phi_{s\rho}^\mu \quad (4.18)$$

whose trace is $\mathcal{F}(\phi_s)$. Recall from (3.28) that $\rho^\mu{}_\nu$ contains the Finsler analogue of the whole Riemann tensor. We will actually not need it in this paper, but it might be useful in further investigations.

Finally we record the perturbative expression for the mean Cartan (3.25)

$$I_\mu^{Ls} = \frac{1}{4}\phi'_{s\mu} + \frac{4-s}{8}\left((D+s-2)\phi_\mu^{s-2} + y_\mu\phi'_{s-2}\right) + \frac{(4-s)(6-s)}{16}(D+s-4)y_\mu\phi_{s-4} \quad (4.19)$$

and for the Landsberg tensor (3.18):

$$L_{\mu\nu\rho}^{Ls} = \frac{1}{4}d\phi_{\mu\nu\rho}^s + \frac{3}{8}(4-s)\left(g_{(\mu\nu}d\phi_{\rho)}^{s-2} + y_{(\mu}d\phi_{\nu\rho)}^{s-2}\right) + \frac{3(4-s)(6-s)}{16}\left(y_{(\mu}y_\nu d\phi_{\rho)}^{s-4} + y_{(\mu}g_{\nu\rho)}d\phi^{s-4}\right) + \frac{(4-s)(6-s)(8-s)}{32}y_\mu y_\nu y_\rho d\phi_{s-6}. \quad (4.20)$$

Again here $I_\nu^L = \sum_{s>2} g_2^{1-s/2} I_\nu^{Ls}$, $L_{\mu\nu\rho}^L = \sum_{s>2} g_2^{1-s/2} L_{\mu\nu\rho}^{Ls}$.

5 Nonlinearities

We now go beyond the linear order in the ϕ_s . This quickly becomes complicated, but we will at least try to make the structure clear in Sec. 5.1. In Sec. 5.2 we make the general results more concrete for ρ_4 and ρ_6 .

5.1 General expansion

We still raise and lower indices with the unperturbed (pseudo-)Riemannian metric, now denoted by $g_{\mu\nu}^0$ to avoid confusion with the fundamental tensor (3.4) of Finsler geometry. We expand the latter and its inverse as in (4.8), (4.12):

$$g_{\mu\nu} = g_{\mu\nu}^0 + \sum_{s>2} g_2^{1-s/2} q_{\mu\nu}^s, \quad g^{\mu\nu} = g_0^{\mu\nu} + \sum_{s>2} g_2^{1-s/2} p_{\mu\nu}^s. \quad (5.1)$$

(4.8) gives

$$q_{\mu\nu}^s = \frac{1}{2}\phi_{\mu\nu}^s - \frac{s-4}{4}(\phi_{s-2}g_{\mu\nu}^0 + 2y_{(\mu}\phi_{\nu)}^{s-2}) + \frac{1}{8}(s-4)(s-6)y_{\mu}y_{\nu}\phi_{s-4}. \quad (5.2)$$

As usual, $\phi_{s<3}$ is understood to be zero; so $q_{\mu\nu}^3 = \frac{1}{2}\phi_{\mu\nu}^3$, $q_{\mu\nu}^4 = \frac{1}{2}\phi_{\mu\nu}^4$, $q_{\mu\nu}^5 = \frac{1}{2}\phi_{\mu\nu}^5 - \frac{1}{4}\phi_3g_{\mu\nu}^0 - \frac{1}{2}y_{(\mu}\phi_{\nu)}^3$, in the usual notation (2.8). The inverse $g^{\mu\nu}$ can be found by inverting (5.1). This is formally similar to several familiar problems in QFT: the answer can be written a bit implicitly as

$$p_s^{\mu\nu} = \sum_{\substack{(s_1, \dots, s_n) \\ \text{partitions of } s-2}} (-1)^n (q_{s_1+2}q_{s_2+2} \dots q_{s_n+2})^{\mu\nu}. \quad (5.3)$$

The parenthesis is explicitly $g_0^{\mu\mu_1} q_{\mu_1\nu_1}^{s_1+2} g_0^{\nu_1\mu_2} q_{\mu_2\nu_2}^{s_2+2} g_0^{\nu_2\mu_3} \dots q_{\mu_n\nu_n}^{s_n+2} g_0^{\nu_n\nu}$; the partitions are to be considered in all possible orders. The first two (5.3) are $p_3^{\mu\nu} = -q_3^{\mu\nu}$, $p_4^{\mu\nu} = -q_4^{\mu\nu} + q_{3\rho}^{\mu} q_3^{\rho\nu}$.

For G^μ , we can work with (4.1) instead of (4.5); it is still more convenient than the original (3.9). The non-linearity in G_s^μ comes from the inverse $g^{\mu\nu}$ in (4.1); in terms of (4.13) we can write

$$G_s^\mu = G_{Ls}^\mu + \sum_{s'>2} p_s^{\mu\nu} G_{L\nu}^{s-s'}, \quad (5.4)$$

The curvature is quite a bit more complicated:

$$\rho_{s\rho}^\mu = \nabla_\rho G_s^\mu - \frac{1}{2}d\partial_{y^\rho} G_s^\mu + \frac{s-4}{4}y_\rho dG_{s-2}^\mu + Q_{s\rho}^\mu. \quad (5.5)$$

The term $Q_{s\rho}^\mu$, quadratic in G_s^μ , is nastier and reads in general

$$\begin{aligned} Q_{s\rho}^\mu &= \sum_{s'=3}^{s-1} \left(-\frac{1}{4}(\partial_{y^\rho} G^\nu)_{s'} (\partial_{y^\nu} G^\mu)_{s-s'+2} + \frac{1}{2}G_{s'}^\nu (\partial_{y^\nu} \partial_{y^\rho} G^\mu)_{s-s'+2} \right); \\ (\partial_{y^\rho} G^\mu)_s &= \partial_{y^\rho} G_s^\mu - \frac{s-4}{2}y_\rho G_{s-2}^\mu, \\ (\partial_{y^\nu} \partial_{y^\rho} G^\mu)_s &= \partial_{y^\nu} \partial_{y^\rho} G_s^\mu - \frac{s-4}{2}(g_{\nu\rho} G_{s-2}^\mu + 2y_{(\nu} \partial_{y^\rho)} G_{s-2}^\mu) + \frac{1}{4}(s-4)(s-6)y_\nu y_\rho G_{s-4}^\mu. \end{aligned} \quad (5.6)$$

The expression (5.5) is only linear and quadratic in G_s^μ , but the p_s appearing in (5.4) are themselves very non-linear in the q_s and hence in the ϕ_s . In particular we see that the non-linearity grows with s .

5.2 Spin-six nonlinear equation

To illustrate the general results above, we now consider the case where only even s are present, and give the first two ρ_s .

The first coefficients of the fundamental tensor and its inverse read

$$\begin{aligned} q_{\mu\nu}^4 &= \frac{1}{2}\phi_{\mu\nu}^4, & q_{\mu\nu}^6 &= \frac{1}{2}\phi_{\mu\nu}^6 - \frac{1}{2}\phi_4 g_{\mu\nu}^0 - y_{(\mu}\phi_{\nu)}^4; \\ p_4^{\mu\nu} &= -\frac{1}{2}\phi_4^{\mu\nu}, & p_6^{\mu\nu} &= q_4^{\mu\rho}q_{4\rho}{}^\nu - q_6^{\mu\nu} = \frac{1}{4}\phi_4^{\mu\rho}\phi_{4\rho}{}^\nu + \frac{1}{2}g_0^{\mu\nu}\phi_4 + y^{(\mu}\phi_4^{\nu)} - \frac{1}{2}\phi_6^{\mu\nu}. \end{aligned} \quad (5.7)$$

The first spray coefficients are

$$G_4^\mu = \frac{1}{2}(d\phi_4^\mu - \nabla^\mu\phi_4), \quad G_6^\mu = \frac{1}{2}(d\phi_6^\mu - \nabla^\mu\phi_6) - \frac{1}{2}y^\mu d\phi_4 - \frac{1}{2}g_4^{\mu\nu}G_{4\nu}. \quad (5.8)$$

In particular, the $s = 4$ result coincides with the linear result (4.13).

As for the curvature, again the $s = 4$ result coincides with the linear (4.17), (4.18), while for $s = 6$ a quadratic term appears:

$$\begin{aligned} \rho_{4\rho}^\mu &= \nabla_\rho G_4^\mu - \frac{1}{2}d\partial_{y^\rho}G_4^\mu = (\rho_4^L)^\mu{}_\rho, \\ \rho_{6\rho}^\mu &= \nabla_\rho G_6^\mu - \frac{1}{2}d\partial_{y^\rho}G_6^\mu + \frac{1}{2}y_\rho dG_4^\mu - \frac{1}{4}\partial_{y^\rho}G_4^\nu\partial_{y^\nu}G_4^\mu + \frac{1}{2}G_4^\nu\partial_{y^\nu}\partial_{y^\rho}G_4^\mu = (\rho_6^L)^\mu{}_\rho + \mathcal{Q}^\mu{}_\rho(\phi_4), \end{aligned} \quad (5.9)$$

where

$$\mathcal{Q}^\mu{}_\rho(\phi_4) = -\frac{1}{2}\nabla_\rho(\phi_4^{\mu\nu}G_\nu^4) + \frac{1}{4}d(\phi_4^{\mu\nu}{}_\rho G_\nu^4 + \phi_4^{\mu\nu}\partial_{y^\rho}G_\nu^4) - \frac{1}{4}\partial_{y^\rho}G_4^\nu\partial_{y^\nu}G_4^\mu + \frac{1}{2}G_4^\nu\partial_{y^\nu}\partial_{y^\rho}G_4^\mu. \quad (5.10)$$

This \mathcal{Q} now denotes the part that is quadratic in ϕ_4 rather than in G_4^μ , so it contains more terms than the Q in (5.6).

Notice that the $\rho_{s\rho}^\mu$ are not symmetric if we lower the first index with $g_{\mu\nu}^0$ (as we are doing in this section). Indeed the symmetric property advertised in 3.2 holds if we lower the index with the full fundamental Finsler tensor, $\rho_{\mu\rho} \equiv g_{\mu\nu}\rho^\nu{}_\rho$. In our spin expansion, the $s = 4$ term of this object reads $g_{\mu\nu}^0\rho_{4\rho}^\nu + \frac{1}{2}\phi_{\mu\nu}^4 R^\nu{}_{\alpha\rho\beta}y^\alpha y^\beta$; this can indeed be shown to be symmetric using (2.6).¹⁰

Finally, taking the trace:

$$\rho_4 = -\frac{1}{2}\mathcal{F}(\phi_4), \quad \rho_6 = -\frac{1}{2}\mathcal{F}(\phi_6) + \frac{1}{4}(D+2)d^2\phi_4 + \mathcal{Q}(\phi_4). \quad (5.11)$$

This signals a cubic interaction among two $s = 4$ and one $s = 6$ fields. (Such an interaction is also present in Vasiliev theories [31].)

6 Challenges for Finsler dynamics

Given the appearance of the Fronsda operator in Sec. 4, we now investigate whether a Finsler action can indeed propagate degrees of freedom with higher spins. Unfortunately this will give mixed results.

¹⁰For the much more laborious $s = 6$ check, I used the software package xAct.

6.1 Equations of motion

For pure GR, Einstein's equations set to zero the Ricci tensor. This suggests to take for pure Finsler gravity

$$\rho = 0, \tag{6.1}$$

sometimes called Rutz equation [32]. For us this has the advantage that we know it already to be related to the Fronsdal operator. A natural variant is $\rho - kF^2 = 0$, for k a constant, which is similar to pure GR with a cosmological constant.

On the other hand, it was argued in [33, 34] that this cannot be obtained from an action. Another natural possibility is to write the closest possible analogue of the Einstein–Hilbert action for pure GR. ρ also contain y variables, and to obtain an analogue of the Ricci scalar we would like to somehow trace over them. This can be achieved by integrating over them, leading to

$$S = \int d^D x d^{D-1} \hat{y} g \frac{\rho}{F^2}. \tag{6.2}$$

(See [20, 11.3.1] and [35].) The measure is the product of the usual $\sqrt{g}d^D x$ and of $\sqrt{g}d^{D-1} \hat{y}$ along the fiber of the sphere tangent bundle SM . The F^2 is included to make the integrand of degree zero, so that it is defined on the fiber of that bundle. The variation of (6.2) gives

$$\rho = \frac{1}{D+2} F^2 g^{\mu\nu} (\partial_{y^\mu} \partial_{y^\nu} \rho + 2D_\mu dI_\nu - 2dI_\mu dI_\nu + 2\partial_{y^\mu} d^2 I_\nu). \tag{6.3}$$

We will see that (6.1) and (6.3) can be analyzed along similar lines.¹¹

In the rest of this section we will analyze the classical solutions of this theory, perturbatively around Minkowski space. In the similar problem for GR, the equations of motion can be approximated by taking (2.2) for $g_{\mu\nu} = \eta_{\mu\nu}$. Crucially, the infinitesimal diffeomorphisms (2.4) can be used to set $\partial^\nu h_{\mu\nu} - \frac{1}{2}\partial_\mu h = 0$, after which the equation of motion reads simply

$$\partial^2 h_{\mu\nu} = 0. \tag{6.4}$$

So there are no massive excitations. Diffeomorphisms can be further used to restrict the massless ones to $\partial^\mu h_{\mu\nu} = u^\mu h_{\mu\nu} = 0$, with u a timelike vector. These results avoid the presence of modes with the wrong sign in the action, which would be problematic for quantization. (While pure GR is ultimately not sensible quantum mechanically because

¹¹A third possibility was proposed in [36], which is however chosen in such a way that its classical solutions coincide with those of ordinary GR. Thus we will not consider it here.

of its non-renormalizability, it can at least be used as a good effective quantum field theory.)

We thus expect the role of gauge transformations to be crucial. The presence of the Fronsdal operator \mathcal{F} in (4.6), (4.14) seems promising in that respect: \mathcal{F} does have an infinite-dimensional kernel, or in other words it has gauge transformations. In the next subsection we examine whether these can be considered gauge transformations for our Finsler equations of motion.

6.2 Broken Fronsdal gauge transformations

The appearance of the Fronsdal operator in (4.14) might give hope that gauge transformations of the form $\delta\phi_s = d\lambda_{s-1}$ can be introduced in Finsler geometry, at least for a small perturbation around flat space $F^2 = 2\eta_2$ (where (2.28) is valid). It is not clear why the F^2 parameterization (4.8) should be invariant under such a transformation.¹² In other words, the gauge transformations of Finsler geometry are only diffeomorphisms. A priori it might be possible, however, that F^2 changes in a way that does not modify the geometry in a physically relevant way. We would then declare such transformations to be additional gauge transformations. This might perhaps be similar to the notion of projective equivalence mentioned in Sec. 4.1.¹³

However, the term $d^2\phi_{s-2}$ in (4.14) should give one pause. Including the Fronsdal term, the total transformation for a Finsler deformation of Minkowski space reads

$$\delta\rho_s^L = d^3 \left(-\frac{1}{4}\lambda'_{s-1} + \alpha_s\lambda_{s-3} \right). \quad (6.5)$$

One might try to cancel the two terms in the parenthesis against each other. The structure looks similar to that of a Stückelberg mechanism, where a mass term locks together the gauge transformations of two fields of different spin. It is also reminiscent of the compensator fields in [26, 37, 38], where an equation of the type $\mathcal{F}(\phi_s) = d^3\mathcal{H}_s$ appears, with $\delta\mathcal{H}_s = \lambda'_{s-1}$.¹⁴

Unfortunately, this combined gauge transformation is nothing but a particular case of the trivial redundancy (4.9), which is only present when the ϕ_s are not traceless. This

¹² [23, (5.9)] notes that the invariance of proper time $\int d\tau F$ can be approximately restored by simultaneously changing the position $x^\mu(\tau)$ in a velocity-dependent fashion.

¹³The infinitesimal counterpart of (4.1) is $\delta G^\mu = \delta P y^\mu$ with $\delta P = (2g_2)^{-1}d\delta F = (2g_2^3)^{-1/2}d\delta F^2$. From (4.3) we now find $\delta\rho = (1 - D/2)d\delta P$; so ρ is not invariant under a projective equivalence.

¹⁴In the AdS case an extra term is present because of (2.39); we will deal with this at the end. For theories of Vasiliev type, a certain boundary condition for the scalar can lead to a Higgs mechanism whose Goldstone field is a bound state of higher spin fields [39].

is easiest to see when we only transform two spins, $\delta\phi_{s_0} = d\lambda_{s_0-1}$, $\delta\phi_{s_0-2} = d\lambda_{s_0-3}$, and the two only contain the same irreducible spin- $(s_0 - 3)$ representation: $\lambda_{s_0-1}^0 = 0$, $\lambda_{s_0-1}'' = 0$, and $\lambda_{s_0-3}' = 0$. Setting $\delta\rho^L$ to zero gives $\lambda_{s_0-3} = \lambda_{s_0-1}'/(6 - D - 2s)$, which upon taking d reduces to (4.9).

A faster way to proceed is to assemble all the λ_s into a single

$$\Lambda = \sum_{s>1} g_2^{1-s/2} \lambda_{s-1}, \quad (6.6)$$

so that now

$$\delta F^2 = d\Lambda. \quad (6.7)$$

Notice that Λ is homogeneous in y of degree one, since F^2 is of degree two. When we perturb around flat (pseudo-)Riemannian space, $F^2 = 2\eta_2$, and use (4.6), the same steps as in (2.28) give

$$\delta\rho = -\frac{1}{4}d^3\Lambda', \quad \Lambda' = \eta^{\mu\nu}\partial_{y^\mu}\partial_{y^\nu}\Lambda. \quad (6.8)$$

The strategy above now corresponds to demanding $\Lambda' = 0$. This is not simply the sum over the individual λ_{s-1}' , because the factors $g_2^{1-s/2}$ in (6.6) also give a contribution when we take the derivatives with respect to y^μ . (This is the subtlety alluded to above (4.6).) Rather:

$$\Lambda' = \sum_{s>1} g_2^{1-s/2} (\lambda_{s-1}' - 4\alpha_s \lambda_{s-3}). \quad (6.9)$$

The coefficients in this expansion are not necessarily traceless, so to set $\Lambda' = 0$ we need to use the logic leading to (4.10). This gives a relation $\lambda_{s-3}^0 + \lambda_{s-1}^1 + \dots = 0$, which upon using (4.10) gives $\Lambda = 0$. Alternatively, the discussion in Sec. 4.2 tells us that the parameterization (6.6) is redundant unless we impose that the λ_{s-1} are irreducible:

$$\lambda_{s-1}' = 0. \quad (6.10)$$

In other words, this choice can be made without loss of generality. Notice that this is the infamous traceless condition on the Fronsdal gauge transformations, which here appears rather naturally. In view of (6.9), $\Lambda' = 0$ implies $\lambda_{s-1} = 0$ for all s .

An even nicer way of seeing the failure of this strategy is to notice that $\Lambda' = \eta^{\mu\nu}\partial_{y^\mu}\partial_{y^\nu}\Lambda = 0$ is demanding Λ to be a harmonic function of the y , while below (6.6) we observed it to be homogeneous of degree one. These two demands are only met by a linear function $\Lambda = \lambda_1$, which corresponds to $\delta g_2 = d\lambda_1$, the familiar diffeomorphisms. Ignoring this possibility as we did above, we recover that $\Lambda' = 0$ implies $\Lambda = 0$.

We have concluded that we cannot set to zero the parenthesis in (6.5), or Λ' in (6.8). In other words, we have tried to find gauge transformations with an algebraic constraint (similar to the traceless constraint for Fronsdal), and we failed. We could still hope that the full (6.5) can somehow be made zero, by playing with the differential operators. However, d^3 has a finite-dimensional kernel, consisting of quadratic polynomials in the x^μ . Thus we also cannot find gauge transformations this way.¹⁵

We now also analyze this possibility using the parameterization in terms of sum over spins. It is lengthier, but instructive, and also a good warm-up for the next subsection. We earlier concluded that we can always fix $\lambda'_{s-1} = 0$. Moreover we recall from Sec. 4.2 that $\phi'_s = 0$ can similarly be assumed without loss of generality. Since $(d\lambda)' = d\lambda' + 2\partial_\mu\lambda^\mu$, we also conclude

$$\partial_\mu\lambda^\mu_{s-1} = 0. \quad (6.11)$$

Setting $\delta\rho^L = 0$ does not necessarily imply that the coefficients $\delta\rho_s^L$ of its g_2 expansion vanish. The latter are given by (6.5) and are not necessarily traceless; thus their trace parts can mix in the total sum $\sum_{s>2} g_2^{1-s/2} \delta\rho_s^L$. As discussed near (4.10), the series can be equivalently rewritten with coefficients $\sum_{j=0}^\infty \delta\rho_{s+2j,j}^L$ (recalling the notation (2.15)), which are now automatically traceless and can be set to zero. Given (6.10), (6.11) and recalling that we are taking $g_2 = \eta_2$, we compute $(d^3\lambda)' = 6d\partial^2\lambda$, $(d^3\lambda)'' = 0$. So gauge invariance of ρ^L requires

$$\left[\alpha_s d^3\lambda_{s-3} - \frac{6\alpha_{s+2}}{D+2s} d\partial^2\lambda_{s-1} \right]_0 = 0. \quad (6.12)$$

One might hope that the two terms can sum to zero; this would lead to a recursive law on the λ_{s-1} . We now show that this cannot happen. The action of d can be diagonalized after going to Fourier transform. Consider first $p^2 < 0$. We can take $p^\mu = (p^0, 0, \dots, 0)$, so

$$d = -iy^0 p_0, \quad d^\dagger = ip_0 \partial_{y^0}. \quad (6.13)$$

(6.11) gives us $\partial_{y^0}\lambda_{s-1} = 0$. The traceless projection in (6.12) adds a slight complication, which is reviewed in App. A. The two terms in (6.12) can be viewed (forgetting the

¹⁵We can apply a similar argument to projective equivalence to flat space. The variation of (4.5) gives $\delta G_L^\mu = \frac{1}{2}d^2\partial_{y_\mu}\Lambda$. From below (4.1) we find $\delta G^\mu = Py^\mu$. If we ignore the finite-dimensional kernel of d^2 , we are led to postulate $\partial_{y_\mu}\Lambda = y^\mu Q$, with $d^2Q = 2P$. This implies a ‘‘purely radial’’ dependence, $\Lambda = \Lambda(\eta_2)$, and the homogeneity property lets us conclude $\Lambda = \lambda_0\sqrt{\eta_2}$, which would correspond to the gauge transformations of a spin-1 field, had we included it in (4.8). This is compatible with [18, Prop. 3.4.8]; notice however that many other Finsler metrics projectively equivalent to flat space are given in that book, not obtained by gauge transformations of Fronsdal type.

constants) as $(zd)^3\lambda_{s-3}$ and $(zd)\lambda_{s-1}$, with $z\lambda \equiv \lambda_0$ the traceless projector. On our y^0 -independent λ 's, the action of $(zd)^j$ is proportional to multiplication by y_0^j . This means that the two terms in (6.12) have y^0 dependence proportional to y_0^3 and y_0 ; thus the two cannot be equal, unless they are both zero. In the case $p^2 = 0$, the second term in (6.12) is directly zero. Moreover, $zd^3 = (zd)^3$ is invertible, as also shown in App. A. Thus there are no solutions to (6.12).

We can also try a similar analysis on a maximally symmetric space such AdS. Here is what changes. First, covariant derivatives don't commute, and that complicates the trace $(d^3\lambda)'$, which is now $(6d(2\nabla^2 - \Delta) - 2[d, \nabla^2])\lambda_{s-1}$. Second, now $\delta\mathcal{F} \neq 0$ even if $\lambda'_{s-1} = 0$, as we saw back in (2.39). Third, it makes sense to consider now the variant of (6.1) discussed below it, $\rho - kF^2 = 0$. This however can be reabsorbed in (2.39) by redefining Λ . All in all (6.12) is changed to

$$\left[-\frac{1}{2}\Lambda\mu_s^2 d\lambda_{s-1} + \alpha_s d^3\lambda_{s-3} - \frac{\alpha_{s+2}}{D+2s} d(6\nabla^2 + \gamma_s)\lambda_{s-1} \right]_0 = 0. \quad (6.14)$$

where $\gamma_s = \Lambda(6(s-1)(D+s-3) + D+2s-3)/(D-1)$. Now we cannot perform a Fourier transform to analyze the resulting condition; we should work in a suitable eigenbasis for AdS. I have not carried out this analysis, but the problem looks similar enough to (6.12) that at small distances (where Fourier transform can be used at least approximately) it should reduce to the one for flat space.

In this subsection we have concluded that in spite of (4.6), (4.14), the Fronsdal gauge transformations $\delta\phi_s = d\lambda_{s-1}$ don't survive in Finsler geometry except as the trivial redundancies (4.9). It is still possible that there is some other gauge transformation, perhaps obtained by complementing the Fronsdal ones by additional terms involving d^\dagger or other operators. In any case, in the next subsection we will sketch a linearized analysis of the Finsler equations of motion.

6.3 Perturbations

Let us first consider (6.1). We use the power series parameterization (4.8), taking $\phi'_s = 0$. We found the linearized ρ^L in (4.14). Just like for the earlier discussion of gauge transformations, $\rho^L = 0$ does not necessarily imply that the ρ_s^L vanish. As discussed near (4.10), the series can be equivalently rewritten with coefficients $\sum_{j=0}^{\infty} \rho_{s+2j,j}^L$, which are now automatically traceless and can be set to zero. Recalling again (2.14), we use (2.25) to compute

$$(d^2\phi_s)'' = -4(\mathcal{F}\phi_s)' = 8(d^\dagger)^2\phi_s, \quad (\mathcal{F}\phi_s)'' = 0. \quad (6.15)$$

So ρ_s^L only has single and double traces, and $\rho_{s+2j,j}^L = 0$ for $j \geq 3$. This leads to the linearized equation of motion

$$\left[-\frac{1}{2}\mathcal{F}\phi_s + \alpha_s d^2\phi_{s-2} + f_s^3(\mathcal{F}\phi_{s+2})' + f_s^4(d^2\phi_s)' \right]_0 = 0, \quad (6.16)$$

where $f_s^3 = (2(D+2s))^{-1}(1 - 4\alpha_{s+4}/(D+2s+2))$, $f_s^4 = -\alpha_{s+2}/(D+2s)$.

We now turn to (6.3). The linearization of the second term is simply ρ' . Among the terms involving I_μ , we can ignore the quadratic one; the others give

$$\begin{aligned} [g^{\mu\nu}(D_\mu dI_\nu + \partial_{y^\mu} d^2 I_\nu)]^L &= (\nabla_\mu d\partial_{y_\mu} + \partial_{y_\mu} d^2 \partial_{y_\mu}) t d F^2 = (2\nabla_\mu d\partial_{y_\mu} + dd^\dagger + d^2 t) t \delta F^2 = \\ &= \left(\mathcal{F} - \frac{1}{2} t d^2 \right) t \delta F^2. \end{aligned} \quad (6.17)$$

We used $1/2[t, d^2] = \nabla^2 + \nabla_\mu d\partial_{y_\mu} + dd^\dagger$, which in turn follows again from (2.25). Overall the linearized version of (6.3) reads

$$\left(\mathcal{F} - \frac{2}{D+2} g_2 (t\mathcal{F} - 4\mathcal{F}t + 2td^2t) \right) \delta F^2 = 0. \quad (6.18)$$

A lengthier computation gives an equation again of the form (6.16), with more complicated coefficients.

Given this qualitative similarity, from now on we will focus on (6.16). We take the pseudo-Riemannian geometry to be flat, $g_2 = \eta_2$. This simplifies $(d^2\phi_s)' = (2\partial^2 + dd^\dagger)\phi_s$. Schematically the equation is now of the form $[(\partial^2 + dd^\dagger)\phi_s + d^2\phi_{s-2} + (d^\dagger)^2\phi_{s+2}]_0 = 0$. This can be clarified further by going to momentum space as in the previous subsection. The simplest case is $p^2 < 0$, corresponding to massive modes. In GR these are gauged away, as reviewed near (6.4), but for Finsler we haven't found a gauge transformation that can achieve a similar result. We can use (6.13) and the explicit decomposition in App. A, which tackles the traceless projection in (6.16). Each ϕ_s is decomposed under the spatial $SO(D-1)$ as a sum $\sum_k Y_{sk} \hat{\phi}_{s,k}^0$, where $\hat{\phi}_{s,k}^0$ is traceless, degree k and y^0 -independent, while Y_{sk} is a polynomial whose highest-order y_0 term is y_0^{s-k} . The operators zd and zd^\dagger (where z is the traceless projector, $z\phi_s = \phi_{s,0}$) are proportional to the identity on each $Y_{sk} \hat{\phi}_{s,k}^0$.

For concreteness let us look at the lowest degrees in (6.16), keeping only even s . For $s = 2$, the equation reads $[(d^\dagger)^2\phi_4]_0 = 0$. The $\hat{\phi}_{4,4}$ and $\hat{\phi}_{4,3}$ don't appear, while $\hat{\phi}_{4,k \leq 2} = 0$. For $s = 4$, schematically we have $[(d^\dagger)^2\phi_6 + (\partial^2 + dd^\dagger)\phi_4]_0 = 0$. Now $\hat{\phi}_{6,6}$ and $\hat{\phi}_{6,5}$ don't appear, while for $k \leq 4$, $\hat{\phi}_{6,k} \propto (\partial^2 + c_{6,k})\phi_{4,k}$, for $c_{6,k}$ some constants; in particular, $\hat{\phi}_{6,k \leq 2} = 0$. Continuing in this fashion we see that $\hat{\phi}_{s,k \leq 2} = 0$, while the $\hat{\phi}_{s,s}$ and $\hat{\phi}_{s,s-1}$ are free, and determine the $\hat{\phi}_{s,k}$ with $3 \leq k \leq s-2$.

Massive modes would then exist for any mass, and they would depend on many free variables. In particular there would be many ϕ_s not transverse to p^μ . This would create issues when trying to quantize this theory. The analysis for massless modes is more complicated, but gives a similar result.

The result of this perturbative analysis is disappointing, but perhaps also expected. Without a gauge transformation, the Finsler equations of motion would be problematic from a perturbative QFT standpoint.

6.4 Discussion

The appearance of Fronsdal operators in the Finsler Ricci curvature is encouraging. But the associated higher spin gauge transformations are broken.¹⁶ Finsler geometry only has diffeomorphisms as gauge transformations.

A pessimistic conclusion would be that the Fronsdal operator appears only for purely mathematical reasons, driven by its similarity with the spin-two situation.

More optimistically, perhaps some sort of gauge transformation can still be found. While the most natural guess for gauge transformations was excluded in Sec. 6.2, I am not excluding that more exotic possibilities might work. Actually the presence of many free parameters in the previous subsection seems to indicate that some sort of gauge transformation should exist. It should be possible to find operators that have those free parameters as their image. We would also like, however, for such a gauge transformation to have some sort of geometrical interpretation. It would be difficult otherwise to imagine that it can survive beyond the linear order. After all, the possibility of some natural interacting structure is the most interesting possible advantage of using Finsler geometry for higher spins, as illustrated in Sec. 5.

While the Finsler Ricci ρ seems important in view of its connection to the Fronsdal operator, another possibility is that it might have to be modified by additional terms. Even more radically, one might have to modify the central tenets of Finsler geometry altogether, as advocated in [24]. It might be helpful to try and recast in terms of Finsler geometry the existing higher spin theories such as those by Vasiliev [8–12], the simpler alternatives in three dimensions [42, 43], or the more recent self-dual theories [44, 45]. In any case, after finding a geometrical quantity that is invariant under a set of infinite transformations that extends diffeomorphism invariance, we would declare such

¹⁶Holography indicates that higher spin gauge transformations are exactly preserved only in the Vasiliev theory for a particular choice of boundary conditions [15]; even their weak breaking at the quantum level is quite constrained [16, 39–41].

transformations to be additional gauge transformations, identifying different Finsler geometries that are equivalent as far as physics is concerned. Time will tell whether any of this can be achieved.

Acknowledgements

It's a pleasure to thank S. Giombi, C. Hull, E. Minguzzi, C. Pfeifer, A. Sagnotti, A. Zaffaroni for very useful discussions and correspondence. My interest in Finsler geometry was rekindled by an ongoing collaboration with B. De Luca, N. De Ponti, A. Mondino, to whom I am very grateful. I would also like to thank the participants in the 2024 Cook's Branch Workshop, to whom a preliminary version of these findings was presented. I am supported in part by INFN and by MIUR-PRIN contract 2022YZ5BA2.

A More details on the symmetrized derivative

In this appendix we give a more explicit discussion of some properties of the operator $d = y^\mu \nabla_\mu$.

We introduce

$$z\phi = (\phi)_0, \quad (\text{A.1})$$

where the $_0$ denotes the traceless part as in (2.15). We now act with d on the decomposition (2.14), and then we take the traceless part of the result. Since $\phi'_{s,k} = 0$, we have $(d\phi_{s,k})' = 2d^\dagger\phi_{s,k}$, $(d\phi_{s,k})'' = 0$. Then

$$(d\phi_s)_0 = (d\phi_{s,0})_0 = d\phi_{s,0} + 2t_{s+1,1}g_2d^\dagger\phi_{s,0}. \quad (\text{A.2})$$

We can also rewrite this as

$$[z, d]\phi_s = 2t_{s+1,1}g_2d^\dagger z\phi_s. \quad (\text{A.3})$$

In particular it follows

$$[z, d]z\phi_s = 0, \quad (\text{A.4})$$

which in turn implies

$$(zd)^k = zd^k, \quad zdzd^\dagger = zdd^\dagger. \quad (\text{A.5})$$

We now specialize to flat space. Working in Fourier transform, there are two cases to consider: p^μ massive or massless. In the former case, by a Lorentz transformation we can always take the momentum to be purely along time, $p^0 \neq 0$, $p^i = 0$, $i = 1, \dots, D-1$.

Then $d = -iy^0 p_0$, $d^\dagger = ip_0 \partial_{y^0}$. A traceless $\phi_{s,0}$ can be decomposed in $\text{SO}(d-1)$ representations as

$$\phi_{s,0} = \sum_k Y_{sk} \hat{\phi}_{s,k}^0(\vec{y}), \quad Y_{sk} = y_0^{s-k} + a_{1sk} \delta_2 y_0^{s-k-2} + \dots \quad (\text{A.6})$$

where $\hat{\phi}_{s,k}^0$ is traceless and degree k ; the s is kept only as a label. The hat denotes dependence on the y^i alone, $\partial_{y^0} \hat{\phi}_{s,k}^0 = 0$, and the coefficients are explicitly

$$a_{1sk} = \frac{(s-k)(s-k-1)}{D+2k-1}, \quad a_{j sk} = \frac{(s-k-2j+2)(s-k-2j+1)}{j(D+2j+2k-3)} a_{j-1,sk}. \quad (\text{A.7})$$

Here $\delta_2 = \frac{1}{2} \delta_{ij} y^i y^j$, and the coefficients are determined by $(\phi_{s,0})' = 0$. (An expansion similar to (A.6) can be found in [2].) One finds

$$(dY_{sk} \hat{\phi}_{sk}^0)_0 = -ip_0 \frac{D+s+k-2}{D+2s-2} Y_{s+1,k} \hat{\phi}_{s,k}^0, \quad (d^\dagger Y_{sk} \hat{\phi}_{sk}^0)_0 = ip_0 (s-k) Y_{s+1,k} \hat{\phi}_{s,k}^0, \quad (\text{A.8})$$

reproducing for $D=4$ formulas in [2]. Using (A.5) one can also find similar expressions for $(d^k Y_{sk} \hat{\phi}_{sk}^0)_0 = (zd)^k \hat{\phi}_{sk}^0$.

For massless momentum we can take $p^- = 0 = p^i$; now $d = -iy^- p_-$. (A.6) is replaced by a decomposition in terms of $\text{SO}(d-2)$ representations:

$$\phi_{s,0} = \sum_k Z_{j_+ sk} \hat{\phi}_{s,k}^0(\vec{y}), \quad Z_{j_+ sk} = y_+^{j_+} y_-^{s-k-j_+} + b_{1sk} \delta_2 y_+^{j_+-1} y_-^{s-k-j_+-1} + \dots, \quad (\text{A.9})$$

where now δ_2 is the transverse $(D-2)$ -dimensional metric, and the coefficients are given by

$$b_{1sk} = \frac{j_+(s-k-j_+)}{D+2k-2}, \quad b_{j sk} = \frac{(j_+-j)(s-k-j_+-j)}{j(D+2j-4+2k)} b_{j-1,sk}. \quad (\text{A.10})$$

(A.8) is replaced by

$$(dZ_{j_+ sk} \hat{\phi}_{sk}^0)_0 = 2ip_- \frac{D+2k+2j_+}{D+2s} Z_{j_++1, s+1, k} \hat{\phi}_{s,k}^0. \quad (\text{A.11})$$

From (A.8), (A.11) it now follows that d has no kernel.

References

- [1] L. P. S. Singh and C. R. Hagen, ‘‘Lagrangian formulation for arbitrary spin. 1. The boson case,’’ *Phys. Rev. D* **9** (1974) 898–909.
- [2] C. Fronsdal, ‘‘Massless fields with integer spin,’’ *Phys. Rev. D* **18** (1978) 3624.

- [3] A. Sagnotti, “Notes on strings and higher spins,” *J. Phys. A* **46** (2013) 214006, [1112.4285](#).
- [4] D. Sorokin, “Introduction to the classical theory of higher spins,” *AIP Conf. Proc.* **767** (2005), no. 1, 172–202, [hep-th/0405069](#).
- [5] D. Ponomarev, “Basic introduction to higher-spin theories,” *Int. J. Theor. Phys.* **62** (2023), no. 7, 146, [2206.15385](#).
- [6] X. Bekaert, S. Cnockaert, C. Iazeolla, and M. A. Vasiliev, “Nonlinear higher spin theories in various dimensions,” in *1st Solvay Workshop on Higher Spin Gauge Theories*, pp. 132–197. 2004. [hep-th/0503128](#).
- [7] S. Giombi, “Higher spin — CFT duality,” in *Theoretical Advanced Study Institute in Elementary Particle Physics: New Frontiers in Fields and Strings*, pp. 137–214. 2017. [1607.02967](#).
- [8] E. S. Fradkin and M. A. Vasiliev, “Cubic interaction in extended theories of massless higher spin fields,” *Nucl. Phys. B* **291** (1987) 141–171.
- [9] E. S. Fradkin and M. A. Vasiliev, “On the gravitational interaction of massless higher spin fields,” *Phys. Lett. B* **189** (1987) 89–95.
- [10] M. A. Vasiliev, “Consistent equation for interacting gauge fields of all spins in (3+1)-dimensions,” *Phys. Lett. B* **243** (1990) 378–382.
- [11] M. A. Vasiliev, “More on equations of motion for interacting massless fields of all spins in (3 + 1) dimensions,” *Phys. Lett. B* **285** (1992) 225–234.
- [12] M. A. Vasiliev, “Properties of equations of motion of interacting gauge fields of all spins in 3 + 1 dimensions,” *Class. Quant. Grav.* **8** (1991) 1387–1417.
- [13] I. R. Klebanov and A. M. Polyakov, “AdS dual of the critical $O(n)$ vector model,” *Phys. Lett. B* **550** (2002) 213–219, [hep-th/0210114](#).
- [14] S. Giombi and X. Yin, “Higher spins in AdS and twistorial holography,” *JHEP* **04** (2011) 086, [1004.3736](#).
- [15] J. Maldacena and A. Zhiboedov, “Constraining conformal field theories with a higher spin symmetry,” *J. Phys. A* **46** (2013) 214011, [1112.1016](#).

- [16] S. Giombi, S. Minwalla, S. Prakash, S. P. Trivedi, S. R. Wadia, and X. Yin, “Chern–Simons theory with vector fermion matter,” *Eur. Phys. J. C* **72** (2012) 2112, [1110.4386](#).
- [17] C.-M. Chang, S. Minwalla, T. Sharma, and X. Yin, “ABJ triality: from higher spin fields to strings,” *J. Phys. A* **46** (2013) 214009, [1207.4485](#).
- [18] S.-S. Chern and Z. Shen, *Riemann–Finsler geometry*, vol. 6. World Scientific Publishing Company, 2005.
- [19] D. Bao, S.-S. Chern, and Z. Shen, *An introduction to Riemann–Finsler geometry*, vol. 200. Springer Science & Business Media, 2000.
- [20] Y.-B. Shen and Z. Shen, *Introduction to modern Finsler geometry*. World Scientific Publishing Company, 2016.
- [21] C. Lämmerzahl and V. Perlick, “Finsler geometry as a model for relativistic gravity,” *Int. J. Geom. Meth. Mod. Phys.* **15** (2018), no. supp01, 1850166, [1802.10043](#).
- [22] C. Pfeifer, “Finsler spacetime geometry in physics,” *Int. J. Geom. Meth. Mod. Phys.* **16** (2019), no. supp02, 1941004, [1903.10185](#).
- [23] B. de Wit and D. Z. Freedman, “Systematics of higher spin gauge fields,” *Phys. Rev. D* **21** (1980) 358.
- [24] C. M. Hull, “ W geometry,” *Commun. Math. Phys.* **156** (1993) 245–275, [hep-th/9211113](#).
- [25] Z.-Q. Guo, “Higher spin theories from Finsler geometry,” [1301.0787](#).
- [26] D. Francia and A. Sagnotti, “Free geometric equations for higher spins,” *Phys. Lett. B* **543** (2002) 303–310, [hep-th/0207002](#).
- [27] C. Aragone and S. Deser, “Consistency problems of hypergravity,” *Phys. Lett. B* **86** (1979) 161–163.
- [28] L. Tamássy, “Relation between metric spaces and Finsler spaces,” *Differential Geometry and its Applications* **26** (2008), no. 5, 483–494.
- [29] E. Minguzzi, “Light cones in Finsler spacetime,” *Commun. Math. Phys.* **334** (2015), no. 3, 1529–1551, [1403.7060](#).

- [30] M. Abate and G. Patrizio, *Finsler metrics – a global approach: with applications to geometric function theory*. Springer, 2006.
- [31] C. Sleight and M. Taronna, “Higher spin interactions from conformal field theory: The complete cubic couplings,” *Phys. Rev. Lett.* **116** (2016), no. 18, 181602, [1603.00022](#).
- [32] S. F. Rutz, “A Finsler generalisation of Einstein’s vacuum field equations,” *General Relativity and Gravitation* **25** (1993) 1139–1158.
- [33] M. Hohmann, C. Pfeifer, and N. Voicu, “Finsler gravity action from variational completion,” *Phys. Rev. D* **100** (2019), no. 6, 064035, [1812.11161](#).
- [34] M. Hohmann, C. Pfeifer, and N. Voicu, “Mathematical foundations for field theories on Finsler spacetimes,” *J. Math. Phys.* **63** (2022), no. 3, 032503, [2106.14965](#).
- [35] C. Pfeifer and M. N. R. Wohlfarth, “Finsler geometric extension of Einstein gravity,” *Phys. Rev. D* **85** (2012) 064009, [1112.5641](#).
- [36] A. Garcia-Parrado and E. Minguzzi, “An anisotropic gravity theory,” *Gen. Rel. Grav.* **54** (2022), no. 11, 150, [2206.09653](#).
- [37] D. Francia and A. Sagnotti, “On the geometry of higher spin gauge fields,” *Class. Quant. Grav.* **20** (2003) S473–S486, [hep-th/0212185](#).
- [38] P. de Medeiros and C. Hull, “Geometric second order field equations for general tensor gauge fields,” *JHEP* **05** (2003) 019, [hep-th/0303036](#).
- [39] L. Girardello, M. Porrati, and A. Zaffaroni, “3-D interacting CFTs and generalized Higgs phenomenon in higher spin theories on AdS,” *Phys. Lett. B* **561** (2003) 289–293, [hep-th/0212181](#).
- [40] J. Maldacena and A. Zhiboedov, “Constraining conformal field theories with a slightly broken higher spin symmetry,” *Class. Quant. Grav.* **30** (2013) 104003, [1204.3882](#).
- [41] O. Aharony, G. Gur-Ari, and R. Yacoby, “ $d = 3$ bosonic vector models coupled to Chern–Simons gauge theories,” *JHEP* **03** (2012) 037, [1110.4382](#).
- [42] A. Campoleoni, S. Fredenhagen, S. Pfenninger, and S. Theisen, “Asymptotic symmetries of three-dimensional gravity coupled to higher-spin fields,” *JHEP* **11** (2010) 007, [1008.4744](#).

- [43] A. Campoleoni, S. Fredenhagen, and S. Pfenninger, “Asymptotic W-symmetries in three-dimensional higher-spin gauge theories,” *JHEP* **09** (2011) 113, [1107.0290](#).
- [44] E. D. Skvortsov, T. Tran, and M. Tsulaia, “Quantum chiral higher spin gravity,” *Phys. Rev. Lett.* **121** (2018), no. 3, 031601, [1805.00048](#).
- [45] K. Krasnov, E. Skvortsov, and T. Tran, “Actions for self-dual higher spin gravities,” *JHEP* **08** (2021) 076, [2105.12782](#).