



Social Neurobiology of Eating Special Issue

A systematic review of implicit attitudes and their neural correlates in eating behaviour

Alessia Gallucci,^{1,2} Lilia Del Mauro,³ Alberto Pisoni,^{2,3} Leonor J Romero Lauro,^{2,3} and Giulia Mattavelli⁴

¹Ph.D. Program in Neuroscience, School of Medicine and Surgery, University of Milano-Bicocca, Via Cadore, 48–20900, Monza, Italy, ²NeuroMi (Neuroscience Center), University of Milano-Bicocca, Piazza dell’Ateneo Nuovo, 1–20126, Milan, Italy, ³Department of Psychology, University of Milano-Bicocca, Piazza dell’Ateneo Nuovo, 1–20126, Milan, Italy and ⁴NETS, School of Advanced Studies, IUSS, Piazza della Vittoria n.15, 27100, Pavia, Italy

Correspondence should be addressed to Giulia Mattavelli, NETS, School of Advanced Studies, IUSS, Palazzo del Broletto, Piazza della Vittoria 15, 27100 Pavia, Italy. E-mail: giulia.mattavelli@iusspavia.it.

Abstract

An increasing number of studies suggest that implicit attitudes towards food and body shape predict eating behaviour and characterize patients with eating disorders (EDs). However, literature has not been previously analysed; thus, differences between patients with EDs and healthy controls and the level of automaticity of the processes involved in implicit attitudes are still matters of debate. The present systematic review aimed to synthesize current evidence from papers investigating implicit attitudes towards food and body in healthy and ED populations. PubMed, EMBASE (Ovid), PsycINFO, Web of Science and Scopus were systematically screened and 183 studies using different indirect paradigms were included in the qualitative analysis. The majority of studies reported negative attitudes towards overweight/obese body images in healthy and ED samples and weight bias as a diffuse stereotypical evaluation. Implicit food attitudes are consistently reported as valid predictors of eating behaviour. Few studies on the neurobiological correlates showed neurostimulation effects on implicit attitudes, but how the brain automatically processes implicit evaluations remains an open area of research. In conclusion, implicit attitudes are relevant measures of eating behaviour in healthy and clinical settings, although evidence about their neural correlates is limited.

Key words: implicit attitudes; food preference; body image evaluation; weight bias; eating disorders

Introduction

Implicit attitudes are behaviours and judgements driven by automatic evaluations, which are triggered independently from

conscious control (Greenwald *et al.*, 1998). In the context of eating behaviour, implicit attitudes towards food and body images are indices of individual preference for different categories of

Received: 28 February 2020; Revised: 17 September 2020; Accepted: 20 November 2020

© The Author(s) 2021. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

food and affective evaluation of bodies in self and other representations. In particular, there is evidence that implicit food attitudes predict the subsequent actual choice for different types of food (e.g. healthy or unhealthy food) and individuals' eating behaviours with modulatory effects of hunger state, craving and eating disorder (ED) symptomatology (Perugini, 2005; Ellis *et al.*, 2014; Richard *et al.*, 2019). On the other hand, implicit attitudes towards underweight or overweight body shape have been related to internalization of thin ideal, body dissatisfaction, drive for thinness and ED symptoms (e.g. Ahern *et al.*, 2008; Cserjési *et al.*, 2010; Heider *et al.*, 2015). The theoretical framework posits that implicit attitudes rely on associative spontaneous processes (in contrast to propositional processes underlying explicit attitudes) that can be triggered automatically and independently from the overt consideration that a person has on the same association in explicit evaluations (Gawronski and Bodenhausen, 2006). Different models have been proposed accounting for the relationship between implicit and explicit attitudes in predicting behaviour, and studies on eating behaviour reported food choice as an example of spontaneous decisions predicted by implicit evaluations (Perugini, 2005; Conner *et al.*, 2007). In addition, strong affective evaluation of food and body image may represent a key aspect of dysfunctional eating and body dissatisfaction in patients with ED (Spring and Bulik, 2014).

However, previous literature on implicit attitudes and eating behaviour is not always consistent since some studies did not replicate findings on the predictive validity of implicit tasks above explicit measures (Ahern and Hetherington, 2006; Ayres *et al.*, 2012), and there are contrasting results from studies assessing implicit attitudes in patients with different ED diagnoses (Roefs *et al.*, 2005b; Khan and Petróczi, 2015; Smith *et al.*, 2018). Crucially, implicit attitudes are measured by indirect tasks, which assess associations between target stimuli and attributes without directly inquiring participants' beliefs and thus are less influenced by social desirability and strategies of self-presentation compared to explicit questionnaires (De Houwer, 2002). This is particularly relevant in clinical or sub-clinical populations with EDs, which tend to have a low level of therapeutic adherence and to mask symptoms as body dissatisfaction, food restraint or craving to avoid interventions and hospitalization (Halmi, 2013). Interestingly, discrepancy between explicit and implicit measures has been reported in individuals with obesity and predicted disinhibited eating (Goldstein *et al.*, 2014; Cserjési *et al.*, 2016). Moreover, implicit attitude towards body images is a widely used measure of weight bias, i.e. the stigmatizing concept that people with obesity or overweight are lazy and lacking in self-control, which predicts prejudice and misbehaviour (O'Brien *et al.*, 2008; Flint *et al.*, 2016). Notably, weight bias is present also in healthcare professionals and in individuals with overweight with relevant clinical consequences (Anselmi *et al.*, 2013; Phelan *et al.*, 2015a; Tomiyama *et al.*, 2015). Indeed, self-directed weight bias in patients with obesity or binge eating disorder (BED), referred as internalization of weight bias, has been related to higher depressive symptoms, perceived stress and worse overall health (Pearl *et al.*, 2013; Hilbert *et al.*, 2014; Phelan *et al.*, 2015a). Considering healthcare professionals, including eating behaviour specialists, there is evidence of weight bias impact on interpersonal relationship, decision-making on treatments and quality of patients' care, although reciprocal influences between implicit and explicit weight bias of patients and healthcare providers are a recent matter of investigation (Forhan and Salas, 2013; Phelan *et al.*, 2015c).

Crucially, studies investigating implicit attitudes typically use indirect tasks as measures of automatic associations and affective evaluations of target stimuli, although it is still an issue of debate which is the level of automaticity of the processes involved in task execution and which are the underpinning brain mechanisms (De Houwer *et al.*, 2009; Forbes *et al.*, 2012). Results from neuroimaging studies and patients with brain lesions have shown that the Implicit Association Tests (IATs; Greenwald *et al.*, 1998), one of the most diffuse tasks to assess implicit attitudes, recruits structural and functional networks involved in cognitive control and inhibition processes as well as automatic self-representation and semantic representation of concepts, including the dorsolateral and ventromedial prefrontal cortex, the anterior cingulate cortex, the insula and the anterior temporal lobe (Chee *et al.*, 2000; Gozzi *et al.*, 2009; Forbes *et al.*, 2012). Moreover, electrophysiological studies with event-related potentials (ERPs) have reported effects related to congruency of blocks and score at the IAT in early time-windows at 90- to 130-ms post-target onset or at the N200 component, supporting the automaticity of brain responses for implicit associations (Forbes *et al.*, 2012; Healy *et al.*, 2015). Whether these automatic processes are common to implicit attitudes in different contexts, or have specific features related to the stimuli eliciting the bias, remains an open question. Research with neuromodulation techniques can be informative on this aspect, providing evidence of a causal relationship between induced modulation or interference in a target region and changes in implicit attitudes. Indeed, it has been shown that transcranial magnetic stimulation (TMS) applied to a different portion of the prefrontal cortex interfered with IAT on gender stereotype or food, whereas parietal stimulation affected implicit religiousness (Cattaneo *et al.*, 2011; Crescentini *et al.*, 2014; Mattavelli *et al.*, 2015). As stated above, implicit attitudes towards food and body are relevant in predicting eating behaviour; thus, it is of interest to investigate the neural bases of these attitudes as potential targets for therapeutic neuromodulation treatments. The reasons for combining neuromodulation and assessment with indirect tasks are twofold: on one hand, implicit measures can represent a feasible alternative to explicit questionnaires to evaluate patients with EDs with difficulties in recognizing/reporting their symptoms; on the other hand, applying neurostimulation on the neural correlates of automatic processes contributing to the maintenance of ED could boost the effect of therapeutic treatment.

Different paradigms have been proposed to evaluate implicit attitudes. The IAT (Greenwald *et al.*, 1998) measures associations of two opposite target categories (e.g. healthy vs unhealthy food) with opposite valence attributes, asking participants to categorize target stimuli and attributes congruently (e.g. healthy-positive and unhealthy-negative) and incongruently paired (e.g. healthy-negative and unhealthy-positive) in subsequent blocks. The Single Category IAT (SC-IAT; Karpinski and Steinman, 2006) consists in the same paradigm with opposite attributes to be associated with one category of stimuli. The affective priming (AP) task (Fazio *et al.*, 1986) is another frequently used paradigm based on reaction times in categorizing positive and negative targets (words or pictures) preceded by prime stimuli (e.g. food stimuli). Responses to targets are informative of participant's affective evaluation of primes, assuming faster responses when prime and target share the same valence (congruent trials). A modified version of the IAT is the Extrinsic Affective Simon Task (EAST; De Houwer, 2003). In this case, participants are asked to categorize valence words presented in white colour as positive/negative words and target stimuli based on their colour

(e.g. blue/green). The EAST is based on the prediction that participants are faster in categorizing with the positive key coloured words that they consider positive. Other indirect tasks are the Implicit Relational Assessment Procedure (IRAP; Barnes-Holmes et al., 2010), which requires to select between two response options (e.g. true/false) for samples and target stimuli presented together (e.g. 'I am' + 'slim', Heider et al., 2015), and the Affective Misattribution Procedure (AMP; Payne et al., 2005), which asks participants to evaluate Chinese characters as pleasant or unpleasant ignoring the preceding stimuli and the difference in responses provide a measure of how the affect evoked by the first image is misattributed to the Chinese character. All these tasks have been used to assess food and body implicit attitudes in healthy and clinical populations.

In the past decade, the assessment of implicit attitudes related to eating behaviour has become widespread, but a review is missing so far. Since the introduction of the IAT by Greenwald et al. (1998) and its early application to assess weight bias and predict eating behaviour (Teachman and Brownell, 2001; Perugini, 2005), different tasks have been proposed and used in healthy, sub-clinical and clinical populations. Previous reviews focused on cognitive bias with food- and body-related stimuli in Stroop or attentional tasks and reported greater bias in patients with EDs compared to control participants (Johansson et al., 2005; Brooks et al., 2011). However, attentional bias is related to the salience of the stimuli, whereas the indirect tasks used to evaluate implicit attitudes involved affective associations and the automatic evaluation in terms of valence of the stimuli (De Houwer, 2002; Brooks et al., 2011).

This study presents a systematic review of studies published up to May 2019 (i.e. when literature search was conducted) concerning implicit attitudes towards food and body in different areas of research. The aim is to analyse and summarize methodological aspects and findings in healthy and clinical populations with EDs. To this purpose, we will provide readers a comprehensive guide of literature in this field giving particular attention to neuroscientific methodological approaches and recent evidence of neurobiological correlates related to implicit attitudes in eating behaviour.

Method

Design and eligibility criteria

The systematic review was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (Moher et al., 2015; Shamseer et al., 2015). Specific eligibility criteria were considered to systematically select and appraise studies (Grant and Booth, 2009). In particular, we included original studies, while systematic reviews, narrative reviews, meta-analysis, conference proceedings, case reports and conference abstracts were excluded. Only peer-reviewed papers published in English were selected. We were interested in studies on implicit attitudes towards food or body images in healthy populations or in patients with EDs. Thus, studies reporting assessment of implicit attitudes towards different targets were excluded (even if sample with EDs was involved). Similarly, studies assessing unconscious processing of food or body images, but with paradigm not related to implicit attitudes, were excluded.

Search strategy and studies selection

The following terms were combined to collect records from PubMed, EMBASE (Ovid), PsycINFO, Web of Science and Scopus:

'implicit attitudes', 'implicit association', 'affective priming', 'eating disorder', 'anorexia nervosa', 'bulimia nervosa', 'binge eating disorder', 'obesity', 'food preference', 'thin ideal' and 'fat phobia' (the search strategies are shown in Appendix 1). After removing duplicates, the records were independently filtered based on title and abstract by three researchers (G.M., A.G. and L.D.M.). To blind the process, we used Rayyan web-based reviews manager (Ouzzani et al., 2016), which allows to screen the records as 'include', 'exclude' or 'maybe'. When records were eligible for exclusion, specific labels were added to justify the reason of exclusion. At the end of the screening, the blind mode was turned off and conflicts were resolved by consensus. Researchers (G.M., A.G. and L.D.M.) independently reviewed the full text of the records in the 'include' and 'maybe' categories. Conflicting decisions were solved based on researchers' consensus.

Out of 633 screened papers, conflicts or unsure decisions for inclusion were 259 (i.e. the two raters gave different decisions of inclusion, exclusion or maybe). Percentage agreement among raters was computed (as suggested in Kottner and Streiner, 2011), resulting in a value of 59.1%, which is considered a moderate agreement (House et al., 1981; Nurjannah and Siwi, 2017). However, indecisions or disagreement were solved by consensus; thus, at the end of the screening procedure, a unanimous agreement on all papers was reached. A detailed presentation of decision percentages is reported in Supplementary Table S1.

Quality assessment

The Cochrane Collaboration's Risk-of-Bias Tool (Higgins et al., 2011) was used to assess the methodological quality of the studies yielded by the search process. The Cochrane Tool allows to rate the following sources of bias as 'high', 'low', 'unclear' or 'not applicable': random sequence generation, allocation concealment, blinding strategy, incomplete outcome data and selective outcome reporting. Three researchers independently evaluated the quality of the records included in the different sections (G.M. studies on neural correlates and food attitudes, A.G. studies on body image and L.D.M. studies on weight bias) by computing, for each Cochrane item, the percentage of the most frequent rating across the retrieved studies. Risk of bias of papers testing implicit attitudes on both food and body was calculated once.

Results

Studies selection and quality assessment

The systematic search retrieved 2840 papers. After removing duplicates, 633 papers were screened based on the abstract and 343 were excluded. The remaining 290 papers were examined as full-text readings and 107 were excluded. The most common reason for exclusion was lack of consistency with the review topic. In total, 183 studies met our inclusion criteria and were included in our review. Figure 1 summarizes the search procedure.

Results are presented on the basis of the main topics that emerged from studies meeting inclusion criteria. A small number of studies investigated the neurobiological correlates of implicit food or body attitudes in healthy individuals or patients with EDs and are described in a specific section ('Studies on neurobiological correlates'). Other studies were grouped depending on whether implicit measures were used to assess attitudes towards food ('Studies on food attitudes'), body image ('Studies on body image') or weight bias ('Studies on weight




Fig. 1. Flow chart showing the selection process of papers.

bias'). Methodological aspects and main results are reported to capture the contribute of the studies in each specific sub-topic. In particular, papers included in body image or weight bias sections were grouped considering the common focus across the studies on the implicit evaluation of different types of body images and shapes or on the assessment of the stigmatizing (and self-stigmatizing) concept of overweight or obesity (i.e. weight bias).

Regarding the quality evaluation, Table 1 reports the risk of bias assessment according to the Cochrane Collaboration's Risk-of-Bias Tool (Higgins *et al.*, 2011). Overall, considering only the applicable Cochrane items, the food topic had the best quality compared to the others.

Studies on neurobiological correlates

Six studies investigated the neurobiological underpinnings of implicit attitudes toward food or body images (Table 2). Three studies were carried out with healthy participants (Mattavelli *et al.*, 2015; Cazzato *et al.*, 2017; Hall *et al.*, 2018a), two involved patients with EDs (Blechert *et al.*, 2011; Mattavelli *et al.*, 2019) and one study assessed patients with Parkinson's disease (PD) treated with dopaminergic replacement (Terenzi *et al.*, 2018). Four of these studies investigated the possibility to modulate IAT performances with non-invasive brain stimulation techniques applied to different cortical regions (Mattavelli *et al.*,

2015, 2019; Cazzato *et al.*, 2017; Hall *et al.*, 2018a). In particular, the study by Mattavelli *et al.* (2015) applied TMS to the medial prefrontal cortex (mPFC) and left parietal cortex while healthy participants were submitted to three different IATs on food, self and flowers/insects. Results demonstrated the causal role of mPFC in monitoring implicit food attitudes and highlighted the impact of individual variability in modulating the effect of neurostimulation. Differently, neuromodulation of dorsolateral prefrontal cortex (dlPFC) by means of a continuous theta burst protocol (cTBS) did not affect the IAT score in the study by Hall *et al.* (2018a). The other study on healthy participants (Cazzato *et al.*, 2017) applied transcranial direct current stimulation (tDCS) on the extrastriate body area (EBA) in the right and left hemispheres and assessed participants' anti-fat bias with valence-IAT and aesthetic-IAT. tDCS was applied in anodal, cathodal and sham mode on each hemisphere. Only the cathodal stimulation (applied to induce a decrease in cortical excitability) of right EBA showed a significant effect in reducing anti-fat bias, measured by the valence-IAT, in male participants. Differently, female participants did not show a reliable anti-fat bias and their performance was not modulated by tDCS. Only one study investigated neurostimulation effects on IAT in patients with ED (Mattavelli *et al.*, 2019) applying anodal or sham tDCS to the mPFC and right EBA and assessing participants' implicit attitudes toward food, body images and flowers/insects. Results showed that the occipito-temporal

Table 1. Risk of bias evaluation of the papers extracted from the systematic review procedure

Studies topic	Cochrane items						
	Selection bias		Performance bias	Detection bias		Attrition bias	Reporting bias
	Random sequence generation	Allocation concealment	Blinding of participants ^a	Blinding of personnel	Blinding of outcome	Incomplete outcome data	Selective reporting
Neural correlates	67% □	83% □	83% ↓	67% □	67% □	100% ↓	100% ↓
Body image	78% □	74% □	89% ↓	81% □	81% □	59% ↓ 41% ↑	96% ↓
Food	58% ↓ 40% □	50% □ 44% ↓	71% ↓	53% □ 29% ↑	51% □ 46% ↑	96% ↓	97% ↓
Weight bias	74% □	74% □	91% ↓	87% □	74% □	74% ↓	92% ↓

↑ = high risk; ↓ = low risk; ■ = unclear; □ = not applicable;

^ablinding of participants was rated as 'low risk' both when participants were randomly assigned to experimental groups and when participants were blinded about the purposes and procedure of the implicit attitudes' assessment.

Table 2. Details of studies on neurobiological correlates of implicit attitudes toward food or body shape

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Method
Blechert et al., 2011	Study 1	23.1 (4.64)	AP	Primes sentences on body shape and weight concern, self-esteem related word target	EEG (study 1)
	AN (20, f = 20)	26.5 (7.78)			
	BN (20, f = 20)	25.4 (4.80)			
	H (20, f = 20)				
	Study 2				
	H-UNRES (21, f = 21)	23.6 (5.02)			
	H-RES (18, f = 18)	22.6 (3.27)			
Cazzato et al., 2017	H (25, f = 13)	f: 22.08 (0.73) m: 22 (0.6)	IAT	Fat, slim body images	tDCS
Hall et al., 2018a	H (37, f = n.r.)	21.6 (n.r.)	IAT	Flavourful snack, Flavourless foods	cTBS
Mattavelli et al., 2015	H (36, f = 21)	23.25 (2.88)	IAT	Tasty and high-fat, tasteless and low-fat food	TMS
Mattavelli et al., 2019	AN/BN/EDNOS (36, f = 36)	25.53 (8.09)	IAT	Tasty high-fat, tasteless low-fat food; Underweight, overweight body images.	tDCS
	H (36, f = 36)	24.03 (3.48)			
Terenzi et al., 2018	PD + BE (16, f = 8)	67.1 (8.2)	AP	Food, non-food	
	PD (15, f = 7)	64.9 (12.9)			
	H (20, f = 10)	68 (6.4)			

AN = anorexia nervosa; AP = affective priming; BN = bulimia nervosa; cTBS = continuous theta burst protocol; EDNOS = eating disorder not otherwise specified; f = female; H = healthy participants; H-RES = healthy restrained eaters; H-UNRES = healthy unrestrained eaters; IAT = Implicit Association Test; n.r. = not reported; PD = Parkinson's disease; PD + BE = Parkinson's disease with binge eating; tDCS = transcranial direct current stimulation; TMS = transcranial magnetic stimulation.

stimulation increased the implicit preference for high-fat tasty food in patients with ED and both mPFC and EBA stimulation increased reaction times in incongruent trials (i.e. tasty food—negative attributes and tasteless food—positive attributes associations). The effect was specific for food attitudes in patients, whereas no modulatory effects resulted for healthy participants and in the other IATs.

Only one study investigated the electrophysiological correlates of AP measuring the association of self-related targets with body shape and weight primes (Blechert et al., 2011). ERPs were recorded in healthy participants and patients with anorexia nervosa (AN) or bulimia nervosa (BN) during the AP task. Results showed a significant congruent-incongruent difference in the N400 amplitude only in the BN group, suggesting that the stronger association of self-evaluation with body shape and

weight concept in ED is encoded in early stage of brain processes in BN patients.

Finally, Terenzi et al. (2018) used an AP on food to evaluate reward sensitivity in patients with PD treated with dopaminergic medications and BED. These patients are at risk of developing impulse control disorder and represent a neurobiological model for striatal dopaminergic anomalous functioning (Dagher and Robbins, 2009). Patients with PD and BED showed reduced priming effect compared to controls for sweet foods, but no differences in explicit rating on liking and wanting the foods.

Studies on food attitudes

Implicit attitudes toward food were investigated in 75 studies (Table 3). Among these, three studies that considered both food

and body stimuli (Spring and Bulik, 2014; Khan and Petróczi, 2015; Moussally et al., 2015) are presented in the body section and also in details in Table 4. Most of these studies (N = 53) measured implicit attitudes in healthy normal-weight participants using the two-categories IAT (N = 26). Ten studies used the SC-IAT, whereas three studies tested participants both with the two-categories IAT and the SC-IAT (Friese et al., 2008; Houben et al., 2010; Guidetti et al., 2012). Five studies used the AP task, one the semantic priming task (Misener and Libben, 2017), three the AMP (Ellis et al., 2014; Woodward and

Treat, 2015; Woodward et al., 2017) and two the EAST or similar variants (Hoefling and Strack, 2008; Veenstra and de Jong, 2010). Four studies combined more than one task, to investigate the role of different implicit measures in predicting behaviour (Roefs et al., 2005a; Conner et al., 2007; Seibt et al., 2007; Genschow et al., 2017). One study assessed recovered and currently diagnosed patients with AN and healthy controls by means of AMP procedure including high- and low-calorie food as well as thin- and fat-related body images (Spring and Bulick, 2014; see Table 4). Nineteen studies evaluated food implicit attitudes in

Table 3. Details of studies on implicit attitudes toward food

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Healthy participants					
Ackermann and Palmer, 2014	H (101, f = 62)	n.r.	IAT	Healthy food, fast food	-
Adams et al., 2017	H (143, f = 134)	22.92 (n.r.)	SC-IAT	Chocolate	Response inhibition training
Alblas et al., 2018	H (125, f = 79)	20.17 (1.88)	IAT	Chocolate, fruits	Evaluative conditioning (health video-games)
Alkozei et al., 2018	H (17, f = 8)	24.53 (4.2)	IAT	High calorie, low-calorie food	Sleep deprivation
Ashby and Stritzke, 2013	H (132, f = 96)	18.67 (3.3)	IAT	High-fat, low-fat food	Priming procedure
Ayres et al., 2012	H (127, f = 97)	18.75 (3.1)	IAT	Chocolate, fruits	-
	H (80, f = 67)	22.1 (7.3)			
Becker et al., 2015	H (98, f = 80)	23 (4.6)	IAT	healthy, unhealthy food	Approach-avoidance training
	H (52, f = 52)	20.47 (2.34)			
	H (104, f = 104)	20.77 (2.94)			
Bongers et al., 2013	H (103, f = 103)	21.94 (3.59)	AP	Healthy, unhealthy food, objects	Emotion induction during milk shake consumption
		20.34 (2.24) and 19.83 (1.83) per condition	SC-IAT	Chocolate, objects	
Conner et al., 2007	H (112, f = 112)	20.34 (2.24) and 19.83 (1.83) per condition	SC-IAT	High-caloric food	-
	H (123, f = 76)	23.7 (5.8)	IAT	Sweets, shapes	
Coricelli et al., 2019	H (104, f = 84)	23.2 (4.9)	EAST	Sweets	-
			IAT	Chocolate, fruits	
			IAT	Natural food, utensils	
Ellis et al., 2014	H (45, f = 22)	23.22 (3.12)	IAT	Transformed food, utensils	-
			AMP	Fruits	
Eschenbeck et al., 2016	H (107, f = 59)	27 (11.9)	IAT	Healthy, unhealthy food	High and low distraction situations
Friese et al., 2008	H (90, f = 79)	21.56 (3.83)	IAT	Chocolate, fruits	Cognitive load or self-regulatory resources during food choice
	H (88, f = 88)	23.19 (4.29)	IAT	Chocolate, fruits	
	69 (f = 69)	22.48 (4.59)	SC-IAT	Chips	
Genschow et al., 2017	48 (f = 0)	24.11 (4.41)	SC-IAT	Beer	Affective or cognitive focus before food choice
	H (91, f = 73)	22.80 (5)	EAT	Chocolate, fruits	
Guidetti et al., 2012	H (85, f = 75)	18.86 (0.97)	AP	Fruits	-
			MT		
			SC-IAT		
Hensels and Baines, 2016	H (95, f = 70)	24.88 (6.16)	IAT	Sweet, savoury snacks	Evaluative conditioning
			IAT	Healthy, unhealthy food	
Hoefling and Strack, 2008	H (66, f = 37)	23.9 (3.7)	EAST	High-calorie, low-calorie food	Food deprived or satiated between groups
Hollands et al., 2011	H (134, f = 101)	24.2 (n.r.)	IAT	Fruits, snacks	Evaluative conditioning
			IAT	Snacks, fruits	
Houben et al., 2010	H (59, f = 59)	31.44 (9.61)	SC-IAT	Snacks	-
			SC-IAT	High-caloric food	
Houben et al., 2012	H (63, f = 63)	34.71 (13.28)	SC-IAT	moderate-caloric food	-
			SC-IAT	Low caloric food	
Kakoschke et al., 2017	H (112, f = 112)	24.88 (5.92)	SC-IAT	Unhealthy food	Approach-avoidance training combined with Go/No-go training

Table 3. (Continued)

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Kraus and Piqueras-Fiszman, 2016	H (108, f = 68)	22.9 (2.9)	IAT-RF	Sandwich, sweets	Hunger state
Lamote et al., 2004	H (26, f = 17) H (29, f = 25)	36.96 (4.03) 19.55 (1.84)	AP AP	Individually positive and negative rated food	(Study 2) Strong or moderate prime condition
Lebens et al., 2011	H (85, f = 85)	E 34.14 (12.87) C 34.23 (13.49)	SC-IAT	Snack	Evaluative conditioning
Mai and Hoffman, 2015	H (203, f = 81)	23.0 (2.58)	IAT	Energy-dense, energy-poor food	-
Mayer et al., 2008	H (50, f = 50)	n.r. (undergraduate students)	IAT	High-caloric food, low-caloric food	Disgusting odour
Maas et al., 2017	H (83, f = 72)	n.r. (undergraduate students)	SC-IAT	High-fat food	-
McConnell et al., 2011	H (56, f = 35)	19.25 (n.r.)	IAT	Apple, chocolate	-
Misener and Libben, 2017	H (115, f = 115)	19.9 (1.47)	SP	ED, non-ED prime-target relation	-
Nederkoon et al., 2010	H (51, f = 51)	19.5 (2.2)	SC-IAT	Snack food	-
Papies et al., 2009	H (91, f = 91) H (100, f = 65)	20.5 (2.2) 20.07 (2.60)	AP AP	High-fat palatable, neutral, unpalatable food	-
Pavlovic et al., 2016	H (89, f = 73) H (40, f = 40)	21.7 (n.r.) 2.4 (n.r.)	IAT	Fruits, sweets	-
Pechey et al., 2015	H (732, f = 371)	51 (14)	SC-IAT	Fruit Chees Cake	-
Perugini, 2005	H (113, f = 62)	25.1 (6.8)	IAT	Snacks, fruits	-
Richard et al., 2019	H (66, f = 66)	20.3 (2.34)	SC-IAT	Chocolate	-
Roefs et al., 2005a	H restrained (32, f = 32) H unrestrained (37, f = 37) H restrained (26, f = 26) H unrestrained (30, f = 30)	19.5 (2) 19.5 (1.8) 19.6 (2.2) 19.3 (1.1)	AP EAST	High-fat, low-fat food High-fat palatable, high-fat Unpalatable, low-fat palatable, low-fat unpalatable food	-
Ragunathan et al., 2006	H (131, f = n.r.)	n.r. (undergraduate students)	IAT	Healthy, unhealthy food	-
Seibt et al., 2007	H (29, f = n.r.) H (74, f = n.r.)	n.r. n.r. (undergraduate students)	IAT EAST	Food, sport Food, flower, non-words	Deprived vs satiated between participant assessment
Stafford and Scheffler, 2008	H (30, f = 24)	30.4 (7.4)	IAT	Food, furniture	Pre-lunch vs post-lunch between participants assessment
Sato et al., 2016	H (34, f = 16)	23.3 (4.5)	AP	Fast food, Japanese diet food	-
Sato et al., 2017	H hungry (28, f = 13) H satiated (28, f = 13)	22.9 (4.4) 23.4 (4.7)	AP	Fast food, Japanese diet food	Hungry/satiated between groups conditions
Schakel et al., 2018	H (120, f = 97)	21.3 (2.4)	IAT	Healthy, unhealthy food	Gamified approach avoidance training and verbal suggestions
Songa and Russo, 2018	H (60, f = 38) H (80, f = 44)	32 (n.r.) 28 (n.r.)	IAT	High-energy, low-energy food	-
Storr and Sparks, 2016	H (183, f = 183)	27.29 (10.84)	IAT	high-calorie, Low-calorie food	Self-affirmation and ego-depletion
Trendel and Werle, 2016	H (283, f = 163) H (142, f = 74)	20.6 (n.r.) 20.6 (n.r.)	IAT	Chocolate, apple	Cognitive load during food choice
Van Dessel et al., 2018	H (389, f = 219) H (184, f = 59)	34 (13) 20 (2)	IAT	Healthy, unhealthy food	Approach avoidance training

Table 3. (Continued)

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Veenstra and de Jong, 2010	H (55, f = 55)	n.r. (undergraduate students)	AST-voice	High-fat, low-fat food	-
Wang et al., 2011	H (100, f = 100)	21.3 (2.4)	SC-IAT	Chocolate	Ego-depletion
Werle et al., 2013	H (94, f = 50)	19.6 (n.r.)	IAT	Healthy, unhealthy food	-
Werntz et al., 2016	H (10115, f = 76.6%) ¹	27.6 (11.4)	IAT	High-fat, low-fat food	-
Woodward et al., 2015	H (238, f = 238)	18.98 (1.72)	AMP	Food	-
Woodward et al., 2017	H (238, f = 238)	19.08 (1.40)	AMP	Food	-
Yen et al., 2010	PMDD (60, f = 60)	23.03 (2.45)	IAT	High-sweet-fat, high-salted-fat food	-
	H (59, f = 59)	22.70 (2.37)			
Overweight/Obesity					
Alabduljader et al., 2018	HW (33, f = 28) OB (20, f = 14)	38.85 (11.36) 40.25 (9.51)	IAT	Unhealthy, healthy food	-
Craeynest et al., 2005	HW (38, f = 21) OB (38, f = 21)	13.53 (2.52) 13.69 (2.63)	EAST	Healthy, unhealthy food; Sedentary, moderate intense, high intense physical activity	-
Craeynest et al., 2006	HW (39, f = 22) OB (39, f = 23)	14.00 (2.40) 14.12 (2.43)	IAT	Fat, non-fat food; Exercise, inactive child	-
Craeynest et al., 2007	HW (40, f = 29) OW (40, f = 29 =	14.83 (0.64) 14.83 (0.75)	IAT	Palatable food, hobby Palatable healthy, palatable unhealthy food	-
Craeynest et al., 2008a	OB (19, f = 12)	12.79 (2.68)	EAST	Healthy, unhealthy food; Sedentary, moderate intense, high intense physical activity	12-month multi-component inpatient programme
Craeynest et al., 2008b	HW (29, f = 16) OW (29, f = 17) HW (29, f = 14) OB (29, f = 17)	14.34 (1.11) 14.59 (1.27) 13.07 (2.09) 13.21 (2.11)	IAT	Fat, lean food	-
Cserjesi et al., 2016	HW (15, f = 7) OB (15, f = 7)	38.4 (9.5) 37.8 (9.5)	AP	Small, medium, large portion of a typical fast food	-
Czyzewska and Graham, 2008	UW (9, f = 9) HW (51, f = 51) OW (12, f = 12) OB (11, f = 11)	21.11 (2.80) 21.61 (3.97) 21.75 (2.30) 24.36 (3.96)	AP	High-calorie non-sweet, high-calorie sweet, low-calorie food, food-related items	-
Ferentzi et al., 2018	OB (129, f = 62)	48 (9.45)	SC-IAT	Unhealthy food	Four sessions of approach-avoidance training
Goldstein et al., 2014	HW/OW (95, f = 95)	19.87 (2.16)	SC-IAT	Chocolate	-
Kemps and Tiggeman, 2015	HW (56, f = 56) OB (56, f = 56)	44.95 (11.82)	IAT	Food (healthy and unhealthy items), non-food	-
McKenna et al., 2016	HW (24, f = 11) OB (25, f = 15) HW (32, f = 16) OB (25, f = 15) HW (42, f = 20) OB (32, f = 16)	20.92 (3.11) 41.57 (8.84) 20.81 (1.36) 41.36 (9.91) 20.76 (3.67) 36.88 (9.76)	IRAP	Healthy, unhealthy	Food restriction
Roefs and Jansen, 2002	HW (31, f = 25) OB (30, f = 24)	40.5 (14.4) 46.3 (14.8)	IAT	High-fat, low-fat food	-
Roefs et al., 2005b	HW (27, f = 27) AN (19, f = 19) HW (27, f = 27) OB (27, f = 27)	20.4 (5.8) 20.6 (6.3) 36.6 (8.7) 36.5 (8.8)	AP	High-fat palatable, low-fat palatable, high-fat unpalatable, low-fat unpalatable food	-
Roefs et al., 2006	HW (26, f = 26) OB (33, f = 33) HW (29, f = 29) OB (27, f = 27)	41.8 (8.1) 41.7 (6.9) 35.8 (10.0) 36.4 (9.9)	AP	High-fat palatable foods, low-fat palatable foods, high-fat unpalatable foods, low-fat unpalatable foods	(i) Focus attention on palatability or health aspects of food (ii) Food craving induction

Table 3. (Continued)

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Sartor et al., 2011	HW (22, f = 15) OW/OB (11, f = 4) Study 2 H (12, f = 7)	23.1 (2.9) 22.2 (1.6) 26 (6)	IAT	Sweet, non-sweet foods and drink	(Study 2) 4 weeks soft drink supplementation
Verbeken et al., 2018	OB (44, f = 23)	12.58 (1.43)	SC-IAT	Unhealthy food	10 sessions of approach-avoidance training
Warschburger et al., 2018	OW/OB (59, f = 33)	13.23 (1.93)	SC-IAT	High energy snacks	Six sessions of approach-avoidance training
Werrij et al., 2009	HW (19, f = 19) OW/OB (24, f = 24) HW (29, f = 29) OB (28, f = 28)	41 (12) 42 (10.3) 37 (8.5) 37 (8.9)	SP	Palatable food, neutral words, disinhibition words Palatable food, neutral words, restraint words	-

AMP = affective misattribution procedure; AP = affective priming; AST = Affective Simon Task; C = control group; E = experimental group; EAST = extrinsic affective Simon test; ED = eating disorder; H = healthy participants; HW = healthy weight; IAT = Implicit Association Test; IAT-RF = recording free variant of IAT; IRAP = Implicit Relational Assessment Procedure; MT = manikin task; n.r. = not reported; OB = obese; OW = overweight; PMDD = premenstrual dysphoric disorder; SC-IAT = Single category IAT; SP = semantic priming; UW = underweight; f = female. ¹ part of a larger web-based data collection on other mental health domains.

Table 4. Details of studies on implicit attitudes toward body image

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Healthy participants					
Ahern et al., 2006	H (86, f = 86)	Range 16–25 years	IAT	Fat and thin silhouettes	-
Ahern et al., 2008	H (105, f = 105)	18.01 (0.15)	IAT	Underweight and normal-weight body images	-
Benas and Gibb, 2011	H (202, f = 202)	18.93 (1.17)	IAT	Fat and thin related words	-
Elran-Barak and Bar-Anan, 2018	H (66 799, f = 47.265)	27.88 (11.9)	IAT	Fat and thin silhouettes	-
Expósito et al., 2015	H (34, f = 34)	23.35 (1.35)	IRAP	Fat and thin women silhouettes	-
Glashouwer et al., 2018	H (72, f = 72)	20.05 (1.41)	RRT	Actual or ideal body image related sentences	-
Heider et al., 2015	H (68, f = 68)	18.72 (2.12)	IRAP	Actual or ideal body image related sentences	-
Heider et al., 2018	H (68, f = 68)	18.72 (2.12)	RRT	Actual or ideal body image related sentences	-
Juarascio et al., 2011	H (80, f = 80)	18.24 (0.68)	IRAP	Fat and thin silhouettes	-
Lydecker et al., 2018	H (657, f = 488)	36.81 (7.96)	IATs	Thin or fat children related words	-
Marini, 2017	H (4.806, f = 3.253/67.7%)	28.32 (11.65)	IAT	Underweight, normal-weight, overweight and obese body images	-
Martijn et al., 2013	Study 1 H (66, f = 66) Study 2 H (39, f = 39)	18.94 (0.99) 21.51 (1.83)	IAT Explicit ratings	Images of supermodels vs normal-sized models	Evaluative conditioning task
Matharu et al., 2014	H (129, f = 91)	25.2 (2.9)	IAT	Fat and thin silhouettes	Standard lecture vs medical humanities intervention
Moussally et al., 2015*	H (121, f = 121)	23.97 (4.80)	AMP	Thin and overweight women picture; permitted and forbidden food	-
Ritzert et al., 2016	H (99, f = 75)	18.6 (1.0)	IRAP	Attractiveness vs disgust/fear related sentences	-
Robstad et al., 2018	H (30, f = 30)	46.53 (n.r.)	IATs	Words related to the concept of obese vs normal body shape	-

Table 4. (Continued)

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Sabin et al., 2015	H (75, f = 41)	48 (-)	IAT	Fat and thin silhouettes	-
Watts et al., 2008	Study 1 H (87, f = 87)	18.91 (2.65) 19.28 (1.83)	AP	Images of body parts and body shapes	-
	Study 2 H (72, f = 72)				
EDs					
Anselmi et al., 2011	OB (43, f = 25) H (331, f = 214)	34.30 (10.01) 26.16 (8.07)	IAT	Faces of thin and fat people	-
Cserjési et al., 2010	AN (35, f = 35) H (35, f = 35)	19.61 (3.42) 20.27 (3.93)	AP	Ultra-thin, average-size, overweight bodies	-
Izquierdo et al., 2019	FP-AN (39, f = 39) NFP-AN (13, f = 13) Low-weight ARFID (10, f = 10) H (32, f = 32)	19.5 (2.5) 18.3 (3.6) 15.2 (3.6) 17.4 (3.1)	IATs	EDI-3 statements; underweight or normal-weight body pictures	-
Keng and Ang, 2019	AN/BN/BED (81, f = 81)	Range 18–55 years	IAT	Body-related words	Mindful breathing exercise vs resting condition
Khan and Petróczi, 2015*	AN (14, f = 14) BN (24, f = 24) EDNOS (16, f = 16). At-risk ED (41, f = 41) H (23, f = 23)	27.00 (12.95) 21.54 (6.19) 20.19 (5.69) 21.56 (7.83) 23.00 (4.97)	IBI-BIAT PBI-BIAT FP-AAT	Normal vs thin silhouettes; High-fat foods vs low-fat foods	-
Parling et al., 2012	AN (12, f = 12) Sub-AN (5, f = 5) H (17, f = 17)	25.33 (6.0) 23.40 (4.4) 24.76 (5.5)	IRAP	'Pro-thin/anti-fat' or 'anti-thin/pro-fat' words	-
Smith et al., 2014	AN (30, f = 30) H (29, f = 29)	20.03 (2.83) 18.93 (1.41)	AP	'Beauty' words (e.g. glamorous, beautiful and attractive), 'ugly' words (e.g. hideous, gross and disgusting), 'neutral' word (e.g. mailbox, cloud and desktop), 'positive' words (e.g. happy, cheerful and elated) in a lexical decision task	Emaciation prime (photos of emaciated looking women) vs thin prime (photos of thin women)
Smith et al., 2018	AN/sub-AN (32, f = 31) BN/sub-BN (37, f = 36) Other ED (23, f = 21) H (85, f = 39)	28.34 (7.88) 32.00 (12.93) 40.35 (12.00) 36.38 (10.75)	AMP	ED-symptom images (emaciation, binge eating, hard exercise and vomiting); body stimuli images (average weight female bodies); eating stimuli images (women eating and words related to eating, e.g. dining, snacking and eating)	-
Spring and Bulik, 2014*	AN (9, f = 9) Recovered AN (14, f = 14) H (29, f = 29)	21.4 (5.79)	AMP	High/low calorie food images and fatness/thinness related images	-

AN = anorexia nervosa; AMP = Affective Misattribution Procedure; AP = affective priming; ARFID = avoidant/restrictive food intake disorder; BED = binge eating disorder; BN = bulimia nervosa; ED = eating disorder; EDNOS = eating disorder not otherwise specified; f = female; FP-AAT = Food Preference Approach Avoidance Task; FP-AN = fat-phobic anorexia nervosa; H = healthy participants; IAT = Implicit Association Test; IBI-BIAT = Ideal Body Image Brief Implicit Association Test; IRAP = Implicit Relational Assessment Procedure; NFP-AN = non-fat phobic anorexia nervosa; n.r. = not reported; OB = obese; OW = overweight; PBI-BIAT = Personal Self Identification Body Image Brief Implicit Association Test; RRT = Relational Responding Task. *Studies measuring implicit attitudes on both body and food (data on food implicit preferences are presented in the food section).

participants with overweight and obesity. With this population, seven studies employed the two-categories IAT, four studies the SC-IAT, three studies the AP and one a semantic priming paradigm (Werrij et al., 2009). The EAST was used in two studies (Craeynest et al., 2005, 2008a) and the IRAP in one (McKenna et al., 2016). The study by Roefs et al. (2005) reported two separate experiments which compared performance at the AP task between patients with AN vs control participants and people with obesity vs control participants, respectively. Concerning the type of stimuli, 47 out of 75 studies used images of food in the implicit tasks, whereas 27 studies employed words of food representing different types of target categories and a study (Becker et al., 2015) used both words or images in different tasks (see Table 3).

Studies investigating food attitudes in healthy normal-weight participants could be grouped on the basis of three main objectives: (i) to assess implicit and explicit measures as predictors of eating behaviour; (ii) to measure differences in implicit food attitudes between individuals or groups with different features in eating behaviour and their relationship with ED symptoms; (iii) to evaluate the impact of manipulations regarding the context, the hunger state or the type of stimuli on implicit food attitudes. Similarly, studies including participants with obesity or overweight aimed at (i) exploring between-group differences in implicit food attitudes or (ii) assessing the effect of manipulations and training programmes on eating behaviour and attitudes.

Implicit attitudes as predictors of eating behaviour. A first study by Perugini (2005) measured implicit and explicit attitudes toward snacks and fruits to test their validity to predict the following behaviour in food choice. Results are discussed as supporting a double dissociation model of the impact of implicit and explicit attitudes on behaviour, since the IAT was significantly related to spontaneous food choice whereas explicit questionnaire was significantly related to self-reported behaviour. Following studies supported this model (Pavlović et al., 2016; Songa and Russo, 2018) and showed the moderating effect of ED symptoms (Ellis et al., 2014) and of individual differences in habit in food behaviour and need for cognition (Conner et al., 2007) on the relative predictive impact of explicit and implicit measures for the subsequent food choice. In line with this evidence, the study by McConnell et al. (2011) reported that both IAT and explicit preference predicted the actual affective experience for eating, but affective forecasting errors were predicted only by the implicit measure suggesting the role of unconscious evaluation in blind affective predictions. Moreover, implicit attitudes have been shown to predict everyday chocolate consumption with interactive patterns of relationship with hunger and craving (Richard et al., 2019). Differently, other studies did not show incremental validity of implicit over explicit measures in predicting food choice, eating behaviour or body mass index (BMI) (Ayres et al., 2012; Ackermann and Palmer, 2014; Woodward and Treat, 2015; Maas et al., 2017; Woodward et al., 2017)

Seven studies investigated how the relationship between implicit and explicit measures and food choice behaviour was influenced by different conditions of cognitive load or self-regulatory resources (Friesen et al., 2008; Eschenbeck et al., 2016; Trendel and Werle, 2016; Wang et al., 2016), positive or negative priming manipulation (Ashby and Stritzke, 2013), emotional induction (Bongers et al., 2013) or manipulation of cognitive or affective focus (Genschow et al., 2017). Six of these studies consistently reported the predictive validity of implicit

measures on the following choice behaviour and also found significant effects of the different manipulations (Friesen et al., 2008; Ashby and Stritzke, 2013; Bongers et al., 2013; Eschenbeck et al., 2016; Trendel and Werle, 2016; Wang et al., 2016). Similarly, Nederkoorn et al. (2010) showed that implicit preference for snack food interacted with response inhibition capacity in predicting weight change over a year. In contrast, the most recent study by Genschow et al. (2017) did not replicate the significant effect of different implicit measures [EAST, AP and manikin task (MT)] in predicting the actual behaviour, neither found significant impact of conditions focusing participants' attention on cognitive or affective aspects.

Another line of research on food perception demonstrated the automaticity of unhealthy-tasty food association and in turn its relationship with body fat (Raghunathan et al., 2006; Mai and Hoffmann, 2015), although the opposite healthy-tasty food association has been reported in a French sample, suggesting intercultural differences in food evaluation (Werle et al., 2013). Moreover, a relevant role of food type has been suggested by Coricelli et al. (2019), who showed different implicit attitudes related to restrictive eating habits and healthiness explicit evaluation, for natural and transformed food matched for calories content. Finally, one study did not assess the actual behaviour, but investigated food attitudes transmission in social contexts testing dyads composed by students with a parent or a friend: results revealed that students' implicit, but not explicit, attitudes toward fruits were associated with those of parents, whereas explicit, but not implicit, attitudes toward snacks were associated with those of friends (Guidetti et al., 2012).

Individual differences and relation with ED symptoms. Five studies compared groups of restrained vs unrestrained eaters using implicit paradigms on high-fat and low-fat foods (Roefs et al., 2005a; Papies et al., 2009; Houben et al., 2010, 2012; Veenstra and de Jong, 2010). Different results are reported depending on stimuli and tasks. Indeed, positive association with high-fat palatable food in both restrained and unrestrained groups is reported in studies using Simon paradigm tasks and AP (Roefs et al., 2005a; Veenstra and de Jong, 2010), although stronger AP in unrestrained eaters is reported by Papies et al. (2009). No positive associations with high-caloric snacks and no group differences emerged using IAT with fruit and snack targets (Study 1 in Houben et al., 2001); in contrast, positive associations with snacks resulted using SC-IAT and this positivity was larger in restrained eaters (Study 2 in Houben et al., 2001). Restrained eaters also showed more positive associations than unrestrained eaters with palatable food in SC-IAT, independently from the caloric food density (Houben et al., 2012).

One study compared women with premenstrual dysphoric disorder (PMDD) and controls assessed with an IAT on high-sweet-fat and high-salted-fat food (Yen et al., 2010). Changes in appetite, overeating and increased food craving are some of PMDD diagnostic criteria; accordingly, the study reported higher implicit positive responses to high-sweet-fat food and higher craving response to food in PMDD group and in women in luteal phase, supporting differences in women's appetite related to menstrual cycle, the relevance of eating assessment in PMDD and the feasibility of implicit measure to evaluate these aspects (Yen et al., 2010).

Four studies investigated the relationship between implicit food preferences and individual variables (Pechey et al., 2015; Sato et al., 2016; Werntz et al., 2016; Misener and Libben, 2017). Sato et al. (2016) showed significant AP effect in evaluating faces

subliminally primed by food or mosaic images, with no differences between high-fat and low-fat food and reported that this effect correlated with individual external eating. In line with this, a semantic priming paradigm showed that priming effect did not differ between ED-related word pairs and non-ED-related word pairs across the whole sample of female students, but higher scores on ED questionnaires and body dissatisfaction were associated with increased priming for ED-related word pairs as compared to non-ED-related word pairs (Misener and Libben, 2017). Moreover, data from website studies showed that implicit preferences for fruit measured with a SC-IAT were significantly related to socioeconomic variables and gender (Pechey et al., 2015), and IAT with high-fat vs low-fat foods and shameful vs acceptable attributes predicted symptoms and concerns related to EDs with stronger high-fat food to shameful association in participants reporting higher score in a EDs questionnaire (Werntz et al., 2016).

Finally, contrasting results come from studies measuring both food and body attitudes: correlations between thin preference and positive evaluation of permitted and forbidden foods are reported by Moussally et al. (2015), in line with results from Spring and Bulik (2014) of higher negative implicit affect toward high-fat food and overweight body images in patients with AN compared to healthy controls; in contrast, Khan and Petróczi (2015) showed that body implicit attitude discriminated between ED and control individuals whereas no differences emerged for food attitudes.

Manipulation effects on implicit attitudes. One study investigated the AP effect manipulating the strength of the positive or negative food prime stimuli, which were individually selected to be extremely or moderately related to positive and negative affects (Lamote et al., 2004). Results showed that the significant AP effect was not moderated by the evaluative extremity of the prime.

Five studies measured implicit food attitudes in participants with different hunger states (Seibt et al., 2007; Hoefling and Strack, 2008; Stafford and Scheffler, 2008; Kraus and Piqueras-Fiszman, 2016; Sato et al., 2017). Administering both motivational and evaluation IATs, Kraus and Piqueras-Fiszman (2016) highlighted that hunger state affected the first, but not the latter implicit attitudes measure. Other studies reported that participants in high vs low hunger state showed increased food preference in an IAT with food and non-food stimuli (Seibt et al., 2007; Stafford and Scheffler, 2008) and increased food preference in supraliminal and subliminal AP tasks with food and mosaic images (Sato et al., 2017). Similarly, hunger state increased the positive association with both high- and low-calorie foods compared to control stimuli in an EAST paradigm (Hoefling and Strack, 2008). This last study compared groups of restrained and unrestrained eaters and reported the same effect of hunger state manipulation in both groups, although restrained eaters had larger positive automatic associations with high-calorie food (Hoefling and Strack, 2008).

One study investigated the effect of sleep restriction on IAT with high- and low-calorie food and reported that food attitudes were not affected by sleep condition in the whole sample, whereas a significant sex by condition interaction revealed that males had greater association of low-calorie food with positive attributes than females in rest condition; no sex difference appeared in sleep restricted condition (Alkozei et al., 2018).

Eleven studies investigated the impact of different conditioning procedures on IAT: evaluative conditioning (EC; Hollands

et al., 2011; Lebens et al., 2011; Hensels and Baines, 2016; Alblas et al., 2018), approach-avoidance training (Becker et al., 2015; Kakoschke et al., 2017; Schakel et al., 2018; Van Dessel et al., 2018), response inhibition training (Adams et al., 2017), ego-depletion and self-affirmation procedures (Storr and Sparks, 2016) and disgusting odour exposure (Mayer et al., 2008). These studies used between groups designs comparing participants randomly assigned to different experimental procedures. EC procedures paired images of unhealthy foods with images of potential adverse health consequences, negative facial expressions or body shapes (Hollands et al., 2011; Lebens et al., 2011; Hensels and Baines, 2016) or used video games to strengthen positive associations with fruits and negative associations with chocolate snacks (Alblas et al., 2018). Similarly, video games or computer tasks were used to train participants to approach healthy food and avoid unhealthy food (Bleckert et al., 2015; Kakoschke et al., 2017; Schakel et al., 2018; Van Dessel et al., 2018). Most of studies assessing implicit attitudes following experimental trainings or EC reported healthier food preference in experimental groups, who received manipulations aimed at increasing positive association to healthy food (Lebens et al., 2011; Hensels and Baines, 2016; Kakoschke et al., 2017; Schakel et al., 2018; Van Dessel et al., 2018). Differently, a study investigating the effect of a response inhibition training on implicit food attitudes reported the absence of manipulation-related modulations (Adams et al., 2017), and one study reported that approach-avoidance training did not affect implicit preferences (Bleckert et al., 2015). Other studies administered the IAT in pre- and post-assessment: Hollands et al. (2011) reported significant interaction between baseline IAT and intervention, with EC reducing preference for snacks vs fruits in individuals with stronger preference at baseline; Alblas et al. (2018) showed that preference for fruit vs chocolate decreased in the control group, but not in the EC group. Disgusting odours spread in the room while participants performed IAT with high- and low-calorie foods did not affect IAT score (Mayer et al., 2008). Finally, Storr and Sparks (2016) reported stronger positive associations with high-calorie foods and negative associations with low-calorie foods in unrestrained compared to restrained eaters, but ego-depletion and self-affirmation procedures did not differently affect restrained and unrestrained eaters in implicit food attitudes.

Group differences in obesity and overweight. Different studies compared healthy-weight participants and participants with obesity for their implicit attitudes towards unhealthy-fat vs healthy-low-fat foods with different types of tasks (see Table 3). Seven of these papers consistently reported more positive attitudes towards healthy food both in healthy-weight adult participants and adult participants with obesity (Roefs and Jansen, 2002; Roefs et al., 2005b; Alabduljader et al., 2018) as well as in children with obesity and healthy-weight (Craeynest et al., 2005, 2006, 2007, 2008b), with no differences between groups in implicit measures. The absence of group differences was also reported in a study showing that both women with obesity and healthy-weight associated high-fat food to restraint concepts (Werrij et al., 2009). On the other hand, participants with obesity responded slower to the high-fat/positive combination than controls in an IAT (Roefs and Jansen, 2002) and showed more positive implicit attitude towards both healthy and unhealthy food in an EAST paradigm (Craeynest et al., 2005). Five other studies reported group differences, with participants with obesity showing greater approach bias to food in a IAT with food

and non-food categories and approach vs avoidance attributes (Kemps and Tiggemann, 2015), higher implicit preference for large fast food portion in an AP paradigm (Cserjesi et al., 2016) and higher preference for sweet than non-sweet food in an IAT (Sartor et al., 2011). Moreover, women with obesity showed significantly more negative attitudes to high-calorie sweet foods and positive attitudes to high-calorie savoury foods compared to healthy- and over-weight participants, with inconsistent results from explicit preference (Czyzewska and Graham, 2008). In line with this result, discrepancy between IAT score and explicit attitudes for chocolate predicted disinhibited eating in healthy- and over-weight participants in a following study (Goldstein et al., 2014).

Effects of manipulations and training on implicit attitudes in obesity. One study reported that manipulation of focus of attention or craving induction affected AP of participants with obesity or lean participants in the same direction with no significant differences between groups (Roefs et al., 2006). Differently, McKenna et al. (2016) reported that participants with obesity and normal-weight participants differed in the automatic responses to healthy and unhealthy food in an IRAP on hunger state and that the groups were differently affected by manipulation of food restriction prior to the assessment. The same paper reported the absence of group differences on IRAP measuring automatic food wanting (McKenna et al., 2016). Five studies used implicit measures as outcome measures following treatments for individuals with obesity (Craeynest et al., 2008a; Sartor et al., 2011; Ferentzi et al., 2018; Verbeken et al., 2018; Warschburger et al., 2018). Craeynest et al. (2008a) reported that children with obesity reduced positive attitude towards healthy and unhealthy food in post-treatment EAST. The other studies showed that IAT score was not affected by intervention of soft drink supplementation (Sartor et al., 2011) and approach avoidant training in adults and children with obesity (Ferentzi et al., 2018; Warschburger et al., 2018; Verbeken et al., 2018).

Studies on body image

Our search retrieved 27 studies about implicit attitudes toward body image (Table 4). Studies using body images to measure weight bias or to investigate neural correlates involved in body representation are reported in the weight bias and neurobiological section, respectively. Eighteen studies involved healthy normal-weight participants, whereas nine studies included samples with EDs. Considering the first groups, most of the studies recruited participants from general population ($N = 14$), whereas the other four studies involved special categories: parents of children with obesity (Lydecker et al., 2006), nurses (Robstad et al., 2018), medical students (Matharu et al., 2014) and clinicians (Sabin et al., 2015). Nine out of 18 studies measured implicit attitudes using the two-categories IAT, whereas in one study participants underwent a multiple-categories IAT (Marini et al., 2017). Two studies (Lydecker et al., 2006; Robstad et al., 2018) combined two different IATs to test both implicit valence attitudes (by using valence related words, i.e. good/bad) and stereotypes (by using judgement-related words, i.e. stupid/smart or lazy/motivated). Four studies used the IRAP (Exposito et al., 2015; Juarascio et al., 2011; Heider et al., 2015; Ritzert et al., 2016), two the Relational Responding Task (Glashouwer et al., 2018; Heider et al., 2018), one the AMP (Moussally et al., 2015) and one the AP (Watts et al., 2008). Concerning the type of stimuli, 10 out of 18 studies compared thin and fat

body images ($N = 7$) or related words ($N = 3$). Two studies used underweight and normal-weight silhouettes (Ahern et al., 2008; Martijn et al., 2013), one study tested implicit attitudes toward underweight, normal-weight and overweight/obese categories (Marini et al., 2017), whereas four studies used sentences to elicit implicit attitudes toward actual and ideal body images (Heider et al., 2015 2018; Glashouwer et al., 2018) or attractiveness vs disgust/fear toward being thin or fat (Ritzert et al., 2016). One study considered images of body parts and of body shape (Watts et al., 2008). The impact of experimental manipulations on implicit attitudes toward body images was explored in two studies (Matharu et al., 2014; Martijn et al., 2013).

Looking at results, 7 out of 18 papers reported negative implicit attitude towards overweight/obese body images (Ahern et al., 2006; Lydecker et al., 2006; Robstad et al., 2018; Elran-Barak and Bar-Anan, 2018; Sabin et al., 2015; Moussally et al., 2015; Watts et al., 2008), when the stimuli were presented both for a short and a long period of time, likely reflecting more automatic or more controlled responses, respectively (Watts et al., 2008). In one study, the self-thin attractive bias was stronger than the self-fat attractive bias (Ritzert et al., 2016). Two studies showed stronger implicit preferences for normal-weight than underweight or overweight/obese body images (Ahern et al., 2008; Marini, 2017), whereas one study did not report a defined bias towards body image as participants showed similar pro-slim and pro-fat implicit attitudes (Exposito et al., 2015). The experimental manipulations were demonstrated to effectively decrease the positive implicit attitude towards underweight people (Martijn et al., 2013) or the negative implicit attitude towards fat people (Matharu et al., 2014). Four studies showed that body implicit attitudes could predict disordered eating, body image dissatisfaction and changes in weight (Juarascio et al., 2011; Heider et al., 2015 2018; Glashouwer et al., 2018). In particular, a study reported that individuals faster in categorizing own actual, but not ideal, body image as fat showed higher body dissatisfaction compared to individuals who represented their actual body image as slim (Glashouwer et al., 2018). One study showed that eating-relevant implicit associations were valid variables to test the negative effects of stereotype (e.g. weight-related peer teasing; Benas and Gibb, 2011). Most of the studies investigating body attitudes in healthy normal-weight participants reported no correlation between implicit and explicit measures. In contrast, four studies showed pro-slim/anti-fat attitudes at both implicit and explicit levels (Robstad et al., 2018; Elran-Barak and Bar-Anan, 2018; Sabin et al., 2015; Benas and Gibb, 2011), and such correlation was confirmed by Matharu et al. (2014) at the baseline measure, but not after experimental manipulation. Three studies did not investigate the correspondence between implicit and explicit attitudes (Watts et al., 2008; Martijn et al., 2013; Moussally et al., 2015). Among the two out of 18 studies testing the predictive value of implicit and explicit attitudes on the behavioural measures, one study reported a negative correlation between implicit and explicit anti-fat bias and the possibility of helping patients with obesity (Robstad et al., 2018). The other study showed that explicit, but not implicit, actual and ideal body image predicted food selection, caloric intake and restraint eating (Glashouwer et al., 2018).

Considering the studies that tested implicit attitudes toward body image in samples of individuals with EDs, most of the studies ($N = 7$) evaluated body implicit attitudes in patients with AN and BN vs healthy controls. One study involved participants with obesity (Anselmi et al., 2011), while another study considered patients with AN, BN and BED (Keng and Ang, 2019). Four studies measured implicit attitudes using the two-categories

IAT (Izquierdo *et al.*, 2019; Anselmi, 2011; Keng and Ang, 2019; Khan and Petr czi, 2015). Among these, Izquierdo *et al.* (2019) added a questionnaire-based IAT using items from the Eating Disorders Inventory (Garner, 2004), whereas Khan and Petr czi (2015) tested both the ideal body image and the personal internalized body image. Two studies used the AP (Cserj si *et al.*, 2010; Smith *et al.*, 2014), other two the AMP (Spring and Bulik, 2014; Smith *et al.*, 2018) and one study applied the Relational Responding Task (Parling *et al.*, 2012). Regarding the type of stimuli, three studies measured implicit attitudes towards underweight or normal-weight silhouettes (Khan and Petr czi, 2015; Izquierdo *et al.*, 2019) or towards thin- or fat-related images (Spring and Bulik, 2014); one study compared thin and fat face images (Anselmi *et al.*, 2011). Parling *et al.* (2012) used words to elicit body shape and weight concerns, while Keng and Ang (2019) used body-related and -unrelated words. The other three studies used ultra-thin, average-size, overweight body images (Cserj si *et al.*, 2010), ED-symptom and body images (Smith *et al.*, 2018) and beauty-related words in a lexical decision task (Smith *et al.*, 2014). Records testing whether an experimental manipulation affected body implicit attitudes showed that exposure to primes of emaciated bodies increased pro-thin bias in women with AN (Smith *et al.*, 2014), while exposure to mindfulness exercises did not alter implicit body dissatisfaction of patients with ED compared to resting condition (Keng and Ang, 2019).

Three studies reported that patients with AN and BN showed pro-thin implicit bias compared to healthy subjects (Izquierdo *et al.*, 2019; Smith *et al.*, 2018; Khan and Petr czi, 2015); similar results were shown by Anselmi *et al.* (2011) in a group of patients with obesity, compared to healthy controls. Conversely, Spring and Bulik (2014) reported that patients with AN, compared to recovered AN individuals and healthy controls, showed negative implicit affect towards overweight stimuli, but not an automatic attraction to thinness. Both pro-thin and anti-fat bias in patients with AN were reported by Parling *et al.* (2012), whereas only anti-fat bias emerged in Cserj si *et al.* (2010), and healthy samples of

both studies were characterized by pro-thin preference, but not by anti-fat preference.

Concerning the correlation between implicit and explicit body attitudes, four studies out of nine reported mixed results (Parling *et al.*, 2012; Khan and Petr czi, 2015; Cserj si *et al.*, 2010; Spring and Bulik, 2014). While one study showed consistent null effect of mindfulness on both implicit and explicit body dissatisfaction (Keng and Ang, 2019), the other four records did not investigate possible correlation between implicit and explicit measures. Among the records on ED samples, only one study reported the predictive value of the implicit attitudes on the ED behaviours (Smith *et al.*, 2018).

Studies on weight bias

Weight bias towards overweight and individuals with obesity has been investigated in 78 studies, using both explicit and implicit measures of preference for 'fat' or 'thin' attributes (Table 5). In order to measure weight stigma, the majority of these studies ($N = 66$) used the two-categories IAT. A minority of studies used either the SC-IAT (Lynagh *et al.*, 2015; Aweidah *et al.*, 2016), the AMP (Pryor *et al.*, 2013; Skinner *et al.*, 2017; Karsay and Schmuck, 2017) or the IRAP (Baker *et al.*, 2017), whereas two studies (Roddy *et al.*, 2010, 2011) paired the two-categories IAT with the IRAP and four papers employed the AP (Degner and Wentura, 2009; Brochu *et al.*, 2011; Glock *et al.*, 2016; Rudolph and Hilbert, 2017). Concerning the type of stimuli employed in the implicit tasks, 27 out of 78 studies used images of thin and fat people, two studies (Anselmi *et al.*, 2013; Penney *et al.*, 2013) used morphed faces of normal-weight and overweight individuals and four studies employed pictures of average-weight and overweight children. Forty-five studies employed words representing different types of target categories whereas one study (Teachman *et al.*, 2003) used both images and words as target categories in different IATs. Regarding the samples, five studies (Carels *et al.*, 2009a, 2009b, 2010, 2011, 2014) recruited adults with overweight

Table 5. Details of studies on weight bias

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Agerstr�m <i>et al.</i> , 2007	Study 2A: Students (88, $f = 58$) Study 2B: Employers (166, $f = 82$)	22.25 (5.73) 43.19 (10.45)	IAT	Obese/normal weight	-
Agerstr�m and Rooth, 2011	Managers (153, $f = 77$)	40.0 (10.41)	IAT	Obese/normal-weight	Manipulation of weight status of job candidates
Anselmi <i>et al.</i> , 2013	UW (53) HW (331) OW (83) OB (43) $f = 327$ of the entire sample	27.06 (8.99)	IAT	Fat/thin people	-
Aweidah <i>et al.</i> , 2016	Diagnostic radiography clinical educators (37, $f = 23$)	n.r.	SC-IAT	Fat	-
Baker <i>et al.</i> , 2017	Five classes of medical students	No demographic data were collected	IRAP	Overweight/obese/slim/thin	-
Bissell and Hays, 2011	H (601, $f = 331$)	n.r.	IAT	Overweight/thin children	Exposure to images of overweight/thin children
Brauhardt <i>et al.</i> , 2014	BED (26, $f = 21$) OB (26, $f = 21$) HW (26, $f = 21$)	34.77 (10.29) 35.19 (11.08) 34.65 (10.70)	Weight Bias-IAT; Self-Esteem-IAT	Thin/fat Self/other	-

Table 5. (Continued)

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Brewis and Wutich, 2012	H from Paraguay (200, f = 200) H from the USA (66, f = 48)	38.9 (13.4) 23.68 (11.2)	IAT	Fat/thin people	-
Brewis et al., 2016	Study 1: HW (116) OW (61) OB (27) f = 103 of the entire sample	23 (n.r.)	IAT	Slim/Fat	-
Brochu and Morrison, 2007	UW (2) AW (75) OW and OB (17) f = 61 of the entire sample	20.11 (4.0)	IAT	Average-weight/overweight people	-
Brochu et al., 2011	University students (80, f = 53)	19.74 (5.05)	AP	Over-weight/normal-weight	-
Carels et al., 2009a	OW and OB (58, f = 52)	47.6 (10.3)	IAT	Fat/thin people	Weight loss intervention
Carels et al., 2009b	OW and OB (58, f = 52)	n.r.	IAT	Fat/thin people	Weight loss intervention
Carels et al., 2010	OW and OB (54, f = 44)	48.7 (11.7)	IAT	Fat/thin people	Weight loss intervention
Carels et al., 2011	OW and OB (53, f = 41)	47.15 (14.1)	Weight Bias-IAT; Self-Esteem-IAT	Fat/thin people Self/other	Weight loss intervention
Carels et al., 2013	OW/OB (42, f = 30) following a weight loss intervention OW/OB (47, f = 38) before a weight loss intervention	46.9 (13.0) 53.7 (13.2)	Stereotype congruent IAT; Stereotype incongruent IAT	Fat/thin people	-
Carels et al., 2014	OW/OB (44, f = 37)	53.2 (13.6)	IAT	Obese/thin people	Weight loss intervention
Cazzato and Makris, 2019	HW (18, f = 11) OW (18, f = 12)	25.28 (0.99) 24 (0.89)	IAT	Fat/slim	-
Chambliss et al., 2004	Undergraduate H (136, f = 57) Graduate in exercise science H (110, f = 53)	Men = 22.7 (2.9) Women = 21.4 (2.9) Men = 25.79 (4.8) Women = 23.5 (2.4)	Weight-attitude IAT; Weight-stereotype IAT	Fat/thin people Fat/thin people	-
Degner and Wentura, 2009	Study 1 (47, f = 24) UW (8) NW (31) OW (8) Study 2 (50, f = 28) UW (5) NW (36) OW (9) Study 3 (65, f = 41) UW (7) NW (39) OW (10) Study 4 (55, f = 55) UW (4) NW (50) OW (1)	25 (n.r.) 21 (n.r.) 23 (n.r.) 47 (n.r.)	AP	Normal-weight/overweight individuals	-
Dimmock et al., 2009	H fitness centre employees (70, f = 40)	27 (9.53)	IAT	Overweight/thin people; Overweight/thin exerciser	-
Domoff et al., 2012	Psychology students (64, f = 42)	20.0 (2.9)	IAT	Fat/thin people	Exposition to a 40-min weight-loss reality shows

Table 5. (Continued)

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Flint et al., 2013	H (28, f = 11)	22.43 (3.59)	IAT	Fat/thin	Counter-conditioning using positive images of obese members of the general public and images of obese celebrities
Flint et al., 2015	H (2380, f = 1767)	27.71 (1.03)	IAT	Fat/thin	-
Flint et al., 2016	Employees (181, f = 74)	38.25 (8.99)	IAT	Fat/thin	Manipulation of weight status of job candidates
Fontana et al., 2013	Physical education teachers (47, f = 28)	37.07 (13.22)	IAT	Fat/thin people	-
Fontana et al., 2017	Professors in physical education departments (94, f = 47)	47.83 (12.18)	IAT	Fat/thin people	-
Gapinski et al., 2006	Undergraduate students (108, f = 108)	19.06 (1.01)	IAT	Fat/thin people	Exposition to a media-based videos intervention
Geier et al., 2003	H (59, f = 59)	19.1 (0.23)	Four IATs	Fat/thin	Exposition to a 'before and after' diet advertisement
Geller and Watkins, 2018	First year medical students H cohort	No demographic data were collected	Weight-attitude IAT; Weight-stereotype IAT	Fat/thin people Fat/thin people	Ethics group session
Glock et al., 2016	Teachers (51, f = 48)	21.12 (6.15)	AP	Words reflecting obesity/thinness	-
Grover et al., 2002	HW (42, f = 22) OW (41, f = 20)	HW men 36.3 (12.7) OW men 32.2 (11.4) HW women 27.8 (11.2) OW women 35.1 (13.8)	Weight-attitude IAT; Weight-identity IAT; gender attitude IAT; gender identity IAT; self-attitude IAT	Light/heavy Self/other Female/male Self/other Self/other	-
Gumble and Carels, 2012	Psychology students (85, f = 47)	19.9 (3.7)	Weight-bias IAT; Weight-identity IAT; Self-esteem IAT; Body image IAT	Fat/thin Self/Other Self/Other Self/Other	-
Halvorson et al., 2019	Hospital attendants (10, f = 7) Pediatric residents (10, f = 7) Pediatric nurses (8, f = 8) Patients (12, f = 7) Parents/caregivers (12, f = 10)	34 (n.r.) 29.0 (n.r.) 31.5 (n.r.) 15 (n.r.) n.r.	IAT	Thin/obese people	-
Hand et al., 2017	Students (302, f = 218)	16 (n.r.)	IAT	n.r.	-
Hart et al., 2016	H African American (207, f = 207) H Non-Hispanic Whites (310, f = 310)	34.45 (10.83) 32.42 (11.17)	Revised-IAT	Overweight/ underweight, obese/underweight, overweight/obese	-

Table 5. (Continued)

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Hilbert and Meyre, 2016	Study 1: University students (128; f = 79) Study 2: Volunteers (128, f = 79)	23.01 (3.24) 35.31 (12.54)	IAT IAT	Thin/fat Thin/fat	Educational inter- vention about the multifactorial aetiology of obesity
Hinman et al., 2015	Students (117, f = 92)	19.3 (1.5)	Stereotype congruent IAT; Stereotype incongruent IAT	Fat/thin Fat/thin	-
Hutchison and Müller, 2018	Children (84, f = 43)	5.89 (1.11)	IAT	Fat/thin children	-
Jiang et al., 2017	104 Asian students UW (23, f = 23) HW (67, f = 67) OW (9, f = 9) OB (5, f = 5)	21.6 (3.3)	IAT	Fat/thin	-
Karsay and Schmuck, 2017	Adolescent (353, f = 176)	17.34 (1.09)	AMP	Obese/non-obese individuals	Exposition to a reality TV show 'The Biggest Loser Teens'
Lund et al., 2018	General practitioners (240, f = 86)	18.5 (10.2)	IAT	Fat/normal weight	-
Lynagh et al., 2015	H enrolled in the health and physical educational curricula for trainee teachers (62, f = 30) H nonspecialist trainee teachers (177, f = 104)	Three age groups: 18–20 years (Special- ist = 38, nonspecial- ist = 106) 21–23 years (Special- ist = 14, nonspecial- ist = 24) >24 years (Special- ist = 7, nonspecial- ist = 16)	SC-IAT	Fat	-
Marini et al., 2013	Volunteers from 71 nations (338.121, f = 238.357)	27.8 (10.64)	IAT	Thin/overweight people	-
Miller et al., 2013	Medical students (310, f = 132)	<25 years = 131 (n.r.) 25–28 years = 145 (n.r.) >28 years = 29 (n.r.)	IAT	Fat/thin	-
O'Brien et al., 2007a	Study1 H (227, f = 140) Study2 H (134, f = 99)	19.98 (2.91) 20.09 (4.199)	IAT	Fat/thin people	-
O'Brien et al., 2007b	H (344, f = 230) First (122) and third (58) year of physical education degree programme; First (95) and third (69) year of psychology programme	First year psychology 18.49 (0.75); First year physical education 18.68 (1.3); Third year psychology 21.8 (4.1); Third year physical education 21.6 (1.9)	Weight-attitude IAT; Weight- stereotype IAT	Fat/thin people Fat/thin people	-
O'Brien et al., 2008	H (104, f = 82)	20.35 (5.04)	Weight-attitude IAT; Weight- stereotype IAT	Fat/thin people Fat/thin people	Manipulation of weight status of job candidates
O'Brien et al., 2010	H (159, f = 135)	20.29 (2.32)	Weight-attitude IAT; Weight- stereotype IAT	Fat/thin people Fat/thin people	Tutorial classes

Table 5. (Continued)

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Penney and Laws, 2013	Students (186, $f = 121$)	19.7 (3.9)	IAT	Thin/obese faces	-
Phelan et al., 2014	Medical students (4,732, $f = 2363$)	23.9 (n.r.)	IAT	Fat/thin people	-
Phelan et al., 2015a	Medical students (4687, $f = 2,344$) HW (3378) UW (163) OW (922) OB (224)	23.9 (2.6)	IAT;	Fat/thin people	-
Phelan et al., 2015b	Medical students (1795, $f = 917$)	$N = 575$ 19–22 years $N = 456$ 23 years $N = 473$ 24–25 years $N = 281$ >26 years	IAT	Fat/thin people	-
Pryor et al., 2013	H (100, $f = 60$)	19.62 (1.42)	AMP	-	Cyberball game
Robertson and Vohra, 2008	Fitness professionals (57, $f = 25$)	29–30 (10.239)	IAT	Fat/thin people	-
Robinson et al., 2014	Health and non-health students (479)	26.2 (7.6)	IAT	Fat/thin people	-
Roddy et al., 2010	Psychology students (80, $f = 58$)	21.1 (3.4)	IAT IRAP	Overweight/average-weight people Overweight/average-weight people	-
Roddy et al., 2011	Students (78, $f = 5$)	20.25 (3.67)	IAT IRAP	Overweight/average-weight people Overweight/average-weight people	-
Rudolph and Hilbert, 2017	Individuals from the community (144)	n.r.	AP	Normal-weight/obese full body pictures	Exposure to health messages
Rukavina et al., 2010	Kinesiology pre-professionals (78, $f = 26$)	21.63 (1.49)	Weight-stereotype IAT	Fat/thin	Classroom and service learning components
Russell-Mayhew et al., 2015	Preservice teachers (30, $f = 25$)	32	IAT	n.r.	Exposure to a professional workshop
Sabin et al., 2012	All test takers (359,261, $f = 220$) Medical doctors (2284, $f = 1285$)	26 (10.7) 33 (12.5)	IAT	Overweight/thin people	-
Schwartz et al., 2003	Professionals engaged in research and/or clinical management of obesity (389, $f = 198$)	n.r.	Weight-attitude IAT; Weight-stereotype IAT	Fat/thin people Fat/thin people	-
Schwartz et al., 2006	UW (128) HW (1756) OW (899) OB (899) EOB (600) $f = \sim 3.555$ of the entire sample (approximately 95% of respondents completed demographic information)	34.6 (n.r.)	Obesity attitude IAT; Obesity stereotype IAT	Fat/thin people Fat/thin people	-

Table 5. (Continued)

Studies	Sample (N)	Mean age (SD)	Implicit measure	Target categories	Manipulation
Scrivano et al., 2017	Students (166)	20.48 (2.31)	IAT	Obese/thin individuals	Exposure to an education about the uncontrollable causes of obesity
Skinner et al., 2017	Children (114, f = 56)	10	AMP	Average-weight/overweight children	-
Solbes and Enesco, 2010	H (120, f = 60)	40 H 6.9 (n.r.) 40 H 8.9 (n.r.) 40 H 10.8 (n.r.)	IAT (child-oriented version)	Fat/thin children	-
Swift et al., 2013	H: CG (21, f = 18) IG (22, f = 18)	21.2 (0.8) 24.6 (7.2)	Weight-attitude IAT; Weight-stereotype IAT	Fat/thin Fat/thin	Anti-stigma films
Teachman and brownell, 2001	Health care specialist (84, f = 24)	48 (9.81)	Weight-attitude IAT; Weight-stereotype IAT	Fat/thin people Fat/thin people	-
Teachman et al., 2003	Study 1: H (144, f = 78) Study 2A: H (90, f = 90) Study 2B: H (63, f = 32)	35 (13.99) 21 (3.87) 42 (16.51)	Weight-attitude IAT; Weight-stereotype IAT Weight-attitude IAT; Weight-stereotype IAT Weight-stereotype IATs	Fat/thin people Fat/thin people Fat/thin people Overweight/underweight people Fat/thin people	Providing information about genetic causes of obesity Written stories of weight discrimination Story of a severe weight discrimination
Thomas et al., 2007	Study 2: Children (94, f = 49)	5	IAT	Fat/thin people	-
Tomiyama et al., 2015	Obesity specialists (232, f = 134)	42.52 (12.82)	Weight-attitude IAT; Weight-stereotype IAT	Fat/thin people Fat/thin people	-
Vallis et al., 2007	Healthcare providers (78, f = 69)	37.39 (10.72)	IAT	Fat/thin people	-
Venturini et al., 2006	Students (45, f = 31)	n.r.	IAT	Normal-weight/fat woman	-
Waller et al., 2012	Nursing H (45, f = 39) Psychology H (45, f = 35)	24.7 (n.r.) 25.60 (n.r.)	IAT	Normal-weight/overweight man and women; normal-weight/overweight man and women in a medical setting	-
Wang et al., 2004	Study 1: HW (64, f = 60) Study 2: OW (48, f = 33)	43.1 (9.4) 48.9 (9.2)	IAT IAT	Fat/thin people Fat/thin people	-
Weinsten et al., 2008	Students (50, f = 35)	n.r.	IAT	Obese/thin	-
Wijayatunga et al., 2019	H (67, f = 64) CG (34, f = 22) IG (33, f = 21)	21.76 (1.43) 21.5 (1.11) 22.03 (1.67)	IAT	Obese/thin people	Learning about uncontrollable cause of obesity and about weight bias

AMP = Affective Misattribution Procedure; AP = affective priming; AW = average weight; BED = binge eating disorder; CG = control group; EOB = extremely obese; f = female; H = healthy participants; HW = healthy weight; IAT = Implicit Association Test; IG = intervention group; IRAP = Implicit Relational Assessment Procedure; n.r. = not reported; OB = obese; OW = overweight; SC-IAT = Single category IAT; UW = underweight.

or obesity engaged in a behavioural weight-loss treatment. Five studies (Solbes and Enesco, 2010; Bissell and Hays, 2010; Hutchison et al., 2018; Skinner et al., 2017; Thomas et al., 2007) involved healthy children, whereas 13 studies involved adult professionals treating obesity. Twenty-six studies analysed the weight bias in healthy university students, whereas two studies (Agerström and Rooth, 2011; Flint et al., 2016) focused on samples of managers and employees. Eight studies included normal-weight, under-weight, over-weight participants or individuals with obesity (Wang et al., 2004; Schwartz et al., 2006; Brochu and Morrison, 2007; Degner and Wentura, 2009; Anselmi et al., 2013; Brauhardt et al., 2014; Brewis et al., 2016; Jiang et al., 2017). One study (Grover et al., 2003) focused on gender differences in weight bias, recruiting normal-weight and overweight males and females. One study (Brewis and Wutich, 2012) involved a sample of healthy women from Paraguay, whereas Hart et al. (2016) recruited two samples of healthy African American and non-Hispanic white women. Marini et al. (2013) recruited volunteers from 71 different nations. Fifteen papers investigated the effect of different manipulations on weight stigma in healthy students and volunteers.

Looking at results, some studies proved evidence for weight stigma at both explicit and implicit level in medical students (Miller et al., 2013; Phelan et al., 2014 2015b; Baker et al., 2017). Weight bias was also found in samples of university students from different fields including psychology, nursing, business and physical education (Agerström et al., 2007; Chambliss et al., 2004; Gumble et al., 2012; Lynagh et al., 2015; Roddy et al., 2010; 2011; Waller et al., 2012). A study showed that weight bias could be transferred to arbitrary stimuli following a training, which asked to associate weight-related and neutral stimuli (Weinstein et al., 2008). The comparison between weight bias held by psychology and physical education students resulted in a stronger implicit anti-fat bias for the latter group (O'Brien et al., 2007b), whereas Robinson et al. (2014) found similar levels of implicit and explicit weight bias among samples of health and non-health discipline students. O'Brien et al. (2007a) found a correlation between the propensity of making physical-appearance-related comparisons with both explicit and implicit anti-fat attitudes. In a different sample of normal-weight and overweight students involved in a weight discrimination task, Cazzato and Makris (2019) showed a better ability to discriminate actions performed by normal-weight as compared to overweight actors. Another group of students was tested in order to explore the relationship between weight prejudice and expressed and actual behaviour. The data collected showed that prejudicial attitudes toward fatness did not necessarily predict discriminatory behaviours (Penney et al., 2013). However, the expression of weight bias was found to be influenced by the motivation to appear non-prejudiced to others and the perception of weight discrimination (Brochu et al., 2011).

Studies examining weight stigma held by youngsters found evidence for weight prejudice among adolescent (Hand et al., 2017) and children (Solbes and Enesco, 2010; Thomas et al., 2007; Hutchison et al., 2018; Skinner et al., 2017), corroborating the hypothesis that the 'thin ideal' occurs in an early stage of life. A study examined media's influence on weight prejudice in young scholars, finding that more time spent watching television was associated with lower levels of anti-fat bias (Bissel and Hays, 2011).

Other studies investigated the weight bias of healthcare professionals. Some researches (Halvorson et al., 2019; Teachman and Brownell, 2001; Sabin et al., 2012; Schwartz et al., 2003; Tomiyama et al., 2015; Lund et al., 2018; Robertson and Vohora,

2008) showed preferences for thinness compared to fatness at both implicit and explicit levels in individuals working in clinical management of overweight individuals, whereas three studies (Vallis et al., 2007; Dimmock et al., 2009; Aweidah et al., 2016) found evidence for weight stigma only at the implicit level. Furthermore, a strong implicit weight bias was found in a group of physical education teachers (Fontana et al., 2013) and in a sample of professors teaching pre-service physical education (Fontana et al., 2017). However, a different research found neutral implicit attitudes toward obesity in sample of pre-service teachers (Glock et al., 2016). An implicit preference to thinness relative to fatness was found in individuals with different BMI (Anselmi et al., 2013) and a stigmatization of obesity has been found in samples of individuals with overweight or obesity and BED (Wang et al., 2004; Brauhardt et al., 2014), with an inverse relation between participants' BMI and the extent of the bias (Schwartz et al., 2006; Brewis et al., 2016). One study found strong evidence for the influence of weight status on the automatic weight evaluation, with overweight participants showing an implicit preference for overweight over normal-weight stimuli (Degner and Wentura, 2009).

Three papers (Agerstrom et al., 2011; O'Brien et al., 2008; Flint et al., 2016) investigated discrimination of people with obesity at workplace, showing implicit and explicit stigmatization towards candidates with obesity in hiring personnel. Moreover, compared to normal-weight people, overweight individuals seem to be associated with positions characterized by restricted interactions with the public (Venturini et al., 2006). Concerning gender differences in weight bias, one study suggested that males hold more negative attitudes toward obesity (Brochu and Morrison, 2007), whereas Grover et al. (2003) showed that anti-fat bias was ubiquitously held by men and women. A different line of research (Carels et al., 2009a, 2009b, 2010, 2011) found evidence for implicit and explicit anti-fat attitudes in adults with overweight/obesity engaged in a behavioural weight-loss treatments, with higher weight stigma correlating with worse treatment outcomes (Carels et al., 2009b).

Since the prevalence of obese and overweight population is growing worldwide, some studies were interested in understanding how weight-related attitudes vary across different cultural settings. Two studies found both implicit and explicit anti-fat attitudes in a sample of UK residents (Flint et al., 2015) and in a group of individuals from 71 nations (Marini et al., 2013). Implicit anti-fat bias was detected in two samples of Asian (Jiang et al., 2017) and African American (Hart et al., 2016) females, whereas explicit anti-fat stigma has been found in a group of Paraguayan women (Brewis and Wutich, 2012). Moreover, two studies (Carels et al., 2013; Hinman et al., 2015) showed that implicit weight bias was significantly greater when obese and thin people were pictured engaging in stereotype congruent than incongruent activities. However, a behavioural weight loss programme resulted in a reduction of the stereotype consistent bias in a sample of individuals with overweight or obesity (Careles et al., 2014). Studies evaluating the impact of different manipulations on weight bias reported heterogeneous results. An exacerbation of weight stigma has been observed after the exposition to a weight-loss reality show (Domoff et al., 2012; Karsay and Schmuck, 2017) and following the administration of a 'before and after' advertisement, typically picturing an overweight person on the left and the new slim version of the same person on the right (Geier et al., 2003). Differently, a reduction of explicit weight bias has been found after informing participants about the uncontrollable causes of obesity (Wijayatunga et al., 2019), after educating about the multifactorial aetiology

of obesity (Hilbert and Meyre, 2016), after a multi-component intervention aimed at reducing weight bias (Rukavina et al., 2010) and after showing anti-stigma films (Swift et al., 2013). Teachman et al. (2003) showed the possibility to reduce implicit weight bias by evoking empathy with stories of discrimination against individuals with obesity, depending on the weight status of participants. A different intervention based on health messages aimed at enhancing physical activity and healthy habits showed a small reduction of implicit anti-fat stigmatization (Rudolph and Hilbert, 2017). Differently, a counter-conditioning intervention (e.g. presentation of positive images of general public or celebrities with obesity) did not result in more positive perception of fatness, at both implicit and explicit levels (Flint et al., 2013). Similarly, weight bias persisted after a media-based conditioning intervention, which relied upon the presentation of videos portraying obese people struggling with their weight status and discrimination (Gapinski et al., 2006). An ethics educational training partially improved attitudes towards obesity (Geller and Watkins, 2018), whereas Russell-Mayhew et al. (2015) showed a reduction of both implicit and explicit weight bias after performing an interactive professional workshop and O'Brien et al. (2010) demonstrated that it was possible to reduce or exacerbate both anti-fat explicit and implicit attitudes, depending on the information provided about causes of obesity. In contrast, Scrivano et al. (2017) found that informing about the causes of obesity had an impact only on the explicit beliefs about the controllability of obesity. In a research that involved an interactive computer game, the Cyberball game, Pryor et al. (2013) focused on the social influence on behavioural expression of weight bias, proving that both explicit and implicit anti-fat attitudes influenced interactions with an overweight player, but only when other players ostracized the overweight subject.

Analyzing the relationship between implicit and explicit measures of weight bias, studies reported heterogeneous results. High levels of implicit bias were coupled with low or completely absent explicit pro-slim/anti-fat preference in some studies (Roddy et al., 2010; Lynagh et al., 2015; Aweidah et al., 2016; Dimmock et al., 2009; Halvorson et al., 2019; Teachman and Brownell, 2001; Carels et al., 2009a; Jiang et al., 2017). Conversely, other studies found similar levels of implicit and explicit weight stigma (Schwartz et al., 2003; Sabin et al., 2012; Marini et al., 2013; Phelan et al., 2014; Tomiyama et al., 2015; Flint et al., 2015, 2016). A minority of data suggested high levels of explicit preference for thinness, coupled with very low levels of implicit weight bias (Brewis and Wutich, 2012).

Discussion

We reviewed studies on implicit attitudes towards food and body in healthy population and in samples of patients with ED. One hundred and eighty three papers were evaluated for bias risks and synthesized. Some main findings emerge from this review. A first evidence is that very few studies explored the neurobiological correlates of implicit attitudes related to eating behaviour. Neuroscientific investigation of eating behaviour has increased in the past decades supporting a brain-based approach to eating disorders and outlining different neurobiological models, which represent the rationale for combining psychotherapy and biological treatments (Val-Laillet et al., 2015; Frank, 2019). Neuroimaging studies reported the involvement of emotional and reward neuronal circuits in monitoring eating behaviour, and neuromodulation treatments have been tested targeting these circuits (Val-Laillet et al., 2015; Hall et al., 2018b).

Therefore, understanding the relationship between neural circuits underpinning eating behaviour and brain mechanisms related to implicit attitudes, which are measures of automatic affective evaluation of food or body (Greenwald and Farnham, 2003; De Houwer et al., 2009), is of great interest. The low number of studies prevents to delineate clear conclusions on this aspect, but it is worth noting that one paper assessing reward sensitivity for food in patients with PD in dopamine replacement treatment reported discrepancies between implicit and explicit food attitudes in patients with PD and binge eating, in line with findings with other samples of disordered eating (Papies et al., 2009; Terenzi et al., 2018). One ERP study showed task-related differences in the N400 component only in patients with BN, supporting the hypothesis that implicit attitudes in EDs could be associated to anomalous brain response at an early stage (Blechert et al., 2011). Neurostimulation studies are consistent in highlighting the crucial impact of individual features in modulating implicit attitudes. Indeed, one study showed that TMS on mPFC affected food attitudes only in a subgroup of participants with low preference for tasty food (Mattavelli et al., 2015), whereas two studies with tDCS on EBA found an effect only in male participants with anti-fat bias, but not in females, and in patients with EDs, but not in control females (Cazzato et al., 2017; Mattavelli et al., 2019). Although using different IATs, these latter studies are consistent in showing the lack of modulatory effect on body images IAT in female participants applying anodal stimulation on EBA. Interestingly, one study stimulating the left dlPFC with cTBS did not find significant effect of neuromodulation on IAT, but reported different correlational patterns between IAT and food consumption in active or sham cTBS condition, supporting a role for left dlPFC in control hunger disinhibition (Hall et al., 2018).

For what concerns food attitudes many different paradigms have been used, with preferences pointing to different directions depending on the type of food categories and evaluative attributes used as stimuli. A general conclusion that can be drawn concerns the fact that most of the studies support the predictive validity of implicit measures on the actual food behaviour. Crucially, the relationship between implicit and explicit measures and behaviour seems to be mediated by individual differences in food habits and ED symptoms (Ellis et al., 2014). In particular, discrepancies between implicit and explicit preferences were more evident in restrained eaters and predicted disinhibited eating with larger validity in individuals with higher level of impulsivity (Papies et al., 2009; Goldstein et al., 2014).

Studies assessing attitudes toward body images show substantially convergent findings of negative implicit attitudes toward overweight body images or stronger preference for normal-weight compared to underweight and overweight bodies in the healthy population (Ahern and Hetherington 2006; Ahern et al., 2008; Lydecker et al., 2006; Watts et al., 2008; Mousally et al., 2015; Sabin et al., 2015; Marini, 2017; Elran-Barak and Bar-Anan, 2018; Robstad et al., 2018), whereas greater pro-thin implicit bias in patients with AN and BN compared to healthy subjects has been reported (Khan and Petróczi, 2015; Smith et al., 2018; Izquierdo et al., 2019). Nevertheless, patients with AN also showed anti-fat bias, rather than attraction to thinness (Spring and Bulik, 2014). Moreover, experimental manipulation could be effectively used to modulate body implicit attitudes in ED samples (Smith et al., 2014), or in healthy subjects (Martijn et al., 2013; Matharu et al., 2014), even though this result was not consistent across the studies (Keng and Ang, 2019). The correspondence between implicit and explicit measures on body image as well as

Appendix 1. Descriptors of search

Search engine	Search algorithm	Alternative keywords
PubMed (May 2019)	<p>Implicit attitudes AND eating disorder: (implicit[All Fields] AND ('attitude'[MeSH Terms] OR 'attitude'[All Fields] OR 'attitudes'[All Fields])) AND ('feeding and eating disorders'[MeSH Terms] OR ('feeding'[All Fields] AND 'eating'[All Fields] AND 'disorders'[All Fields]) OR 'feeding and eating disorders'[All Fields] OR ('eating'[All Fields] AND 'disorder'[All Fields]) OR 'eating disorder'[All Fields])</p> <p>Implicit attitudes AND anorexia nervosa: (implicit[All Fields] AND ('attitude'[MeSH Terms] OR 'attitude'[All Fields] OR 'attitudes'[All Fields])) AND ('anorexia nervosa'[MeSH Terms] OR ('anorexia'[All Fields] AND 'nervosa'[All Fields]) OR 'anorexia nervosa'[All Fields])</p> <p>Implicit attitudes AND bulimia nervosa: (implicit[All Fields] AND ('attitude'[MeSH Terms] OR 'attitude'[All Fields] OR 'attitudes'[All Fields])) AND ('bulimia nervosa'[MeSH Terms] OR ('bulimia'[All Fields] AND 'nervosa'[All Fields]) OR 'bulimia nervosa'[All Fields])</p> <p>Implicit attitudes AND binge eating disorder: (implicit[All Fields] AND ('association'[MeSH Terms] OR 'association'[All Fields])) AND ('binge-eating disorder'[MeSH Terms] OR ('binge-eating'[All Fields] AND 'disorder'[All Fields]) OR 'binge-eating disorder'[All Fields] OR ('binge'[All Fields] AND 'eating'[All Fields] AND 'disorder'[All Fields]) OR 'binge eating disorder'[All Fields])</p> <p>Implicit attitudes AND obesity: (implicit[All Fields] AND ('attitude'[MeSH Terms] OR 'attitude'[All Fields] OR 'attitudes'[All Fields])) AND ('obesity'[MeSH Terms] OR 'obesity'[All Fields])</p> <p>Implicit attitudes AND food preference: (implicit[All Fields] AND ('attitude'[MeSH Terms] OR 'attitude'[All Fields] OR 'attitudes'[All Fields])) AND ('food preferences'[MeSH Terms] OR ('food'[All Fields] AND 'preferences'[All Fields]) OR 'food preferences'[All Fields] OR ('food'[All Fields] AND 'preference'[All Fields]) OR 'food preference'[All Fields])</p> <p>Implicit attitudes AND thin idea: (implicit[All Fields] AND ('attitude'[MeSH Terms] OR 'attitude'[All Fields] OR 'attitudes'[All Fields])) AND (thin[All Fields] AND ('IDEA J Law Technol'[Journal] OR 'idea'[All Fields]))</p> <p>Implicit attitudes AND thin ideal: (implicit[All Fields] AND ('attitude'[MeSH Terms] OR 'attitude'[All Fields] OR 'attitudes'[All Fields])) AND (thin[All Fields] AND ideal[All Fields])</p> <p>Implicit attitudes AND fat phobia: (implicit[All Fields] AND ('association'[MeSH Terms] OR 'association'[All Fields])) AND (fat[All Fields] AND ('phobic disorders'[MeSH Terms] OR ('phobic'[All Fields] AND 'disorders'[All Fields]) OR 'phobic disorders'[All Fields] OR 'phobia'[All Fields]))</p>	Implicit association; Affective priming
EMBASE (May 2019)	<p>Implicit attitudes AND eating disorder: implicit AND attitudes AND eating AND disorder</p> <p>Implicit attitudes AND anorexia nervosa: implicit AND attitudes AND anorexia AND nervosa</p> <p>Implicit attitudes AND bulimia nervosa: implicit AND attitudes AND bulimia AND nervosa</p> <p>Implicit attitudes AND binge eating disorder: implicit AND attitudes AND binge AND eating AND disorder</p> <p>Implicit attitudes AND obesity: implicit AND attitudes AND obesity</p> <p>Implicit attitudes AND food preference: implicit AND attitudes AND food AND preference</p> <p>Implicit attitudes AND thin idea: implicit AND attitudes AND thin AND idea</p> <p>Implicit attitudes AND thin ideal: implicit AND attitudes AND thin AND ideal</p> <p>Implicit attitudes AND fat phobia: implicit AND attitudes AND fat AND phobia</p> <p>Implicit attitudes AND eating disorder: noft("implicit attitudes") AND noft("eating disorder")</p>	Implicit association; Affective priming
PsychINFO (May 2019)	<p>Implicit attitudes AND eating disorder: noft("implicit attitudes") AND noft("eating disorder")</p> <p>Implicit attitudes AND anorexia nervosa: noft("implicit attitudes") AND noft("anorexia nervosa")</p> <p>Implicit attitudes AND bulimia nervosa: noft("implicit attitudes") AND noft("bulimia nervosa")</p> <p>Implicit attitudes AND binge eating disorder: noft("implicit attitudes") AND noft("binge eating disorder")</p> <p>Implicit attitudes AND obesity: noft("implicit attitudes") AND noft("obesity")</p> <p>Implicit attitudes AND food preference: noft("implicit attitudes") AND noft("food preference")</p> <p>Implicit attitudes AND thin idea: noft("implicit attitudes") AND noft("thin idea")</p> <p>Implicit attitudes AND thin ideal: noft("implicit attitudes") AND noft("thin ideal")</p> <p>Implicit attitudes AND fat phobia: noft("implicit attitudes") AND noft("fat phobia")</p>	Implicit association; Affective priming

(Continued)

Search engine	Search algorithm	Alternative keywords
SCOPUS (May 2019)	Implicit attitudes AND eating disorder: TITLE-ABS-KEY (implicit AND attitudes AND eating AND disorder)	Implicit association; Affective priming
	Implicit attitudes AND anorexia nervosa: TITLE-ABS-KEY (implicit AND attitudes AND anorexia AND nervosa)	
	Implicit attitudes AND bulimia nervosa: TITLE-ABS-KEY (implicit AND attitudes AND bulimia AND nervosa)	
	Implicit attitudes AND binge eating disorder: TITLE-ABS-KEY (implicit AND attitudes AND binge AND eating AND disorder)	
	Implicit attitudes AND obesity: TITLE-ABS-KEY (implicit AND attitudes AND obesity)	
	Implicit attitudes AND food preference: TITLE-ABS-KEY (implicit AND attitudes AND food AND preference)	
	Implicit attitudes AND thin idea: TITLE-ABS-KEY (implicit AND attitudes AND thin AND idea)	
	Implicit attitudes AND thin ideal: TITLE-ABS-KEY (implicit AND attitudes AND thin AND ideal)	
	Implicit attitudes AND fat phobia: TITLE-ABS-KEY (implicit AND attitudes AND fat AND phobia)	

In each search algorithm, the 'Implicit attitudes' keyword have been replaced with the alternatives in the right column; noft = research in all filed excluding entire manuscripts due to the exceeding number of retrieved records; TITLE-ABS-KEY = research in title, abstract and keywords (excluding entire manuscripts) due to the exceeding number of retrieved records.

References

- Ackermann, C.L., Palmer, A. (2014). The contribution of implicit cognition to the Theory of Reasoned Action Model: a study of food preferences. *Journal of Marketing Management*, 30(5–6), 529–50.
- Adams, R.C., Lawrence, N.S., Verbruggen, F., Chambers, C.D. (2017). Training response inhibition to reduce food consumption: mechanisms, stimulus specificity and appropriate training protocols. *Appetite*, 109, 11–23.
- Agerström, J., Carlsson, R., Rooth, D.O. (2007). Ethnicity and obesity: evidence of implicit work performance stereotypes in Sweden. *EconStor*, 20.
- Agerström, J., Rooth, D.-O. (2011). The role of automatic obesity stereotypes in real hiring discrimination. *The Journal of Applied Psychology*, 96, 790–805.
- Ahern, A.L., Bennett, K.M., Hetherington, M.M. (2008). Internalization of the ultra-thin ideal: positive implicit associations with underweight fashion models are associated with drive for thinness in young women. *Eating Disorders*, 16(4), 294–307.
- Ahern, A.L., Hetherington, M.M. (2006). The thin ideal and body image: an experimental study of implicit attitudes. *Psychology of Addictive Behaviors*, 20(3), 338.
- Alabduljader, K., Cliffe, M., Sartor, F., Papini, G., Cox, W.M., Kubis, H.P. (2018). Ecological momentary assessment of food perceptions and eating behavior using a novel phone application in adults with or without obesity. *Eating Behaviors*, 30(May), 35–41.
- Alblas, E.E., Folkvord, F., Anschutz, D.J., et al. (2018). Investigating the impact of a health game on implicit attitudes towards food and food choice behaviour of young adults. *Appetite*, 128, 294–302.
- Alkozei, A., Killgore, W.D.S., Smith, R., et al. (2018). Chronic sleep restriction differentially affects implicit biases toward food among men and women: preliminary evidence. *Journal of Sleep Research*, 27(4), 1–4.
- Anselmi, P., Vianello, M., Robusto, E. (2013). Preferring thin people does not imply derogating fat people. A Rasch analysis of the implicit weight attitude. *Obesity*, 21, 261–65.
- Anselmi, P., Vianello, M., Robusto, E. (2011). Positive associations primacy in the IAT. *Experimental Psychology*, 58, 376–84.
- Ashby, C.R., Stritzke, W.G.K. (2013). Is sensitivity to reward associated with the malleability of implicit inclinations toward high-fat food? *Emotion*, 13(4), 711–23.
- Aweidah, L., Robinson, J., Cumming, S., Lewis, S. (2016). Australian diagnostic radiographers' attitudes and perceptions of imaging obese patients: a study of self, peers and students. *Radiography*, 22, e258–63.
- Ayres, K., Conner, M.T., Prestwich, A., Smith, P. (2012). Do implicit measures of attitudes incrementally predict snacking behaviour over explicit affect-related measures? *Appetite*, 58(3), 835–41.
- Baker, T.K., Smith, G.S., Jacobs, N.N., et al. (2017). A deeper look at implicit weight bias in medical students. *Advances in Health Sciences Education*, 22, 889–900.
- Barnes-Holmes, D., Barnes-Holmes, Y., Stewart, I., Boles, S. (2010). A sketch of the implicit relational assessment procedure (IRAP) and the relational elaboration and coherence (REC) model. *Psychological Record*, 60(3), 527–42.
- Becker, D., Jostmann, N.B., Wiers, R.W., Holland, R.W. (2015). Approach avoidance training in the eating domain: testing the effectiveness across three single session studies. *Appetite*, 85, 58–65.
- Benas, J.S., Gibb, B.E. (2011). Childhood teasing and adult implicit cognitive biases. *Cognitive Therapy and Research*, 35(6), 491–96.
- Bissell, K., Hays, H. (2010). Understanding anti-fat bias in children: the role of media and appearance anxiety in third to

- sixth graders' implicit and explicit attitudes toward obesity. *Mass Communication and Society*, 14(1), 113–40.
- Blechert, J., Ansoorge, U., Beckmann, S., Tuschen-Caffier, B. (2011). The undue influence of shape and weight on self-evaluation in anorexia nervosa, bulimia nervosa and restrained eaters: a combined ERP and behavioral study. *Psychological Medicine*, 41(1), 185–94.
- Bongers, P., Jansen, A., Houben, K., Roefs, A. (2013). Happy eating: the single target implicit association test predicts overeating after positive emotions. *Eating Behaviors*, 14(3), 348–55.
- Brauhardt, A., Rudolph, A., Hilbert, A. (2014). Implicit cognitive processes in binge-eating disorder and obesity. *Journal of Behavior Therapy and Experimental Psychiatry*, 45(2), 285–90.
- Brewis, A., Brennhof, S., van Wodern, I., Bruening, M. (2016). Weight stigma and eating behaviors on a college campus: are students immune to stigma's effects? *Preventive Medicine Reports*, 4, 578–84.
- Brewis, A.A., Wutich, A. (2012). Explicit versus implicit fat-stigma. *American Journal of Human Biology*, 24, 332–38.
- Brochu, P.M., Gawronski, B., Esses, V.M. (2011). The integrative prejudice framework and different forms of weight prejudice: an analysis and expansion. *Group Processes & Intergroup Relations*, 14(3), 429–44.
- Brochu, P.M., Morrison, M.A. (2007). Implicit and explicit prejudice toward overweight and average-weight men and women: testing their correspondence and relation to behavioral intentions. *The Journal of Social Psychology*, 147(6), 681–706.
- Brooks, S.J., O'Daly, O.G., Uher, R., et al. (2011). Differential neural responses to food images in women with bulimia versus anorexia nervosa. *PLoS One*, 6(7), e22259.
- Carels, R.A., Hinman, N.G., Hoffmann, D.A., et al. (2014). Implicit bias about weight and weight loss treatment outcomes. *Eating Behaviors*, 15(4), 648–53.
- Carels, R.A., Hinman, N.G., Burmeister, J.M., Hoffmann, D.A., Ashrafioun, L., Koball, A.M. (2013). Stereotypical images and implicit weight bias in overweight/obese people. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*, 18(4), 441–5.
- Carels, R.A., Hinman, N., Koball, A., Oehlhof, M.W., Gumble, A., Young, K.M. (2011). The self-protective nature of implicit identity and its relationship to weight bias and short-term weight loss. *Obesity Facts*, 4, 278–83.
- Carels, R.A., Wott, C.B., Young, K.M., Gumble, A., Koball, A., Oehlhof, M.W. (2010). Implicit, explicit, and internalized weight bias and psychosocial maladjustment among treatment-seeking adults. *Eating Behaviors*, 11, 180–5.
- Carels, R.A., Young, K.M., Wott, C.B., et al. (2009a). Internalized weight stigma and its ideological correlates among weight loss treatment seeking adults. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 14(2–3), e92–7.
- Carels, R.A., Young, K.M., Wott, C.B., et al. (2009b). Weight bias and weight loss treatment outcomes in treatment-seeking adults. *Annals of Behavioral Medicine*, 37(3), 350–5.
- Cattaneo, Z., Mattavelli, G., Platania, E., Papagno, C. (2011). The role of the prefrontal cortex in controlling gender-stereotypical associations: a TMS investigation. *NeuroImage*, 56, 1839–46.
- Cazzato, V., Makris, S. (2019). Implicit preference towards slim bodies and weight-stigma modulate the understanding of observed familiar actions. *Psychological Research*, 83, 1825–35.
- Cazzato, V., Makris, S., Urgesi, C. (2017). Cathodal transcranial direct current stimulation of the extrastriate visual cortex modulates implicit anti-fat bias in male, but not female, participants. *Neuroscience*, 359, 92–104.
- Chambliss, H.O., Finley, C.E., Blair, S.N. (2004). Attitudes toward obese individuals among exercise science students. *Medicine and Science in Sports and Exercise*, 36, 468–74.
- Chee, M.W., Sriram, N., Soon, C.S., Lee, K.M. (2000). Dorsolateral prefrontal cortex and the implicit association of concepts and attributes. *Neuroreport*, 11(1), 135–40.
- Conner, M.T., Perugini, M., O'Gorman, R., Ayres, K., Prestwich, A. (2007). Relations between implicit and explicit measures of attitudes and measures of behavior: evidence of moderation by individual difference variables. *Personality and Social Psychology Bulletin*, 33(12), 1727–40.
- Coricelli, C., Feroni, F., Osimo, S.A., Rumiati, R.I. (2019). Implicit and explicit evaluations of foods: the natural and transformed dimension. *Food Quality and Preference*, 73, 143–53.
- Craeynest, M., Crombez, G., Deforche, B., Tanghe, A., De Bourdeaudhuij, I. (2008a). The role of implicit attitudes towards food and physical activity in the treatment of youth obesity. *Eating Behaviors*, 9(1), 41–51.
- Craeynest, M., Crombez, G., Koster, E.H., Haerens, L., De Bourdeaudhuij, I. (2008b). Cognitive-motivational determinants of fat food consumption in overweight and obese youngsters: the implicit association between fat food and arousal. *Journal of Behavior Therapy and Experimental Psychiatry*, 39(3), 354–68.
- Craeynest, M., Crombez, G., Haerens, L., De Bourdeaudhuij, I. (2007). Do overweight youngsters like food more than lean peers? Assessing their implicit attitudes with a personalized Implicit Association Task. *Food Quality and Preference*, 18(8), 1077–84.
- Craeynest, M., Crombez, G., De Houwer, J., Deforche, B., De Bourdeaudhuij, I. (2006). Do children with obesity implicitly identify with sedentariness and fat food? *Journal of Behavior Therapy and Experimental Psychiatry*, 37(4), 347–57.
- Craeynest, M., Crombez, G., De Houwer, J., Deforche, B., Tanghe, A., De Bourdeaudhuij, I. (2005). Explicit and implicit attitudes towards food and physical activity in childhood obesity. *Behaviour Research and Therapy*, 43(9), 1111–20.
- Crescentini, C., Aglioti, S.M., Fabbro, F., Urgesi, C. (2014). Virtual lesions of the inferior parietal cortex induce fast changes of implicit religiousness/spirituality. *Cortex*, 54, 1–15.
- Cserjesi, R., De Vos, I., Deroost, N. (2016). Discrepancy between implicit and explicit preferences for food portions in obesity. *International Journal of Obesity*, 40(9), 1464–7.
- Cserjesi, R., Vermeulen, N., Luminet, O., et al. (2010). Explicit vs. implicit body image evaluation in restrictive anorexia nervosa. *Psychiatry Research*, 175(1–2), 148–53.
- Czyzewska, M., Graham, R. (2008). Implicit and explicit attitudes to high- and low-calorie food in females with different BMI status. *Eating Behaviors*, 9(3), 303–12.
- Dagher, A., Robbins, T.W. (2009). Personality, addiction, dopamine: insights from Parkinson's disease. *Neuron*, 61(4), 502–10.
- Degner, J., Wentura, D. (2009). Not everybody likes the thin and despises the fat: one's weight matters in the automatic activation of weight-related social evaluations. *Social Cognition*, 27(2), 202–21.
- De Houwer, J., Teige-Mocigemba, S., Spruyt, A., Moors, A. (2009). Implicit measures: a normative analysis and review. *Psychological Bulletin*, 135(3), 347–68.
- De Houwer, J. (2003). The Extrinsic Affective Simon Task. *Experimental Psychology*, 50(2), 77–85.
- De Houwer, J. (2002). The Implicit Association Test as a tool for studying dysfunctional associations in psychopathology: strengths and limitations. *Journal of Behavior Therapy and Experimental Psychiatry*, 33(2), 115–33.

- Dimmock, J.A., Hallett, B.E., Grove, R.J. (2009). Attitudes toward overweight individuals among fitness center employees: an examination of contextual effects. *Research Quarterly for Exercise and Sport*, 80(3), 641–7.
- Domoff, S.E., Hinman, N.G., Koball, A.M., et al. (2012). The effects of reality television on weight bias: an examination of the biggest loser. *Obesity*, 20, 993–8.
- Ellis, E.M., Kiviniemi, M.T., Cook-Cottone, C. (2014). Implicit affective associations predict snack choice for those with low, but not high levels of eating disorder symptomatology. *Appetite*, 77, 124–32.
- Elran-Barak, R., Bar-Anan, Y. (2018). Implicit and explicit anti-fat bias: the role of weight-related attitudes and beliefs. *Social Science & Medicine*, 204, 117–24.
- Eschenbeck, H., Heim-Dreger, U., Steinhilber, A., Kohlmann, C.-W. (2016). Self-regulation of healthy nutrition: automatic and controlled processes. *BMC Psychology*, 4(1), 1–8.
- Expósito, P.M., López, M.H., Valverde, M.R. (2015). Assessment of implicit anti-fat and pro-slim attitudes in young women using the implicit relational assessment procedure. *International Journal of Psychology and Psychological Therapy*, 15(1), 17–32.
- Fazio, R.H., Sanbonmatsu, D.M., Powell, M.C., Kardes, F.R. (1986). On the automatic activation of attitudes. *Journal of Personality and Social Psychology*, 50(2), 229–38.
- Ferentzi, H., Scheibner, H., Wiers, R., et al. (2018). Retraining of automatic action tendencies in individuals with obesity: a randomized controlled trial. *Appetite*, 126, 66–72.
- Flint, S.W., Čadek, M., Codreanu, S.C., Ivić, V., Zomer, C., Gomoiu, A. (2016). Obesity discrimination in the recruitment process: “you’re not hired!”. *Frontiers in Psychology*, 7, 647.
- Flint, S.W., Hudson, J., Lavallee, D. (2015). UK adults’ implicit and explicit attitudes towards obesity: a cross-sectional study. *BMC Obesity*, 2(1), 31.
- Flint, S.W., Hudson, J., Lavallee, D. (2013). Counter-conditioning as an intervention to modify anti-fat attitudes. *Health Psychology Research*, 1, e24.
- Fontana, F., Furtado, O., Jr, Mazzardo, O., Hong, D., de Campos, W. (2017). Anti-fat bias by professors teaching physical education majors. *European Physical Education Review*, 23(1), 127–38.
- Fontana, F.E., Furtado, O., Marston, R., Mazzardo, O., Gallagher, J. (2013). Anti-fat bias among physical education teachers and majors. *Physical Educator*, 70(1), 15.
- Forbes, C.E., Cameron, K.A., Grafman, J., et al. (2012). Identifying temporal and causal contributions of neural processes underlying the Implicit Association Test (IAT). *Frontiers in Human Neuroscience*, 6(November 2012), 1–18.
- Forhan, M., Salas, X.R. (2013). Inequities in healthcare: a review of bias and discrimination in obesity treatment. *Canadian Journal of Diabetes*, 37(3), 205–9.
- Frank, G.K.W. (2019). Editorial to the virtual issue highlighting neuroscience based research in eating disorders to mark the 49th Society for Neuroscience Annual Meeting. *International Journal of Eating Disorders*, 52(11), 1332–5.
- Friese, M., Hofmann, W., Wänke, M. (2008). When impulses take over: moderated predictive validity of explicit and implicit attitude measures in predicting food choice and consumption behaviour. *British Journal of Social Psychology*, 47(3), 397–419.
- Gapinski, K.D., Schwartz, M.B., Brownell, K.D. (2006). Can television change anti-fat attitudes and behavior? *Journal of Applied Biobehavioral Research*, 11(1), 1–28.
- Garner, D.M. (2004). *Eating Disorder Inventory-3 (EDI-3)*. Professional Manual. Odessa, FL: Psychological Assessment Resources.
- Gawronski, B., Bodenhausen, G.V. (2006). Associative and propositional processes in evaluation: an integrative review of implicit and explicit attitude change. *Psychological Bulletin*, 132(5), 692–731.
- Geier, A.B., Schwartz, M.B., Brownell, K.D. (2003). “Before and after” diet advertisements escalate weight stigma. *Eating and Weight Disorders*, 8, 282–8.
- Geller, G., Watkins, P.A. (2018). Addressing medical students’ negative bias toward patients with obesity through ethics education. *AMA Journal of Ethics*, 20, e948–59.
- Genschow, O., Demanet, J., Hersche, L., Brass, M. (2017). An empirical comparison of different implicit measures to predict consumer choice. *PLoS One*, 12(8), 1–13.
- Glashouwer, K.A., Bennik, E.C., de Jong, P.J., Spruyt, A. (2018). Implicit measures of actual versus ideal body image: relations with self-reported body dissatisfaction and dieting behaviors. *Cognitive Therapy and Research*, 42(5), 622–35.
- Glock, S., Beverborg, A.O.G., Müller, B.C.N. (2016). Pre-service teachers’ implicit and explicit attitudes toward obesity influence their judgments of students. *Social Psychology of Education*, 19(1), 97–115.
- Goldstein, S.P., Forman, E.M., Meiran, N., Herbert, J.D., Juarascio, A.S., Butryn, M.L. (2014). The discrepancy between implicit and explicit attitudes in predicting disinhibited eating. *Eating Behaviors*, 15(1), 164–70.
- Gozzi, M., Raymont, V., Solomon, J., Koenigs, M., Grafman, J. (2009). Dissociable effects of prefrontal and anterior temporal cortical lesions on stereotypical gender attitudes. *Neuropsychologia*, 47, 2125–32.
- Grant, M.J., Booth, A. (2009). A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Information & Libraries Journal*, 26(2), 91–108.
- Greenwald, A.G., Nosek, B.A., Banaji, M.R. (2003). Understanding and using the Implicit Association Test: I. An improved scoring algorithm. *Journal of Personality and Social Psychology*, 85(2), 197–216.
- Greenwald, A.G., McGhee, D.E., Schwartz, J.L.K. (1998). Measuring individual differences in implicit cognition: the Implicit Association Test. *Journal of Personality and Social Psychology*, 74(6), 1464–80.
- Grover, V.P., Keel, P.K., Mitchell, J.P. (2003). Gender differences in implicit weight identity. *International Journal of Eating Disorders*, 34, 125–35.
- Guidetti, M., Conner, M., Prestwich, A., Cavazza, N. (2012). The transmission of attitudes towards food: twofold specificity of similarities with parents and friends. *British Journal of Health Psychology*, 17(2), 346–61.
- Gumble, A., Carels, R. (2012). The harmful and beneficial impacts of weight bias on well-being: the moderating influence of weight status. *Body Image*, 9, 101–7.
- Hall, P.A., Lowe, C.J., Safati, A.B., Li, H., Klassen, E.B., Burhan, A.M. (2018a). Effects of left dlPFC modulation on social cognitive processes following food sampling. *Appetite*, 126, 73–9.
- Hall, P., Vincent, C.M., Burhan, A.M. (2018b). Non-invasive brain stimulation for food cravings, consumption, and disorders of eating: A review of methods, findings and controversies. *Appetite*, 124, 78–88.
- Halmi, K.A. (2013). Perplexities of treatment resistance in eating disorders. *BMC Psychiatry*, 13(1), 292.
- Halvorson, E.E., Curley, T., Wright, M., Skelton, J.A. (2019). Weight bias in pediatric inpatient care. *Academic Paediatrics*, 19, 780–6.
- Hand, W.B., Robinson, J.C., Stewart, M.W., Zhang, L., Hand, S.C. (2017). The identity threat of weight stigma in adolescents. *Western Journal of Nursing Research*, 39(8), 991–1007.

- Hart, E.A., Sbrocco, T., Carter, M.M. (2016). Ethnic identity and implicit anti-fat bias: similarities and differences between African American and Caucasian women. *Ethnicity & Disease*, 26, 69–76.
- Healy, G.F., Boran, L., Smeaton, A.F. (2015). Neural patterns of the implicit association test. *Frontiers in Human Neuroscience*, 9(Nov), 1–16.
- Heider, N., Spruyt, A., De Houwer, J. (2018). Body dissatisfaction revisited: on the importance of implicit beliefs about actual and ideal body image. *Psychologica Belgica*, 57(4), 158.
- Heider, N., Spruyt, A., De Houwer, J. (2015). Implicit beliefs about ideal body image predict body image dissatisfaction. *Frontiers in Psychology*, 6, 1402.
- Hensels, I.S., Baines, S. (2016). Changing ‘gut feelings’ about food: an evaluative conditioning effect on implicit food evaluations and food choice. *Learning and Motivation*, 55, 31–44.
- Higgins, J.P., Altman, D.G., Gøtzsche, P.C., et al. (2011). The Cochrane Collaboration’s tool for assessing risk of bias in randomised trials. *BMJ*, 343, d5928.
- Hilbert, A., Meyre, D. (2016). Weight stigma reduction and genetic determinism. *PLoS One*, 11(9), e0162993.
- Hilbert, A., Braehler, E., Haeuser, W., Zenger, M. (2014). Weight bias internalization, core self-evaluation, and health in overweight and obese persons. *Obesity*, 22, 79–85.
- Hinman, N.G., Burmeister, J.M., Kiefner, A.E., Borushok, J., Carels, R.A. (2015). Stereotypical portrayals of obesity and the expression of implicit weight bias. *Body Image*, 12, 32–5.
- Hoefling, A., Strack, F. (2008). The tempting effect of forbidden foods. High calorie content evokes conflicting implicit and explicit evaluations in restrained eaters. *Appetite*, 51(3), 681–9.
- Hollands, G.J., Prestwich, A., Marteau, T.M. (2011). Using aversive images to enhance healthy food choices and implicit attitudes: an experimental test of evaluative conditioning. *Health Psychology*, 30(2), 195–203.
- Houben, K., Roefs, A., Jansen, A. (2012). Guilty pleasures II: restrained eaters’ implicit preferences for high, moderate and low-caloric food. *Eating Behaviors*, 13(3), 275–7.
- Houben, K., Roefs, A., Jansen, A. (2010). Guilty pleasures. Implicit preferences for high calorie food in restrained eating. *Appetite*, 55(1), 18–24.
- House, A.E., House, B.J., Campbell, M.B. (1981). Measures of inter-observer agreement: calculation formulas and distribution effects. *Journal of Behavioral Assessment*, 3(1), 37–57.
- Hutchison, S.M., Müller, U. (2018). Explicit and implicit measures of weight stigma in young children. *Merrill-Palmer Quarterly*, 64(4), 427–58.
- Izquierdo, A., Plessow, F., Becker, K.R., et al. (2019). Implicit attitudes toward dieting and thinness distinguish fat-phobic and non-fat-phobic anorexia nervosa from avoidant/restrictive food intake disorder in adolescents. *International Journal of Eating Disorders*, 52(4), 419–27.
- Jiang, W., Tan, J., Fassnacht, D.B. (2017). Implicit and explicit anti-fat bias among Asian females. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 22, 457–65.
- Johansson, L., Ghaderi, A., Andersson, G. (2005). Stroop interference for food- and body-related words: a meta-analysis. *Eating Behaviors*, 6(3), 271–81.
- Juarascio, A.S., Forman, E.M., Timko, C.A., Herbert, J.D., Butryn, M., Lowe, M. (2011). Implicit internalization of the thin ideal as a predictor of increases in weight, body dissatisfaction, and disordered eating. *Eating Behaviors*, 12(3), 207–13.
- Kakoschke, N., Kemps, E., Tiggemann, M. (2017). The effect of combined avoidance and control training on implicit food evaluation and choice. *Journal of Behavior Therapy and Experimental Psychiatry*, 55, 99–105.
- Karpinski, A., Steinman, R.B. (2006). The Single Category Implicit Association Test as a measure of implicit social cognition. *Journal of Personality and Social Psychology*, 91, 16–32.
- Karsay, K., Schmuck, D. (2019). “Weak, sad, and lazy fatties”: adolescents’ explicit and implicit weight bias following exposure to weight loss reality TV shows. *Media Psychology*, 22(1), 60–81.
- Kemps, E., Tiggemann, M. (2015). Approach bias for food cues in obese individuals. *Psychology & Health*, 30(3), 370–80.
- Keng, S.-L., Ang, Q. (2019). Effects of mindfulness on negative affect, body dissatisfaction, and disordered eating urges. *Mindfulness*, 10(9), 1779–91.
- Khan, S., Petróczy, A. (2015). Stimulus-response compatibility tests of implicit preference for food and body image to identify people at risk for disordered eating: a validation study. *Eating Behaviors*, 16, 54–63.
- Kottner, J., Streiner, D.L. (2011). The difference between reliability and agreement. *Journal of Clinical Epidemiology*, 64(6), 701–2.
- Kraus, A.A., Piqueras-Fiszman, B. (2016). Sandwich or sweets? An assessment of two novel implicit association tasks to capture dynamic motivational tendencies and stable evaluations towards foods. *Food Quality and Preference*, 49, 11–9.
- Lamote, S., Hermans, D., Baeyens, F., Eelen, P. (2004). An exploration of affective priming as an indirect measure of food attitudes. *Appetite*, 42(3), 279–86.
- Lebens, H., Roefs, A., Martijn, C., Houben, K., Nederkoorn, C., Jansen, A. (2011). Making implicit measures of associations with snack foods more negative through evaluative conditioning. *Eating Behaviors*, 12(4), 249–53.
- Lund, T.B., Brodersen, J., Sandøe, P. (2018). A study of anti-fat bias among Danish general practitioners and whether this bias and general practitioners’ lifestyle can affect treatment of tension headache in patients with obesity. *Obesity Facts*, 11(6), 501–13.
- Lydecker, J.A., O’Brien, E., Grilo, C.M. (2018). Parents have both implicit and explicit biases against children with obesity. *Journal of Behavioral Medicine*, 41(6), 784–91.
- Lynagh, M., Cliff, K., Morgan, P.J. (2015). Attitudes and beliefs of nonspecialist and specialist trainee health and physical education teachers toward obese children: evidence for “anti-fat” bias. *The Journal of School Health*, 9, 595–603.
- Maas, J., Woud, M.L., Keijsers, G.P.J., Rinck, M., Becker, E.S., Wiers, R.W. (2017). The attraction of sugar: an association between body mass index and impaired avoidance of sweet snacks. *Journal of Experimental Psychopathology*, 8(1), 40–54.
- Mai, R., Hoffmann, S. (2015). How to combat the unhealthy = tasty intuition: the influencing role of health consciousness. *Journal of Public Policy & Marketing*, 34(1), 63–83.
- Marini, M. (2017). Underweight vs. Overweight/obese: which weight category do we prefer? Dissociation of weight-related preferences at the explicit and implicit level. *Obesity Science & Practice*, 3(4), 390–8.
- Marini, M., Sriram, N., Schnabel, K., et al. (2013). Overweight people have low levels of implicit weight bias, but overweight nations have high levels of implicit weight bias. *PLoS One*, 8, e83543.
- Martijn, C., Sheeran, P., Wesseldijk, L.W., et al. (2013). Evaluative conditioning makes slim models less desirable as standards for comparison and increases body satisfaction. *Health Psychology*, 32(4), 433.
- Martin, L.E., Holsen, L.M., Chambers, R.J., et al. (2010). Neural mechanisms associated with food motivation in obese and healthy weight adults. *Obesity*, 18(2), 254–60.

- Matharu, K., Shapiro, J.F., Hammer, R.R., Kravitz, R.L., Wilson, M.D., Fitzgerald, F.T. (2014). Reducing obesity prejudice in medical education. *Education for Health: Change in Learning and Practice*, 27(3), 231–7.
- Mattavelli, G., Gallucci, A., Schiena, G., et al. (2019). Transcranial direct current stimulation modulates implicit attitudes towards food in eating disorders. *International Journal of Eating Disorders*, 52(5), 576–81.
- Mattavelli, G., Zuglian, P., Dabroi, E., Gaslini, G., Clerici, M., Papagno, C. (2015). Transcranial magnetic stimulation of medial prefrontal cortex modulates implicit attitudes towards food. *Appetite*, 89, 70–6.
- Mayer, B., Bos, A.E.R., Muris, P., Huijding, J., Vlieland, M. (2008). Does disgust enhance eating disorder symptoms? *Eating Behaviors*, 9(1), 124–7.
- McConnell, A.R., Dunn, E.W., Austin, S.N., Rawn, C.D. (2011). Blind spots in the search for happiness: implicit attitudes and nonverbal leakage predict affective forecasting errors. *Journal of Experimental Social Psychology*, 47(3), 628–34.
- McKenna, I., Hughes, S., Barnes-Holmes, D., De Schryver, M., Yoder, R., O'Shea, D. (2016). Obesity, food restriction, and implicit attitudes to healthy and unhealthy foods: lessons learned from the implicit relational assessment procedure. *Appetite*, 100, 41–54.
- Miller, D.P., Spangler, J.G., Vitolins, M.Z., et al. (2013). Are medical students aware of their anti-obesity bias? *Academic Medicine*, 88(7), 978.
- Misener, K., Libben, M. (2017). Risk for eating disorders modulates interpretation bias in a semantic priming task. *Body Image*, 21, 103–6.
- Moher, D., Shamseer, L., Clarke, M., et al. (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews*, 4(1), 1.
- Mohr, H.M., Röder, C., Zimmermann, J., Hummel, D., Negele, A., Grabhorn, R. (2011). Body image distortions in bulimia nervosa: investigating body size overestimation and body size satisfaction by fMRI. *NeuroImage*, 56(3), 1822–31.
- Moussally, J.M., Billieux, J., Mobbs, O., Rothen, S., Van der Linden, M. (2015). Implicitly assessed attitudes toward body shape and food: the moderating roles of dietary restraint and disinhibition. *Journal of Eating Disorders*, 3(1), 47.
- Nederkoorn, C., Houben, K., Hofmann, W., Roefs, A., Jansen, A. (2010). Control yourself or just eat what you like? Weight gain over a year is predicted by an interactive effect of response inhibition and implicit preference for snack foods. *Health Psychology*, 29(4), 389.
- Nurjannah, I., Siwi, S.M. (2017). Guidelines for analysis on measuring interrater reliability of nursing outcome classification. *International Journal of Research in Medical Sciences*, 5(4), 1169–75.
- O'Brien, K.S., Puhl, R.M., Latner, J.D., Mir, A.S., Hunter, J.A. (2010). Reducing anti-fat prejudice in preservice health students: a randomized trial. *Obesity*, 18, 2138–44.
- O'Brien, K.S., Latner, J.D., Halberstadt, J., Hunter, J.A., Anderson, J., Caputi, P. (2008). Do antifat attitudes predict anti-fat behaviors? *Obesity*, 16, s87–92.
- O'Brien, K.S., Hunter, J.A., Halberstadt, J., Anderson, J. (2007a). Body image and explicit and implicit anti-fat attitudes: the mediating role of physical appearance comparisons. *Body Image*, 4, 249–56.
- O'Brien, K.S., Hunter, J.A., Banks, M. (2007b). Implicit anti-fat bias in physical educators: physical attributes, ideology and socialization. *International Journal of Obesity*, 31, 308–14.
- Ouzzani, M., Hammady, H., Fedorowicz, Z., Elmagarmid, A. (2016). Rayyan—a web and mobile app for systematic reviews. *Systematic reviews*, 5(1), 210.
- Payne, B.K., Cheng, C.M., Govorun, O., Stewart, B.D. (2005). An inkblot for attitudes: affect misattribution as implicit measurement. *Journal of Personality and Social Psychology*, 89(3), 277–93.
- Papies, E.K., Stroebe, W., Aarts, H. (2009). Who likes it more? Restrained eaters' implicit attitudes towards food. *Appetite*, 53(3), 279–87.
- Parling, T., Cernvall, M., Stewart, I., Barnes-Holmes, D., Ghaderi, A. (2012). Using the implicit relational assessment procedure to compare implicit pro-thin/anti-fat attitudes of patients with anorexia nervosa and non-clinical controls. *Eating Disorders*, 20(2), 127–43.
- Pavlović, M., Žeželj, I., Marinković, M., Sućević, J. (2016). Implicit preference of sweets over fruit as a predictor of their actual consumption. *British Food Journal*, 118(10), 2567–80.
- Pearl, R.L., White, M.A., Grilo, C.M. (2013). Weight bias internalization, depression, and self-reported health among overweight binge eating disorder patients. *Obesity*, 22, 142–8.
- Pechey, R., Monsivais, P., Ng, Y.-L., Marteau, T.M. (2015). Why don't poor men eat fruit? Socioeconomic differences in motivations for fruit consumption. *Appetite*, 84, 271–9.
- Penney, E., Laws, C. (2013). Application of the MODE model to implicit weight prejudice and its influence on expressed and actual behavior among college students. *Journal of Applied Social Psychology*, 43, E229–36.
- Perugini, M. (2005). Predictive models of implicit and explicit attitudes. *British Journal of Social Psychology*, 44(1), 29–45.
- Phelan, S.M., Dovidio, J.F., Puhl, R.M., et al. (2014). Implicit and explicit weight bias in a national sample of 4,732 medical students: the medical student CHANGES study. *Obesity*, 4, 1201–8.
- Phelan, S.M., Burgess, D.J., Puhl, R., et al. (2015a). The adverse effect of weight stigma on the well-being of medical students with overweight or obesity: findings from a national survey. *Journal of General Internal Medicine*, 30(9), 1251–8.
- Phelan, S.M., Puhl, R.M., Burke, S.E., et al. (2015b). The mixed impact of medical school on medical students' implicit and explicit weight bias. *Medical Education*, 49, 983–92.
- Phelan, S.M., Burgess, D.J., Yeazel, M.W., Hellerstedt, W.L., Griffin, J.M., van Ryn, M. (2015c). Impact of weight bias and stigma on quality of care and outcomes for patients with obesity. *Obesity Reviews*, 16(4), 319–26.
- Pryor, J.B., Reeder, G.D., Wesselmann, E.D., Williams, K.D., Wirth, J.H. (2013). The influence of social norms upon behavioral expressions of implicit and explicit weight-related stigma in an interactive game. *The Yale Journal of Biology and Medicine*, 86, 189–201.
- Ragunathan, R., Naylor, R.W., Hoyer, W.D. (2006). The unhealthy = tasty intuition and its effects on taste inferences, enjoyment, and choice of food products. *Journal of Marketing*, 70(4), 170–84.
- Richard, A., Meule, A., Blechert, J. (2019). Implicit evaluation of chocolate and motivational need states interact in predicting chocolate intake in everyday life. *Eating Behaviors*, 33, 1–6.
- Ritzert, T.R., Anderson, L.M., Reilly, E.E., Gorrell, S., Forsyth, J.P., Anderson, D.A. (2016). Assessment of weight/shape implicit bias related to attractiveness, fear, and disgust. *The Psychological Record*, 66(3), 405–17.

- Robertson, N., Vohora, R. (2008). Fitness vs. fatness: implicit bias towards obesity among fitness professionals and regular exercisers. *Psychology of Sport and Exercise*, 9(4), 547–57.
- Robinson, E.L., Ball, L.E., Leveritt, M.D. (2014). Obesity bias among health and non-health students attending an Australian university and their perceived obesity education. *Journal of Nutrition Education and Behavior*, 46(5), 390–5.
- Roddy, S., Stewart, I., Barnes-Holmes, D. (2011). Facial reactions reveal that slim is good but fat is not bad: implicit and explicit measures of body-size bias. *European Journal of Social Psychology*, 41(6), 688–94.
- Roddy, S., Stewart, I., Barnes-Holmes, D. (2010). Anti-fat, pro-slim, or both? Using two reaction-time based measures to assess implicit attitudes to the slim and overweight. *Journal of Health Psychology*, 15(3), 416–25.
- Roefs, A., Quaedackers, L., Werrij, M.Q., et al. (2006). The environment influences whether high-fat foods are associated with palatable or with unhealthy. *Behaviour Research and Therapy*, 44(5), 715–36.
- Roefs, A., Herman, C.P., MacLeod, C.M., Smulders, F.T.Y., Jansen, A. (2005a). At first sight: how do restrained eaters evaluate high-fat palatable foods? *Appetite*, 44(1), 103–14.
- Roefs, A., Stapert, D., Isabella, L.A.S., Wolters, G., Wojciechowski, F., Jansen, A. (2005b). Early associations with food in anorexia nervosa patients and obese people assessed in the affective priming paradigm. *Eating Behaviors*, 6(2), 151–63.
- Roefs, A., Jansen, A. (2002). Implicit and explicit attitudes toward high-fat foods in obesity. *Journal of Abnormal Psychology*, 111(3), 517–21.
- Robstad, N., Siebler, F., Söderhamn, U., Westergren, T., Fegran, L. (2018). Design and psychometric testing of instruments to measure qualified intensive care nurses' attitudes toward obese intensive care patients. *Research in Nursing & Health*, 41(6), 525–34.
- Rudolph, A., Hilbert, A. (2017). The effects of obesity-related health messages on explicit and implicit weight bias. *Frontiers in Psychology*, 7, 2064.
- Rukavina, P.B., Li, W., Shen, B., Sun, H. (2010). A service learning based project to change implicit and explicit bias toward obese individuals in kinesiology pre-professionals. *Obesity Facts*, 3, 117–26.
- Russell-Mayhew, S., Nutter, S., Ireland, A., et al. (2015). Pilot testing a professional development model for preservice teachers in the area of health and weight: feasibility, utility, and efficacy. *Advances in School Mental Health Promotion*, 8(3), 176–86.
- Sabin, J.A., Moore, K., Noonan, C., Lallemand, O., Buchwald, D. (2015). Clinicians' implicit and explicit attitudes about weight and race and treatment approaches to overweight for American Indian children. *Childhood Obesity*, 11(4), 456–65.
- Sabin, J.A., Marini, M., Nosek, B.A. (2012). Implicit and explicit anti-fat bias among a large sample of medical doctors by BMI, race/ethnicity and gender. *PLoS One*, 7, e48448.
- Sartor, F., Donaldson, L.F., Markland, D.A., Loveday, H., Jackson, M.J., Kubis, H.P. (2011). Taste perception and implicit attitude toward sweet related to body mass index and soft drink supplementation. *Appetite*, 57(1), 237–46.
- Sato, W., Sawada, R., Kubota, Y., Toichi, M., Fushiki, T. (2017). Homeostatic modulation on unconscious hedonic responses to food. *BMC Research Notes*, 10(1), 1–6.
- Sato, W., Sawada, R., Kubota, Y., Toichi, M., Fushiki, T. (2016). Unconscious affective responses to food. *PLoS One*, 11(8), e0160956.
- Schakel, L., Veldhuijzen, D.S., Van Middendorp, H., et al. (2018). The effects of a gamified approach avoidance training and verbal suggestions on food outcomes. *PLoS One*, 13(7), 7–9.
- Schwartz, M.B., Vartanian, L.R., Nosek, B.A., Brownell, K.D. (2006). The influence of one's own body weight on implicit and explicit anti-fat bias. *Obesity*, 14, 440–7.
- Schwartz, M.B., Chambliss, H.O., Brownell, K.D., Blair, S.N., Billington, C. (2003). Weight bias among health professionals specializing in obesity. *Obesity Research*, 11, 1033–9.
- Scrivano, R.M., Scisco, J.L., Giumetti, G.W. (2017). The impact of applicants' weight and education about obesity on applicant ratings. *Psi Chi Journal of Psychological Research*, 22(4), 278–85.
- Seibt, B., Häfner, M., Deutsch, R. (2007). Prepared to eat: how immediate affective and motivational responses to food cues are influenced by food deprivation. *European Journal of Social Psychology*, 37(2), 359–79.
- Shamseer, L., Moher, D., Clarke, M., et al. (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*, 349; g7647.
- Skinner, A.C., Payne, K., Perrin, A.J., et al. (2017). Implicit weight bias in children age 9 to 11 years. *Pediatrics*, 140(1), e20163936.
- Smith, A.R., Forrest, L.N., Velkoff, E.A., Ribeiro, J.D., Franklin, J. (2018). Implicit attitudes toward eating stimuli differentiate eating disorder and non-eating disorder groups and predict eating disorder behaviors. *International Journal of Eating Disorders*, 51(4), 343–51.
- Smith, A.R., Joiner, T.E., Jr, Dodd, D.R. (2014). Examining implicit attitudes toward emaciation and thinness in anorexia nervosa. *International Journal of Eating Disorders*, 47(2), 138–47.
- Solbes, I., Enesco, I. (2010). Explicit and implicit anti-fat attitudes in children and their relationships with their body images. *Obesity Facts*, 3, 23–32.
- Songa, G., Russo, V. (2018). IAT, consumer behaviour and the moderating role of decision-making style: an empirical study on food products. *Food Quality and Preference*, 64, 205–20.
- Spring, V.L., Bulik, C.M. (2014). Implicit and explicit affect toward food and weight stimuli in anorexia nervosa. *Eating Behaviors*, 15(1), 91–4.
- Stafford, L.D., Scheffler, G. (2008). Hunger inhibits negative associations to food but not auditory biases in attention. *Appetite*, 51(3), 731–4.
- Storr, S.M., Sparks, P. (2016). Does self-affirmation following ego depletion moderate restrained eaters' explicit preferences for, and implicit associations with, high-calorie foods? *Psychology and Health*, 31(7), 840–56.
- Striegel-Moore, R.H., Rosselli, F., Perrin, N., et al. (2009). Gender difference in the prevalence of eating disorder symptoms. *International Journal of Eating Disorders*, 42(5), 471–4.
- Swift, J.A., Tischler, V., Markham, S., et al. (2013). Are anti-stigma films a useful strategy for reducing weight bias among trainee healthcare professionals? Results of a pilot randomized control trial. *Obesity Facts*, 6, 101–7.
- Teachman, B.A., Jeyaram, S., Gapinski, K.D., Brownell, K.D., Rawlins, M. (2003). Demonstrations of implicit anti-fat bias: the impact of providing causal information and evoking empathy. *Health Psychology*, 22, 68–78.
- Teachman, B.A., Brownell, K.D. (2001). Implicit anti-fat bias among health professionals: is anyone immune? *International Journal of Obesity and Related Metabolic Disorders*, 25, 1525–31.

- Terenzi, D., Rumiati, R.I., Catalan, M., et al. (2018). Reward sensitivity in Parkinson's patients with binge eating. *Parkinsonism & Related Disorders*, 51, 79–84.
- Thomas, S.R., Burton Smith, R., Ball, P.J. (2007). Implicit attitudes in very young children: an adaptation of the IAT. *Current Research in Social Psychology*, 13(7), 75–85.
- Tomiyaama, A.J., Finch, L.E., Incollingo Belsky, A.C., et al. (2015). Weight bias in 2001 versus 2013: contradictory attitudes among obesity researchers and health professionals. *Obesity*, 23, 46–53.
- Trendel, O., Werle, C.O.C. (2016). Distinguishing the affective and cognitive bases of implicit attitudes to improve prediction of food choices. *Appetite*, 104, 33–43.
- Van Dessel, P., Hughes, S., De Houwer, J. (2018). Consequence-based approach-avoidance training: a new and improved method for changing behavior. *Psychological Science*, 29(12), 1899–910.
- Val-Laillet, D., Aarts, E., Weber, B., et al. (2015). Neuroimaging and neuromodulation approaches to study eating behavior and prevent and treat eating disorders and obesity. *NeuroImage: Clinical*, 8, 1–31.
- Vallis, T.M., Currie, B., Lawlor, D., Ransom, T. (2007). Health-care professional bias against the obese: how do we know if we have a problem? *Canadian Journal of Diabetes*, 31(4), 365–70.
- Veenstra, E.M., de Jong, P.J. (2010). Restrained eaters show enhanced automatic approach tendencies towards food. *Appetite*, 55(1), 30–6.
- Venturini, B., Castelli, L., Tomelleri, S. (2006). Not all jobs are suitable for fat people: experimental evidence of a link between being fat and “out-of-sight” jobs. *Social Behavior and Personality*, 34(4), 389–98.
- Verbeken, S., Boendermaker, W.J., Loeyts, T., et al. (2018). Feasibility and effectiveness of adding an approach avoidance training with game elements to a residential childhood obesity treatment—a pilot study. *Behaviour Change*, 35(2), 91–107.
- Waller, T., Lampman, C., Lupfer-Johnson, G. (2012). Assessing bias against overweight individuals among nursing and psychology students: an implicit association test. *Journal of Clinical Nursing*, 21(23–24), 3504–12.
- Wang, Y., Zhu, J., Hu, Y., et al. (2016). The effect of implicit preferences on food consumption: moderating role of ego depletion and impulsivity. *Frontiers in Psychology*, 7, 1699.
- Wang, S.S., Brownell, K.D., Wadden, T.A. (2004). The influence of the stigma of obesity on overweight individuals. *International Journal of Obesity*, 28, 1333–7.
- Warschburger, P., Gmeiner, M., Morawietz, M., Rinck, M. (2018). Battle of plates: a pilot study of an approach-avoidance training for overweight children and adolescents. *Public Health Nutrition*, 21(2), 426–34.
- Watts, K., Cranney, J., Gleitzman, M. (2008). Automatic evaluation of body-related images. *Body Image*, 5(4), 352–64.
- Weinstein, J.H., Wilson, K.G., Drake, C.E., Kellum, K.K. (2008). A relational frame theory contribution to social categorization. *Behavior and Social Issues*, 17(1), 40–65.
- Werle, C.O., Trendel, O., Ardito, G. (2013). Unhealthy food is not tastier for everybody: the “healthy = tasty” French intuition. *Food Quality and Preference*, 28(1), 116–21.
- Werntz, A.J., Steinman, S.A., Glenn, J.J., Nock, M.K., Teachman, B.A. (2016). Characterizing implicit mental health associations across clinical domains. *Journal of Behavior Therapy and Experimental Psychiatry*, 52, 17–28.
- Werrij, M.Q., Roefs, A., Janssen, I., et al. (2009). Early associations with palatable foods in overweight and obesity are not disinhibition related but restraint related. *Journal of Behavior Therapy and Experimental Psychiatry*, 40(1), 136–46.
- Wijayatunga, N.N., Kim, Y., Butsch, W.S., Dhurandhar, E.J. (2019). The effects of a teaching intervention on weight bias among kinesiology undergraduate students. *International Journal of Obesity*, 43, 2273–81.
- Woodward, H.E., Treat, T.A., Cameron, C.D., Yegorova, V. (2017). Valence and arousal-based affective evaluations of foods. *Eating Behaviors*, 24, 26–33.
- Woodward, H.E., Treat, T.A. (2015). Unhealthy how?: Implicit and explicit affective evaluations of different types of unhealthy foods. *Eating Behaviors*, 17, 27–32.
- Yen, J.Y., Chang, S.J., Ko, C.H., et al. (2010). The high-sweet-fat food craving among women with premenstrual dysphoric disorder: emotional response, implicit attitude and rewards sensitivity. *Psychoneuroendocrinology*, 35(8), 1203–12.