

Article

Lipschitz Continuity Results for a Class of Parametric Variational Inequalities and Applications to Network Games

Mauro Passacantando ¹  and Fabio Raciti ^{2,*} 

¹ Department of Business and Law, University of Milan-Bicocca, Via degli Arcimboldi 8, 20126 Milan, Italy; mauro.passacantando@unimib.it

² Department of Mathematics and Computer Science, University of Catania, Viale A. Doria 6, 95125 Catania, Italy

* Correspondence: fabio.raciti@unict.it

Abstract: We consider a class of finite-dimensional variational inequalities where both the operator and the constraint set can depend on a parameter. Under suitable assumptions, we provide new estimates for the Lipschitz constant of the solution, which considerably improve previous ones. We then consider the problem of computing the mean value of the solution with respect to the parameter and, to this end, adapt an algorithm devised to approximate a Lipschitz function whose analytic expression is unknown, but can be evaluated in arbitrarily chosen sample points. Finally, we apply our results to a class of Nash equilibrium problems, and generalized Nash equilibrium problems on networks.

Keywords: parametric variational inequality; Lipschitz continuity; worst case error; network games; Nash equilibrium; generalized Nash equilibrium

1. Introduction

In many optimization or variational inequality (VI) problems, some data can depend on parameters, and it is important to analyze the regularity of the solution with respect to those parameters. For a detailed account of some classical results on the topic of parametric optimization, the reader can refer to [1]. The investigation of the Lipschitz behavior of solutions to convex minimization problems, from the point of view of set valued analysis, has also been carried out in the influential paper [2], and recently extended to abstract quasi-equilibrium problems in [3]. On a more applied context, in the case of linear programming, sharp Lipschitz constants for feasible and optimal solutions of some perturbed linear programs were derived in [4,5]. Another influential paper focusing on the estimate of Lipschitz constants for the solutions of a perturbed system of equalities and inequalities is [6], while in [7], the estimate of the Lipschitz constant is connected with of the role played by the norm of the inverse of a non-singular matrix in bounding the perturbation of the solution of a system of equations in terms of a right-hand side perturbation. The topic of parametric sensitivity analysis for variational inequalities has been developed by many authors in recent decades, and the excellent monograph [8] contains many related results. Let us notice, however, that while the number of papers dealing with local parametric differentiability for variational inequalities is very large, the investigation of parametric variational inequalities under the mere Lipschitz assumptions on data have received less attention, in particular with respect to the problem of estimating the global Lipschitz constant of the solution.

The local Lipschitz continuity of the solution of a variational inequality with parametric constraint set was investigated in [9], while more recently, the authors [10] derived an estimate of the global Lipschitz constant of the unique solution of a parametrized variational inequality, by making an extensive use of Lagrange multipliers techniques. Specifically, the Lagrange multiplier computations were used to derive the contribution of the parametric variation in the constraint set to the corresponding variation in the solution of the variational inequality. Interestingly enough, similar results were obtained in [11] with purely geometric methods.



Citation: Passacantando, M.; Raciti, F. Lipschitz Continuity Results for a Class of Parametric Variational Inequalities and Applications to Network Games. *Algorithms* **2023**, *16*, 458. <https://doi.org/10.3390/a16100458>

Academic Editor: Sorin-Mihai Grad

Received: 31 August 2023

Revised: 21 September 2023

Accepted: 25 September 2023

Published: 26 September 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

In [12], the authors improved the above-mentioned estimate but realized that, due to the complexity of the problem, their estimates were also far from being optimal.

In this paper, we derive new estimates for the Lipschitz constant of the solution of a parametric variational inequality, which considerably improve the previous ones. We then combine our results with a variant of the algorithm proposed in [13,14] to provide an approximation, on a given interval, of a univariate Lipschitz function whose analytic expression is unknown, but can be computed in arbitrarily chosen sample points (such a function is often called a black box-function). It turns out that the same algorithm also provides an approximation of the integral of the black-box function on the interval under consideration. Thus, we can use our estimate of the Lipschitz constant of the solution, to approximate its mean value on a given interval. Our results are then applied to investigate a class of Nash and generalized Nash equilibrium problems on networks.

The specific class of Nash equilibrium problems known as Network Games (or games played on networks) was investigated for the first time in the paper by Ballester et al. [15]. Players are modeled as nodes of a graph, and the social or economic relationship between any two players is represented by a link connecting the two corresponding nodes. Two players connected with a link are called neighbors. Considering linear-quadratic utility functions, the authors were able to express the unique Nash equilibrium as a series expansion involving the powers of the adjacency matrix of the graph, thus showing that, although players only interact with their neighbors, the equilibrium also depends on indirect contacts (i.e., neighbors of neighbors, neighbors of neighbors of neighbors, and so on). A wealth of social and economic processes may be modeled using the theory of Network Games; see, for instance, the papers [16–19], the survey [20] and the monograph [21]. The solution methods used in all the above-mentioned references combine the best response approach with some results of Graph Theory (see, e.g., [22]). It is worth noticing that although the connection between variational inequalities and Nash equilibrium problems was proven about forty years ago [23], very few papers have dealt with the variational inequality approach to Network Games and, in this regard, the interested reader can find a detailed investigation in [24]. Furthermore, the case of *generalized* network Nash equilibrium problems with shared constraints has not yet been investigated within the above-mentioned framework, with the only exception of the paper [25].

The paper is organized as follows. In Section 2, we provide estimates of the Lipschitz constant of the solution of our parametric VI in a general case, as well as in four different cases where the dependence on the parameter is more specific. In Section 3, we describe the algorithm used to approximate the solution components and their mean values. Section 4 is devoted to the application of our results to a class of Network Games, and is structured in three subsections. Specifically, in Section 4.1 we provide the basic concepts on Network Games, in Section 4.2 we describe the quadratic reference model of a network Nash equilibrium problem, while in Section 4.3 we first recall the definition of generalized Nash equilibrium problem with shared constraints, and the concept of variational solution, and then consider the quadratic reference model in this new setting. In Section 5, we apply our estimates and algorithm to some parametric network games. We then summarize our work and outline some future research perspectives in a small conclusion section. The paper ends with an appendix, where we derive some exact results that can be obtained in the special case where the Lipschitz constant of the operator of the variational inequality equals its strong monotonicity constant.

2. Problem Formulation and Lipschitz Constant Estimates

Let $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ and, for each $t \in [0, T]$, $K(t)$ be a nonempty, closed and convex subset of \mathbb{R}^n . We denote with $\langle \cdot, \cdot \rangle$ the scalar product in \mathbb{R}^n . The variational inequality on \mathbb{R}^n , with parameter $t \in [0, T]$, is the following problem: for each $t \in [0, T]$, find $x(t) \in K(t)$ such that:

$$\langle F(t, x(t)), y - x(t) \rangle \geq 0, \quad \forall y \in K(t). \quad (1)$$

In the case where, for each $t \in [0, T]$ the solution of (1) is unique, we will prove, assuming the Lipschitz continuity of F and of the parametric functions describing $K(t)$, the Lipschitz continuity of the solution map $t \mapsto x(t)$ and derive an a priori estimate of its Lipschitz constant Λ . Specifically, we estimate Λ in the general case where the variables t and x cannot be separated, as well as in the separable case where $F(t, x) = G(x) + H(t)$. Moreover, in the special circumstance where $K(t) = K$, for any $t \in [0, T]$, two simplified formulas are derived. We then focus on computing approximations for the mean value of the components $x_j : [0, T] \rightarrow \mathbb{R}, j = 1, \dots, n$:

$$\frac{1}{T} \int_0^T x_j(t) dt. \tag{2}$$

We recall here some useful monotonicity properties.

Definition 1. A map $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is monotone on a set $A \subset \mathbb{R}^n$ if and only if

$$\langle F(x) - F(y), x - y \rangle \geq 0, \quad \forall x, y \in A.$$

If the equality holds only when $x = y$, F is said to be strictly monotone.

A stronger type of monotonicity is given by the following definition.

Definition 2. $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is α -strongly monotone on A if and only if $\exists \alpha > 0$:

$$\langle F(x) - F(y), x - y \rangle \geq \alpha \|x - y\|^2, \quad \forall x, y \in A.$$

The concepts of strict and strong monotonicity coincide for linear operators and they are equivalent to the positive definiteness of the corresponding matrix. When F also depends on a parameter, the following property is useful.

Definition 3. $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is uniformly α -strongly monotone on A if and only if $\exists \alpha > 0$:

$$\langle F(t, x) - F(t, y), x - y \rangle \geq \alpha \|x - y\|^2 \quad \forall x, y \in A, \forall t \in [0, T].$$

Sufficient conditions for the existence and uniqueness of the solution of a variational inequality problem can be found in [8]. We mention here the following classical theorem which, in the parametric case, has to be applied for each $t \in [0, T]$.

Theorem 1. Let $K \subset \mathbb{R}^n$ be a closed and convex set and $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ continuous on K . If K is bounded, then the variational inequality problem $VI(F, K)$ admits at least one solution. If K is unbounded, then the existence of a solution is guaranteed by the following coercivity condition:

$$\lim_{\|x\| \rightarrow +\infty} \frac{\langle F(x) - F(x_0), x - x_0 \rangle}{\|x - x_0\|} = +\infty,$$

for $x \in K$ and some $x_0 \in K$. Furthermore, if F is α -strongly monotone on K , then the solution exists and is unique.

Let us now prove two theorems, along with three corresponding corollaries, which constitute the main contribution of the paper. Such results, at different levels of generality, ensure that the unique solution of (1) is Lipschitz continuous, and also provide an estimate of the Lipschitz constant of the solution.

Theorem 2 (General case). Assume that $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is uniformly α -strongly monotone on \mathbb{R}^n and F is Lipschitz continuous on $[0, T] \times \mathbb{R}^n$ with constant L , i.e.,

$$\|F(t_1, x_1) - F(t_2, x_2)\| \leq L(|t_1 - t_2| + \|x_1 - x_2\|), \quad \forall x_1, x_2 \in \mathbb{R}^n, \forall t_1, t_2 \in [0, T].$$

Moreover, we assume that $K(t)$ is a closed and convex set for any $t \in [0, T]$, and there exists $M \geq 0$ such that

$$\|p_{K(t_1)}(x) - p_{K(t_2)}(x)\| \leq M|t_1 - t_2|, \quad \forall x \in \mathbb{R}^n, \forall t_1, t_2 \in [0, T], \tag{3}$$

where $p_{K(t)}(x)$ denotes the projection of x on the closed convex set $K(t)$.

Then, for any $t \in [0, T]$, there exists a unique solution $x(t)$ of the VI($F, K(t)$) and $x(t)$ is Lipschitz continuous on $[0, T]$, with an estimated constant equal to

$$\Lambda_1 = \begin{cases} \inf_{z \in (0, 2\tilde{\alpha})} \left[\frac{M}{1-s} + \frac{z(1+z)}{s(1-s)} \right], & \text{if } \tilde{\alpha} < 1, \\ M + 2\sqrt{2M} + 1, & \text{if } \tilde{\alpha} = 1, \end{cases}$$

where $\tilde{\alpha} = \alpha/L$ and $s = \sqrt{z^2 - 2\tilde{\alpha}z + 1}$.

Proof. The existence and uniqueness of the solution $x(t)$ follows from Theorem 1. Given $t_1, t_2 \in [0, T]$, we denote $x_1 := x(t_1)$ and $x_2 := x(t_2)$. We know that

$$x_1 = p_{K(t_1)}(x_1 - \lambda F(t_1, x_1)), \quad x_2 = p_{K(t_2)}(x_2 - \lambda F(t_2, x_2)),$$

hold for any $\lambda > 0$, due to the characterization of the projection onto a closed convex set and the Brower fixed point theorem (see, e.g., [8]). Hence, we obtain the following estimate:

$$\begin{aligned} \|x_2 - x_1\| &= \|p_{K(t_2)}(x_2 - \lambda F(t_2, x_2)) - p_{K(t_1)}(x_1 - \lambda F(t_1, x_1))\| \\ &\leq \|p_{K(t_2)}(x_2 - \lambda F(t_2, x_2)) - p_{K(t_1)}(x_2 - \lambda F(t_2, x_2))\| \\ &\quad + \|p_{K(t_1)}(x_2 - \lambda F(t_2, x_2)) - p_{K(t_1)}(x_1 - \lambda F(t_1, x_1))\| \\ &\leq M|t_1 - t_2| + \|(x_2 - \lambda F(t_2, x_2)) - (x_1 - \lambda F(t_1, x_1))\|, \end{aligned} \tag{4}$$

where the second inequality follows from assumption (3) and the non-expansiveness of the projection map. Moreover, we have

$$\begin{aligned} &\|(x_2 - \lambda F(t_2, x_2)) - (x_1 - \lambda F(t_1, x_1))\|^2 \\ &= \|x_2 - x_1\|^2 + \lambda^2 \|F(t_2, x_2) - F(t_1, x_1)\|^2 - 2\lambda \langle x_2 - x_1, F(t_2, x_2) - F(t_1, x_1) \rangle \\ &\leq \|x_2 - x_1\|^2 + \lambda^2 L^2 (\|x_2 - x_1\| + |t_2 - t_1|)^2 - 2\lambda \langle x_2 - x_1, F(t_2, x_2) - F(t_1, x_1) \rangle \\ &= \|x_2 - x_1\|^2 + \lambda^2 L^2 (\|x_2 - x_1\| + |t_2 - t_1|)^2 - 2\lambda \langle x_2 - x_1, F(t_2, x_2) - F(t_2, x_1) \rangle \\ &\quad - 2\lambda \langle x_2 - x_1, F(t_2, x_1) - F(t_1, x_1) \rangle \\ &\leq \|x_2 - x_1\|^2 + \lambda^2 L^2 (\|x_2 - x_1\| + |t_2 - t_1|)^2 - 2\lambda \alpha \|x_2 - x_1\|^2 \\ &\quad + 2\lambda \|x_2 - x_1\| \|F(t_2, x_1) - F(t_1, x_1)\| \\ &= \|x_2 - x_1\|^2 + \lambda^2 L^2 (\|x_2 - x_1\| + |t_2 - t_1|)^2 - 2\lambda \alpha \|x_2 - x_1\|^2 \\ &\quad + 2\lambda L \|x_2 - x_1\| |t_2 - t_1| \\ &= (1 + \lambda^2 L^2 - 2\lambda \alpha) \|x_2 - x_1\|^2 + \lambda^2 L^2 |t_2 - t_1|^2 \\ &\quad + 2\lambda L (1 + \lambda L) \|x_2 - x_1\| |t_2 - t_1|. \end{aligned} \tag{5}$$

It is well-known that $\alpha \leq L$, since the assumptions guarantee that the following inequalities

$$\alpha \|x - y\|^2 \leq \langle F(t, x) - F(t, y), x - y \rangle \leq \|F(t, x) - F(t, y)\| \|x - y\| \leq L \|x - y\|^2$$

hold for any $x, y \in \mathbb{R}^n$ and $t \in [0, T]$. Thus, we obtain

$$1 + \lambda^2 L^2 - 2\lambda\alpha \geq 1 + \lambda^2 L^2 - 2\lambda L = (1 - \lambda L)^2 \geq 0,$$

hence $s := \sqrt{1 + \lambda^2 L^2 - 2\lambda\alpha}$ is well defined. Notice that if $\alpha < L$, then $s > 0$ for any $\lambda > 0$, while if $\alpha = L$ then $s > 0$ for any $\lambda \in (0, 1/L) \cup (1/L, +\infty)$. Moreover, we have

$$1 + \lambda L = \sqrt{1 + \lambda^2 L^2 + 2\lambda L} > \sqrt{1 + \lambda^2 L^2 - 2\lambda\alpha} = s,$$

thus, if $s > 0$, we have

$$\lambda^2 L^2 |t_2 - t_1|^2 \leq \left[\lambda L \left(\frac{1 + \lambda L}{s} \right) |t_2 - t_1| \right]^2. \tag{6}$$

It follows from (5) and (6) that

$$\begin{aligned} & \| (x_2 - \lambda F(t_2, x_2)) - (x_1 - \lambda F(t_1, x_1)) \|^2 \\ & \leq (s \|x_2 - x_1\|)^2 + \left[\lambda L \left(\frac{1 + \lambda L}{s} \right) |t_2 - t_1| \right]^2 + 2\lambda L(1 + \lambda L) \|x_2 - x_1\| |t_2 - t_1| \\ & = \left[s \|x_2 - x_1\| + \lambda L \left(\frac{1 + \lambda L}{s} \right) |t_2 - t_1| \right]^2, \end{aligned}$$

hence

$$\| (x_2 - \lambda F(t_2, x_2)) - (x_1 - \lambda F(t_1, x_1)) \| \leq s \|x_2 - x_1\| + \lambda L \left(\frac{1 + \lambda L}{s} \right) |t_2 - t_1|. \tag{7}$$

It follows from (4) and (7) that

$$(1 - s) \|x_2 - x_1\| \leq \left(M + \frac{\lambda L(1 + \lambda L)}{s} \right) |t_2 - t_1|.$$

Notice that $1 - s > 0$ if and only if $\lambda \in (0, 2\alpha/L^2)$. Therefore, in the case $\alpha < L$, we obtain

$$\|x_2 - x_1\| \leq \left[\frac{M}{1 - s} + \frac{\lambda L(1 + \lambda L)}{s(1 - s)} \right] |t_2 - t_1|, \quad \forall \lambda \in \left(0, \frac{2\alpha}{L^2} \right), \tag{8}$$

while in the case $\alpha = L$, we obtain

$$\|x_2 - x_1\| \leq \left[\frac{M}{1 - s} + \frac{\lambda L(1 + \lambda L)}{s(1 - s)} \right] |t_2 - t_1|, \quad \forall \lambda \in \left(0, \frac{1}{L} \right) \cup \left(\frac{1}{L}, \frac{2}{L} \right). \tag{9}$$

If we set $z = \lambda L$ and $\tilde{\alpha} = \alpha/L$, then we can write $s = \sqrt{z^2 - 2\tilde{\alpha}z + 1}$ and the estimate (8), which, in case $\tilde{\alpha} < 1$, reads as

$$\|x_2 - x_1\| \leq \inf_{z \in (0, 2\tilde{\alpha})} \left[\frac{M}{1 - s} + \frac{z(1 + z)}{s(1 - s)} \right] |t_2 - t_1|.$$

In case $\tilde{\alpha} = 1$, we have $s = |z - 1|$, and the estimate (9) reads as

$$\|x_2 - x_1\| \leq \left[\inf_{z \in (0, 1) \cup (1, 2)} f(z) \right] |t_2 - t_1|,$$

where

$$f(z) = \frac{M}{1 - |z - 1|} + \frac{z(1 + z)}{|z - 1|(1 - |z - 1|)} = \begin{cases} \frac{M}{z} + \frac{1 + z}{1 - z} & \text{if } z \in (0, 1), \\ \frac{M}{2 - z} + \frac{z(1 + z)}{(z - 1)(2 - z)} & \text{if } z \in (1, 2). \end{cases}$$

An analytical study of the function f (see Appendix A.1) provides

$$\inf_{z \in (0,1)} f(z) = M + 2\sqrt{2M} + 1,$$

and

$$\inf_{z \in (1,2)} f(z) = \frac{M^2 + 7M + 8 + 2\sqrt{2M + 12}}{M + 8 - 2\sqrt{2M + 12}}.$$

Moreover, it is easy to check that the following inequality

$$\inf_{z \in (1,2)} f(z) \geq \inf_{z \in (0,1)} f(z)$$

holds for any $M \geq 0$. Therefore, we obtain

$$\inf_{z \in (0,1) \cup (1,2)} f(z) = \inf_{z \in (0,1)} f(z) = M + 2\sqrt{2M} + 1,$$

thus the thesis follows. \square

Corollary 1 (General operator and constant feasible region). *Assume that F satisfies the same assumptions of Theorem 2 and the feasible region K does not depend on t . Then, the solution $x(t)$ of the VI(F, K) is Lipschitz continuous on $[0, T]$ with estimated constant equal to $\Lambda_1 = L/\alpha$.*

Proof. Since K does not depend on t , inequality (3) holds with $M = 0$. Hence, Theorem 2 guarantees that $x(t)$ is Lipschitz continuous on $[0, T]$ with a constant equal to

$$\begin{cases} \inf_{z \in (0,2\tilde{\alpha})} \frac{z(1 + z)}{s(1 - s)}, & \text{if } \tilde{\alpha} < 1, \\ 1, & \text{if } \tilde{\alpha} = 1, \end{cases}$$

where $\tilde{\alpha} = \alpha/L$ and $s = \sqrt{z^2 - 2\tilde{\alpha}z + 1}$. It is sufficient to prove that if $\tilde{\alpha} < 1$, then

$$\inf_{z \in (0,2\tilde{\alpha})} \frac{z(1 + z)}{s(1 - s)} = \frac{1}{\tilde{\alpha}}.$$

Since $\tilde{\alpha} > 0$, we have

$$(1 + \tilde{\alpha})z^2 - 2\tilde{\alpha}z + 1 + \tilde{\alpha} > 0, \quad \forall z \in \mathbb{R},$$

thus

$$(1 + \tilde{\alpha})z^2 \left[(1 + \tilde{\alpha})z^2 - 2\tilde{\alpha}z + 1 + \tilde{\alpha} \right] > 0, \quad \forall z > 0,$$

On the other hand, we have

$$\begin{aligned} & (1 + \tilde{\alpha})z^2 \left[(1 + \tilde{\alpha})z^2 - 2\tilde{\alpha}z + 1 + \tilde{\alpha} \right] \\ &= (1 + \tilde{\alpha})^2 z^4 - 2\tilde{\alpha}(1 + \tilde{\alpha})z^3 + (1 + \tilde{\alpha})^2 z^2 \\ &= (1 + \tilde{\alpha})^2 z^4 + \tilde{\alpha}^2 z^2 + 1 - 2\tilde{\alpha}(1 + \tilde{\alpha})z^3 + 2(1 + \tilde{\alpha})z^2 - 2\tilde{\alpha}z - z^2 + 2\tilde{\alpha}z - 1 \\ &= [(1 + \tilde{\alpha})z^2 - \tilde{\alpha}z + 1]^2 - z^2 + 2\tilde{\alpha}z - 1 \\ &= [(1 + \tilde{\alpha})z^2 - \tilde{\alpha}z + 1]^2 - s^2, \end{aligned}$$

hence

$$[(1 + \tilde{\alpha})z^2 - \tilde{\alpha}z + 1]^2 > s^2, \quad \forall z > 0.$$

Since $s > 0$ and $(1 + \tilde{\alpha})z^2 - \tilde{\alpha}z + 1 > 0$ hold for any $z > 0$, we obtain

$$(1 + \tilde{\alpha})z^2 - \tilde{\alpha}z + 1 > s, \quad \forall z > 0.$$

The latter inequality is equivalent to

$$\tilde{\alpha}z(1 + z) > s - z^2 + 2\tilde{\alpha}z - 1 = s(1 - s), \quad \forall z > 0.$$

Since $s > 0$ and $1 - s > 0$ hold for any $z \in (0, 2\tilde{\alpha})$, we obtain

$$\frac{z(1 + z)}{s(1 - s)} > \frac{1}{\tilde{\alpha}}, \quad \forall z \in (0, 2\tilde{\alpha}).$$

On the other hand, we have

$$\lim_{z \rightarrow 0} \frac{z(1 + z)}{s(1 - s)} = \lim_{z \rightarrow 0} \frac{z(1 + z)(1 + s)}{s(1 - s^2)} = \lim_{z \rightarrow 0} \frac{z(1 + z)(1 + s)}{sz(2\tilde{\alpha} - z)} = \lim_{z \rightarrow 0} \frac{(1 + z)(1 + s)}{s(2\tilde{\alpha} - z)} = \frac{1}{\tilde{\alpha}}.$$

Therefore, the proof is complete. \square

Theorem 3 (Separable operator). Assume that $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is separable, i.e.,

$$F(t, x) = G(x) + H(t), \quad \forall t \in [0, T], x \in \mathbb{R}^n,$$

where $G : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is α -strongly monotone on \mathbb{R}^n and Lipschitz continuous on \mathbb{R}^n with constant L_x , and $H : [0, T] \rightarrow \mathbb{R}^n$ is Lipschitz continuous on \mathbb{R}^n with constant L_t . Moreover, we assume that $K(t)$ is a closed and convex set for any $t \in [0, T]$, and there exists $M \geq 0$, such that (3) holds.

Then, for any $t \in [0, T]$, there exists a unique solution $x(t)$ of the VI($F, K(t)$), and $x(t)$ is Lipschitz continuous on $[0, T]$ with an estimated constant equal to

$$\Lambda_2 = \begin{cases} \inf_{z \in (0, 2\hat{\alpha})} \left[\frac{M}{1 - \hat{s}} + \frac{\hat{L}z(1 + z)}{\hat{s}(1 - \hat{s})} \right], & \text{if } \hat{\alpha} < 1, \\ M + 2\sqrt{2M\hat{L}} + \hat{L}, & \text{if } \hat{\alpha} = 1, \end{cases}$$

where $\hat{\alpha} = \alpha / L_x$, $\hat{L} = L_t / L_x$ and $\hat{s} = \sqrt{z^2 - 2\hat{\alpha}z + 1}$.

Proof. The existence and uniqueness of the solution $x(t)$ follows from Theorem 1. Given $t_1, t_2 \in [0, T]$, we denote $x_1 := x(t_1)$ and $x_2 := x(t_2)$. For any $\lambda > 0$, we have the following estimate:

$$\begin{aligned} \|x_2 - x_1\| &= \|p_{K(t_2)}(x_2 - \lambda F(t_2, x_2)) - p_{K(t_1)}(x_1 - \lambda F(t_1, x_1))\| \\ &\leq \|p_{K(t_2)}(x_2 - \lambda F(t_2, x_2)) - p_{K(t_1)}(x_2 - \lambda F(t_2, x_2))\| \\ &\quad + \|p_{K(t_1)}(x_2 - \lambda F(t_2, x_2)) - p_{K(t_1)}(x_1 - \lambda F(t_1, x_1))\| \\ &\leq M|t_1 - t_2| + \|(x_2 - \lambda F(t_2, x_2)) - (x_1 - \lambda F(t_1, x_1))\|. \end{aligned} \tag{10}$$

Moreover, we have

$$\begin{aligned}
 & \| (x_2 - \lambda F(t_2, x_2)) - (x_1 - \lambda F(t_1, x_1)) \|^2 \\
 &= \|x_2 - x_1\|^2 + \lambda^2 \|F(t_2, x_2) - F(t_1, x_1)\|^2 - 2\lambda \langle x_2 - x_1, F(t_2, x_2) - F(t_1, x_1) \rangle \\
 &= \|x_2 - x_1\|^2 + \lambda^2 \|G(x_2) - G(x_1) + H(t_2) - H(t_1)\|^2 - 2\lambda \langle x_2 - x_1, G(x_2) - G(x_1) \\
 &\quad - 2\lambda \langle x_2 - x_1, H(t_2) - H(t_1) \rangle \\
 &\leq \|x_2 - x_1\|^2 + \lambda^2 (L_x \|x_2 - x_1\| + L_t |t_2 - t_1|)^2 - 2\lambda \langle x_2 - x_1, G(x_2) - G(x_1) \\
 &\quad - 2\lambda \langle x_2 - x_1, H(t_2) - H(t_1) \rangle \\
 &\leq \|x_2 - x_1\|^2 + \lambda^2 (L_x \|x_2 - x_1\| + L_t |t_2 - t_1|)^2 - 2\lambda \alpha \|x_2 - x_1\|^2 \\
 &\quad + 2\lambda \|x_2 - x_1\| \|H(t_2) - H(t_1)\| \\
 &\leq \|x_2 - x_1\|^2 + \lambda^2 (L_x \|x_2 - x_1\| + L_t |t_2 - t_1|)^2 - 2\lambda \alpha \|x_2 - x_1\|^2 \\
 &\quad + 2\lambda L_t \|x_2 - x_1\| |t_2 - t_1| \\
 &= (1 + \lambda^2 L_x^2 - 2\lambda \alpha) \|x_2 - x_1\|^2 + \lambda^2 L_t^2 |t_2 - t_1|^2 \\
 &\quad + 2\lambda L_t (1 + \lambda L_x) \|x_2 - x_1\| |t_2 - t_1|. \tag{11}
 \end{aligned}$$

Since $\alpha \leq L_x$, we have

$$1 + \lambda^2 L_x^2 - 2\lambda \alpha \geq 1 + \lambda^2 L_x^2 - 2\lambda L_x = (1 - \lambda L_x)^2 \geq 0,$$

hence $\hat{s} := \sqrt{1 + \lambda^2 L_x^2 - 2\lambda \alpha}$ is well defined. Notice that if $\alpha < L_x$, then $s > 0$ for any $\lambda > 0$, while if $\alpha = L_x$, then $s > 0$ for any $\lambda \in (0, 1/L_x) \cup (1/L_x, +\infty)$. Moreover, we have

$$1 + \lambda L_x = \sqrt{1 + \lambda^2 L_x^2 + 2\lambda L_x} > \sqrt{1 + \lambda^2 L_x^2 - 2\lambda \alpha} = \hat{s},$$

thus, if $\hat{s} > 0$, we have

$$\lambda^2 L_t^2 |t_2 - t_1|^2 \leq \left[\lambda L_t \left(\frac{1 + \lambda L_x}{\hat{s}} \right) |t_2 - t_1| \right]^2. \tag{12}$$

It follows from (11) and (12) that

$$\begin{aligned}
 & \| (x_2 - \lambda F(t_2, x_2)) - (x_1 - \lambda F(t_1, x_1)) \|^2 \\
 &\leq (\hat{s} \|x_2 - x_1\|)^2 + \left[\lambda L_t \left(\frac{1 + \lambda L_x}{\hat{s}} \right) |t_2 - t_1| \right]^2 + 2\lambda L_t (1 + \lambda L_x) \|x_2 - x_1\| |t_2 - t_1| \\
 &= \left[\hat{s} \|x_2 - x_1\| + \lambda L_t \left(\frac{1 + \lambda L_x}{\hat{s}} \right) |t_2 - t_1| \right]^2,
 \end{aligned}$$

hence

$$\| (x_2 - \lambda F(t_2, x_2)) - (x_1 - \lambda F(t_1, x_1)) \| \leq \hat{s} \|x_2 - x_1\| + \lambda L_t \left(\frac{1 + \lambda L_x}{\hat{s}} \right) |t_2 - t_1|. \tag{13}$$

It follows from (10) and (13) that

$$(1 - \hat{s}) \|x_2 - x_1\| \leq \left(M + \frac{\lambda L_t (1 + \lambda L_x)}{\hat{s}} \right) |t_2 - t_1|.$$

Notice that $1 - \hat{s} > 0$ if and only if $\lambda \in (0, 2\alpha/L_x^2)$. Therefore, in the case $\alpha < L_x$, we obtain

$$\|x_2 - x_1\| \leq \frac{M\hat{s} + \lambda L_t (1 + \lambda L_x)}{\hat{s}(1 - \hat{s})} |t_2 - t_1|, \quad \forall \lambda \in \left(0, \frac{2\alpha}{L_x^2} \right), \tag{14}$$

while in the case $\alpha = L_x$, we obtain

$$\|x_2 - x_1\| \leq \frac{M\hat{s} + \lambda L_t(1 + \lambda L_x)}{\hat{s}(1 - \hat{s})} |t_2 - t_1|, \quad \forall \lambda \in \left(0, \frac{1}{L_x}\right) \cup \left(\frac{1}{L_x}, \frac{2}{L_x}\right). \quad (15)$$

If we set $z = \lambda L_x$, $\hat{\alpha} = \alpha/L_x$ and $\hat{L} = L_t/L_x$, then we can write $\hat{s} = \sqrt{z^2 - 2\hat{\alpha}z + 1}$, and the estimate (14), in the case $\hat{\alpha} < 1$, reads as

$$\|x_2 - x_1\| \leq \inf_{z \in (0, 2\hat{\alpha})} \left[\frac{M}{1 - \hat{s}} + \frac{\hat{L}z(1 + z)}{\hat{s}(1 - \hat{s})} \right] |t_2 - t_1|.$$

In the case $\hat{\alpha} = 1$, we have $\hat{s} = |z - 1|$ and the estimate (15) reads as

$$\|x_2 - x_1\| \leq \left[\inf_{z \in (0,1) \cup (1,2)} f(z) \right] |t_2 - t_1|,$$

where

$$f(z) = \frac{M}{1 - |z - 1|} + \frac{\hat{L}z(1 + z)}{|z - 1|(1 - |z - 1|)} = \begin{cases} \frac{M}{z} + \frac{\hat{L}(1 + z)}{1 - z} & \text{if } z \in (0, 1), \\ \frac{M}{2 - z} + \frac{\hat{L}z(1 + z)}{(z - 1)(2 - z)} & \text{if } z \in (1, 2). \end{cases}$$

An analytical study of the function f (see Appendix A.2) provides

$$\inf_{z \in (0,1)} f(z) = M + 2\sqrt{2M\hat{L}} + \hat{L},$$

and

$$\inf_{z \in (1,2)} f(z) = \frac{M^2 + 7M\hat{L} + 8\hat{L}^2 + 2\hat{L}\sqrt{2M\hat{L} + 12\hat{L}^2}}{M + 8\hat{L} - 2\sqrt{2M\hat{L} + 12\hat{L}^2}}.$$

Moreover, it is easy to check that the following inequality

$$\inf_{z \in (1,2)} f(z) \geq \inf_{z \in (0,1)} f(z)$$

holds for any $M \geq 0$ and $\hat{L} > 0$. Therefore, we obtain

$$\inf_{z \in (0,1) \cup (1,2)} f(z) = \inf_{z \in (0,1)} f(z) = M + 2\sqrt{2M\hat{L}} + \hat{L},$$

thus the thesis follows. \square

Corollary 2 (Separable operator and constant feasible region). *Assume that F satisfies the assumptions of Theorem 3 and the feasible region K does not depend on t . Then, the solution $x(t)$ of the VI(F, K) is Lipschitz continuous on $[0, T]$ with estimated constant equal to $\Lambda_2 = L_t/\alpha$.*

Proof. Theorem 3 guarantees that $x(t)$ is Lipschitz continuous on $[0, T]$ with a constant equal to

$$\begin{cases} \inf_{z \in (0, 2\hat{\alpha})} \frac{\hat{L}z(1 + z)}{\hat{s}(1 - \hat{s})}, & \text{if } \hat{\alpha} < 1, \\ \hat{L} = L_t/\alpha, & \text{if } \hat{\alpha} = 1, \end{cases}$$

where $\hat{\alpha} = \alpha/L_x$, $\hat{L} = L_t/L_x$ and $\hat{s} = \sqrt{z^2 - 2\hat{\alpha}z + 1}$. Moreover, similarly to the proof of Corollary 1, where $\tilde{\alpha}$ is replaced by $\hat{\alpha}$, it can be proved that if $\hat{\alpha} < 1$, then

$$\inf_{z \in (0, 2\hat{\alpha})} \frac{\hat{L}z(1+z)}{\hat{s}(1-\hat{s})} = \frac{\hat{L}}{\hat{\alpha}} = \frac{L_t}{\alpha}.$$

□

In the special case where F does not depend on t , we obtain the same estimate proved in [12].

Corollary 3 (Constant operator). *Assume that F and K satisfy the assumptions of Theorem 3 and F does not depend on t , i.e., $H(t) \equiv 0$. Then, the solution $x(t)$ of the VI(F, K) is Lipschitz continuous on $[0, T]$ with estimated constant equal to*

$$\Lambda_2 = \frac{M}{1 - \sqrt{1 - \left(\frac{\alpha}{L}\right)^2}}.$$

Proof. If $H(t) \equiv 0$, then $L_t = 0$, and Theorem 3 guarantees that $x(t)$ is Lipschitz continuous on $[0, T]$ with a constant equal to

$$\begin{cases} \inf_{z \in (0, 2\hat{\alpha})} \frac{M}{1 - \hat{s}}, & \text{if } \hat{\alpha} < 1, \\ M, & \text{if } \hat{\alpha} = 1, \end{cases}$$

where $\hat{\alpha} = \alpha/L_x = \alpha/L$ and $\hat{s} = \sqrt{z^2 - 2\hat{\alpha}z + 1}$.

Moreover, it is easy to check that

$$\inf_{z \in (0, 2\hat{\alpha})} \frac{M}{1 - \hat{s}} = \frac{M}{1 - \sqrt{1 - \hat{\alpha}^2}} = \frac{M}{1 - \sqrt{1 - \left(\frac{\alpha}{L}\right)^2}},$$

thus the thesis follows. □

Table 1 summarizes the estimate of the Lipschitz constant of $x(t)$ depending on the features of the map F and the feasible set K .

Table 1. Estimate of the Lipschitz constant of $x(t)$ depending on the features of the map F and the feasible set K .

	General K	Constant K
General F	$\begin{cases} \inf_{z \in (0, 2\tilde{\alpha})} \left[\frac{M}{1-s} + \frac{z(1+z)}{s(1-s)} \right], & \text{if } \tilde{\alpha} < 1, \\ M + 2\sqrt{2M} + 1, & \text{if } \tilde{\alpha} = 1, \end{cases}$ where $\tilde{\alpha} = \frac{\alpha}{L}, s = \sqrt{z^2 - 2\tilde{\alpha}z + 1}$	$\Lambda_1 = \frac{L}{\alpha}$
Separable F	$\begin{cases} \inf_{z \in (0, 2\hat{\alpha})} \left[\frac{M}{1-\hat{s}} + \frac{\hat{L}z(1+z)}{\hat{s}(1-\hat{s})} \right], & \text{if } \hat{\alpha} < 1, \\ M + 2\sqrt{2M\hat{L}} + \hat{L}, & \text{if } \hat{\alpha} = 1, \end{cases}$ where $\hat{\alpha} = \frac{\alpha}{L_x}, \hat{L} = \frac{L_t}{L_x}, \hat{s} = \sqrt{z^2 - 2\hat{\alpha}z + 1}$	$\Lambda_2 = \frac{L_t}{\alpha}$
Constant F	$\Lambda_2 = \frac{M}{1 - \sqrt{1 - \left(\frac{\alpha}{L}\right)^2}}$	–

Theorems 2 and 3, along with Corollaries 1, 2 and 3, do not provide the solution $x(t)$ of the parametric VI in closed form, but guarantee that it is a Lipschitz continuous function of the parameter t and provide an estimate of the corresponding constant. The knowledge of the Lipschitz constant is a key tool to provide lower and upper approximations of both the solution $x(t)$ and its mean value on $[0, T]$, as the algorithm described in the next section shows.

3. Approximation Algorithm

Let $x_j : [0, T] \rightarrow \mathbb{R}$ be any component of the solution vector x of (1); assume that x_j is a Lipschitz continuous function and Λ is an estimate of its Lipschitz constant, i.e.,

$$|x_j(t_1) - x_j(t_2)| \leq \Lambda |t_1 - t_2|, \quad \forall t_1, t_2 \in [0, T]. \tag{16}$$

Furthermore, let I be the integral of x_j over $[0, T]$ and, to begin with, assume that we only know the value of x_j at the two endpoints $0, T$. We wish to construct an approximation of x_j in the interval $[0, T]$ and also obtain an estimate \hat{I} of the value of I , together with an estimate of the error $|I - \hat{I}|$.

The Lipschitz continuity of x_j allow us to localize its graph by drawing four lines passing trough the points $(0, x_j(0))$ and $(T, x_j(T))$, with slopes Λ and $-\Lambda$. We thus construct a parallelogram $P_{0,T}$ containing the unknown graph of the function (see Figure 1, left). Specifically, the two upper and the two lower sides of the parallelogram (dashed lines) correspond, respectively, to an *upper bound* x_j^u and a *lower bound* x_j^l for x_j . The functions x_j^u and x_j^l can be written as follows:

$$\begin{aligned} x_j^u(t) &= \min\{x_j(0) + \Lambda t, x_j(T) - \Lambda(t - T)\}, \\ x_j^l(t) &= \max\{x_j(0) - \Lambda t, x_j(T) + \Lambda(t - T)\}. \end{aligned}$$




Figure 1. Upper and lower bounds (dashed lines) and best approximation (solid line) of a Lipschitz continuous function evaluated at $t = 0$ and $t = T$ (left), and same curves after the evaluation at the midpoint $t = T/2$.

Denote now with $\hat{x}_j(t)$ an approximation of the value of x_j at some $t \in [0, T]$, with $\delta(t) = |x_j(t) - \hat{x}_j(t)|$ the error on the estimate, and with $e_w(t) = \max\{x_j^u(t) - \hat{x}_j(t), \hat{x}_j(t) - x_j^l(t)\}$ the largest possible absolute error on the estimate, whence:

$$\delta(t) \leq e_w(t),$$

where the equality sign holds when x_j coincides with its upper or lower bound. The case where $\delta(t) = e_w(t)$ can be called the *worst case*, and the value of $\hat{x}_j(t)$ that minimizes $e_w(t)$ is the mean value

$$\hat{x}_j(t) = \frac{x_j^u(t) + x_j^l(t)}{2}, \tag{17}$$

which yields

$$e_w(t) = \frac{x_j^u(t) - x_j^l(t)}{2}. \tag{18}$$

The graph of $\hat{x}_j(t)$, in the worst case, is represented by the solid line in Figure 1, on the left. In order to obtain a global measure of the error in the interval $[0, T]$, in the worst case, it is then natural to define:

$$E_w(0, T) = \int_0^T e_w(t) dt = \int_0^T \frac{x_j^u(t) - x_j^l(t)}{2} dt, \tag{19}$$

which is equal to the half of the area of $P_{0,T}$, denoted by $A(0, T)$, and thus $E_w(0, T) = A(0, T)/2$.

We thus have an approximation \hat{x}_j of the function x_j , together with the estimates e_w and E_w of the *worst case* local and total error, respectively. The average worst case error is defined as $E_w(0, T)/T$.

We want now to approximate the integral I of x_j over $[0, T]$. Notice that lower and upper bounds for I are given, respectively, by

$$I^l = \int_0^T x_j^l(t) dt = \frac{(x_j(0) - x_j(T))^2}{4\Lambda} - \frac{\Lambda}{4}T^2 + \frac{T(x_j(0) + x_j(T))}{2},$$

$$I^u = \int_0^T x_j^u(t) dt = -\frac{(x_j(0) - x_j(T))^2}{4\Lambda} + \frac{\Lambda}{4}T^2 + \frac{T(x_j(0) + x_j(T))}{2}.$$

The estimate of I that differs the less from the two extreme I^l and I^u is the average value

$$\hat{I} = \frac{I^l + I^u}{2} = \frac{T(x_j(0) + x_j(T))}{2}.$$

For this choice of \hat{I} , the error $|I - \hat{I}|$ can reach the value of $E_I = (I^u - I^l)/2$ in the worst case.

It follows from (17) that \hat{I} is equal to the integral of the approximating function \hat{x}_j . We remark from (19) that the total worst case error E_w coincides with the estimate of the error E_I on the integral \hat{I} . Hence, both for the estimate of the function and of its integral, the quantity that measures the corresponding errors is the same, and it is exactly half the area of the parallelogram $P_{0,T}$. It is easy to check (see [13,14]) that, for any pair of points t_1, t_2 (with $t_1 < t_2$), the area of parallelogram P_{t_1,t_2} is

$$A(t_1, t_2) = \int_{t_1}^{t_2} (x_j^u(t) - x_j^l(t)) dt = \frac{\Lambda^2(t_1 - t_2)^2 - (x_j(t_1) - x_j(t_2))^2}{2\Lambda}. \tag{20}$$

Now, we want to analyze how the evaluation of x_j in a further point improves the approximation procedure. For this, let $t_1 = 0, t_2 = T$, let t_3 be an arbitrary point in $(0, T)$ and repeat the previous procedure in the intervals $[0, t_3]$ and $[t_3, T]$, respectively. The graph of x_j is now contained in the union of parallelograms P_{0,t_3} and $P_{t_3,T}$. Therefore, a new sample point allows to better locate the graph of x_j . In this case, the bounding functions are computed as follows:

$$x_j^u(t) = \begin{cases} \min\{x_j(0) + \Lambda't, x_j(t_3) - \Lambda'(t - t_3)\} & \text{if } t \in [0, t_3], \\ \min\{x_j(t_3) + \Lambda''(t - t_3), x_j(T) - \Lambda''(t - T)\} & \text{if } t \in [t_3, T], \end{cases}$$

$$x_j^l(t) = \begin{cases} \max\{x_j(0) - \Lambda' t, x_j(t_3) + \Lambda'(t - t_3)\} & \text{if } t \in [0, t_3], \\ \max\{x_j(t_3) - \Lambda''(t - t_3), x_j(T) + \Lambda''(t - T)\} & \text{if } t \in [t_3, T], \end{cases}$$

where Λ' and Λ'' denote the update estimates of the Lipschitz constant of x_j in the intervals $[0, t_3]$ and $[t_3, T]$, respectively.

For the two sub-intervals, we can provide approximations of x_j and of its integral, and estimates of the corresponding errors $E_w(t_1, t_3)$ and $E_w(t_3, t_2)$.

We observe that the smaller the sum of the areas of the new parallelograms is, the better the new estimate of the function is. Hence, with the evaluation of x_j at new points and repeating the above procedure, we can further improve the approximation x_j and its integral, as specified by the following algorithm.

Step 0. (Initialization)

Fix a positive constant ε_{tot} , which represents the desired maximum error on the integral of x_j (or total worst case error in the estimate of x_j itself). Create a list of evaluation points, which initially contains the two endpoints 0 and T , and the corresponding list of function values $x_j(0), x_j(T)$. Calculate the area $A(0, T)$ of parallelogram $P_{0,T}$ by means of (20) and initialize a list of parallelograms' areas.

Step 1. (New evaluation)

Locate the parallelogram having the largest area, and the corresponding interval, say $[t_1, t_2]$ (at the first step, $[t_1, t_2]$ coincides with the whole interval $[0, T]$). Add the new evaluation point $p = (t_1 + t_2)/2$, and update the lists of points and function values accordingly. Compute the two Lipschitz constants in the intervals $[t_1, p]$ and $[p, t_2]$. Update the parallelograms' areas list by removing the area of parallelogram P_{t_1,t_2} and inserting the areas of the two new parallelograms $P_{t_1,p}$ and P_{p,t_2} .

Step 2. (Stopping rule)

Compute the worst case error E_{tot} over the interval $[0, T]$:

$$E_{tot} = (\text{sum of the areas of all parallelograms}) / 2.$$

If $E_{tot} \leq \varepsilon_{tot}$, then stop the procedure, otherwise go to Step 1.

At the k -th iteration of the algorithm, we thus have divided the interval $[0, T]$ with $k + 2$ points, which we reorder as $\{t_1 = 0, t_2, \dots, t_{k+1}, t_{k+2} = T\}$, and we denote with Λ_i the estimate of the Lipschitz constant of x_j in $[t_i, t_{i+1}]$. The bounding functions in $[t_i, t_{i+1}]$ are then given by:

$$X_j^u(t)_{|[t_i,t_{i+1}]} = \min_{t \in [t_i,t_{i+1}]} \{x_j(t_i) + \Lambda_i(t - t_i), x_j(t_{i+1}) - \Lambda_i(t - t_{i+1})\},$$

$$X_j^l(t)_{|[t_i,t_{i+1}]} = \max_{t \in [t_i,t_{i+1}]} \{x_j(t_i) - \Lambda_i(t - t_i), x_j(t_{i+1}) + \Lambda_i(t - t_{i+1})\}.$$

In the next result, we provide a complexity analysis of the algorithm.

Theorem 4. *After k iterations of the algorithm, the worst case error E_{tot} is, at most, equal to*

$$\frac{E_0}{2^{\lfloor \log_2(k+1) \rfloor}} \tag{21}$$

where

$$E_0 = \frac{\Lambda^2 T^2 - (x_j(0) - x_j(T))^2}{4\Lambda}$$

is the worst case error before the algorithm starts. Therefore, the algorithm stops after at most $\lceil 2E_0/\varepsilon_{tot} \rceil$ iterations.

Proof. At each iteration, we need to compute the two new estimates of the Lipschitz constant in the intervals $[t_1, p]$ and $[p, t_2]$. Such estimates are less than or equal to the

Lipschitz estimate Λ in the interval $[0, T]$. Since the area of each parallelogram given by Formula (20) (and thus the error on the integral) is an increasing function of Λ , the worst case occurs when the two estimates coincide with Λ . Hence, the worst case complexity analysis of the algorithm reduces to that of the algorithm where the Lipschitz estimate is not updated at any iteration. Therefore, following the same argument as in [13], we can prove the inequality

$$\frac{A_k}{A_0} \leq \frac{1}{2^i}, \tag{22}$$

where A_k denotes the sum of the areas of all parallelograms generated by the algorithm after k iterations, A_0 is the area of the parallelogram before the algorithm starts, and i is the integer such that $2^i \leq k + 1 < 2^{i+1}$. Since the worst case error E_{tot} after k iterations is $E_{tot} = A_k/2$, $E_0 = A_0/2$, and $i = \lfloor \log_2(k + 1) \rfloor$, we obtain the following inequality from (22):

$$E_{tot} = \frac{A_k}{2} \leq \frac{A_0/2}{2^i} = \frac{E_0}{2^{\lfloor \log_2(k+1) \rfloor}},$$

thus (21) holds.

If we denote $\bar{k} = \lceil 2E_0/\varepsilon_{tot} \rceil$, then

$$\bar{k} + 1 > \left\lceil \frac{2E_0}{\varepsilon_{tot}} \right\rceil \geq \frac{2E_0}{\varepsilon_{tot}},$$

thus, we obtain

$$\varepsilon_{tot} > \frac{2E_0}{\bar{k} + 1} = \frac{E_0}{2^{\log_2(\bar{k}+1)-1}} \geq \frac{E_0}{2^{\lfloor \log_2(\bar{k}+1) \rfloor}},$$

hence (21) guarantees that the worst case error after \bar{k} iterations is less than ε_{tot} ; that is, the algorithm stops after at most \bar{k} iterations. \square

4. Application to Parametric Network Games

The theoretical results proved in Section 2 and the approximation algorithm described in Section 3 can be applied to any parametric VI in the form (1). It is well known that parametric VIs in the form (1) can be viewed as the dynamic equilibrium models, in the case where the problem data depend on a time parameter t . Several applications of these models have been studied in recent years (see, e.g., [26,27] and references therein). On the other hand, network games are an interesting class of non-cooperative games with an underlying network structure, which can be modeled through VIs [24]. A number of applications of network games have been analyzed in the literature, from social networks to juvenile delinquency (see, e.g., [20,21] and references therein).

In this section, we apply the results of Sections 2 and 3 to a class of parametric network games, which can be modeled as parametric VIs in the form (1), where both the players' utility functions and their strategy sets depend on a time parameter. Specifically, in Section 4.1, a brief recall of Graph Theory and Game Theory concepts is given together with the definition of parametric network games. Then, we focus on the linear-quadratic model, both in the framework of the network Nash equilibrium problem (Section 4.2) and the network generalized Nash equilibrium problem (Section 4.3).

4.1. Basics of Network Games

We begin with a recapitulation of foundational Graph Theory concepts. We define a graph g as a pair of sets (V, E) , where V represents the set of nodes, while E represents the set of arcs composed of pairs of nodes (v, w) . Arcs sharing identical terminal nodes are categorized as parallel arcs, while arcs in the form of (v, v) are designated as loops. For the context at hand, we exclusively consider simple graphs, which are characterized by the absence of both parallel arcs and loops. Within our framework, the players are represented by the n nodes of the graph. Furthermore, we operate within the framework of undirected

graphs, wherein arcs (v, w) and (w, v) are considered equivalent. Two nodes v and w are adjacent if they are connected by the arc (v, w) . The information regarding the adjacency of nodes can be stored in the form of the adjacency matrix G , where the elements g_{ij} assume a value of 1 when (v_i, v_j) is an arc and 0 otherwise. Notice that matrix G is symmetric and with diagonal elements equal to 0. For a specific node v , the nodes that share a direct link with it through arcs are referred to as the neighbors of v . These neighbors are collated within the set $N_v(g)$. The cardinality of $N_v(g)$ is the degree of v .

A walk is defined as a sequence $v_0, e_1, v_1, e_2, \dots, v_k$ comprising both graph nodes v_i and graph edges e_i , for any $1 \leq i \leq k$, where the arc e_i has endpoints v_{i-1} and v_i . Notice that within a walk, the possibility of revisiting a node or traversing an arc multiple times is permitted. The length of a walk is equal to the number of its arcs. The indirect connections between any two nodes within the graph are described through the utilization of the powers of the adjacency matrix G . Indeed, it can be proved that the element $g_{ij}^{[k]}$ of G^k provides the number of walks of length k between the nodes v_i and v_j .

We now recall some concepts of Game Theory, which will then be adapted to our specific framework. A strategic game (in normal form) is a model of interaction among decision makers, which are usually called players. Each player has to make a choice among various possible actions, but the outcome of her choice also depends on the choice made by the other players. The elements of the game are:

- A set of n players;
- A set A_i , for each player $i \in \{1, \dots, n\}$, called the strategy (or action) set of player i ; each element $(x_1, x_2, \dots, x_n) \in A := A_1 \times A_2 \times \dots \times A_n$ is called a strategy profile;
- For each player $i \in \{1, \dots, n\}$, a set of preferences, according to which they can order the various profiles;
- For each player $i \in \{1, \dots, n\}$, a utility (or payoff) function, $u_i : A \rightarrow \mathbb{R}$, which represents their preferences such that to higher values there correspond better outcomes. Let us notice that the utility function of player i does not merely depend on x_i , but on the whole strategy profile $x = (x_1, x_2, \dots, x_n)$; that is, the interaction with the other players cannot be neglected in their decision processes.

In the case of more than two players, it is important to distinguish the action of player i from the action of all the other players. To this end, we will use the notation x_{-i} to denote the subvector $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$, and write $x = (x_i, x_{-i}) = (x_1, \dots, x_n)$. Since each player i wishes to maximize their utility, and only controls her action variable x_i , a useful solution concept is that of Nash equilibrium of the game.

Definition 4. A Nash equilibrium is a vector $x^* \in A$ such that:

$$u_i(x_i^*, x_{-i}^*) \geq u_i(x_i, x_{-i}^*), \quad \forall i \in \{1, \dots, n\}, \forall x_i \in A_i. \tag{23}$$

The interested reader can find a comprehensive theoretical treatment of Game Theory in the book [28], along with various applications.

We now adapt the above concepts to our parametric network problem and, for simplicity, still denote the set of players by $\{1, 2, \dots, n\}$ instead of $\{v_1, v_2, \dots, v_n\}$. Thus, consider an interval $[0, T]$, and for each $t \in [0, T]$, denote with $A_i(t) \subset \mathbb{R}$ the parametric strategy space of player i , while $A(t) = A_1(t) \times \dots \times A_n(t)$ is called the parametric space of action profiles. Each player i is endowed with a payoff function $u_i : [0, T] \times A(t) \rightarrow \mathbb{R}$ that he/she wishes to maximize. The notation $u_i(t, x, G)$ is often utilized when one wants to emphasize the dependence on the graph structure. We wish to find a Nash equilibrium of the parametric game; that is, for each $t \in [0, T]$, we seek an element $x^*(t) \in A(t)$ such that for each $i \in \{1, \dots, n\}$:

$$u_i(t, x_i^*(t), x_{-i}^*(t)) \geq u_i(t, x_i, x_{-i}^*(t)), \quad \forall x_i \in A_i(t). \tag{24}$$

A characteristic of network games is that the vector x_{-i} is only made up of components x_j such that $j \in N_i(g)$; that is, j is a neighbor of i . For modeling purposes, and also for simplifying the analysis, it is common to make further assumptions on how variations in the actions of a player’s neighbors affect her marginal utility. We assume that the type of interaction does not change in $[0, T]$ and, in the case where $u_i(t, \cdot) \in C^2(\mathbb{R}^n)$, we obtain the following definitions.

Definition 5. We say that the network game has the property of strategic substitutes if for each player i the following condition holds:

$$\frac{\partial^2 u_i(t, x_i, x_{-i})}{\partial x_j \partial x_i} < 0, \quad \forall (i, j) : g_{ij} = 1, \forall t \in [0, T], \forall x \in A(t).$$

Definition 6. We say that the network game has the property of strategic complements if for each player i the following condition holds:

$$\frac{\partial^2 u_i(t, x_i, x_{-i})}{\partial x_j \partial x_i} > 0, \quad \forall (i, j) : g_{ij} = 1, \forall t \in [0, T], \forall x \in A(t).$$

We will use the variational inequality approach, pioneered by [23] (in the non-parametric case), to solve Nash equilibrium problems. If, for each $t \in [0, T]$, the u_i are continuously differentiable functions on \mathbb{R}^n , and $u_i(t, \cdot, x_{-i})$ are concave, the Nash equilibrium problem is equivalent to the variational inequality $VI(F, A(t))$:

For each $t \in [0, T]$, find $x^*(t) \in A(t)$ such that

$$\langle F(t, x^*(t)), x - x^*(t) \rangle \geq 0, \quad \forall x \in A(t), \tag{25}$$

where

$$[F(t, x)]^\top := - \left(\frac{\partial u_1}{\partial x_1}(t, x), \dots, \frac{\partial u_n}{\partial x_n}(t, x) \right) \tag{26}$$

is also called the *pseudo-gradient* of the game, according to the terminology introduced by Rosen [29].

In the following subsection, we introduce the parametric linear-quadratic utility functions, which will be used both for the Nash equilibrium problem and the generalized Nash equilibrium problem. This simple functional form as been extensively used in the literature (see, e.g, [20]) because, in the case on non-negative strategy space, it allows solutions in closed form. We consider instead the case where the strategy space of each player has a (parameter-dependent) upper bound.

4.2. The Linear-Quadratic Network Nash Equilibrium Problem

Let $A_i(t) = [0, U_i(t)]$ for any $i \in \{1, \dots, n\}$, $t \in [0, T]$, where, for each i , $U_i(t)$ is a positive and Lipschitz continuous function on $[0, T]$. The payoff of player i is given by:

$$u_i(t, x) = k(t)x_i - \frac{1}{2}x_i^2 + \phi \sum_{j=1}^n g_{ij}x_i x_j, \tag{27}$$

In this model $k : [0, T] \rightarrow \mathbb{R}$ is a positive Lipschitz function, and $\phi > 0$ describes the interaction between a player and their neighbors, which corresponds to the case of strategic complements. Moreover, since $k(t)$ is the same for all players, they only differ because of their position in the network. The pseudo-gradient’s components are given by:

$$F_i(t, x) = -k(t) + x_i - \phi \sum_{j=1}^n g_{ij}x_j, \quad i \in \{1, \dots, n\}, \tag{28}$$

which can be written in compact form as:

$$F(t, x) = (I - \phi G)x - k(t)\mathbf{1},$$

where $\mathbf{1} = (1, \dots, 1)^\top \in \mathbb{R}^n$. We will seek Nash equilibrium points by solving the variational inequality: for each $t \in [0, T]$, find $x^*(t) \in A(t)$ such that

$$\langle (I - \phi G)x^*(t) - k(t)\mathbf{1}, x - x^*(t) \rangle \geq 0, \quad \forall x \in A(t). \tag{29}$$

Lemma 1. *F is uniformly strongly monotone if $\phi\rho(G) < 1$, where $\rho(G)$ is the spectral radius of G.*

Proof. The symmetric matrix $I - \phi G$ is positive definite if and only if its minimum eigenvalue $\lambda_{\min}(I - \phi G)$ is positive. On the other hand, $\lambda_{\min}(I - \phi G) = 1 - \phi\lambda_{\max}(G)$. Since G is a symmetric non-negative matrix, the Perron–Frobenius Theorem guarantees that $\lambda_{\max}(G) = \rho(G)$, hence $I - \phi G$ is positive definite if and only if $\phi\rho(G) < 1$. This condition does not depend on $t \in [0, T]$. \square

Notice that, when $\phi\rho(G) < 1$, the map F in (28) satisfies the assumptions of Theorems 2 and 3 with constants $\alpha = 1 - \phi\rho(G)$, $L_x = \|I - \phi G\|_2$, $L_t = L_k\sqrt{n}$, where L_k is the Lipschitz constant of $k(t)$, and $L = \max\{L_x, L_t\}$. Moreover, the feasible region $A(t)$ satisfies assumption (3) with constant $M = \|(L_1, \dots, L_n)\|_2$, where L_i is the Lipschitz constant of $U_i(t)$ for any $i = 1, \dots, n$ (see [11]). Therefore, it follows from Theorems 2 and 3 that the Nash equilibrium $x^*(t)$ is a Lipschitz continuous function with estimated constant equal to $\min\{\Lambda_1, \Lambda_2\}$.

Remark 1. *The game under consideration also falls, for each $t \in [0, T]$, in the class of potential games according to the definition introduced by Monderer and Shapley [30]. Indeed, a potential function is given by:*

$$P(t, x) = \sum_{i=1}^n u_i(t, x) - \frac{\phi}{2} \sum_{i=1}^n \sum_{j=1}^n g_{ij}x_i x_j. \tag{30}$$

Monderer and Shapley have proven that, in general, the solutions of the problem

$$\max_{x \in A(t)} P(t, x)$$

form a subset of the solution set of the Nash game. Because both problems have a unique solution under the condition $\phi\rho(G) < 1$, it follows that the two problems share the same solution.

4.3. The Linear-Quadratic Network Generalized Nash Equilibrium Problem

We start this subsection with a recall of generalized Nash equilibrium problems (GNEPs) with shared constraints and their variational solutions. More details can be found in the already-mentioned paper by Rosen, and in the more recent papers [31–38]. We directly introduce the parametric case, and all the following definitions and properties hold for any fixed $t \in [0, T]$.

In GNEPs, the strategy set of each player is dependent on the strategies of their rivals. In this context, we examine a simplified scenario where the variable x is constrained to be non-negative ($x \geq 0$). Moreover, we consider a function $g : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^m$, which characterizes the collective constraints shared among the players. For each t , and for each move x_{-i} of the other players, the strategy set of player i is then given by:

$$K_i(t, x_{-i}) = \{x_i \in \mathbb{R}_+ : g(t, x) = g(t, x_i, x_{-i}) \leq 0\}.$$

Definition 7. The GNEP is the problem of finding, for each $t \in [0, T]$, $x^*(t) \in \mathbb{R}^n$ such that, for any $i \in \{1, \dots, n\}$, $x_i^*(t) \in K_i(t, x_{-i}^*(t))$ and

$$u_i(t, x_i^*(t), x_{-i}^*(t)) \geq u_i(x_i, x_{-i}^*(t)), \quad \forall t \in [0, T], \forall x_i \in K_i(t, x_{-i}^*(t)). \quad (31)$$

We assume that that, for each fixed $t \in [0, T]$, $x_i \in K_i(t, x_{-i}^*)$, the functions $u_i(t, \cdot, x_{-i}^*)$ are concave and continuously differentiable, and the components of $g(t, \cdot)$ are convex and continuously differentiable. As a consequence, a necessary and sufficient condition for $x_i^*(t) \in K_i(t, x_{-i}^*)$ to satisfy (31) is

$$-\frac{\partial u_i(t, x_i^*(t), x_{-i}^*(t))}{\partial x_i}(x_i - x_i^*(t)) \geq 0, \quad \forall t \in [0, T], \forall x_i \in K_i(t, x_{-i}^*(t)). \quad (32)$$

Thus, if we define $F(t, x)$ as in (26), and

$$K(t, x) = K_1(t, x_{-1}) \times \dots \times K_n(t, x_{-n}),$$

it follows that $x^*(t)$ is a GNE if and only if, $\forall t \in [0, T]$, $x^*(t) \in K(t, x^*(t))$ and

$$\langle F(t, x^*(t)), x - x^*(t) \rangle \geq 0, \quad \forall x \in K(t, x^*(t)). \quad (33)$$

The above problem, where the feasible set also depends on the solution, is called a quasi-variational inequality (with parameter t), and its solution is as difficult as the original GNEP. Let us now remark that even if the strict monotonicity assumption on F is satisfied for each t , the above problem (equivalent to the original GNEP) has infinite solutions. We can select a specific solution by studying its Lagrange multipliers.

Thus, assume that, for each $t \in [0, T]$, $x^*(t)$ is a solution of GNEP. Hence, for each i , $x_i^*(t)$ solves the maximization problem

$$\max_{x_i} \{u_i(t, x_i, x_{-i}^*(t)) : g(t, x_i, x_{-i}^*(t)) \leq 0, x_i \geq 0\}.$$

Under some standard constraint qualification, we can then write the KKT conditions for each maximization problem. We then introduce the Lagrange multiplier $\lambda^i(t) \in \mathbb{R}^m$ associated with the constraint $g(t, x_i, x_{-i}^*) \leq 0$ and the multiplier $\mu_i(t) \in \mathbb{R}$ associated with the nonnegativity constraint $x_i \geq 0$. The Lagrangian function for each player i reads as:

$$L_i(t, x_i, x_{-i}^*(t), \lambda^i(t), \mu_i(t)) = u_i(t, x_i, x_{-i}^*(t)) - \langle g(t, x_i, x_{-i}^*(t)), \lambda^i(t) \rangle + \mu_i(t)x_i$$

and the KKT conditions for all players are given by:

$$\nabla_{x_i} L_i(t, x_i^*(t), x_{-i}^*(t), \lambda^{i*}(t), \mu_i^*(t)) = 0, \quad i = 1, \dots, n, \quad (34)$$

$$\lambda_j^{i*}(t)g_j(t, x^*(t)) = 0, \lambda_j^{i*}(t) \geq 0, g_j(t, x^*(t)) \leq 0, \quad i = 1, \dots, n, j = 1, \dots, m \quad (35)$$

$$\mu_i^*(t)x_i^*(t) = 0, \mu_i^*(t) \geq 0, x_i^*(t) \geq 0, \quad i = 1, \dots, n. \quad (36)$$

Conversely, under the assumptions made, if, for any fixed t , $(x^*(t), \lambda(t), \mu^*(t))$, where $\lambda^*(t) = (\lambda^{1*}(t), \dots, \lambda^{n*}(t))$ and $\mu^*(t) = (\mu_1^*(t), \dots, \mu_n^*(t))$, satisfy the KKT system (34)–(36), then $x^*(t)$ is a GNE. We are now in position to classify equilibria.

Definition 8. Let $x^*(t)$ be a GNE which, together with the Lagrange multipliers $\lambda^*(t) = (\lambda^{1*}(t), \dots, \lambda^{n*}(t))$ and $\mu^*(t) = (\mu_1^*(t), \dots, \mu_n^*(t))$, satisfies the KKT system of all players. We call $x^*(t)$ a normalized equilibrium if there exists a vector $r(t) \in \mathbb{R}_{++}^n$ and a vector $\bar{\lambda}(t) \in \mathbb{R}_+^m$ such that

$$\lambda^{i*}(t) = \frac{\bar{\lambda}(t)}{r_i(t)}, \quad \forall i = 1, \dots, n,$$

which means that, for a normalized equilibrium, the multipliers of the constraints shared by all players are proportional to a common multiplier. In the special case where, for some t , $r_i(t) = 1$ for any i , i.e., the multipliers coincide for each player, $x^*(t)$ is called variational equilibrium (VE). Rosen [29] proved that if the feasible set, which in our case is:

$$K(t) = \{x \in \mathbb{R}_+^n : g(t, x) \leq 0\}$$

is compact and convex, then there exists a normalized equilibrium for each $r(t) \in \mathbb{R}_{++}^n$.

Now, let us define, for each $r(t) \in \mathbb{R}_{++}^n$, the vector function $F^{r(t)} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ as follows:

$$[F^{r(t)}(t, x)]^\top := -\left(r_1(t) \frac{\partial u_1}{\partial x_1}(t, x), \dots, r_n(t) \frac{\partial u_n}{\partial x_n}(t, x)\right).$$

The variational inequality approach for finding the normalized equilibria of the GNEP is expressed by the following theorem, which can be viewed as a special case of Proposition 3.2 in [34] in a parametric setting.

Theorem 5.

1. Suppose that $x^*(t)$ is a solution of $VI(F^{r(t)}, K(t))$, where $r(t) \in \mathbb{R}_{++}^n$, a constraint qualification holds at $x^*(t)$ and $(\bar{\lambda}(t), \bar{\mu}(t)) \in \mathbb{R}^m \times \mathbb{R}^n$ are the multipliers associated with $x^*(t)$. Then, $x^*(t)$ is a normalized equilibrium such that the multipliers $(\lambda^{i*}(t), \mu_i^*(t))$ of each player i satisfy the following conditions:

$$\lambda^{i*}(t) = \frac{\bar{\lambda}(t)}{r_i(t)}, \quad \mu_i^*(t) = \frac{\bar{\mu}_i(t)}{r_i(t)}, \quad \forall i = 1, \dots, n.$$

2. If $x^*(t)$ is a normalized equilibrium such that the multipliers $(\lambda^{i*}(t), \mu_i^*(t))$ of each player i satisfy the following conditions:

$$\lambda^{i*}(t) = \frac{\bar{\lambda}(t)}{r_i(t)}, \quad \forall i = 1, \dots, n,$$

for some vector $\bar{\lambda}(t) \in \mathbb{R}_+^m$ and $r(t) \in \mathbb{R}_{++}^n$, then $x^*(t)$ is a solution of $VI(F^{r(t)}, K(t))$ and $(\bar{\lambda}(t), r_1(t)\mu_1^*(t), \dots, r_n(t)\mu_n^*(t))$ are the corresponding multipliers.

The variational equilibria can then be computed by solving, for any $t \in [0, T]$, the following variational inequality: find $x^*(t) \in K(t)$ such that

$$\langle F(t, x^*(t)), x - x^*(t) \rangle \geq 0, \quad \forall x \in K(t). \tag{37}$$

As an example of network GNEP, we consider the same payoff functions defined as in (27), while the strategy set of player i is given by the usual individual constraint $x_i \geq 0$ and an additional constraint, shared by all the players, on the total quantity of activities of all players, that is

$$K_i(t, x_{-i}) = \left\{ x_i \in \mathbb{R}_+ : \sum_{i=1}^n x_i \leq C(t) \right\}, \quad i = 1, \dots, n, \quad t \in [0, T],$$

where $C : [0, T] \rightarrow \mathbb{R}$ is a given positive and Lipschitz continuous function. Depending on the specific application, the additional constraint can have the meaning of a collective budget upper bound or of a limited availability of a certain commodity. If $\phi\rho(G) < 1$, then

the variational equilibrium can thus be found by solving $VI(F, K(t))$, where F is the same as in (28) and

$$K(t) = \left\{ x \in \mathbb{R}_+^n : \sum_{i=1}^n x_i \leq C(t) \right\}. \tag{38}$$

Notice that $K(t)$ satisfies Assumption (3), with constant M equal to the Lipschitz constant of the function C (see [11]). Therefore, Theorems 2 and 3 guarantee that the variational equilibrium $x^*(t)$ is a Lipschitz continuous function with estimated constant equal to $\min\{\Lambda_1, \Lambda_2\}$.

5. Numerical Experiments

In this section, some numerical experiments show the performance of the algorithm described in Section 3 based on the new Lipschitz estimates provided in Section 2.

First, we compare the new Lipschitz estimate Λ_1 provided in Theorem 2 with that provided in [12] from the numerical point of view on a set of sample parameters. Then, we compare the performance of the algorithm described in Section 3 based on the Lipschitz estimate used. The algorithm and the computation of Lipschitz estimates were implemented in MATLAB R2023a.

5.1. Comparison between Λ_1 and the Estimate in [12]

Under the same assumptions of Theorem 2, the authors in [12] proved that the solution $x(t)$ of the $VI(F, K(t))$ is Lipschitz continuous with a constant equal to

$$\Lambda_0 = \inf_{(\varepsilon, \lambda)} \left\{ \frac{M + \sqrt{\frac{\lambda^2 + \lambda}{\varepsilon} + \lambda^2}}{1 - \sqrt{1 + \lambda^2(1 + \varepsilon) - \lambda(2\tilde{\alpha} - \varepsilon)}} : 0 < \varepsilon < 2\tilde{\alpha}, 0 < \lambda(1 + \varepsilon) + \varepsilon < 2\tilde{\alpha} \right\}, \tag{39}$$

where $\tilde{\alpha} = \alpha/L$. Since it is not easy to write the estimates Λ_0 and Λ_1 in closed form, we decided to compare them numerically. We considered the following grid for the parameters $(\tilde{\alpha}, M)$: $\tilde{\alpha}$ varies from 0.01 to 1 with a step equal to 0.01, while M varies from 0 to 10 with step equal to 0.01. For any value of $(\tilde{\alpha}, M)$ in the considered grid, we computed the values of Λ_0 and Λ_1 : the minimum value in Λ_0 was numerically approximated (due to the non-convexity of the objective function and constraints), while the minimum value in Λ_1 was computed by exploiting the MATLAB function `fminbnd`. The results show that the new estimate Λ_1 definitely improves Λ_0 . In fact, Λ_1 is always less than Λ_0 and the relative improvement in Λ_1 compared to Λ_0 , i.e., $(\Lambda_0 - \Lambda_1)/\Lambda_0$, is about 42% on average with a minimum of 13% and a maximum of 99%.

5.2. Performance of the Algorithm

We now show the performance of the algorithm described in Section 3 to solve a linear-quadratic network Nash equilibrium problem (Example 1) and a linear-quadratic network GNEP (Example 2). We implemented three different versions of the algorithm according to the Lipschitz estimate used. We considered the estimate Λ_0 provided in [12], and the new estimates Λ_1 and Λ_2 introduced in Theorems 2 and 3, respectively.

Example 1. We consider a linear-quadratic network Nash equilibrium problem as in the framework of Section 4.2, with a network of 8 nodes (players) as shown in Figure 2.




Figure 2. Network topology of Example 1.

The spectral radius of the adjacency matrix G is $\rho(G) \simeq 3.1019$. We set the function $k(t) = t + 5$, with $t \in [0, 15]$. We set two different values for parameter ϕ : $\phi = 0.1/\rho(G)$ or $\phi = 0.5/\rho(G)$, so that there exists a unique Nash equilibrium in both cases. We set the upper bounds $U_i(t) = \min\{2i + (t + 5)/2, 18\}$ for any $i = 1, \dots, 8$. We recall that the map F in (28) satisfies the assumptions of Theorems 2 and 3 with constants $\alpha = 1 - \phi\rho(G)$, $L_x = \|I - \phi G\|_2$, $L_t = \sqrt{8}$, $L = \max\{L_x, L_t\}$, and the feasible region

$$A(t) = \prod_{i=1}^8 [0, U_i(t)]$$

satisfies Assumption (3) with constant $M = \|(L_1, \dots, L_8)\|_2$, where L_i is the Lipschitz constant of the function U_i for any $i = 1, \dots, 8$. We run the algorithm to approximate the fifth component of the solution vector $x(t)$, due to the special position of the player 5 in the network. To evaluate the function $x_5(t)$ during the execution of the algorithm, i.e., to find Nash equilibria of the problem, we exploited the ad hoc algorithm proposed in [39].

Table 2 shows the number of iterations performed (or Nash equilibria computed) by the three versions of the algorithm (based on the estimates Λ_0 , Λ_1 and Λ_2) for different values of ϕ and the error tolerance ϵ_{tot} . We set the maximum number of iterations equal to 10^6 . Results show that the algorithm based on the estimate Λ_2 outperforms the other two versions based on the estimates Λ_0 and Λ_1 . Table 3 shows the convergence of the approximate mean value of $x_5(t)$ (computed by the algorithm based on Λ_2) for different values of ϕ .

Table 2. Example 1 with a moving feasible region: number of iterations of the algorithm based on the Lipschitz estimate used.

ϵ_{tot}	$\phi = 0.1/\rho(G)$			$\phi = 0.5/\rho(G)$		
	Alg with Λ_0	Alg with Λ_1	Alg with Λ_2	Alg with Λ_0	Alg with Λ_1	Alg with Λ_2
1	5107	1799	640	15,383	5246	1714
0.5	10,213	3597	1277	30,765	10,490	3426
0.2	26,069	8554	3260	78,858	26,431	7995
0.1	52,136	17,107	6518	157,715	52,860	15,989
0.05	104,270	34,212	13,034	315,428	105,719	31,977
0.02	246,321	92,250	30,788	815,228	249,976	85,721
0.01	492,640	184,499	61,575	$>1 \times 10^6$	499,950	171,441
Lipschitz estimate	$\Lambda_0 = 88.46$	$\Lambda_1 = 33.18$	$\Lambda_2 = 11.18$	$\Lambda_0 = 276.49$	$\Lambda_1 = 96.82$	$\Lambda_2 = 30.31$

Table 3. Example 1 with a moving feasible region: approximate mean value of $x_5(t)$.

ϵ_{tot}	$\phi = 0.1/\rho(G)$	$\phi = 0.5/\rho(G)$
1	13.0189685363	15.2238272767
0.5	13.0189711251	15.2238276377
0.2	13.0189724192	15.2238277070
0.1	13.0189727295	15.2238277328
0.05	13.0189728137	15.2238277376
0.02	13.0189728309	15.2238277386
0.01	13.0189728372	15.2238277390

Moreover, we run the algorithm to solve the problem in the case the feasible region $A(t)$ is constant, with $U_i(t) = 18$ for any t and $i = 1, \dots, 8$, and the map F is the same as before. Tables 4 and 5 show the number of iterations of the algorithm and the approximate mean value of $x_5(t)$, respectively. Notice that, in this case, the Lipschitz estimates Λ_1 and Λ_2 coincide, since $L = L_t$ and the algorithm based on such an estimate is much more efficient than the one based on Λ_0 .

Table 4. Example 1 with a constant feasible region: number of iterations of the algorithm based on the Lipschitz estimate used.

ϵ_{tot}	$\phi = 0.1/\rho(G)$			$\phi = 0.5/\rho(G)$		
	Alg with Λ_0	Alg with Λ_1	Alg with Λ_2	Alg with Λ_0	Alg with Λ_1	Alg with Λ_2
1	2835	179	179	7985	343	343
0.5	5668	357	357	15,968	684	684
0.2	13,983	881	881	43,207	1701	1701
0.1	27,965	1760	1760	86,413	3401	3401
0.05	55,928	3519	3519	172,824	6800	6800
0.02	128,807	8111	8111	433,801	15,862	15,862
0.01	257,613	16,220	16,220	867,601	31,722	31,722
Lipschitz estimate	$\Lambda_0 = 45.07$	$\Lambda_1 = 3.14$	$\Lambda_2 = 3.14$	$\Lambda_0 = 138.59$	$\Lambda_1 = 5.66$	$\Lambda_2 = 5.66$

Table 5. Example 1 with a constant feasible region: approximate mean value of $x_5(t)$.

ϵ_{tot}	$\phi = 0.1/\rho(G)$	$\phi = 0.5/\rho(G)$
1	13.0336960026	15.9007292403
0.5	13.0337000695	15.9007436640
0.2	13.0337020507	15.9007521737
0.1	13.0337029357	15.9007529259
0.05	13.0337033781	15.9007531658
0.02	13.0337033949	15.9007532022
0.01	13.0337034020	15.9007532066

Example 2. We now consider a linear-quadratic network generalized Nash equilibrium problem, as in the framework of Section 4.3. We assume that the player network is the same as Example 1. We set the function $k(t) = t + 5$, with $t \in [0, 15]$, $\phi = 0.1/\rho(G)$ or $\phi = 0.5/\rho(G)$, and $K(t)$ defined as in (38) with $C(t) = \min\{13 + t, 18\}$. The map F in (28) satisfies the assumptions of Theorems 2 and 3 with constants $\alpha = 1 - \phi\rho(G)$, $L_x = \|I - \phi G\|_2$, $L_t = \sqrt{8}$, $L = \max\{L_x, L_t\}$, and the feasible region satisfies assumption (3) with constant $M = 1$. In order to find the unique variational equilibrium $x(t)$ of the problem, we found the maximizer of the potential function (30) by means of the MATLAB function `quadprog`. As in Example 1, we run the algorithm to approximate the fifth component of the solution vector $x(t)$.

Table 6 shows the number of iterations performed by the three versions of the algorithm (based on the estimates Λ_0 , Λ_1 and Λ_2) for different values of ϕ and ϵ_{tot} . Results show again that the algorithm based on the estimate Λ_2 outperforms the other two versions based on the estimates Λ_0 and Λ_1 . Table 7 shows the convergence of the approximate mean value of $x_5(t)$ for different values of ϕ .

Table 6. Example 2 with a moving feasible region: number of iterations of the algorithm based on the Lipschitz estimate used.

ϵ_{tot}	$\phi = 0.1/\rho(G)$			$\phi = 0.5/\rho(G)$		
	Alg with Λ_0	Alg with Λ_1	Alg with Λ_2	Alg with Λ_0	Alg with Λ_1	Alg with Λ_2
1	3353	497	302	10,118	1210	637
0.5	6703	992	602	20,233	2419	1272
0.2	16,053	2498	1438	51,723	6282	3222
0.1	32,104	4994	2875	103,444	12,561	6441
0.05	64,206	9986	5747	206,886	25,120	12,880
0.02	163,601	25,447	15,288	501,923	60,264	33,256
0.01	327,200	50,892	30,574	$>1 \times 10^6$	120,525	66,511
Lipschitz estimate	$\Lambda_0 = 79.14$	$\Lambda_1 = 26.80$	$\Lambda_2 = 9.93$	$\Lambda_0 = 246.59$	$\Lambda_1 = 76.16$	$\Lambda_2 = 25.65$

Table 7. Example 1 with a moving feasible region: approximate mean value of $x_5(t)$.

ϵ_{tot}	$\phi = 0.1/\rho(G)$	$\phi = 0.5/\rho(G)$
1	2.0960942274	1.9169266928
0.5	2.0960948096	1.9169268259
0.2	2.0960949552	1.9169268675
0.1	2.0960949916	1.9169268696
0.05	2.0960950007	1.9169268701
0.02	2.0960950030	1.9169268702
0.01	2.0960950035	1.9169268703

Similarly to Example 1, Table 8 and 9 show the number of iterations of the algorithm and the approximate mean value of $x_5(t)$ in the case where the feasible region $K(t)$ is constant, with $C(t) = 18$ for any t , and the map F is the same as before.

Table 8. Example 2 with a constant feasible region: number of iterations of the algorithm based on the Lipschitz estimate used.

ϵ_{tot}	$\phi = 0.1/\rho(G)$			$\phi = 0.5/\rho(G)$		
	Alg with Λ_0	Alg with Λ_1	Alg with Λ_2	Alg with Λ_0	Alg with Λ_1	Alg with Λ_2
1	2836	200	200	7985	358	358
0.5	5671	399	399	15,969	714	714
0.2	13,988	944	944	43,210	1755	1755
0.1	27,975	1887	1887	86,419	3509	3509
0.05	55,948	3773	3773	172,837	7017	7017
0.02	128,837	9392	9392	433,825	16,141	16,141
0.01	257,673	18,783	18,783	867,648	32,281	32,281
Lipschitz estimate	$\Lambda_0 = 45.07$	$\Lambda_1 = 3.14$	$\Lambda_2 = 3.14$	$\Lambda_0 = 138.59$	$\Lambda_1 = 5.66$	$\Lambda_2 = 5.66$

Table 9. Example 2 with a constant feasible region: approximate mean value of $x_5(t)$.

ϵ_{tot}	$\phi = 0.1/\rho(G)$	$\phi = 0.5/\rho(G)$
1	2.197847188363	2.009981572725
0.5	2.197847188363	2.009981572725
0.2	2.197847188363	2.009981572725
0.1	2.197847188363	2.009981572725
0.05	2.197847188363	2.009981572725
0.02	2.197847188363	2.009981572725
0.01	2.197847188364	2.009981572727

6. Conclusions

In this paper, we improved previous estimates of the global Lipschitz constant of the unique solution of a parametric variational inequality. Having in mind various applications where the parameter represents time, we utilized the estimated constant to approximate the

mean value of the solution by modifying an algorithm previously derived to approximate black box Lipschitz functions. We then applied our results to two class of games played on networks, where we assumed that some terms of the players’ utility functions as well as the constraint set could vary with time. In future work, we plan to investigate the case where also the adjacency matrix of the graph is time-dependent.

Author Contributions: M.P. and F.R. have contributed equally to this work. All authors have read and agreed to the published version of the manuscript.

Funding: This research was partially supported by the research project “Programma ricerca di ateneo UNICT 2020-22 linea 2-OMNIA” of the University of Catania. This support is gratefully acknowledged.

Data Availability Statement: The data presented in this study are available in the article.

Acknowledgments: The authors are members of the Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA—National Group for Mathematical Analysis, Probability and their Applications) of the Istituto Nazionale di Alta Matematica (INdAM—National Institute of Higher Mathematics).

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Appendix A.1. Special Case $\tilde{\alpha} = 1$ in Theorem 2.

The function

$$f(z) = \frac{M}{z} + \frac{1+z}{1-z}$$

is strictly convex in $(0, 1)$ and

$$f'(z) = \frac{2z^2 - Mz^2 + 2Mz - M}{z^2(1-z)^2}.$$

If $M = 0$, then $\inf_{z \in (0,1)} f(z) = 1$. If $M > 0$, then the minimizer of f is

$$\bar{z} = \begin{cases} \frac{-M + \sqrt{2M}}{2-M} & \text{if } M < 2, \\ \frac{1}{2} & \text{if } M = 2, \\ \frac{M - \sqrt{2M}}{M-2} & \text{if } M > 2. \end{cases}$$

Therefore, we have

$$\begin{aligned} \inf_{z \in (0,1)} f(z) &= \begin{cases} 7 & \text{if } M = 2, \\ \frac{M}{\frac{\sqrt{2M}-M}{2-M}} + \frac{1 + \frac{\sqrt{2M}-M}{2-M}}{1 - \frac{\sqrt{2M}-M}{2-M}} & \text{if } M \neq 2, \end{cases} \\ &= \begin{cases} 7 & \text{if } M = 2, \\ 1 + M + 2\sqrt{2M} & \text{if } M \neq 2, \end{cases} \\ &= 1 + M + 2\sqrt{2M}. \end{aligned}$$

The function

$$f(z) = \frac{M}{2-z} + \frac{z(1+z)}{(z-1)(2-z)} = \frac{z^2 + z(M+1) - M}{-z^2 + 3z - 2}$$

is strictly convex in $(1, 2)$ and

$$f'(z) = \frac{(4+M)z^2 - 2z(M+2) + M-2}{(-z^2 + 3z - 2)^2}.$$

Thus, the minimum is reached in

$$\bar{z} = \frac{M + 2 + \sqrt{2M + 12}}{M + 4},$$

which belongs to the interval (1, 2). Therefore, we have

$$\begin{aligned} \inf_{z \in (1,2)} f(z) &= \frac{4M + 24 + (M^2 + 7M + 8)\sqrt{2M + 12}}{(M + 8)\sqrt{2M + 12} - 4M - 24} \\ &= \frac{\sqrt{8}\sqrt{M + 6} + M^2 + 7M + 8}{M + 8 - \sqrt{8}\sqrt{M + 6}} \\ &= \frac{M^2 + 7M + 8 + 2\sqrt{2M + 12}}{M + 8 - 2\sqrt{2M + 12}}. \end{aligned}$$

Appendix A.2. Special Case $\tilde{\alpha} = 1$ in Theorem 3.

The function

$$f(z) = \frac{M}{z} + \hat{L} \frac{1+z}{1-z}$$

is strictly convex in (0, 1) and

$$f'(z) = \frac{2Lz^2 - M(1-z)^2}{z^2(1-z)^2}.$$

If $M = 0$, then $\inf_{z \in (0,1)} f(z) = \hat{L}$. If $M > 0$, then the minimizer of f is

$$\bar{z} = \begin{cases} \frac{-M + \sqrt{2\hat{L}M}}{2\hat{L} - M} & \text{if } 2\hat{L} - M > 0, \\ \frac{-M + \sqrt{2\hat{L}M}}{2\hat{L} - M} & \text{if } 2\hat{L} - M < 0, \\ \frac{1}{2} & \text{if } 2\hat{L} - M = 0. \end{cases}$$

Therefore, we have

$$\begin{aligned} \inf_{z \in (0,1)} f(z) &= \begin{cases} 7\hat{L} & \text{if } M = 2\hat{L}, \\ \frac{M(2\hat{L} - M)}{\sqrt{2\hat{L}M - M}} + \hat{L} \frac{1 + \frac{\sqrt{2\hat{L}M - M}}{2\hat{L} - M}}{1 - \frac{\sqrt{2\hat{L}M - M}}{2\hat{L} - M}} & \text{if } M \neq 2\hat{L}, \end{cases} \\ &= \begin{cases} 7\hat{L} & \text{if } M = 2\hat{L}, \\ M + 2\sqrt{2\hat{L}M} + \hat{L} & \text{if } M \neq 2\hat{L}, \end{cases} \\ &= M + 2\sqrt{2\hat{L}M} + \hat{L}. \end{aligned}$$

The function

$$f(z) = \frac{M}{2-z} + \hat{L} \frac{z(1+z)}{(z-1)(2-z)} = \frac{\hat{L}z^2 + z(M + \hat{L}) - M}{-z^2 + 3z - 2}$$

is strictly convex in (1, 2) and

$$f'(z) = \frac{(4\hat{L} + M)z^2 - 2(2\hat{L} + M)z + M - 2\hat{L}}{(-z^2 + 3z - 2)^2}.$$

The minimum point is

$$\bar{z} = \frac{M + 2\hat{L} + \sqrt{2\hat{L}M + 12\hat{L}^2}}{4\hat{L} + M}$$

which belongs to $(1, 2)$. Moreover, after some algebra, we obtain:

$$\begin{aligned} \inf_{z \in (1,2)} f(z) &= \frac{(M^2 + 7\hat{L}M + 8\hat{L}^2)\sqrt{12\hat{L}^2 + 2\hat{L}M + 4M\hat{L}^2 + 24\hat{L}^3}}{(M + 8\hat{L})\sqrt{12\hat{L}^2 + 2\hat{L}M} - 4\hat{L}M - 24\hat{L}^2} \\ &= \frac{M^2 + 7\hat{L}M + 8\hat{L}^2 + \hat{L}\sqrt{8\hat{L}\sqrt{M + 6\hat{L}}}}{M + 8\hat{L} - \sqrt{8\hat{L}\sqrt{M + 6\hat{L}}}} \\ &= \frac{M^2 + 7M\hat{L} + 8\hat{L}^2 + 2\hat{L}\sqrt{2M\hat{L} + 12\hat{L}^2}}{M + 8\hat{L} - 2\sqrt{2M\hat{L} + 12\hat{L}^2}}. \end{aligned}$$

References

- Bank, B.; Guddat, J.; Klatte, D.; Kummer, B.; Tammer, K. *Non-Linear Parametric Optimization*; Springer: Berlin/Heidelberg, Germany, 1983.
- Aubin, J.-P. Lipschitz behavior of solutions to convex minimization problems. *Math. Oper. Res.* **1984**, *9*, 87–111.
- Mansour, M.A.; Bahraoui, M.A.; Adham, E.B. Approximate solutions of quasi-equilibrium problems: Lipschitz dependence of solutions on parameters. *J. Appl. Numer. Optim* **2021**, *3*, 297–314.
- Li, W. The sharp Lipschitz constants for feasible and optimal solutions of a perturbed linear program. *Linear Algebra Appl.* **1993**, *187*, 15–40.
- Li, W. Sharp Lipschitz constants for basic optimal solutions and basic feasible solutions of linear programs. *SIAM J. Control Optim.* **1994**, *32*, 140–153.
- Klatte, D.; Gisbert, T. A note of Lipschitz constants for solutions of linear inequalities and equations. *Linear Algebra Appl.* **1996**, *244*, 365–374.
- Mangasarian, O.L.; Shiau, T.H. Lipschitz continuity of solutions of linear inequalities, programs and complementarity problems. *SIAM J. Control Optim.* **1986**, *25*, 583–595.
- Facchinei, F.; Pang, J.-S. *Finite-Dimensional Variational Inequalities and Complementarity Problems*; Springer: Berlin, Germany, 2003.
- Yen, N.D. Lipschitz Continuity of Solutions of Variational Inequalities with a Parametric Polyhedral Constraint. *Math. Oper. Res.* **1995**, *20*, 695–708.
- Maugeri, A.; Scrimali, L. Global Lipschitz Continuity of Solutions to Parameterized Variational Inequalities. *Boll. Dell'Unione Mat. Ital.* **2009**, *2*, 45–69.
- Causa, A.; Raciti, F. Lipschitz Continuity Results for a Class of Variational Inequalities and Applications: A Geometric Approach. *J. Optim. Theory Appl.* **2010**, *145*, 235–248.
- Falsaperla, P.; Raciti, F. Global Approximation of Solutions of Time-Dependent Variational Inequalities. *Numer. Funct. Anal. Optim.* **2014**, *35*, 1018–1042.
- Zabinsky, Z.B.; Smith, R.L.; Kristinsdottir, B.P. Optimal estimation of univariate black-box Lipschitz functions with upper and lower error bounds. *Comput. Oper. Res.* **2003**, *30*, 1539–1553.
- Baran, I.; Demain, E.D.; Katz, D.A. Optimally Adaptive Integration of Univariate Lipschitz Functions. *Algorithmica* **2008**, *50*, 255–278.
- Ballester, C.; Calvo-Armengol, A.; Zenou, Y. Who's Who in Networks. Wanted: The Key Player. *Econometrica* **2006**, *74*, 1403–1417.
- Belhaj, M.; Bramoullé Y.; Deroian F. Network games under strategic complementarities. *Games Econ. Behav.* **2014**, *88*, 310–319.
- Calvo-Armengol, A.; Patacchini, E.; Zenou, Y. Peer effects and social networks in education. *Rev. Econ. Stud.* **2009**, *76*, 1239–1267.
- Helsley, R.; Zenou, Y. Social networks and interactions in cities. *J. Econ. Theory* **2014**, *150*, 426–466.
- Liu, X.; Patacchini, E.; Zenou, Y. Endogenous peer effects: Local aggregate or local average? *J. Econ. Behav. Organ.* **2014**, *103*, 39–59.
- Jackson, M.O.; Zenou, Y. Games on Networks. In *Handbook of Game Theory with Economic Applications*; Elsevier: Amsterdam, The Netherlands, 2015; pp. 95–163.
- Goyal, S. *Connections: An Introduction to the Economics of Networks*; Princeton University Press: Princeton, NJ, USA, 2007.
- Bonacich, P. Power and centrality: A family of measures. *Am. J., Sociol.* **1987**, *92*, 1170–1182.
- Gabay, D.; Moulin, H. On the uniqueness and stability of Nash equilibria in noncooperative games. In *Applied Stochastic Control in Econometrics and Management Science*; Bensoussan A., Kleindorfer P., Tapiero C.S., Eds.; North-Holland: Amsterdam, The Netherlands, 1980; pp. 271–294.
- Parise, F.; Ozdaglar, A. A variational inequality framework for network games: Existence, uniqueness, convergence and sensitivity analysis. *Games Econ. Behav.* **2019**, *114*, 47–82.
- Passacantando, M.; Raciti, F. A note on generalized Nash games played on networks. In *Nonlinear Analysis, Differential Equations, and Applications*; Rassias, T.M., Ed.; Optimization and Its Applications; Springer: Berlin/Heidelberg, Germany, 2021; Volume 173, pp. 365–380.
- Daniele, P. *Dynamic Networks and Evolutionary Variational Inequalities*; Edward Elgar Publishing: Cheltenham, UK, 2006.
- Pang, J.-S.; Stewart, D.E. Differential variational inequalities, *Math. Program.* **2008**, *113*, 345–424.
- Osborne, M.J. *Introduction to Game Theory: International Edition*; OUP Catalogue, Oxford University Press: Oxford, UK, 2009.
- Rosen, J.B. Existence and uniqueness of equilibrium points for concave n person games. *Econometrica* **1965**, *33*, 520–534.
- Monderer, D.; Shapley, L.S. Potential games. *Games Econ. Behav.* **1996**, *14*, 124–143.

31. Facchinei, F.; Fischer, A.; Piccialli, V. On generalized Nash games and variational inequalities. *Oper. Res. Lett.* **2007**, *35*, 159–164.
32. Facchinei, F.; Kanzov, C. Generalized Nash equilibrium problems. *Ann. Oper. Res.* **2010**, *175*, 177–211.
33. Nabetani, K. Variational Inequality Approaches to Generalized Nash Equilibrium Problems, Master's Thesis, Department of Applied Mathematics and Physics Graduate School of Informatics, Kyoto University, Kyoto, Japan, 2008.
34. Nabetani, K.; Tseng, P.; Fukushima, M. Parametrized variational inequality approaches to generalized Nash equilibrium problems with shared constraints. *Comput. Optim. Appl.* **2011**, *48*, 423–452.
35. Nagurney, A.; Alvarez Flores, E.; Soylu, C. A Generalized Nash Equilibrium network model for post-disaster humanitarian relief. *Transp. Res. E Logist. Transp. Rev.* **2026**, *95*, 1–18.
36. Aussel, D.; Dutta, J. Generalized Nash equilibrium problem, variational inequality and quasiconvexity. *Oper. Res. Lett.* **2011**, *36*, 461–464.
37. Aussel, D.; Cao Van, K.; Salas, D. Existence Results for Generalized Nash Equilibrium Problems under Continuity-Like Properties of Sublevel Sets. *Siam J. Optim.* **2021**, *31*, 2784–2806.
38. Krilašević, S., Grammatico, S. Learning generalized Nash equilibria in multi-agent dynamical systems via extremum seeking control. *Automatica* **2021**, *133*, 109846.
39. Passacantando, M.; Raciti, F. A finite convergence algorithm for solving linear-quadratic network games with strategic complements and bounded strategies. *Optim. Methods Softw.* **2023**, 1–24. <https://doi.org/10.1080/10556788.2023.2205644>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.