




Abdominal drainage after elective colorectal surgery: propensity score-matched retrospective analysis of an Italian cohort

Stefano Guadagni¹ , Marco Catarci^{2,3,*} , Francesco Masedu⁴, Mohammad Ehsanul Karim^{5,6} , Marco Clementi¹, Giacomo Ruffo⁷, Massimo Giuseppe Viola⁸, Felice Borghi^{9,10}, Gianandrea Baldazzi^{11,12}, Marco Scatizzi¹³, Felice Pirozzi¹⁴, Paolo Delrio¹⁵, Gianluca Garulli¹⁶, Pierluigi Marini¹⁷, Roberto Campagnacci¹⁸, Raffaele De Luca¹⁹, Ferdinando Ficari²⁰, Giuseppe Sica²¹, Stefano Scabini²², Andrea Liverani²³, Marco Caricato²⁴ and Alberto Patriti²⁵, the Italian ColoRectal Anastomotic Leakage (iCral) study group

- ¹General Surgery Unit, University of L'Aquila, L'Aquila, Italy
²General Surgery Unit, Sandro Pertini Hospital, ASL Roma 2, Roma, Italy
³General Surgery Unit, 'C.&G. Mazzoni' Hospital, Ascoli Piceno, Italy
⁴Department of Applied Clinical Sciences and Biotechnology, University of L'Aquila, L'Aquila, Italy
⁵School of Population and Public Health, The University of British Columbia, Vancouver, BC, Canada
⁶Centre for Health Evaluation and Outcome Sciences, St.Paul's Hospital, Vancouver, BC, Canada
⁷General Surgery Unit, IRCCS Sacro Cuore Don Calabria Hospital, Negrar di Valpolicella (VR), Italy
⁸General Surgery Unit, Cardinale G. Panico Hospital, Tricase, Italy
⁹Oncologic Surgery Unit, Candiolo Cancer Institute, FPO-IRCCS, Candiolo, Italy
¹⁰General & Oncologic Surgery Unit, Department of Surgery, Santa Croce e Carle Hospital, Cuneo, Italy
¹¹General Surgery Unit, ASST Ovest Milanese, Legnano, Italy
¹²General Surgery Unit, ASST Nord Milano, Sesto San Giovanni, Italy
¹³General Surgery Unit, Santa Maria Annunziata & Serristori Hospital, Firenze, Italy
¹⁴General Surgery Unit, ASL Napoli 2 Nord, Pozzuoli, Italy
¹⁵Colorectal Surgical Oncology, Istituto Nazionale per lo Studio e la Cura dei Tumori, 'Fondazione Giovanni Pascale IRCCS-Italia', Napoli, Italy
¹⁶General Surgery Unit, Infermi Hospital, Rimini, Italy
¹⁷General & Emergency Surgery Unit, San Camillo-Forlanini Hospital, Roma, Italy
¹⁸General Surgery Unit, 'C. Urbani' Hospital, Jesi, Italy
¹⁹Department of Surgical Oncology, IRCCS Istituto Tumori 'Giovanni Paolo II', Bari, Italy
²⁰General Surgery and IBD Unit, Careggi University Hospital, Firenze, Italy
²¹Minimally Invasive Surgery Unit, Policlinico Tor Vergata University Hospital, Roma, Italy
²²General & Oncologic Surgery Unit, IRCCS 'San Martino' National Cancer Center, Genova, Italy
²³General Surgery Unit, Regina Apostolorum Hospital, Albano Laziale, Italy
²⁴Colorectal Surgery Unit, Policlinico Campus BioMedico, Roma, Italy
²⁵Department of Surgery, Marche Nord Hospital, Pesaro e Fano, Italy

*Correspondence to: Marco Catarci, UOC Chirurgia Generale, Ospedale Sandro Pertini, ASL Roma 2, Via dei Monti Tiburtini 385, 00157 Roma, Italy (e-mail: marco.catarci@aslroma2.it)

Abstract

Background: In Italy, surgeons continue to drain the abdominal cavity in more than 50 per cent of patients after colorectal resection. The aim of this study was to evaluate the impact of abdominal drain placement on early adverse events in patients undergoing elective colorectal surgery.

Methods: A database was retrospectively analysed through a 1:1 propensity score-matching model including 21 covariates. The primary endpoint was the postoperative duration of stay, and the secondary endpoints were surgical site infections, infectious morbidity rate defined as surgical site infections plus pulmonary infections plus urinary infections, anastomotic leakage, overall morbidity rate, major morbidity rate, reoperation and mortality rates. The results of multiple logistic regression analyses were presented as odds ratios (OR) and 95 per cent c.i.

Results: A total of 6157 patients were analysed to produce two well-balanced groups of 1802 patients: group (A), no abdominal drain(s) and group (B), abdominal drain(s). Group A versus group B showed a significantly lower risk of postoperative duration of stay >6 days (OR 0.60; 95 per cent c.i. 0.51–0.70; $P < 0.001$). A mean postoperative duration of stay difference of 0.86 days was detected between groups. No difference was recorded between the two groups for all the other endpoints.

Conclusion: This study confirms that placement of abdominal drain(s) after elective colorectal surgery is associated with a non-clinically significant longer (0.86 days) postoperative duration of stay but has no impact on any other secondary outcomes, confirming that abdominal drains should not be used routinely in colorectal surgery.

Introduction

More than 100 years after the statements of Robert Lawson Tait 'When in doubt, drain' and of William Stewart Halsted 'No drainage at all is better than the ignorant employment of it'¹, the assumption that the placement of peritoneal drains after elective colorectal surgery can provide diagnostic and therapeutic benefit through prevention and early detection of anastomotic leak or other intraperitoneal collections is debated^{2,3}. Evidence suggests that drains can stimulate serous fluid production and may lead to an increased risk of surgical site infection (SSI)⁴ and adhesions, and prolonged hospital length of stay (LOS), impacting on postoperative pain control, mobility^{4,5}, increased perceived discomfort and anxiety⁶. The Enhanced Recovery After Surgery (ERAS) Society⁷, the American Society of Colon and Rectal Surgeons and the Society of American Gastrointestinal and Endoscopic Surgeons⁸, French⁹ and Italian¹⁰ guidelines, based on RCTs^{11,12}, older^{13,14} and more recent^{15,16} meta-analyses or systematic reviews of RCTs, strongly recommend that pelvic and peritoneal drains should not be used routinely in colorectal surgery. However, this strong recommendation is based on moderate-quality evidence^{8,17} (all the RCTs showed a bias of surgeon blinding, and some of them had a bias of allocation concealment and sequence randomization method¹², systematic reviews/meta-analyses included a large number of infra-promontory anastomoses in which a pelvic drain is almost always placed) and mainly on data observed before the widespread application of minimally invasive surgery. Conversely, many surgeons, particularly in Europe and China¹⁸, still believe that prophylactic drainage may remove collected fluid, thus reducing the risk of intra-abdominal infection, favouring early detection of postoperative complications such as intra-abdominal bleeding or anastomotic leakage, and minimize their severity, possibly avoiding reoperation^{19,20}.

Despite the above-mentioned recommendations, recent large observational studies in Italy, Spain and Europe^{21–25} report an abdominal drain placement rate after colorectal resection ranging from 40 to 70 per cent, reaching 90 per cent in a recent survey among German and Austrian surgeons²⁶, whereas these rates are generally reported below 15 per cent in North America^{27,28}.

The aim of the present study was to address the existing gap in knowledge by evaluating the impact of the omission of abdominal drains on early adverse events in patients who underwent elective colorectal surgery. Data were used from two prospective open-label observational multicentre studies of the Italian ColoRectal Anastomotic Leakage (iCral) study group^{24,25}.

Methods

Study design

This was a retrospective propensity score-matched analysis (PSMA) of patients who had undergone colorectal surgery for malignant and benign diseases enrolled in two consecutive studies upon explicit inclusion/exclusion criteria, in 78 surgical centres in Italy from January 2019 to September 2021: iCral2²⁴ and iCral3²⁵.

Patient population and data collection

The inclusion criteria were: ASA class I, II or III; elective or delayed urgency setting (defined as >48 h from admission in iCral2 and >24 h from admission in iCral3); patient's written informed consent for inclusion in the study and processing of sensitive data. The exclusion criteria were pregnancy, hyperthermic chemotherapy (HIPEC) for carcinomatosis and incomplete data.

The iCral2 study excluded patients with a protective stoma proximal to the anastomosis; conversely, these patients were included in the iCral3 study. Both studies were conducted in accordance with the Declaration of Helsinki and guidelines for good clinical practice E6 (R2). The study protocols were approved by the ethics committee of the coordinating centre (Marche Regional Ethics Committee (CERM) 2018/334 released on 28 November, 2018 for iCral2 and 2020/192 released on 30 July, 2020 for iCral3) and registered at clinicaltrials.gov (NCT03771456 for iCral2 and NCT04397627 for iCral3). Subsequently, all other centres were authorized to participate by their local ethics committees. Due to the retrospective nature of the current analysis, no specific authorization was requested.

To control for data imbalance derived from several treatment confounders, the present PSMA study included 6157 patients (73.7 per cent) out of 8359 in the parent studies, based on explicit exclusion criteria: any anastomosis located <10 cm from the anal verge, any anastomosis protected by a proximal stoma, delayed urgency, neo-adjuvant therapy, perioperative steroids and dialysis (Fig. 1). The variables and outcomes recorded in the PSMA study population are shown in Tables 1 and 2. To optimize the effectiveness of PSMA by reducing the number of unmatched cases, continuous variables were categorized according to their median values.

Outcomes

All enrolled patients were followed up for 8 weeks after surgery by local investigators, who were left free to manage the perioperative interval according to their usual local criteria, including any additional exam and time to discharge. Any adverse event was recorded and graded according to Clavien-Dindo³⁰ and the Japanese Clinical Oncology Group (JCOG) extended criteria³¹ as well as any reoperation, readmission or death. Anastomotic leakage (AL) was defined according to the international consensus³². All the outcomes were calculated at 60 days after surgery.

The primary endpoint was the duration of postoperative hospital stay (LOS, inclusive of any readmission) either dichotomized according to its median value or considered as a continuous variable. The secondary endpoints were: superficial and/or deep surgical site infections (s-d-SSI), defined as drain-specific complications including purulent drainage from superficial incisions, positive culture of fluid or tissue from superficial incisions, pain or tenderness, localized swelling, redness, heat, and/or infections involving deep fascial and muscle layers without dehiscence³³; deep wound dehiscence; abdominal collection/abscess defined as intraperitoneal postoperative collections that altered the normal postoperative course, requiring either medical, radiological, endoscopic or surgical intervention³³; SSI defined as s-d-SSI plus abdominal collection/abscess plus deep wound dehiscence; infectious morbidity rate defined as SSI plus pulmonary infections plus urinary infections; AL; overall morbidity rate (any adverse event); major morbidity rate (any adverse event grade > II); reoperation (any unplanned operation) rates; mortality (any death) rates.

Statistical analysis

This was a retrospective PSMA of two prospective cohorts, with sample sizes calculated and reported in the respective core papers^{24,25}. Events per variable guideline were followed³⁴. There were no missing data in the database of 6157 patients. The target of estimand was represented by the average treatment effect in the true population of interest (ATT).




Fig. 1 Study flow chart according to the reporting and guidelines in propensity score analysis²⁹

iCral, Italian ColoRectal Anastomotic Leakage study group; ERAS, enhanced recovery after surgery; SMD, standardized mean difference.

A propensity score-matching model^{35,36} was used for the analysis (Fig. 1). An adjusted logistic regression was used to estimate the propensity scores of the treatment and control groups. The exposure variable was a treatment that implied no abdominal drain(s) placement in elective colorectal surgery, and 21 covariates, potentially affecting the treatment³⁷, were selected: age, sex, ASA class, BMI, diabetes, chronic renal failure, chronic liver disease, nutritional status measured through the Mini Nutritional Assessment—Short Form (MNA-SF)³⁸, surgery for malignancy, centre volume, hospital type (academic/metropolitan versus local/regional), surgical unit type (general versus oncologic/colorectal), mini-invasive surgery, standard surgical procedure, operation length (minutes), intra- or extracorporeal anastomosis, stapled versus handsewn anastomosis, end-to-end anastomosis, preoperative blood transfusion(s), intra- and/or postoperative blood transfusion(s), and overall ERAS pathway adherence rates.

To ensure that the treatment groups were balanced³⁹, a PSMA using the software ‘R’[®] (Version 4.2.2, The R Foundation[®] for Statistical Computing, Vienna, Austria, 2022) was performed. A

nearest neighbour approach with a logit distance metric and a caliper of 0.1 to minimize differences between the groups was used as well as adjusted logistic regression to estimate the association between the treatment variable and outcomes.

Balance in the matched groups was assessed by calculating the standardized mean difference (SMD), using a threshold of 0.1 (an SMD less than 0.1 typically indicates a negligible difference between the means of the groups) and the general variance ratio (a variance ratio close to 1 indicates that variances are equal in the two groups). For outcome modelling, an adjusted logistic regression was performed based on a treatment variable represented by no abdominal drain placement in elective colorectal surgery and on the same 21 covariates selected for the PSMA⁴⁰, presenting odds ratios (OR) and 95 per cent c.i. The eventual effect of any unobserved confounder was tested through a sensitivity analysis⁴¹, using the library ‘SensitivityR5’ of the software R[®] (Version 4.2.2, The R Foundation[®] for Statistical Computing, Vienna, Austria, 2022) and presenting the Γ values (each 0.1 increment of Γ values

Table 1 Descriptive analysis of the variables considered in the 6157 patients evaluated by the Italian ColoRectal Anastomotic Leakage study group (iCral)

	Overall (n = 6157)	No drain(s) (n = 2168)	Drain(s) (n = 3989)	P*
Age (years)				
< 70	3133 (50.9)	1112 (51.3)	2021 (50.1)	0.640
≥ 70	3024 (49.1)	1056 (48.7)	1968 (49.3)	
Sex				
Male	3205 (52.0)	1059 (48.2)	2146 (53.8)	<0.010
Female	2952 (48.0)	1109 (51.2)	1843 (46.2)	
ASA class				
I-II	3916 (63.6)	1455 (67.1)	2461 (61.7)	<0.010
III	2241 (36.4)	713 (32.9)	1528 (38.3)	
BMI (kg/m²)				
≤ 25.25	3092 (50.2)	1138 (52.5)	1954 (49.0)	0.010
> 25.25	3065 (49.8)	1030 (47.5)	2035 (51.0)	
Diabetes				
Yes	917 (14.9)	282 (13.0)	635 (15.9)	<0.010
No	5240 (85.1)	1886 (87.0)	3354 (84.1)	
Chronic renal failure				
Yes	256 (4.2)	93 (4.3)	163 (4.1)	0.700
No	5901 (95.8)	2075 (95.7)	3826 (95.9)	
Chronic liver disease				
Yes	66 (1.1)	16 (0.7)	50 (1.2)	0.060
No	6091 (98.9)	2152 (99.3)	3939 (98.8)	
MNA-SF				
≤ 12	3282 (53.3)	1025 (47.3)	2257 (56.6)	<0.010
> 12	2875 (46.7)	1143 (52.7)	1732 (43.4)	
Surgery for malignancy				
Yes	4496 (73.0)	1655 (76.3)	2841 (71.2)	<0.010
No	1661 (27.0)	513 (23.7)	1148 (28.8)	
Diverticular disease	882 (53.1)	290 (13.4)	592 (14.8)	
Endometriosis	45 (2.7)	2 (0.1)	43 (1.1)	
Polyps	318 (19.2)	141 (6.5)	177 (4.4)	
IBD	180 (10.8)	32 (1.5)	148 (3.7)	
Other	236 (14.2)	48 (2.2)	188 (4.7)	
Mini-invasive surgery				
No	913 (14.8)	134 (6.2)	779 (19.5)	<0.010
Yes	5244 (85.2)	2034 (93.8)	3210 (80.5)	
Laparoscopic	4441 (84.7)	1815 (83.7)	2626 (65.8)	
Robotic	508 (9.7)	178 (8.2)	330 (8.3)	
Converted	295 (5.6)	41 (1.9)	254 (6.4)	
Standard procedure				
Yes	5192 (84.3)	1940 (89.5)	3252 (81.5)	<0.010
Right colectomy	2852 (54.9)	1177 (54.3)	1675 (42.0)	
Left colectomy	2029 (39.1)	684 (31.6)	1345 (33.7)	
Anterior resection	311 (6.0)	79 (3.6)	232 (5.8)	
No	965 (15.7)	228 (10.5)	737 (18.5)	
Transverse colectomy	154 (16.0)	45 (2.1)	109 (2.7)	
Splenic flexure colectomy	218 (22.6)	72 (3.3)	146 (3.7)	
Hartmann reversal	149 (15.4)	24 (1.1)	125 (3.1)	
(Sub) total colectomy	120 (12.4)	24 (1.1)	96 (2.4)	
Other	324 (33.6)	63 (2.9)	261 (6.5)	
Anastomosis 1				
Intracorporeal	3964 (64.4)	1779 (82.1)	2185 (54.3)	<0.010
Extracorporeal	2193 (35.6)	389 (17.9)	1804 (45.2)	
Anastomosis 2				
Stapled	5460 (88.7)	2043 (94.2)	3417 (85.7)	<0.010
Handsewn	697 (11.3)	125 (5.8)	572 (14.3)	
Anastomosis 3				
End-to-end	2467 (40.1)	779 (35.9)	1688 (42.3)	<0.010
Other shape	3690 (59.9)	1389 (64.1)	2301 (57.7)	
Operation length (min)				
≤ 170	3169 (51.5)	1265 (58.3)	1904 (47.7)	<0.010
> 170	2988 (48.5)	903 (41.7)	2085 (52.3)	
Hospital type				
Met./ac.	4012 (65.2)	1459 (67.3)	2553 (64.0)	0.010
Local/regional	2145 (34.8)	709 (32.7)	1436 (36.0)	
Unit type				
Colorectal/oncologic	1107 (18.0)	372 (17.2)	735 (18.4)	0.220
General	5050 (82.0)	1796 (82.8)	3254 (81.6)	

(continued)

Table 1 (continued)

	Overall (n = 6157)	No drain(s) (n = 2168)	Drain(s) (n = 3989)	P*
Centre volume				
< 4 patients/month	1822 (29.6)	577 (26.6)	1245 (31.2)	
≥ 4 patients/month	4335 (70.4)	1591 (73.4)	2744 (68.8)	<0.010
Preoperative BT(s)				
Yes	374 (6.1)	127 (5.9)	247 (6.2)	
No	5783 (93.9)	2041 (94.1)	3742 (93.8)	0.600
Intra-/postoperative BT(s)				
Yes	417 (6.8)	114 (5.3)	303 (7.6)	<0.010
No	5740 (93.2)	2054 (94.7)	3686 (92.4)	
Overall ERAS adherence (%)				
≤ 75.0	3161 (51.3)	668 (30.8)	2493 (62.5)	
> 75.0	2996 (48.7)	1500 (69.2)	1496 (37.5)	<0.010
Nutritional screening	4170 (67.7)	1628 (75.1)	2542 (63.7)	
Prehabilitation	2386 (38.8)	1097 (50.6)	1289 (32.3)	
Counselling	4073 (66.2)	1716 (79.2)	2357 (59.1)	
Immune enhancing nutrition	1830 (29.7)	854 (39.4)	976 (24.5)	
Antithrombotic prophylaxis	5607 (91.1)	2023 (93.3)	3584 (89.9)	
Antibiotic prophylaxis	5771 (93.7)	2061 (95.1)	3710 (93.0)	
No mechanical bowel preparation	4257 (69.1)	1784 (82.3)	2473 (62.0)	
Preoperative carbohydrates load	3449 (56.0)	1520 (70.1)	1929 (48.4)	
No preanaesthesia	4739 (77.0)	1857 (85.7)	2882 (72.3)	
Standard anaesthesia protocol	4936 (80.2)	1862 (85.9)	3074 (77.1)	
Normothermia	5588 (90.8)	2039 (94.1)	3549 (89.0)	
Goal-directed or restrictive fluid therapy	4738 (77.0)	1816 (83.8)	2922 (73.3)	
Postoperative nausea/vomit prophylaxis	5253 (85.3)	1927 (88.9)	3326 (83.4)	
Multimodal analgesia	5434 (88.3)	2048 (94.5)	3386 (84.9)	
No nasogastric tube	5145 (83.6)	2064 (95.2)	3081 (77.2)	
Minimally invasive surgery	5244 (85.2)	2034 (93.8)	3210 (80.5)	
Urinary catheter < 24–48 h	4746 (77.1)	1971 (90.9)	2775 (69.6)	
Early mobilization	3501 (56.9)	1593 (73.5)	1908 (47.8)	
Early oral feeding	3243 (52.7)	1574 (72.6)	1669 (41.8)	
Preadmission check	4916 (79.8)	2025 (93.4)	2891 (72.5)	

Values are n (%) unless otherwise stated. *Chi square independence test with one degree of freedom; MNA-SF, Mini Nutritional Assessment—Short Form; IBD, inflammatory bowel disease; Intracorporeal, anastomosis performed under visual control through the scope; Extracorporeal, anastomosis performed under direct visual control through an open access; Met./ac., metropolitan/academic; BT, blood transfusion; ERAS, enhanced recovery after surgery items.

representing a 10 per cent odds of differential assignment to treatment due to any unobserved variable).

Results

A total of 8359 patients who underwent colorectal resection with anastomosis were enrolled in two consecutive studies upon explicit inclusion/exclusion criteria, in 78 surgical centres in Italy from January 2019 to September 2021: iCral2²⁴ and iCral3²⁵.

The overall rate of abdominal drain placement after elective colorectal surgery was 64.8 per cent (3989 of 6157 patients). Tables 1 and 2 provide descriptions of the study covariates and, regarding univariable outcome analysis, drain omission was significantly associated with a lower risk of s-d-SSI, SSI, overall morbidity rate, mortality rate and LOS >6 days. The prevalence characteristics of the 3989 patients in whom abdominal drain(s) were placed are reported in Table 2. Drain(s) placement was significantly prevalent in males, ASA III, BMI >25.25 kg/m², diabetes, MNA-SF ≤12, surgery for benign disease open surgery, non-standard procedures (transverse colectomy, splenic flexure colectomy, Hartmann reversal, (sub) total colectomy, other) in comparison to standard procedures (right colectomy, left colectomy, anterior resection), extracorporeal anastomosis, handsewn anastomosis, end-to-end anastomosis, operation length >170 min, local/regional hospitals in comparison to metropolitan/academic hospitals, centre volume <4 patients/month, intra/postoperative blood transfusion(s), overall ERAS adherence <75 per cent.

For the PSMA, 3604 patients were included, and two groups of 1802 patients were generated (Fig. 1): group A (no abdominal drain(s), true population of interest), and group B (abdominal drain(s), control population). This population of 3604 patients included data deriving from 77 (98.7 per cent) of the original 78 centres: group A included data deriving from 60 (77.9 per cent) centres and group B from 75 (97.4 per cent) centres. A good balance between the two groups was achieved, SMD within 0.1 (Table 3 and Fig. 2), with a model variance ratio of 1.0843.

Group A versus group B showed a significantly lower risk of LOS >6 days (408 (22.6 per cent) versus 575 (31.9 per cent) events; OR 0.60; 95 per cent c.i. 0.51–0.70; P < 0.001). Sensitivity analysis for LOS calculated a Γ of 1.5 (P upper bound = 0.090), meaning that assuming the probabilities of assignment to the two treatment groups to be different because of unknown and/or unmeasured confounding variables, 50 per cent of patients should have been treated by drain(s) placement instead of omission to alter the significant association between drain(s) omission and LOS <6 days. The overall mean(standard deviation (s.d.)) LOS was 5.77(5.77) days in group A versus 6.63(5.70) days in group B (P < 0.0001; two tailed Student's t test with equal variances), with a mean difference of 0.86 days in favour of group A.

No difference was recorded between the two groups regarding all the other endpoints: s-d-SSI (OR 0.98; 95 per cent c.i. 0.64–1.48; P = 0.900); deep wound dehiscence (OR 2.20; 95 per cent c.i. 0.52–9.30; P = 0.280); abdominal collection/abscess (OR 1.13; 95 per cent c.i. 0.64–1.99; P = 0.670); SSI (OR 1.15; 95 per cent c.i. 0.82–

Table 2 Descriptive analysis of the outcomes considered in the 6157 patients evaluated by the Italian ColoRectal Anastomotic Leakage

	Overall	No drain(s)	Drain(s)	OR (95% c.i.)*
s-d-SSI				
Yes	208 (3.4)	52 (2.4)	156 (3.9)	0.60 (0.44–0.83) P <0.010
No	5949 (96.6)	2116 (97.6)	3833 (96.1)	Reference
Deep wound dehiscence				
Yes	14 (0.2)	6 (0.3)	8 (0.2)	1.38 (0.48–3.99) P = 0.550
No	6143 (99.8)	2162 (99.7)	3981 (99.8)	Reference
Abdominal collection/abscess				
Yes	87 (1.4)	29 (1.3)	58 (1.6)	0.92 (0.59–1.44) P = 0.710
No	6070 (98.6)	2139 (98.7)	3931 (98.5)	Reference
SSI				
Yes	290 (4.7)	84 (3.9)	206 (5.2)	0.74 (0.57–0.96) P = 0.020
No	5867 (95.3)	2084 (96.1)	3783 (94.8)	Reference
Infectious morbidity rate				
Yes	401 (6.5)	125 (5.8)	276 (6.9)	0.82 (0.66–1.02) P = 0.080
No	5756 (93.5)	2043 (94.2)	3713 (93.1)	Reference
Reoperation on				
Yes	284 (4.6)	101 (4.7)	183 (4.6)	1.02 (0.79–1.30) P = 0.900
No	5873 (95.5)	2067 (95.3)	3806 (95.4)	Reference
LOS				
≤ 6 days	3966 (64.4)	1683 (77.6)	2283 (57.2)	Reference
> 6 days	2191 (35.6)	485 (22.4)	1706 (42.8)	0.39 (0.34–0.43) P <0.010
Anastomotic leakage				
Yes	211 (3.4)	67 (3.1)	144 (3.6)	0.85 (0.63–1.14) P = 0.280
No	5946 (96.6)	2101 (96.9)	3845 (96.4)	Reference
Overall morbidity rate				
Yes	1666 (27.1)	542 (25.0)	1124 (28.2)	0.85 (0.75–0.96) P = 0.010
No	4491 (72.9)	1626 (75.0)	2865 (71.8)	Reference
Major morbidity rate				
Yes	331 (5.4)	108 (5.0)	223 (5.6)	0.89 (0.70–1.12) P = 0.310
No	5826 (94.6)	2060 (95.0)	3766 (94.4)	Reference
Mortality rate				
Yes	56 (0.9)	10 (0.5)	46 (1.2)	0.40 (0.20–0.79) P = 0.010
No	6101 (99.1)	2158 (99.5)	3943 (98.8)	Reference
LOS (days)	6.89(6.08)† (6 (4–7)‡)	5.67(5.57)† (4 (3–6)‡)	7.55(6.24)† (6 (5–8)‡)	

Values are n (%) unless otherwise stated. *Univariate ORs estimation with Wolf valuation of the c.i.; †Mean(s.d.). ‡Median (i.q.r.). s-d-SSI, superficial and/or deep surgical site infections; SSI, s-d-SSI plus deep wound dehiscence plus abdominal collection/abscess; Infectious morbidity rate, s-d-SSI plus deep wound dehiscence plus abdominal collection/abscess plus pulmonary infections plus urinary infections; LOS, length of postoperative hospital stay.

1.62; P = 0.420); infectious morbidity rate (OR 1.21; 95 per cent c.i. 0.90–1.62; P = 0.190); AL (OR 0.99; 95 per cent c.i. 0.67–1.46; P = 0.950); overall morbidity rate (OR 1.06; 95 per cent c.i. 0.90–1.24; P = 0.480); major morbidity rate (OR 1.11; 95 per cent c.i. 0.81–1.52; P = 0.500); reoperation rate (OR 1.19; 95 per cent c.i. 0.85–1.66; P = 0.300); mortality rate (OR 0.67; 95 per cent c.i. 0.27–1.68; P = 0.390).

Discussion

This study presents data on a retrospective PSMA of a prospective multicentre database comparing drain(s) versus no drain(s) placement after elective colorectal surgery. This study involved 78 surgical centres, representing a snapshot of real-life clinical practice in Italy. Abdominal drain(s) placement after elective colorectal surgery was performed in 64.8 per cent of 6157 patients, and the univariable analysis of this population demonstrated a statistically significant association between drain(s) placement and a higher risk of s-d-SSI, SSI, overall morbidity rate, mortality rate and prolonged LOS, confirming the observations of previous studies^{4,7,8,11–16}. Conversely, our PSMA showed that omission of drain(s) placement after elective colorectal surgery was significantly associated with a lower risk of LOS >6 days, albeit with a small and not clinically significant reduction of 0.86 days mean difference. No statistically significant association was detected for secondary outcomes.

The main aim of the present analysis was to identify any reason supporting the use of drains by Italian (and European) surgeons following elective colorectal resections; there was no single reason to support their use. While LOS is an important outcome for hospital managers and for costs associated with the care of patients with colorectal diseases, it is of relatively little interest to patients and surgeons compared with other endpoints such as AL, major morbidity rate, reoperation rate and quality of life. This study did not demonstrate any difference in the risk of AL, major adverse events and reoperations. This disproves the possible role of abdominal drain(s) on earlier diagnosis and treatment of AL, for which we have highlighted the role of the joint use of clinical scores, C-reactive protein and procalcitonin⁴². The use of abdominal and pelvic drain(s) will continue to exist in a minority (for example, <20 per cent) of selected patients (low rectal anastomoses, immunocompromised and/or frail patients, heavily contaminated or dirty procedures, excessive blood loss and/or intraoperative complications). However, the routine placement is not supported⁴³, and a progressive de-implementation strategy should be actively sought at organizational and surgeon levels⁴⁴.

A recent retrospective PSMA of a prospective international cohort²³ on the same topic used a 'full matching' model, which may result in bias as some observations may not have suitable matches.

Table 3 Variables distribution in treatment and control groups before and after propensity score-matching

Covariates	Before propensity score-matching				After propensity score-matching			
	No drain(s) n = 2168 (35.2%)	Drain(s) n = 3989 (64.8%)	P*	SMD	No drain(s) n = 1802 (50.0%)	Drain(s) n = 1802 (50.0%)	P*	SMD
Age								
< 70 years	1112	2021	0.660	-0.01	903	919	0.620	0.02
≥ 70 years	1056	1968	0.660	0.01	899	883	0.620	-0.02
Sex								
Male	1059	2146	<0.010	0.01	903	921	0.570	0.02
Female	1109	1843	<0.010	-0.01	899	881	0.570	-0.02
ASA class								
I-II	1455	2461	<0.010	-0.11	1192	1188	0.920	-0.005
III	713	1528	<0.010	0.11	610	614	0.920	0.005
BMI (kg/m²)								
≤ 25.25	1138	1954	<0.010	-0.07	892	898	0.870	0.01
> 25.25	1030	2035	<0.010	0.07	910	904	0.870	-0.01
Diabetes								
Yes	282	635	<0.010	0.08	257	241	0.470	-0.03
No	1886	3354	<0.010	-0.08	1545	1561	0.470	0.03
Chronic renal failure								
Yes	93	163	0.750	-0.01	78	73	0.740	-0.01
No	2075	3826	0.750	0.01	1724	1729	0.740	0.01
Chronic liver disease								
Yes	16	50	0.080	0.05	15	11	0.550	-0.03
No	2152	3939	0.080	-0.05	1787	1791	0.550	0.03
MNA-SF								
≤ 12	1025	2257	<0.010	0.19	901	908	0.840	0.01
> 12	1143	1732	<0.010	-0.19	901	894	0.840	-0.01
Surgery for malignancy								
Yes	1655	2841	<0.010	-0.12	1344	1344	1.000	0.00
No	513	1148	<0.010	0.12	458	458	1.000	0.00
Mini-invasive surgery								
Yes	2034	3210	<0.010	-0.41	1672	1656	0.350	-0.03
No	134	779	<0.010	0.41	130	146	0.350	0.03
Standard procedures								
Yes	1940	3252	<0.010	-0.23	1596	1565	0.130	-0.05
No	228	737	<0.010	0.23	206	237	0.130	0.05
Anastomosis 1								
Intracorporeal	1779	2185	<0.010	-0.61	1422	1396	0.310	-0.03
Extracorporeal	389	1804	<0.010	0.61	380	406	0.310	0.03
Anastomosis 2								
Stapled	2043	3417	<0.010	-0.29	1681	1675	0.740	-0.01
Handsewn	125	572	<0.010	0.29	121	127	0.740	0.01
Anastomosis 3								
End-to-end	779	1688	<0.010	0.13	704	700	0.920	-0.004
Other shape	1389	2301	<0.010	-0.13	1098	1102	0.920	0.004
Operation length								
≤ 170 min	1265	1904	<0.010	-0.21	1024	986	0.210	-0.04
> 170 min	903	2085	<0.010	0.21	778	816	0.210	0.04
Hospital type								
Met./ac.	1459	2553	0.010	-0.07	1169	1193	0.420	0.03
Local/regional	709	1436	0.010	0.07	633	609	0.420	-0.03
Unit type								
Col/onc	372	735	0.230	0.03	327	324	0.930	-0.004
General	1796	3254	0.230	-0.03	1475	1478	0.930	0.004
Centre volume								
Low	577	1245	<0.010	0.10	513	432	<0.010	-0.10
High	1591	2744	<0.010	-0.10	1289	1370	<0.010	0.10
Preoperative BT								
Yes	127	247	0.640	0.01	112	108	0.830	-0.01
No	2041	3742	0.640	-0.01	1690	1694	0.830	0.01
Intrapostoperative BT								
Yes	114	303	<0.010	0.10	106	104	0.940	-0.005
No	2054	3686	<0.010	-0.10	1696	1698	0.940	0.005
Overall ERAS adherence								
≤ 75.0%	668	2493	<0.010	0.67	656	657	1.000	0.001
> 75.0%	1500	1496	<0.010	-0.67	1146	1145	1.000	-0.001

*Student's test for proportions. SMD, standardized mean difference; MNA-SF, Mini Nutritional Assessment-Short Form; Intracorporeal, anastomosis performed under visual control through the scope; Extracorporeal, anastomosis performed under direct visual control through an open access; Met./ac., metropolitan/academic; Col/onc: colorectal/oncologic; BT, blood transfusion; ERAS, enhanced recovery after surgery items.




Fig. 2 Love plot of covariate standardized mean differences between treatment and control groups before and after matching; the vertical lines represent the interval of ± 0.1 within which balance is considered acceptable ERAS, enhanced recovery after surgery.

The main strength of this large sample size study is that it followed rigorous guidelines for applying PSMA^{29,45}, being based on the following items: rigorous patient selection from the parent population, performed upon explicit criteria: to limit data imbalance, several potential confounders related to the surgical procedure (delayed urgency, operations without any abdominal incision/trans-anal procedures) or exclusively impacting on a subgroup of patients (anastomosis located <10 cm from the anal verge, neo-adjuvant therapy, proximal protective stoma, administration of perioperative steroids, patients treated by dialysis) were excluded; a reasoned inclusion of 21 conditioning variables (covariates): hospital type, surgical unit type and centre volume to account for the potential imbalance of multicentre, clustered data; adherence to the ERAS pathway items to account for the potential imbalance of medical, anaesthetic and surgical perioperative management; resections for benign and malignant diseases, mini-invasive or open surgery, standard and non-standard procedures²⁴, intracorporeal (anastomosis performed under visual control through the scope) or extracorporeal (anastomosis performed under direct visual control through an open access) anastomoses, stapled or handsewn anastomoses, end-to-end or different fashion anastomoses, and operation length, in relation to the imbalance of the surgical treatment; pre- and intrapostoperative blood

transfusion(s) to account for transfusion-related morbidity rate⁴⁶; age, sex, ASA class, body mass index, diabetes, chronic renal failure, chronic liver disease, and Mini Nutritional Assessment–Short Form, to account for patient imbalance; evaluation of the treatment effect through an adjusted multiple regression model including the same 21 covariates used for matching⁴⁰; a clear, sheer and restrictive balance algorithm (Fig. 1); a sensitivity analysis for unmeasured confounders.

Another strength of this study was the large number of enrolled patients in a well-defined time-lapse in a large number of centres, representing a very wide sample of surgical units performing colorectal resections in Italy. Although the multicentre nature of the considered data may be a definite source of clustering bias, it is undoubtedly representative of real-life data.

However, this study has several limitations, and its results should be interpreted with caution. First, several controversial risk factors were not measured or recorded in the parent studies: single surgeon's experience⁴⁷, material, type and time to removal of drain(s)⁴⁸, and indication (routine or selective) for drain(s) placement²³. Second, although a sensitivity analysis of unmeasured confounders has been conducted, potential residual unknown factors and the inability to rule out potential measurement errors by the participating investigators, may have had an impact on the results.

This study confirms that abdominal drain(s) placement after elective colorectal surgery is linked to a slightly prolonged non-clinically relevant LOS, without influencing anastomotic leakage, major morbidity rate and reoperation rate. Abdominal drains should not be routinely used in elective colorectal surgery.

Collaborators

Assistance with the study: iCral study group co-investigators: Stefano Mancini²⁶, Gian Luca Baiocchi²⁷, Roberto Santoro²⁸, Walter Siquini²⁹, Gianluca Guercioni³, Massimo Basti³⁰, Corrado Pedrazzani³¹, Mauro Totis³², Alessandro Carrara³³, Andrea Lucchi³⁴, Maurizio Pavanello³⁵, Andrea Muratore³⁶, Stefano D'Ugo³⁷, Alberto Di Leo³⁸, Giusto Pignata³⁹, Ugo Elmore⁴⁰, Gabriele Anania⁴¹, Massimo Carlini⁴², Francesco Corcione⁴³, Nereo Vettoretto⁴⁴, Graziano Longo⁴⁵, Mario Sorrentino⁴⁶, Antonio Giuliani⁴⁷, Giovanni Ferrari⁴⁸, Lucio Taglietti⁴⁹, Augusto Verzelli⁵⁰, Mariantonietta Di Cosmo⁵¹, Davide Cavaliere⁵², Marco Milone⁵³, Stefano Rauseri⁵⁴, Giovanni Ciaccio⁵⁵, Giovanni Tebala⁵⁶, Giuseppe Brisinda⁵⁷, Stefano Berti⁵⁸, Paolo Millo⁵⁹, Luigi Boni⁶⁰, Mario Guerrieri⁶¹, Roberto Persiani⁶², Dario Parini⁶³, Antonino Spinelli⁶⁴, Michele Genna⁶⁵, Vincenzo Bottino⁶⁶, Andrea Coratti⁶⁷, Dario Scala⁶⁸, Umberto Rivolta⁶⁹, Micaela Piccoli⁷⁰, Carlo Talarico⁷¹, Franco Roviello⁷², Alessandro Anastasi⁷³, Giuseppe Maria Ettore⁷⁴, Mauro Montuori⁷⁵, Pierpaolo Mariani⁷⁶, Nicolò de Manzini⁷⁷, Annibale Donini⁷⁸, Mariano Fortunato Armellino⁷⁹, Carlo Feo⁸⁰, Silvio Guerrieri⁸¹, Andrea Costanzi⁸², Federico Marchesi⁸³, Moreno Cicetti⁸⁴, Paolo Ciano², Michele Benedetti², Leonardo Antonio Montemurro², Maria Sole Mattei², Elena Belloni², Daniela Apa², Matteo Di Carlo², Elisa Bertocchi⁷, Gaia Masini⁷, Amedeo Altamura⁸, Francesco Rubichi⁸, Desirée Cianflocca¹⁰, Marco Migliore¹⁰, Diletta Cassini^{11,12}, Lorenzo Pandolfini¹³, Alessandro Falsetto¹³, Antonio Sciuto¹⁴, Ugo Pace¹⁵, Andrea Fares Bucci¹⁵, Francesco Monari¹⁶, Grazia Maria Attinà¹⁷, Angela Maurizi¹⁸, Michele Simone¹⁹, Francesco Giudici²⁰, Fabio Cianchi²⁰, Bruno Sensi²¹, Alessandra Aprile²², Domenico Soriero²², Andrea Scarinci²³, Gabriella Teresa Capolupo²⁴, Valerio Sisti²⁵, Marcella Lodovica Ricci²⁵, Andrea Sagnotta²⁶, Sarah Molfino²⁷, Pietro Amodio²⁸, Alessandro Cardinali²⁹, Simone Cicconi³, Irene Marziali³, Diletta Frazzini³⁰, Cristian Conti³¹, Nicolò Tamini³², Marco Braga³², Michele Motter³³, Giuseppe Tirone³³, Giacomo Martorelli³⁴, Alban Cacurri³⁴, Carlo Di Marco³⁵, Patrizia Marsanic³⁶, Nicoletta Sveva Pipitone Federico³⁶, Marcello Spampinato³⁷, Lorenzo Crepaz³⁸, Jacopo Andreuccetti³⁹, Iliara Canfora³⁹, Giulia Maggi⁴⁰, Matteo Chiozza⁴¹, Domenico Spoleitini⁴², Rosa Marcellinaro⁴², Umberto Bracale⁴³, Roberto Peltrini⁴³, Maria Michela Di Nuzzo⁴³, Emanuele Botteri⁴⁴, Simone Santoni⁴⁵, Massimo Stefanoni⁴⁶, Giovanni Del Vecchio⁴⁷, Carmelo Magistro⁴⁸, Silvia Ruggiero⁴⁹, Arianna Birindelli⁴⁹, Andrea Budassi⁵⁰, Daniele Zigiotta⁵¹, Leonardo Solaini⁵², Giorgio Ercolani⁵², Giovanni Domenico De Palma⁵³, Silvia Tenconi⁵⁴, Paolo Locurto⁵⁵, Antonio Di Cintio⁵⁶, Maria Michela Chiarello⁵⁷, Maria Cariati⁵⁷, Andrea Gennai⁵⁸, Manuela Grivon⁵⁹, Elisa Cassinotti⁶⁰, Monica Ortenzi⁶¹, Alberto Biondi⁶², Maurizio De Luca⁶³, Francesco Carrano⁶⁴, Francesca Fior⁶⁵, Antonio Ferronetti⁶⁶, Giuseppe Giuliani⁶⁷, Graziella Marino⁶⁸, Camillo Leonardo Bertoglio⁶⁹, Francesca Pecchini⁷⁰, Vincenzo Greco⁷¹, Roberto Piagnerelli⁷², Giuseppe Canonico⁷³, Marco Colasanti⁷⁴, Enrico Pinotti⁷⁵, Roberta Carminati⁷⁶, Edoardo Osenda⁷⁷, Luigina Graziosi⁷⁸, Ciro De Martino⁷⁹, Giovanna Ioia⁷⁹, Fioralba Pindozi⁸⁰, Lorenzo Organetti⁸¹, Michela Monteleone⁸², Giorgio Dalmonte⁸³, Gabriele La Gioia⁸⁴.

From: ²⁶General & Oncologic Surgery Unit, San Filippo Neri Hospital, ASL Roma 1; ²⁷General Surgery Unit 3, Department of Clinical and Experimental Sciences, University of Brescia; ²⁸General Oncologic Surgery Unit, Belcolle Hospital, Viterbo; ²⁹General Surgery Unit, S. Lucia Hospital, Macerata; ³⁰General Surgery Unit, Spirito Santo Hospital, Pescara; ³¹General & HPB Surgery Unit, University Hospital, Verona; ³²Colorectal Surgery Unit, San Gerardo Hospital, ASST Monza; ³³1st General Surgery Unit, S. Chiara Hospital, Trento; ³⁴General Surgery Unit, 'Ceccarini' Hospital, Riccione (RN); ³⁵General Surgery Unit, AULSS2 Marca Trevigiana, Conegliano Veneto (TV); ³⁶General Surgery Unit, 'E. Agnelli' Hospital, Pinerolo (TO); ³⁷General Surgery Unit, 'V. Fazzi' Hospital, Lecce; ³⁸General and Minimally Invasive Surgery Unit, San Camillo Hospital, Trento; ³⁹2nd General Surgery Unit 2, Spedali Civili di Brescia; ⁴⁰Gastroenterologic Surgery Unit, IRCCS S. Raffaele Hospital, Milano; ⁴¹General & Laparoscopic Surgery Unit, University Hospital, Ferrara; ⁴²General Surgery Unit, S. Eugenio Hospital, ASL Roma 2; ⁴³General Oncologic and Mininvasive Surgery Unit, 'Federico II' University, Napoli; ⁴⁴General Surgery Unit, Spedali Civili of Brescia, Montichiari (BS); ⁴⁵General Surgery Unit, Policlinico Casilino, Roma; ⁴⁶General Surgery Unit, Latisana-Palmanova Hospital, Friuli Centrale University (UD); ⁴⁷General Surgery Unit, S. Carlo Hospital, Potenza; ⁴⁸General Oncologic and Mininvasive Surgery Unit, Great Metropolitan Niguarda Hospital, Milano; ⁴⁹General Surgery Unit, ASST Valcamonica, Esine (BS); ⁵⁰General Surgery Unit, Profili Hospital, Fabriano (AN); ⁵¹General & Upper GI Surgery Unit, University Hospital, Verona; ⁵²General & Oncologic Surgery Unit, AUSL Romagna, Forlì (FC); ⁵³General & Endoscopic Surgery Unit, 'Federico II' University, Napoli; ⁵⁴General Surgery Unit, Gallarate Hospital (VA); ⁵⁵General Surgery Unit, S. Elia Hospital, Caltanissetta; ⁵⁶General Surgery Unit, S. Maria Hospital, Terni; ⁵⁷General Surgery Unit, San Giovanni di Dio Hospital, Crotone; ⁵⁸General Surgery Unit, ASL 5 Liguria POLL, La Spezia; ⁵⁹General Surgery Unit, 'U. Parini' Regional Hospital, Aosta; ⁶⁰General Surgery Unit, Fondazione IRCCS Ca' Granda, Policlinico Maggiore Hospital, Milano; ⁶¹Surgical Clinic, Torrette Hospital, University of Ancona; ⁶²General Surgery Unit, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Roma; ⁶³General Surgery Unit, S. Maria della Misericordia Hospital, Rovigo; ⁶⁴Colorectal Surgery Unit, Humanitas University, Rozzano (MI); ⁶⁵General & Bariatric Surgery Unit, University Hospital, Verona; ⁶⁶General & Oncologic Surgery Unit, Evangelico Betania Hospital, Napoli; ⁶⁷General Surgery Unit, Misericordia Hospital, Grosseto; ⁶⁸Abdominal Oncologic Surgery Unit, Basilicata Oncologic Hospital, Rionero in Vulture (PZ); ⁶⁹General Surgery Unit, Fornaroli Hospital, ASST Ovest Milanese, Magenta (MI); ⁷⁰General Surgery Unit, Civil Hospital, Baggiovara (MO); ⁷¹General Surgery Unit, Villa dei Gerani Hospital, Vibo Valentia (VV); ⁷²Surgical Clinic, University of Siena; ⁷³General Surgery Unit, San Giovanni di Dio Hospital, Firenze; ⁷⁴General & Transplant Surgery Unit, San Camillo-Forlanini Hospital, Roma; ⁷⁵General & Mininvasive Surgery Unit, S. Pietro Hospital, Ponte San Pietro (BG); ⁷⁶General Surgery Unit, Pesenti Fenaroli Hospital, Alzano Lombardo (BG); ⁷⁷Surgical Clinic, University of Trieste; ⁷⁸General & Emergency Surgery Unit, University of Perugia; ⁷⁹General & Emergency Surgery Unit, S. Giovanni di Dio e Ruggi d'Aragona Hospital, Salerno; ⁸⁰General Surgery Unit, Delta Hospital, Lagosanto (FE); ⁸¹General Surgery Unit, 'F. Murri' Hospital, Fermo; ⁸²General Surgery Unit, S. Leopoldo Hospital, Merate (LC); ⁸³Surgical Clinic, University of Parma; ⁸⁴General Surgery Unit, S. Maria della Misericordia Hospital, Urbino (PU); Italy.

Funding

The authors have no funding to declare. Medtronic SI® Italy provided unconditional support for the organization of three iCral2 study investigator meetings, held in Rome, Italy—October 2018, Matera, Italy—June 2019 and Bologna, Italy—October 2019.

Acknowledgements

M.Cat. is the study group coordinator.

Disclosure

M.Cat. reports personal fees from Baxter Spa outside the submitted work. The authors declare no other conflict of interest.

Data availability

Data are available upon reasonable request from M.Cat., iCral Study Group coordinator (e-mail: marco.catarci@aslroma2.it).

Author contributions

Stefano Guadagni (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Resources, Software, Validation, Visualization, Writing—original draft, Writing—review & editing), Marco Catarci (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing—original draft, Writing—review & editing), Francesco Masedu (Data curation, Formal analysis, Investigation, Supervision, Validation, Visualization, Writing—original draft, Writing—review & editing), Mohammad Karim (Data curation, Formal analysis, Supervision, Visualization, Writing—review & editing), Marco Clementi (Data curation, Investigation, Validation, Visualization, Writing—review & editing), Giacomo Ruffo (Conceptualization, Data curation, Investigation, Validation, Visualization, Writing—review & editing), Massimo Viola (Data curation, Investigation, Validation, Visualization, Writing—review & editing), Felice Borghi (Conceptualization, Data curation, Formal analysis, Supervision, Validation, Visualization, Writing—review & editing), Gianandrea Baldazzi (Data curation, Investigation, Validation, Writing—review & editing), Marco Scatizzi (Conceptualization, Data curation, Investigation, Supervision, Validation, Visualization, Writing—review & editing), Felice Pirozzi (Data curation, Investigation, Validation, Visualization, Writing—review & editing), Paolo Delrio (Conceptualization, Data curation, Validation, Visualization, Writing—review & editing), Gianluca Garulli (Data curation, Formal analysis, Investigation, Validation, Visualization, Writing—review & editing), Pierluigi Marini (Data curation, Formal analysis, Investigation, Validation, Visualization, Writing—review & editing), Roberto Campagnacci (Conceptualization, Data curation, Investigation, Supervision, Writing—review & editing), Raffaele De Luca (Data curation, Investigation, Validation, Visualization, Writing—review & editing), Ferdinando Ficari (Data curation, Investigation, Supervision, Validation, Visualization, Writing—review & editing), Giuseppe Sica (Data curation, Investigation, Validation, Visualization, Writing—review & editing), Stefano Scabini (Data curation, Investigation, Validation, Visualization, Writing—review &

editing), Andrea Liverani (Data curation, Investigation, Visualization, Writing—review & editing), Marco Caricato (Conceptualization, Data curation, Investigation, Supervision, Validation, Visualization, Writing—review & editing) and Alberto Patriti (Conceptualization, Data curation, Investigation, Visualization, Writing—review & editing).

References

- Smith SR, Gilmore OJ. Surgical drainage. *Br J Hosp Med* 1985;**33**: 308–315
- Memon MA, Memon MI, Donohue JH. Abdominal drains: a brief historical review. *Ir Med J* 2001;**94**:164–166
- Puleo F, Mishra N, Hall J. Use of intra-abdominal drains. *Clin Colon Rectal Surg* 2013;**26**:174–177
- Mujagic E, Zeindler J, Coslovsky M, Hoffmann H, Soysal SD, Mechera R et al. The association of surgical drains with surgical site infections—a prospective observational study. *Am J Surg* 2019;**217**:17–23
- Tsujinaka S, Konishi F. Drain vs no drain after colorectal surgery. *Indian J Surg Oncol* 2011;**2**:3–8
- Findik UY, Topcu SY, Vatansver O. Effects of drains on pain. Comfort and anxiety in patients undergone surgery. *Int J Caring Sci* 2013;**6**:412–419
- Gustafsson UO, Scott MJ, Hubner M, Nygren J, Demartines N, Francis N et al. Guidelines for perioperative care in elective colorectal surgery: Enhanced Recovery After Surgery (ERAS®) Society Recommendations: 2018. *World J Surg* 2019;**43**:659–695
- Irani JL, Hedrick TL, Miller TE, Lee L, Steinhagen E, Shogan BD et al. Clinical practice guidelines for enhanced recovery after colon and rectal surgery from the American Society of Colon and Rectal Surgeons and the Society of American Gastrointestinal and Endoscopic Surgeons. *Surg Endosc* 2023;**37**:5–30
- Alfonsi P, Slim K, Chauvin M, Mariani P, Faucheron JL, Fletcher D et al. Réhabilitation rapide après une chirurgie colorectale programmée [Guidelines for enhanced recovery after elective colorectal surgery]. *Ann Fr Anesth Reanim* 2014;**33**:370–384
- Ficari F, Borghi F, Catarci M, Scatizzi M, Alagna V, Bachini I et al. Enhanced recovery pathways in colorectal surgery: a consensus paper by the Associazione Chirurghi Ospedalieri Italiani (ACOI) and the PeriOperative Italian Society (POIS). *G Chir* 2019;**40**:1–40
- Sagar PM, Couse N, Kerin M, May J, MacFie J. Randomized trial of drainage of colorectal anastomosis. *Br J Surg* 1993;**80**:769–771
- Merad F, Yahouchi E, Hay JM, Fingerhut A, Laborde Y, Langlois-Zantain O. Prophylactic abdominal drainage after elective colonic resection and suprapromontory anastomosis: a multicenter study controlled by randomization. *Arch Surg* 1998;**133**:309–314
- Jesus EC, Karliczek A, Matos D, Castro AA, Atallah AN. Prophylactic anastomotic drainage for colorectal surgery. *Cochrane Database Syst Rev* 2004;**2004**:CD002100
- Petrowsky H, Demartines N, Rousson V, Clavien PA. Evidence-based value of prophylactic drainage. *Ann Surg* 2004;**240**:1074–1085
- Zhang HY, Zhao CL, Xie J, Ye YW, Sun JF, Ding ZH et al. To drain or not to drain in colorectal anastomosis: a meta-analysis. *Int J Colorectal Dis* 2016;**31**:951–960
- Podda M, Di Saverio S, Davies RJ, Atzeni J, Balestra F, Viridis F et al. Prophylactic intra-abdominal drainage following colorectal anastomoses. A systematic review and meta-analysis of randomized controlled trials. *Am J Surg* 2020;**219**:164–174
- Guyatt G, Guterma D, Baumann MH, Addrizzo-Harris D, Hylek EM, Phillips B et al. Grading strength of recommendations and quality of evidence in clinical guidelines: report from an American College of Chest Physicians Task Force. *Chest* 2006;**129**:174–181

18. van Rooijen SJ, Jongen ACHM, Wu ZG, Ji JF, Slooter GD, Roumen RMH et al. Definition of colorectal anastomotic leakage: a consensus survey among Dutch and Chinese colorectal surgeons. *World J Gastroenterol* 2017;**23**:6172–6180
19. Solaini L, Cavaliere D, Pecchini F, Perna F, Avanzolini A, Vitali G et al. The use of intra-abdominal drain in minimally invasive right colectomy: a propensity score-matched analysis on postoperative outcomes. *Int J Colorectal Dis* 2019;**34**:2137–2141
20. Taflampas P, Christodoulakis M, Tsiftsis DD. Anastomotic leakage after low anterior resection for rectal cancer: facts, obscurity, and fiction. *Surg Today* 2009;**39**:183–188
21. Ripollés-Melchor J, Ramírez-Rodríguez JM, Casans-Francés R, Aldecoa C, Abad-Motos A, Logroño-Egea M et al. Association between use of enhanced recovery after surgery protocol and postoperative complications in colorectal surgery: the postoperative outcomes within enhanced recovery after surgery protocol (POWER) study. *JAMA Surg* 2019;**154**:725–736
22. Ripollés-Melchor J, Abad-Motos A, Cecconi M, Pearse R, Jaber S, Slim K et al. Association between use of enhanced recovery after surgery protocols and postoperative complications in colorectal surgery in Europe: the EuroPOWER international observational study. *J Clin Anesth* 2022;**80**:110752
23. EuroSurg Collaborative. Intraperitoneal drain placement and outcomes after elective colorectal surgery: international matched, prospective, cohort study. *Br J Surg* 2022;**109**:520–529
24. Catarci M, Ruffo G, Viola MG, Pirozzi F, Delrio P, Borghi F et al. ERAS program adherence-institutionalization, major morbidity and anastomotic leakage after elective colorectal surgery: the iCral2 multicenter prospective study. *Surg Endosc* 2022; **36**:3965–3984
25. Italian ColoRectal Anastomotic Leakage (iCral) study group. Patient-reported outcomes, return to intended oncological therapy and enhanced recovery pathways after colorectal surgery: a prospective multicenter observational investigation by the Italian ColoRectal Anastomotic Leakage (iCral 3) study group. *Ann Surg Open* 2023;**4**:e267
26. Willis MA, Keller PS, Sommer N, Koch F, Ritz JP, Beyer K et al. Adherence to fast track measures in colorectal surgery—a survey among German and Austrian surgeons. *Int J Colorectal Dis* 2023;**38**:80
27. Nelson G, Kiyang LN, Crumley ET, Chuck A, Nguyen T, Paris P et al. Implementation of Enhanced Recovery After Surgery (ERAS) across a provincial healthcare system: the ERAS Alberta colorectal surgery experience. *World J Surg* 2016;**40**:1092–1103
28. Aarts MA, Rotstein OD, Pearsall EA, Victor JC, Okrainec A, McKenzie M et al. Postoperative ERAS interventions have the greatest impact on optimal recovery: experience with implementation of ERAS across multiple hospitals. *Ann Surg* 2018;**267**:992–997
29. Yao XI, Wang X, Speicher PJ, Hwang ES, Cheng P, Harpole DH et al. Reporting and guidelines in propensity score analysis: a systematic review of cancer and cancer surgical studies. *J Natl Cancer Inst* 2017;**109**:djw323
30. Dindo D, Demartines N, Clavien PA. Classification of surgical complications. A new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;**240**: 205–213
31. Katayama H, Kurokawa Y, Nakamura K, Ito H, Kanemitsu Y, Masuda N et al. Extended Clavien-Dindo classification of surgical complications: Japan Clinical Oncology Group postoperative complications criteria. *Surg Today* 2016;**46**:668–685
32. Rahbari NN, Weitz J, Hohenberger W, Heald RJ, Moran B, Ulrich A et al. Definition and grading of anastomotic leakage following anterior resection of the rectum: a proposal by the International Study Group of Rectal Cancer. *Surgery* 2010;**147**:339–351
33. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control* 2008;**36**:309–332
34. Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 1996;**49**:1373–1379
35. Austin PC. An introduction to propensity score methods for reducing the effects of confounding in observational studies. *Multivariate Behav Res* 2011;**46**:399–424
36. Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects. *Biometrika* 1983;**70**:41–55
37. Brookhart MA, Schneeweiss S, Rothman KJ, Glynn RJ, Avorn J, Stürmer T. Variable selection for propensity score models. *Am J Epidemiol* 2006;**163**:1149–1156
38. Kaiser MJ, Bauer JM, Ramsch C, Uter W, Guigoz Y, Cederholm T et al. Validation of the Mini Nutritional Assessment Short-Form (MNA-SF): a practical tool for identification of nutritional status. *J Nutr Health Aging* 2009;**13**:782–788
39. Austin PC. Optimal caliper widths for propensity-score matching when estimating differences in means and differences in proportions in observational studies. *Pharm Stat* 2011;**10**:150–161
40. Ho DE, Imai K, King G, Stuart EA. Matching as nonparametric preprocessing for reducing model dependence in parametric causal inference. *Polit Anal* 2007;**15**:199–236
41. Rosenbaum PR. *Design of Observational studies* (2nd edn). Springer Series in Statistics. New York, NY: Springer Nature Switzerland AG, 2020, 317–336
42. Italian ColoRectal Anastomotic Leakage (iCral) Study Group. Anastomotic leakage after elective colorectal surgery: a prospective multicentre observational study on use of the Dutch leakage score, serum procalcitonin and serum C-reactive protein for diagnosis. *BJS Open* 2020;**4**:499–507
43. Catarci M, Ruffo G, Viola MG, Pirozzi F, Delrio P, Borghi F et al. High adherence to enhanced recovery pathway independently reduces major morbidity and mortality rates after colorectal surgery: a reappraisal of the iCral2 and iCral3 multicenter prospective studies. *G Chir* 2023;**43**:2
44. Heus P, van Dulmen SA, Weenink JW, Naaktgeboren CA, Takada T, Verkerk EW et al. What are effective strategies to reduce low-value care? An analysis of 121 randomized deimplementation studies. *J Healthc Qual* 2023;**45**:261–271
45. Simoneau G, Pellegrini F, Debray TPA, Rouette J, Muñoz J, Platt RW et al. Recommendations for the use of propensity score methods in multiple sclerosis research. *Mult Scler* 2022;**28**:1467–1480
46. Catarci M, Guadagni S, Masedu F, Montemurro LA, Ciano P, Benedetti M et al. Blood transfusions and adverse events after colorectal surgery: a propensity-score-matched analysis of a hen-egg issue. *Diagnostics (Basel)* 2023;**13**:952
47. García-Granero E, Navarro F, Cerdán Santacruz C, Frasson M, García-Granero A, Marinello F et al. Individual surgeon is an independent risk factor for leak after double-stapled colorectal anastomosis: an institutional analysis of 800 patients. *Surgery* 2017;**162**:1006–1016
48. Yeh CY, Changchien CR, Wang JY, Chen J-S, Chen HH, Chiang J-M et al. Pelvic drainage and other risk factors for leakage after elective anterior resection in rectal cancer patients: a prospective study of 978 patients. *Ann Surg* 2005; **241**:9–13