



*Monotonicity methods for asymptotics of solutions to
Schrödinger equations near isolated singularities of the
electromagnetic potential*

Veronica Felli

Dipartimento di Matematica ed Applicazioni

University of Milano–Bicocca

`veronica.felli@unimib.it`

joint work with **Alberto Ferrero** and **Susanna Terracini**

Schrödinger operators with singular potentials

In quantum mechanics, the hamiltonian of a non-relativistic charged particle in an **electromagnetic** field has the form

$$(-i\nabla + \mathcal{A})^2 + V$$

$$\mathcal{A}: \mathbb{R}^N \rightarrow \mathbb{R}^N$$

magnetic potential associated to the magnetic field $B = \text{curl } \mathcal{A}$.

$$V: \mathbb{R}^N \rightarrow \mathbb{R}$$

electric potential.

Schrödinger operators with singular potentials

In quantum mechanics, the hamiltonian of a non-relativistic charged particle in an **electromagnetic** field has the form

$$(-i\nabla + \mathcal{A})^2 + V$$

$\mathcal{A}: \mathbb{R}^N \rightarrow \mathbb{R}^N$ **magnetic potential** associated to the magnetic field $B = \text{curl } \mathcal{A}$.

$V: \mathbb{R}^N \rightarrow \mathbb{R}$ **electric potential**.

For $N \geq 2$, we consider **singular homogeneous electromagnetic potentials** which make the operator *invariant* by scaling

$$\mathcal{A}(x) = \frac{\mathbf{A}\left(\frac{x}{|x|}\right)}{|x|}$$

$$\mathbf{A} \in C^1(\mathbb{S}^{N-1}, \mathbb{R}^N)$$

and

$$V(x) = -\frac{a\left(\frac{x}{|x|}\right)}{|x|^2}$$

$$a \in L^\infty(\mathbb{S}^{N-1}, \mathbb{R})$$

A prototype in dimension 2

Aharonov-Bohm magnetic potentials are associated to thin solenoids: if the radius of the solenoid tends to zero while the flux through it remains constant, then the particle is subject to a δ -type magnetic field, which is called **Aharonov-Bohm** field. An associated vector potential in \mathbb{R}^2 is

$$\mathcal{A}(x_1, x_2) = \alpha \left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2} \right), \quad (x_1, x_2) \in \mathbb{R}^2,$$

with α = circulation of \mathcal{A} around the solenoid.

A prototype in dimension 2

Aharonov-Bohm magnetic potentials are associated to thin solenoids: if the radius of the solenoid tends to zero while the flux through it remains constant, then the particle is subject to a δ -type magnetic field, which is called **Aharonov-Bohm** field. An associated vector potential in \mathbb{R}^2 is

$$\mathcal{A}(x_1, x_2) = \alpha \left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2} \right), \quad (x_1, x_2) \in \mathbb{R}^2,$$

with α = circulation of \mathcal{A} around the solenoid.

Aharonov-Bohm vector potentials are

- singular at 0,

A prototype in dimension 2

Aharonov-Bohm magnetic potentials are associated to thin solenoids: if the radius of the solenoid tends to zero while the flux through it remains constant, then the particle is subject to a δ -type magnetic field, which is called **Aharonov-Bohm** field. An associated vector potential in \mathbb{R}^2 is

$$\mathcal{A}(x_1, x_2) = \alpha \left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2} \right), \quad (x_1, x_2) \in \mathbb{R}^2,$$

with α = circulation of \mathcal{A} around the solenoid.

Aharonov-Bohm vector potentials are

- singular at 0,
- homogeneous of degree -1

A prototype in dimension 2

Aharonov-Bohm magnetic potentials are associated to thin solenoids: if the radius of the solenoid tends to zero while the flux through it remains constant, then the particle is subject to a δ -type magnetic field, which is called **Aharonov-Bohm field**. An associated vector potential in \mathbb{R}^2 is

$$\mathbf{A}(x_1, x_2) = \alpha \left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2} \right), \quad (x_1, x_2) \in \mathbb{R}^2,$$

with α = circulation of \mathbf{A} around the solenoid.

Aharonov-Bohm vector potentials are

- singular at 0,
- homogeneous of degree -1

- transversal, i.e.

$$\mathbf{A}(\theta) \cdot \theta = 0 \quad \text{for all } \theta \in \mathbb{S}^{N-1} \quad \text{(TC)}$$

Singular homogeneous electric potentials

which scale as the laplacian arise in nonrelativistic molecular physics, see [J. M. Lévy-Leblond, *Phys. Rev.* (1967)]. The potential describing the interaction between an electric charge and the **dipole moment** $\mathbf{D} \in \mathbb{R}^N$ of a molecule has the form

$$V(x) = -\frac{\lambda (x \cdot \mathbf{d})}{|x|^3} \quad \text{in } \mathbb{R}^N,$$

where $\lambda \propto$ **magnitude** of the dipole moment \mathbf{D}

$\mathbf{d} = \mathbf{D}/|\mathbf{D}| =$ **orientation** of \mathbf{D} .

Singular homogeneous electric potentials

which scale as the laplacian arise in nonrelativistic molecular physics, see [J. M. Lévy-Leblond, Phys. Rev. (1967)]. The potential describing the interaction between an electric charge and the **dipole moment** $\mathbf{D} \in \mathbb{R}^N$ of a molecule has the form

$$V(x) = -\frac{\lambda (x \cdot \mathbf{d})}{|x|^3} \quad \text{in } \mathbb{R}^N,$$

where $\lambda \propto$ **magnitude** of the dipole moment \mathbf{D}

$\mathbf{d} = \mathbf{D}/|\mathbf{D}| =$ orientation of \mathbf{D} .

Schrödinger operators with dipole-type potentials $\frac{a(x/|x|)}{|x|^2}$ are studied in

[Terracini, Adv. Diff. Equations (1996)]

[F.-Marchini-Terracini, Discr. Contin. Dyn. Syst. (2008)]

[F.-Marchini-Terracini, Indiana Univ. Math. J. (2009)]

Problem:

describe the *asymptotic behavior at the singularity* of solutions to equations associated to Schrödinger operators with singular electromagnetic potentials of type

$$\mathcal{L}_{\mathbf{A},a} := \left(-i \nabla + \frac{\mathbf{A}\left(\frac{x}{|x|}\right)}{|x|} \right)^2 - \frac{a\left(\frac{x}{|x|}\right)}{|x|^2}$$

in a domain $\Omega \subset \mathbb{R}^N$ containing either the origin or a neighborhood of ∞ .

Problem:

describe the *asymptotic behavior at the singularity* of solutions to equations associated to Schrödinger operators with singular electromagnetic potentials of type

$$\mathcal{L}_{\mathbf{A},a} := \left(-i \nabla + \frac{\mathbf{A}\left(\frac{x}{|x|}\right)}{|x|} \right)^2 - \frac{a\left(\frac{x}{|x|}\right)}{|x|^2}$$

in a domain $\Omega \subset \mathbb{R}^N$ containing either the origin or a neighborhood of ∞ .

- **Linear perturbation of $\mathcal{L}_{\mathbf{A},a}$:**

$$\mathcal{L}_{\mathbf{A},a} u = h(x) u$$

with $h \in L_{\text{loc}}^\infty(\Omega \setminus \{0\})$ negligible with respect to $\frac{1}{|x|^2}$ near the singularity

Problem:

describe the *asymptotic behavior at the singularity* of solutions to equations associated to Schrödinger operators with singular electromagnetic potentials of type

$$\mathcal{L}_{\mathbf{A},a} := \left(-i \nabla + \frac{\mathbf{A}\left(\frac{x}{|x|}\right)}{|x|} \right)^2 - \frac{a\left(\frac{x}{|x|}\right)}{|x|^2}$$

in a domain $\Omega \subset \mathbb{R}^N$ containing either the origin or a neighborhood of ∞ .

- **Linear perturbation of $\mathcal{L}_{\mathbf{A},a}$:**

$$\mathcal{L}_{\mathbf{A},a}u = h(x)u$$

with $h \in L_{\text{loc}}^\infty(\Omega \setminus \{0\})$ negligible with respect to $\frac{1}{|x|^2}$ near the singularity

- **Semilinear equations of type**

$$\mathcal{L}_{\mathbf{A},a}u = f(x, u(x))$$

with f having at most critical growth.

References

Regularity properties of solutions to Schrödinger equations with less singular magnetic and electric potentials:

[Kurata, Math. Z., (1997)]: $N \geq 3$, *local boundedness* (and *continuity*) if the electric potential and the square of the magnetic one belong to the *Kato class*.

References

Regularity properties of solutions to Schrödinger equations with less singular magnetic and electric potentials:

[Kurata, Math. Z., (1997)]: $N \geq 3$, *local boundedness* (and *continuity*) if the electric potential and the square of the magnetic one belong to the *Kato class*.

[Kurata, Proc. AMS, 125 (1997)]: $N \geq 3$, *unique continuation* (*Kato class*).

References

Regularity properties of solutions to Schrödinger equations with less singular magnetic and electric potentials:

[Kurata, Math. Z., (1997)]: $N \geq 3$, *local boundedness* (and *continuity*) if the electric potential and the square of the magnetic one belong to the *Kato class*.

[Kurata, Proc. AMS, 125 (1997)]: $N \geq 3$, *unique continuation* (*Kato class*).

[Chabrowski-Szulkin, Topol. Methods Nonlinear Anal., (2005)]: *boundedness and decay at ∞* of solutions for $N \geq 3$, L^2_{loc} magnetic potentials, and *electric potentials* with $L^{N/2}$ negative part.

References

Behavior of solutions to Schrödinger equations with singular inverse square electric potentials for $\mathbf{A} = 0$ (i.e. no magnetic vector potential):

[F.-Schneider, Adv. Nonl. Studies (2003)] : *Hölder continuity results for degenerate elliptic equations with singular weights; include asymptotics of solutions near the pole for potentials $\frac{\lambda}{|x|^2}$.*

References

Behavior of solutions to Schrödinger equations with singular inverse square electric potentials for $\mathbf{A} = 0$ (i.e. no magnetic vector potential):

[F.-Schneider, Adv. Nonl. Studies (2003)] : Hölder continuity results for *degenerate elliptic equations with singular weights*; include asymptotics of solutions near the pole for potentials $\frac{\lambda}{|x|^2}$.

[F.-Marchini-Terracini, Discrete Contin. Dynam. Systems (2008)]: exact asymptotics of solutions near the pole for anisotropic inverse-square singular potentials $\frac{\alpha(x/|x|)}{|x|^2}$, through *separation of variables and comparison methods*.

References

Behavior of solutions to Schrödinger equations with singular inverse square electric potentials for $\mathbf{A} = 0$ (i.e. no magnetic vector potential):

[F.-Schneider, Adv. Nonl. Studies (2003)] : Hölder continuity results for *degenerate elliptic equations with singular weights*; include asymptotics of solutions near the pole for potentials $\frac{\lambda}{|x|^2}$.

[F.-Marchini-Terracini, Discrete Contin. Dynam. Systems (2008)]: exact asymptotics of solutions near the pole for anisotropic inverse-square singular potentials $\frac{\alpha(x/|x|)}{|x|^2}$, through *separation of variables and comparison methods*.

[Pinchover, Ann. IHP Anal. Nonlinaire (1994)]: existence of the limit at the singularity of any quotient of two positive solutions in some linear and semilinear cases.

Remark

Comparison and maximum principles play a crucial role both in [F.-Marchini-Terracini (2008)] and [Pinchover (1994)].

In the presence of a **singular magnetic potential**, comparison methods are no more available, preventing us from a direct extension of the aforementioned results.

We overcome this difficulty by a **Almgren type monotonicity formula** and **blow-up methods**.

The angular operator

We aim to describe the **rate** and the **shape** of the singularity of solutions, by relating them to the **eigenvalues** and the **eigenfunctions** of a Schrödinger operator on the sphere \mathbb{S}^{N-1} corresponding to the angular part of $\mathcal{L}_{\mathbf{A},a}$:

$$L_{\mathbf{A},a} := \left(-i \nabla_{\mathbb{S}^{N-1}} + \mathbf{A} \right)^2 - a$$

The angular operator

We aim to describe the **rate** and the **shape** of the singularity of solutions, by relating them to the **eigenvalues** and the **eigenfunctions** of a Schrödinger operator on the sphere \mathbb{S}^{N-1} corresponding to the angular part of $\mathcal{L}_{\mathbf{A},a}$:

$$L_{\mathbf{A},a} := \left(-i \nabla_{\mathbb{S}^{N-1}} + \mathbf{A} \right)^2 - a$$

For $a \in L^\infty(\mathbb{S}^{N-1}, \mathbb{R})$ and $\mathbf{A} \in C^1(\mathbb{S}^{N-1}, \mathbb{R}^N)$, the operator $L_{\mathbf{A},a}$ on \mathbb{S}^{N-1} admits a diverging sequence of real eigenvalues

$$\mu_1(\mathbf{A}, a) \leq \mu_2(\mathbf{A}, a) \leq \cdots \leq \mu_k(\mathbf{A}, a) \leq \cdots$$

The angular operator

We aim to describe the **rate** and the **shape** of the singularity of solutions, by relating them to the **eigenvalues** and the **eigenfunctions** of a Schrödinger operator on the sphere \mathbb{S}^{N-1} corresponding to the angular part of $\mathcal{L}_{\mathbf{A},a}$:

$$L_{\mathbf{A},a} := \left(-i \nabla_{\mathbb{S}^{N-1}} + \mathbf{A} \right)^2 - a$$

For $a \in L^\infty(\mathbb{S}^{N-1}, \mathbb{R})$ and $\mathbf{A} \in C^1(\mathbb{S}^{N-1}, \mathbb{R}^N)$, the operator $L_{\mathbf{A},a}$ on \mathbb{S}^{N-1} admits a diverging sequence of real eigenvalues

$$\mu_1(\mathbf{A}, a) \leq \mu_2(\mathbf{A}, a) \leq \cdots \leq \mu_k(\mathbf{A}, a) \leq \cdots$$

Positivity of the quadratic form associated to $\mathcal{L}_{\mathbf{A},a}$ is ensured by

$$\mu_1(\mathbf{A}, a) > - \left(\frac{N-2}{2} \right)^2 \quad \text{(PD)}$$

The functional setting

$\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C}) :=$ completion of $C_c^\infty(\mathbb{R}^N \setminus \{0\}, \mathbb{C})$ with respect to

$$\|u\|_{\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C})} := \left(\int_{\mathbb{R}^N} \left(|\nabla u(x)|^2 + \frac{|u(x)|^2}{|x|^2} \right) dx \right)^{1/2}.$$

The functional setting

$\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C}) :=$ completion of $C_c^\infty(\mathbb{R}^N \setminus \{0\}, \mathbb{C})$ with respect to

$$\|u\|_{\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C})} := \left(\int_{\mathbb{R}^N} \left(|\nabla u(x)|^2 + \frac{|u(x)|^2}{|x|^2} \right) dx \right)^{1/2}.$$

$$\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C}) = \left\{ u \in L_{\text{loc}}^1(\mathbb{R}^N \setminus \{0\}, \mathbb{C}) : \frac{u}{|x|}, \nabla u \in L^2(\mathbb{R}^N, \mathbb{C}^N) \right\}.$$

The functional setting

$\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C}) :=$ completion of $C_c^\infty(\mathbb{R}^N \setminus \{0\}, \mathbb{C})$ with respect to

$$\|u\|_{\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C})} := \left(\int_{\mathbb{R}^N} \left(|\nabla u(x)|^2 + \frac{|u(x)|^2}{|x|^2} \right) dx \right)^{1/2}.$$

$$\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C}) = \left\{ u \in L_{\text{loc}}^1(\mathbb{R}^N \setminus \{0\}, \mathbb{C}) : \frac{u}{|x|}, \nabla u \in L^2(\mathbb{R}^N, \mathbb{C}^N) \right\}.$$

Under assumptions **(TC)** and **(PD)**, $\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C}) = \mathcal{D}_{\mathbf{A},a}^{1,2}(\mathbb{R}^N)$, where $\mathcal{D}_{\mathbf{A},a}^{1,2}(\mathbb{R}^N)$ is the completion of $C_c^\infty(\mathbb{R}^N \setminus \{0\}, \mathbb{C})$ with respect to

$$\|u\|_{\mathcal{D}_{\mathbf{A},a}^{1,2}(\mathbb{R}^N)} := \left(\int_{\mathbb{R}^N} \left[\left| \left(\nabla + i \frac{\mathbf{A}(x/|x|)}{|x|} \right) u(x) \right|^2 - \frac{a(x/|x|)}{|x|^2} |u(x)|^2 \right] dx \right)^{1/2}$$

Moreover the norms $\|\cdot\|_{\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C})}$ and $\|\cdot\|_{\mathcal{D}_{\mathbf{A},a}^{1,2}(\mathbb{R}^N)}$ are equivalent.

The functional setting

$$N \geq 3$$

Hardy's inequality

$$\left(\frac{N-2}{2}\right)^2 \int_{\mathbb{R}^N} \frac{|u|^2}{|x|^2} dx \leq \int_{\mathbb{R}^N} |\nabla u|^2 dx$$

The functional setting

$$N \geq 3$$

Hardy's inequality

$$\left(\frac{N-2}{2}\right)^2 \int_{\mathbb{R}^N} \frac{|u|^2}{|x|^2} dx \leq \int_{\mathbb{R}^N} |\nabla u|^2 dx$$

↓

$$\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C}) = \mathcal{D}^{1,2}(\mathbb{R}^N) = \overline{C_c^\infty(\mathbb{R}^N, \mathbb{C})}^{\|\cdot\|_{\mathcal{D}^{1,2}(\mathbb{R}^N)}}$$

$$\|u\|_{\mathcal{D}^{1,2}(\mathbb{R}^N)} := \left(\int_{\mathbb{R}^N} |\nabla u(x)|^2 dx \right)^{1/2}$$

and the norms $\|\cdot\|_{\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C})}$ and $\|\cdot\|_{\mathcal{D}^{1,2}(\mathbb{R}^N)}$ are equivalent.

The functional setting If $N = 2$, **(TC)** holds (i.e. $\mathbf{A}(\theta) \cdot \theta = 0$), and

$$\Phi_{\mathbf{A}} := \frac{1}{2\pi} \int_0^{2\pi} \alpha(t) dt \notin \mathbb{Z} \quad \textbf{(ND)}$$

where $\alpha(t) := \mathbf{A}(\cos t, \sin t) \cdot (-\sin t, \cos t)$

The functional setting If $N = 2$, (TC) holds (i.e. $\mathbf{A}(\theta) \cdot \theta = 0$), and

$$\Phi_{\mathbf{A}} := \frac{1}{2\pi} \int_0^{2\pi} \alpha(t) dt \notin \mathbb{Z} \quad (\mathbf{ND})$$

where $\alpha(t) := \mathbf{A}(\cos t, \sin t) \cdot (-\sin t, \cos t)$

↓ [Laptev-Weidl (1999)]

$C_c^\infty(\mathbb{R}^N \setminus \{0\}, \mathbb{C})$ functions satisfy the following **Hardy inequality**:

$$\underbrace{\left(\min_{k \in \mathbb{Z}} |k - \Phi_{\mathbf{A}}| \right)^2}_{\text{optimal}} \int_{\mathbb{R}^2} \frac{|u(x)|^2}{|x|^2} dx \leq \int_{\mathbb{R}^2} \left| \nabla u(x) + i \frac{\mathbf{A}(x/|x|)}{|x|} u(x) \right|^2 dx$$

The functional setting If $N = 2$, (TC) holds (i.e. $\mathbf{A}(\theta) \cdot \theta = 0$), and

$$\Phi_{\mathbf{A}} := \frac{1}{2\pi} \int_0^{2\pi} \alpha(t) dt \notin \mathbb{Z} \quad (\text{ND})$$

where $\alpha(t) := \mathbf{A}(\cos t, \sin t) \cdot (-\sin t, \cos t)$

↓ [Laptev-Weidl (1999)]

$C_c^\infty(\mathbb{R}^N \setminus \{0\}, \mathbb{C})$ functions satisfy the following **Hardy inequality**:

$$\underbrace{\left(\min_{k \in \mathbb{Z}} |k - \Phi_{\mathbf{A}}| \right)^2}_{\text{optimal}} \int_{\mathbb{R}^2} \frac{|u(x)|^2}{|x|^2} dx \leq \int_{\mathbb{R}^2} \left| \nabla u(x) + i \frac{\mathbf{A}(x/|x|)}{|x|} u(x) \right|^2 dx$$

$$\mathcal{D}_*^{1,2}(\mathbb{R}^N, \mathbb{C}) = \mathcal{D}_{\mathbf{A}}^{1,2}(\mathbb{R}^2) = \text{completion w.r.t. } \left\| \nabla u + i \frac{\mathbf{A}(x/|x|)}{|x|} u \right\|_{L^2(\mathbb{R}^2, \mathbb{C})}$$

The functional setting

In an open bounded domain $\Omega \subset \mathbb{R}^N$, $0 \in \Omega$, let

$H_*^1(\Omega, \mathbb{C}) =$ completion of $\left\{ \begin{array}{l} u \in H^1(\Omega, \mathbb{C}) \cap C^\infty(\Omega, \mathbb{C}) : \\ u \text{ vanishes in a neighborhood of } 0 \end{array} \right\}$ w.r.t.

$$\|u\|_{H_*^1(\Omega, \mathbb{C})} = \left(\|\nabla u\|_{L^2(\Omega, \mathbb{C}^N)}^2 + \|u\|_{L^2(\Omega, \mathbb{C})}^2 + \left\| \frac{u}{|x|} \right\|_{L^2(\Omega, \mathbb{C})}^2 \right)^{1/2}.$$

The functional setting

In an open bounded domain $\Omega \subset \mathbb{R}^N$, $0 \in \Omega$, let

$H_*^1(\Omega, \mathbb{C}) =$ completion of $\left\{ \begin{array}{l} u \in H^1(\Omega, \mathbb{C}) \cap C^\infty(\Omega, \mathbb{C}) : \\ u \text{ vanishes in a neighborhood of } 0 \end{array} \right\}$ w.r.t.

$$\|u\|_{H_*^1(\Omega, \mathbb{C})} = \left(\|\nabla u\|_{L^2(\Omega, \mathbb{C}^N)}^2 + \|u\|_{L^2(\Omega, \mathbb{C})}^2 + \left\| \frac{u}{|x|} \right\|_{L^2(\Omega, \mathbb{C})}^2 \right)^{1/2}.$$

$$H_*^1(\Omega, \mathbb{C}) = \left\{ u \in H^1(\Omega, \mathbb{C}) : \frac{u}{|x|} \in L^2(\Omega, \mathbb{C}) \right\}$$

The functional setting

In an open bounded domain $\Omega \subset \mathbb{R}^N$, $0 \in \Omega$, let

$$H_*^1(\Omega, \mathbb{C}) = \text{completion of } \left\{ \begin{array}{l} u \in H^1(\Omega, \mathbb{C}) \cap C^\infty(\Omega, \mathbb{C}) : \\ u \text{ vanishes in a neighborhood of } 0 \end{array} \right\} \text{ w.r.t.}$$

$$\|u\|_{H_*^1(\Omega, \mathbb{C})} = \left(\|\nabla u\|_{L^2(\Omega, \mathbb{C}^N)}^2 + \|u\|_{L^2(\Omega, \mathbb{C})}^2 + \left\| \frac{u}{|x|} \right\|_{L^2(\Omega, \mathbb{C})}^2 \right)^{1/2}.$$

$$H_*^1(\Omega, \mathbb{C}) = \left\{ u \in H^1(\Omega, \mathbb{C}) : \frac{u}{|x|} \in L^2(\Omega, \mathbb{C}) \right\}$$

- If $N \geq 3$, $H_*^1(\Omega) = H^1(\Omega, \mathbb{C})$ and their norms are equivalent.
- If $N = 2$, $H_*^1(\Omega)$ is strictly smaller than $H^1(\Omega, \mathbb{C})$.

The Almgren frequency function

Studying regularity of area-minimizing surfaces of codimension ≥ 1 , in 1979 Almgren introduced the *frequency function*

$$N(r) = \frac{r \int_{B_r} |\nabla u|^2 dx}{\int_{\partial B_r} u^2}$$

and observed that, if u is harmonic, then $N \nearrow$ in r .

The Almgren frequency function

Studying regularity of area-minimizing surfaces of codimension ≥ 1 , in 1979 Almgren introduced the *frequency function*

$$N(r) = \frac{r \int_{B_r} |\nabla u|^2 dx}{\int_{\partial B_r} u^2}$$

and observed that, if u is harmonic, then $N \nearrow$ in r .

“*frequency*”: if u is a harmonic function in \mathbb{R}^2 homogeneous of degree k ($u_k(r, \theta) = a_k r^k \sin(k\theta)$), then $N(r) = k$.

The Almgren frequency function

Studying regularity of area-minimizing surfaces of codimension ≥ 1 , in 1979 Almgren introduced the *frequency function*

$$N(r) = \frac{r \int_{B_r} |\nabla u|^2 dx}{\int_{\partial B_r} u^2}$$

and observed that, if u is harmonic, then $N \nearrow$ in r .

The Almgren monotonicity formula was used in

- [Garofalo-Lin, Indiana Univ. Math. J. (1986)]: generalization to variable coefficient elliptic operators in divergence form (unique continuation)
- [Athanasopoulos-Caffarelli-Salsa, Amer. J. Math. (2008)]: regularity of the free boundary in obstacle problems.
- [Caffarelli-Lin, J. AMS (2008)] regularity of free boundary of the limit components of singularly perturbed elliptic systems.

The Almgren type frequency function

In an open bounded $\Omega \ni 0$, let u be a $H_*^1(\Omega)$ -weak **solution** to $\mathcal{L}_{\mathbf{A},a}u = h(x)u$, with h satisfying

$$h \in L_{\text{loc}}^\infty(\Omega \setminus \{0\}, \mathbb{C}), \quad |h(x)| = O(|x|^{-2+\varepsilon}) \text{ as } |x| \rightarrow 0 \quad (\mathbf{H}_0)$$

The Almgren type frequency function

In an open bounded $\Omega \ni 0$, let u be a $H_*^1(\Omega)$ -weak solution to $\mathcal{L}_{\mathbf{A},a}u = h(x)u$, with h satisfying

$$h \in L_{\text{loc}}^\infty(\Omega \setminus \{0\}, \mathbb{C}), \quad |h(x)| = O(|x|^{-2+\varepsilon}) \text{ as } |x| \rightarrow 0 \quad (\mathbf{H}_0)$$

For small $r > 0$ define

$$D(r) = \frac{1}{r^{N-2}} \int_{B_r} \left[\left| \nabla u + i \frac{\mathbf{A}(\frac{x}{|x|})}{|x|} u \right|^2 - \frac{a(\frac{x}{|x|})}{|x|^2} |u|^2 - (\Re h) |u|^2 \right] dx,$$

$$H(r) = \frac{1}{r^{N-1}} \int_{\partial B_r} |u|^2 dS.$$

The Almgren type frequency function

In an open bounded $\Omega \ni 0$, let u be a $H_*^1(\Omega)$ -weak **solution** to $\mathcal{L}_{\mathbf{A},a}u = h(x)u$, with h satisfying

$$h \in L_{\text{loc}}^\infty(\Omega \setminus \{0\}, \mathbb{C}), \quad |h(x)| = O(|x|^{-2+\varepsilon}) \text{ as } |x| \rightarrow 0 \quad (\mathbf{H}_0)$$

For small $r > 0$ define

$$D(r) = \frac{1}{r^{N-2}} \int_{B_r} \left[\left| \nabla u + i \frac{\mathbf{A}(\frac{x}{|x|})}{|x|} u \right|^2 - \frac{a(\frac{x}{|x|})}{|x|^2} |u|^2 - (\Re h) |u|^2 \right] dx,$$

$$H(r) = \frac{1}{r^{N-1}} \int_{\partial B_r} |u|^2 dS.$$

If **(PD)** holds and $u \not\equiv 0$,
 $\Rightarrow H(r) > 0$ for small $r > 0$

\rightsquigarrow

Almgren type frequency function

$$\mathcal{N}(r) = \mathcal{N}_{u,h}(r) = \frac{D(r)}{H(r)}$$

is well defined in a suitably small interval $(0, \bar{r})$.

The Almgren type frequency function

$\mathcal{N} \in W_{\text{loc}}^{1,1}(0, \bar{r})$ and, in a distributional sense and for a.e. $r \in (0, \bar{r})$,

$$\mathcal{N}'(r) = \frac{2r \left[\left(\int_{\partial B_r} \left| \frac{\partial u}{\partial \nu} \right|^2 dS \right) \left(\int_{\partial B_r} |u|^2 dS \right) - \left(\int_{\partial B_r} \Re \left(u \frac{\partial \bar{u}}{\partial \nu} \right) dS \right)^2 \right]}{\left(\int_{\partial B_r} |u|^2 dS \right)^2} + \frac{\alpha_h(r)}{\int_{\partial B_r} |u|^2 dS}$$

where $\alpha_h(r) = 2 \left[\int_{B_r} \Re(h(x) \overline{u(x)} (x \cdot \nabla u(x))) dx \right.$
 $\left. + \frac{N-2}{2} \int_{B_r} (\Re h(x)) |u(x)|^2 dx - \frac{r}{2} \int_{\partial B_r} (\Re h(x)) |u(x)|^2 dS \right]$

The Almgren type frequency function

$\mathcal{N} \in W_{\text{loc}}^{1,1}(0, \bar{r})$ and, in a distributional sense and for a.e. $r \in (0, \bar{r})$,

$$\mathcal{N}'(r) = \frac{2r \left[\left(\int_{\partial B_r} \left| \frac{\partial u}{\partial \nu} \right|^2 dS \right) \left(\int_{\partial B_r} |u|^2 dS \right) - \left(\int_{\partial B_r} \Re \left(u \frac{\partial \bar{u}}{\partial \nu} \right) dS \right)^2 \right]}{\left(\int_{\partial B_r} |u|^2 dS \right)^2} + \frac{\alpha_h(r)}{\int_{\partial B_r} |u|^2 dS}$$

Schwarz's inequality

\leq
0

where $\alpha_h(r) = 2 \left[\int_{B_r} \Re(h(x) \overline{u(x)} (x \cdot \nabla u(x))) dx \right.$

$$\left. + \frac{N-2}{2} \int_{B_r} (\Re h(x)) |u(x)|^2 dx - \frac{r}{2} \int_{\partial B_r} (\Re h(x)) |u(x)|^2 dS \right]$$

The Almgren type frequency function

$\mathcal{N} \in W_{\text{loc}}^{1,1}(0, \bar{r})$ and, in a distributional sense and for a.e. $r \in (0, \bar{r})$,

$$\mathcal{N}'(r) = \frac{2r \left[\left(\int_{\partial B_r} \left| \frac{\partial u}{\partial \nu} \right|^2 dS \right) \left(\int_{\partial B_r} |u|^2 dS \right) - \left(\int_{\partial B_r} \Re \left(u \frac{\partial \bar{u}}{\partial \nu} \right) dS \right)^2 \right]}{\left(\int_{\partial B_r} |u|^2 dS \right)^2} + \frac{\alpha_h(r)}{\int_{\partial B_r} |u|^2 dS}$$

Schwarz's inequality

\leq
0

where $\alpha_h(r) = 2 \left[\int_{B_r} \Re(h(x) \overline{u(x)} (x \cdot \nabla u(x))) dx \right.$

$$\left. + \frac{N-2}{2} \int_{B_r} (\Re h(x)) |u(x)|^2 dx - \frac{r}{2} \int_{\partial B_r} (\Re h(x)) |u(x)|^2 dS \right]$$

$$(H_0) \implies \left| \frac{\alpha_h(r)}{\int_{\partial B_r} |u|^2 dS} \right| \leq \text{const } r^{-1+\varepsilon}$$

The Almgren type frequency function

$\mathcal{N} \in W_{\text{loc}}^{1,1}(0, \bar{r})$ and, in a distributional sense and for a.e. $r \in (0, \bar{r})$,

$$\mathcal{N}'(r) = \underbrace{\frac{2r \left[\left(\int_{\partial B_r} \left| \frac{\partial u}{\partial \nu} \right|^2 dS \right) \left(\int_{\partial B_r} |u|^2 dS \right) - \left(\int_{\partial B_r} \Re \left(u \frac{\partial \bar{u}}{\partial \nu} \right) dS \right)^2 \right]}{\left(\int_{\partial B_r} |u|^2 dS \right)^2}}_{\text{Schwarz's inequality}} + \underbrace{\frac{\alpha_h(r)}{\int_{\partial B_r} |u|^2 dS}}_{\text{integrable}}$$

where $\alpha_h(r) = 2 \left[\int_{B_r} \Re(h(x) \overline{u(x)} (x \cdot \nabla u(x))) dx \right. \\ \left. + \frac{N-2}{2} \int_{B_r} (\Re h(x)) |u(x)|^2 dx - \frac{r}{2} \int_{\partial B_r} (\Re h(x)) |u(x)|^2 dS \right]$

\implies the limit $\gamma := \lim_{r \rightarrow 0^+} \mathcal{N}(r)$ exists and is finite.

Blow-up: set $w^\lambda(x) = \frac{u(\lambda x)}{\sqrt{H(\lambda)}}$, so that $\int_{\partial B_1} |w^\lambda|^2 dS = 1$.

$\{w^\lambda\}_{\lambda \in (0, \bar{\lambda})}$ is bounded in $H_*^1(B_1) \implies$ for any $\lambda_n \rightarrow 0^+$, $w^{\lambda_{n_k}} \rightharpoonup w$ in $H_*^1(B_1)$ along a subsequence $\lambda_{n_k} \rightarrow 0^+$, and $\int_{\partial B_1} |w|^2 dS = 1$.

Blow-up: set $w^\lambda(x) = \frac{u(\lambda x)}{\sqrt{H(\lambda)}}$, so that $\int_{\partial B_1} |w^\lambda|^2 dS = 1$.

$\{w^\lambda\}_{\lambda \in (0, \bar{\lambda})}$ is bounded in $H_*^1(B_1) \implies$ for any $\lambda_n \rightarrow 0^+$, $w^{\lambda_{n_k}} \rightharpoonup w$ in $H_*^1(B_1)$ along a subsequence $\lambda_{n_k} \rightarrow 0^+$, and $\int_{\partial B_1} |w|^2 dS = 1$.

$$(E_k) \quad \mathcal{L}_{\mathbf{A}, a} w^{\lambda_{n_k}}(x) = \lambda_{n_k}^2 h(\lambda_{n_k} x) w^{\lambda_{n_k}}(x) \quad \xrightarrow[\text{limit}]{\text{weak}} \quad (E) \quad \mathcal{L}_{\mathbf{A}, a} w(x) = 0 \text{ in } B_1$$

Bootstrap and classical regularity theory \implies

$$w^{\lambda_{n_k}} \rightarrow w \quad \text{in } C_{\text{loc}}^{1, \tau}(B_1 \setminus \{0\}), \quad \tau \in (0, 1), \quad H^1(B_r, \mathbb{C}), \quad H_*^1(B_r), \quad r \in (0, 1).$$

Blow-up: set $w^\lambda(x) = \frac{u(\lambda x)}{\sqrt{H(\lambda)}}$, so that $\int_{\partial B_1} |w^\lambda|^2 dS = 1$.

$\{w^\lambda\}_{\lambda \in (0, \bar{\lambda})}$ is bounded in $H_*^1(B_1) \implies$ for any $\lambda_n \rightarrow 0^+$, $w^{\lambda_{n_k}} \rightharpoonup w$ in $H_*^1(B_1)$ along a subsequence $\lambda_{n_k} \rightarrow 0^+$, and $\int_{\partial B_1} |w|^2 dS = 1$.

$$(E_k) \quad \mathcal{L}_{\mathbf{A}, a} w^{\lambda_{n_k}}(x) = \lambda_{n_k}^2 h(\lambda_{n_k} x) w^{\lambda_{n_k}}(x) \quad \xrightarrow[\text{limit}]{\text{weak}} \quad (E) \quad \mathcal{L}_{\mathbf{A}, a} w(x) = 0 \text{ in } B_1$$

Bootstrap and classical regularity theory \implies

$$w^{\lambda_{n_k}} \rightarrow w \quad \text{in } C_{\text{loc}}^{1, \tau}(B_1 \setminus \{0\}), \quad \tau \in (0, 1), \quad H^1(B_r, \mathbb{C}), \quad H_*^1(B_r), \quad r \in (0, 1).$$

If $\mathcal{N}_k(r)$ the Almgren frequency function associated to (E_k) and $\mathcal{N}_w(r)$ is the Almgren frequency function associated to (E) , then

$$\lim_{k \rightarrow \infty} \mathcal{N}_k(r) = \mathcal{N}_w(r) \quad \text{for all } r \in (0, 1).$$

Blow-up:

By scaling $\mathcal{N}_k(r) = \mathcal{N}(\lambda_{n_k} r)$

\Downarrow

$$\mathcal{N}_w(r) = \lim_{k \rightarrow \infty} \mathcal{N}(\lambda_{n_k} r) = \gamma \quad \forall r \in (0, 1)$$

Blow-up:

By scaling $\mathcal{N}_k(r) = \mathcal{N}(\lambda_{n_k} r)$

\Downarrow

$$\mathcal{N}_w(r) = \lim_{k \rightarrow \infty} \mathcal{N}(\lambda_{n_k} r) = \gamma \quad \forall r \in (0, 1)$$

Then \mathcal{N}_w is constant in $(0, 1)$ and hence $\mathcal{N}'_w(r) = 0$ for any $r \in (0, 1)$

\Downarrow

$$\left(\int_{\partial B_r} \left| \frac{\partial w}{\partial \nu} \right|^2 dS \right) \cdot \left(\int_{\partial B_r} |w|^2 dS \right) - \left(\int_{\partial B_r} \Re \left(w \frac{\partial \bar{w}}{\partial \nu} \right) dS \right)^2 = 0$$

Therefore w and $\frac{\partial w}{\partial \nu}$ are parallel as vectors in $L^2(\partial B_r, \mathbb{C})$, i.e. \exists a real valued function $\eta = \eta(r)$ such that $\frac{\partial w}{\partial \nu}(r, \theta) = \eta(r)w(r, \theta)$ for $r \in (0, 1)$.

Blow-up:

By scaling $\mathcal{N}_k(r) = \mathcal{N}(\lambda_{n_k} r)$

\Downarrow

$$\mathcal{N}_w(r) = \lim_{k \rightarrow \infty} \mathcal{N}(\lambda_{n_k} r) = \gamma \quad \forall r \in (0, 1)$$

Then \mathcal{N}_w is constant in $(0, 1)$ and hence $\mathcal{N}'_w(r) = 0$ for any $r \in (0, 1)$

\Downarrow

$$\left(\int_{\partial B_r} \left| \frac{\partial w}{\partial \nu} \right|^2 dS \right) \cdot \left(\int_{\partial B_r} |w|^2 dS \right) - \left(\int_{\partial B_r} \Re \left(w \frac{\partial \bar{w}}{\partial \nu} \right) dS \right)^2 = 0$$

Therefore w and $\frac{\partial w}{\partial \nu}$ are parallel as vectors in $L^2(\partial B_r, \mathbb{C})$, i.e. \exists a real valued function $\eta = \eta(r)$ such that $\frac{\partial w}{\partial \nu}(r, \theta) = \eta(r)w(r, \theta)$ for $r \in (0, 1)$.

After integration we obtain

$$w(r, \theta) = e^{\int_1^r \eta(s) ds} w(1, \theta) = \varphi(r) \psi(\theta), \quad r \in (0, 1), \quad \theta \in \mathbb{S}^{N-1}.$$

Blow-up:

$$w(r, \theta) = \varphi(r)\psi(\theta)$$

Rewriting equation (E) $\mathcal{L}_{\mathbf{A},a}w(x) = 0$ in polar coordinates we obtain

$$\left(-\varphi''(r) - \frac{N-1}{r}\varphi'(r)\right)\psi(\theta) + r^{-2}\varphi(r)L_{\mathbf{A},a}\psi(\theta) = 0.$$

Blow-up:

$$w(r, \theta) = \varphi(r)\psi(\theta)$$

Rewriting equation (E) $\mathcal{L}_{\mathbf{A},a}w(x) = 0$ in polar coordinates we obtain

$$\left(-\varphi''(r) - \frac{N-1}{r}\varphi'(r)\right)\psi(\theta) + r^{-2}\varphi(r)L_{\mathbf{A},a}\psi(\theta) = 0.$$

Then ψ is an eigenfunction of the operator $L_{\mathbf{A},a}$.

Let $\mu_{k_0}(\mathbf{A}, a)$ be the corresponding eigenvalue $\implies \varphi(r)$ solves

$$-\varphi''(r) - \frac{N-1}{r}\varphi'(r) + r^{-2}\mu_{k_0}(\mathbf{A}, a)\varphi(r) = 0.$$

Blow-up:

$$w(r, \theta) = \varphi(r)\psi(\theta)$$

Rewriting equation (E) $\mathcal{L}_{\mathbf{A},a}w(x) = 0$ in polar coordinates we obtain

$$\left(-\varphi''(r) - \frac{N-1}{r}\varphi'(r)\right)\psi(\theta) + r^{-2}\varphi(r)L_{\mathbf{A},a}\psi(\theta) = 0.$$

Then ψ is an eigenfunction of the operator $L_{\mathbf{A},a}$.

Let $\mu_{k_0}(\mathbf{A}, a)$ be the corresponding eigenvalue $\implies \varphi(r)$ solves

$$-\varphi''(r) - \frac{N-1}{r}\varphi'(r) + r^{-2}\mu_{k_0}(\mathbf{A}, a)\varphi(r) = 0.$$

Then $\varphi(r) = c_1r^{\sigma^+} + c_2r^{\sigma^-}$ with $\sigma^\pm = -\frac{N-2}{2} \pm \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_{k_0}(\mathbf{A}, a)}$.

$$|x|^{\sigma^-}\psi\left(\frac{x}{|x|}\right) \notin H_*^1(B_1) \rightsquigarrow c_2 = 0, \quad \varphi(1) = 1 \rightsquigarrow c_1 = 1$$

\Downarrow

$$w(r, \theta) = r^{\sigma^+}\psi(\theta)$$

From $\mathcal{N}_w(r) \equiv \gamma$, we deduce that $\gamma = \sigma^+$.

Asymptotics at the singularity

Step 1: any $\lambda_n \rightarrow 0^+$ admits a subsequence $\{\lambda_{n_k}\}_{k \in \mathbb{N}}$ s.t.

$$\frac{u(\lambda_{n_k} x)}{\sqrt{H(\lambda_{n_k})}} \rightarrow |x|^\gamma \psi\left(\frac{x}{|x|}\right)$$

weakly in $H^1(B_1)$

strongly in $H^1(B_r)$ for all $r \in (0, 1)$

in $C_{\text{loc}}^{1,\tau}(B_1 \setminus \{0\})$ for all $\tau \in (0, 1)$

$$\gamma = -\frac{N-2}{2} + \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_{k_0}(\mathbf{A}, a)},$$

ψ eigenfunction associated to μ_{k_0} .

Asymptotics at the singularity

Step 1: any $\lambda_n \rightarrow 0^+$ admits a subsequence $\{\lambda_{n_k}\}_{k \in \mathbb{N}}$ s.t.

$$\frac{u(\lambda_{n_k} x)}{\sqrt{H(\lambda_{n_k})}} \rightarrow |x|^\gamma \psi\left(\frac{x}{|x|}\right)$$

weakly in $H^1(B_1)$

strongly in $H^1(B_r)$ for all $r \in (0, 1)$

in $C_{\text{loc}}^{1,\tau}(B_1 \setminus \{0\})$ for all $\tau \in (0, 1)$

$$\gamma = -\frac{N-2}{2} + \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_{k_0}(\mathbf{A}, a)},$$

ψ eigenfunction associated to μ_{k_0} .

Step 2: $\lim_{r \rightarrow 0^+} \frac{H(r)}{r^{2\gamma}}$ is finite and > 0

(Step 1 + separation of variables)

Step 3: Let $m = \text{multiplicity of } \mu_{k_0}(\mathbf{A}, a)$ and $\{\psi_i\}_{i=j_0, \dots, j_0+m-1}$ be an $L^2(\mathbb{S}^{N-1}, \mathbb{C})$ -orthonormal basis for the associated eigenspace.

Step 1 $\implies \exists \{\beta_i\}_{i=j_0, \dots, j_0+m-1} \subset \mathbb{R}, \beta_i \neq 0$ for some i , s.t.

$$\lambda_{n_k}^{-\gamma} u(\lambda_{n_k} \theta) \rightarrow \sum_{i=j_0}^{j_0+m-1} \beta_i \psi_i(\theta) \quad \text{in } C^{1,\tau}(\mathbb{S}^{N-1})$$

$$\lambda_{n_k}^{1-\gamma} \nabla u(\lambda_{n_k} \theta) \rightarrow \sum_{i=j_0}^{j_0+m-1} \beta_i (\gamma \psi_i(\theta) \theta + \nabla_{\mathbb{S}^{N-1}} \psi_i(\theta)) \quad \text{in } C^{0,\tau}(\mathbb{S}^{N-1})$$

Step 3: Let $m = \text{multiplicity of } \mu_{k_0}(\mathbf{A}, a)$ and $\{\psi_i\}_{i=j_0, \dots, j_0+m-1}$ be an $L^2(\mathbb{S}^{N-1}, \mathbb{C})$ -orthonormal basis for the associated eigenspace.

Step 1 $\implies \exists \{\beta_i\}_{i=j_0, \dots, j_0+m-1} \subset \mathbb{R}, \beta_i \neq 0$ for some i , s.t.

$$\lambda_{n_k}^{-\gamma} u(\lambda_{n_k} \theta) \rightarrow \sum_{i=j_0}^{j_0+m-1} \beta_i \psi_i(\theta) \quad \text{in } C^{1,\tau}(\mathbb{S}^{N-1})$$

$$\lambda_{n_k}^{1-\gamma} \nabla u(\lambda_{n_k} \theta) \rightarrow \sum_{i=j_0}^{j_0+m-1} \beta_i (\gamma \psi_i(\theta) \theta + \nabla_{\mathbb{S}^{N-1}} \psi_i(\theta)) \quad \text{in } C^{0,\tau}(\mathbb{S}^{N-1})$$

Fix $R > 0$ such that $\overline{B_R} \subset \Omega$ and expand $u(\lambda \theta) = \sum_{k=1}^{\infty} \varphi_k(\lambda) \psi_k(\theta)$, then

$$\begin{aligned} \beta_i &= \lim_{k \rightarrow \infty} \lambda_{n_k}^{-\gamma} \varphi_i(\lambda_{n_k}) \\ &= \int_{\mathbb{S}^{N-1}} \left[R^{-\gamma} u(R\theta) + \int_0^R \frac{h(s\theta)u(s\theta)}{2\gamma + N - 2} \left(s^{1-\gamma} - \frac{s^{\gamma+N-1}}{R^{2\gamma+N-2}} \right) ds \right] \overline{\psi_i(\theta)} dS(\theta) \end{aligned}$$

depends neither on the sequence $\{\lambda_n\}_{n \in \mathbb{N}}$ nor on its subsequence $\{\lambda_{n_k}\}_{k \in \mathbb{N}}$

\implies **the convergences actually hold as $\lambda \rightarrow 0^+$.**

Asymptotics at the singularity

Theorem 1 [F.-Ferrero-Terracini, JEMS, to appear]

Let $\Omega \ni 0$ be a bounded open set in \mathbb{R}^N , $N \geq 2$, **(TC)**, **(PD)**, and **(H₀)** hold. If $u \not\equiv 0$ weakly solves $\mathcal{L}_{\mathbf{A},a}u = h(x)u$ in Ω , then $\exists k_0 \in \mathbb{N}$, $k_0 \geq 1$, s. t.

$$\gamma = \lim_{r \rightarrow 0^+} \mathcal{N}_{u,h}(r) = -\frac{N-2}{2} + \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_{k_0}(\mathbf{A}, a)}.$$

Furthermore, for any $\tau \in (0, 1)$, as $\lambda \rightarrow 0^+$,

$$\lambda^{-\gamma}u(\lambda\theta) \quad \rightarrow \quad \sum_{i=j_0}^{j_0+m-1} \beta_i \psi_i(\theta) \quad \text{in } C^{1,\tau}(\mathbb{S}^{N-1})$$

$$\lambda^{1-\gamma}\nabla u(\lambda\theta) \quad \rightarrow \quad \sum_{i=j_0}^{j_0+m-1} \beta_i (\gamma\psi_i(\theta)\theta + \nabla_{\mathbb{S}^{N-1}}\psi_i(\theta)) \quad \text{in } C^{0,\tau}(\mathbb{S}^{N-1}).$$

Asymptotics at the singularity

Corollary. Under the same assumptions as Theorem 1, let u be a weak $H_*^1(\Omega)$ -solution to $\mathcal{L}_{\mathbf{A},a}u = h(x)u$.

- (i) If $u(x) = O(|x|^k)$ as $|x| \rightarrow 0$ for all $k \in \mathbb{N}$, then $u \equiv 0$ in Ω .
- (ii) If $0 < \gamma < 1$ then $u \in C_{\text{loc}}^{0,\gamma}(\Omega)$.
- (iii) If $\gamma \geq 1$ then u is locally Lipschitz continuous in Ω .

Asymptotics at the singularity

Corollary. Under the same assumptions as Theorem 1, let u be a weak $H_*^1(\Omega)$ -solution to $\mathcal{L}_{\mathbf{A},a}u = h(x)u$.

- (i) If $u(x) = O(|x|^k)$ as $|x| \rightarrow 0$ for all $k \in \mathbb{N}$, then $u \equiv 0$ in Ω .
- (ii) If $0 < \gamma < 1$ then $u \in C_{\text{loc}}^{0,\gamma}(\Omega)$.
- (iii) If $\gamma \geq 1$ then u is locally Lipschitz continuous in Ω .

(i) is a ***strong unique continuation property***. It extends to singular homogeneous magnetic potentials the unique continuation property proved by **Kurata** for electromagnetic potentials in the Kato class.

Further results

- **Semilinear equations** $\mathcal{L}_{\mathbf{A},a}u = f(x, u(x))$ in $\Omega \ni 0$ with $f : \Omega \times \mathbb{C} \rightarrow \mathbb{C}$ a Carathéodory function satisfying

$$\left| \frac{f(x, z)}{z} \right| \leq \begin{cases} C(1 + |z|^{2^*-2}), & \text{if } N \geq 3, \\ C(1 + |z|^{p-2}) \text{ for some } p > 2, & \text{if } N = 2, \end{cases}$$

for a.e. $x \in \Omega$ and for all $z \in \mathbb{C} \setminus \{0\}$, where $2^* = \frac{2N}{N-2}$.

Further results

- **Semilinear equations** $\mathcal{L}_{\mathbf{A},a}u = f(x, u(x))$ in $\Omega \ni 0$ with $f : \Omega \times \mathbb{C} \rightarrow \mathbb{C}$ a Carathéodory function satisfying

$$\left| \frac{f(x, z)}{z} \right| \leq \begin{cases} C(1 + |z|^{2^*-2}), & \text{if } N \geq 3, \\ C(1 + |z|^{p-2}) \text{ for some } p > 2, & \text{if } N = 2, \end{cases}$$

for a.e. $x \in \Omega$ and for all $z \in \mathbb{C} \setminus \{0\}$, where $2^* = \frac{2N}{N-2}$.

Under the further assumption $\mu_1(0, a) > -\left(\frac{N-2}{2}\right)^2$ a **Brezis-Kato type iteration argument** provides an **upper bound** for solutions and then reduces the semilinear problem to a linear one with enough control on the perturbing potential at the singularity to apply Theorem 1 to recover the **exact asymptotic behavior of solutions at the singularity**.

Further results

- **Semilinear equations** $\mathcal{L}_{\mathbf{A},a}u = f(x, u(x))$ in $\Omega \ni 0$ with $f : \Omega \times \mathbb{C} \rightarrow \mathbb{C}$ a Carathéodory function satisfying

$$\left| \frac{f(x, z)}{z} \right| \leq \begin{cases} C(1 + |z|^{2^*-2}), & \text{if } N \geq 3, \\ C(1 + |z|^{p-2}) \text{ for some } p > 2, & \text{if } N = 2, \end{cases}$$

for a.e. $x \in \Omega$ and for all $z \in \mathbb{C} \setminus \{0\}$, where $2^* = \frac{2N}{N-2}$.

Under the further assumption $\mu_1(0, a) > -\left(\frac{N-2}{2}\right)^2$ a **Brezis-Kato type iteration argument** provides an **upper bound** for solutions and then reduces the semilinear problem to a linear one with enough control on the perturbing potential at the singularity to apply Theorem 1 to recover the **exact asymptotic behavior of solutions at the singularity**.

- invariance by the Kelvin transform \implies **asymptotics at ∞** for solutions in external domains

Example: Aharonov-Bohm magnetic potentials in $\dim N = 2$

$$\mathbf{A}(\cos t, \sin t) = \alpha(-\sin t, \cos t), \alpha \in \mathbb{R} \setminus \mathbb{Z}, \quad a(\cos t, \sin t) = a_0, a_0 \in \mathbb{R}$$

$$\left(-i \nabla + \alpha \left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2} \right) \right)^2 u - \frac{a_0}{|x|^2} u = h u,$$

with $x = (x_1, x_2) \in \Omega \subset \mathbb{R}^2$,
 Ω bounded, $0 \in \Omega$,
and h verifying (H_0) .

Eigenvalues: $\{(\alpha - j)^2 - a_0 : j \in \mathbb{Z}\} \Rightarrow \mu_1(\mathbf{A}, a) = (\text{dist}(\alpha, \mathbb{Z}))^2 - a_0$

If $\text{dist}(\alpha, \mathbb{Z}) \neq \frac{1}{2}$, then all eigenvalues are simple and the eigenspace associated to the first eigenvalue is generated by $\psi(\cos t, \sin t) = e^{-ijt}$.

If $\text{dist}(\alpha, \mathbb{Z}) = \frac{1}{2}$, then all eigenvalues have multiplicity 2.

Example: Aharonov-Bohm magnetic potentials in $\dim N = 2$

$$\mathbf{A}(\cos t, \sin t) = \alpha(-\sin t, \cos t), \alpha \in \mathbb{R} \setminus \mathbb{Z}, \quad a(\cos t, \sin t) = a_0, a_0 \in \mathbb{R}$$

$$\left(-i \nabla + \alpha \left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2} \right) \right)^2 u - \frac{a_0}{|x|^2} u = h u,$$

with $x = (x_1, x_2) \in \Omega \subset \mathbb{R}^2$,
 Ω bounded, $0 \in \Omega$,
 and h verifying (H_0) .

$$\text{Eigenvalues: } \{(\alpha - j)^2 - a_0 : j \in \mathbb{Z}\} \Rightarrow \mu_1(\mathbf{A}, a) = (\text{dist}(\alpha, \mathbb{Z}))^2 - a_0$$

If $\text{dist}(\alpha, \mathbb{Z}) \neq \frac{1}{2}$, then all eigenvalues are simple and the eigenspace associated to the first eigenvalue is generated by $\psi(\cos t, \sin t) = e^{-ij t}$.

If $\text{dist}(\alpha, \mathbb{Z}) = \frac{1}{2}$, then all eigenvalues have multiplicity 2.

If $a_0 < (\text{dist}(\alpha, \mathbb{Z}))^2$, $\text{dist}(\alpha, \mathbb{Z}) \neq \frac{1}{2} \Rightarrow \exists j_0 \in \mathbb{Z}, \beta \in \mathbb{C}$ s.t.

$$\lambda^{-\sqrt{(\alpha - j_0)^2 - a_0}} u(\lambda \cos t, \lambda \sin t) \xrightarrow{\lambda \rightarrow 0^+} \beta e^{-ij_0 t} \quad \text{in } C^{1,\tau}(0, 2\pi)$$

Example: Aharonov-Bohm magnetic potentials in dim $N = 2$

$$\mathbf{A}(\cos t, \sin t) = \alpha(-\sin t, \cos t), \alpha \in \mathbb{R} \setminus \mathbb{Z}, \quad a(\cos t, \sin t) = a_0, a_0 \in \mathbb{R}$$

$$\left(-i \nabla + \alpha \left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2} \right) \right)^2 u - \frac{a_0}{|x|^2} u = h u, \quad \text{with } x = (x_1, x_2) \in \Omega \subset \mathbb{R}^2, \\ \Omega \text{ bounded, } 0 \in \Omega, \\ \text{and } h \text{ verifying } (H_0).$$

$$\text{Eigenvalues: } \{(\alpha - j)^2 - a_0 : j \in \mathbb{Z}\} \Rightarrow \mu_1(\mathbf{A}, a) = (\text{dist}(\alpha, \mathbb{Z}))^2 - a_0$$

If $\text{dist}(\alpha, \mathbb{Z}) \neq \frac{1}{2}$, then all eigenvalues are simple and the eigenspace associated to the first eigenvalue is generated by $\psi(\cos t, \sin t) = e^{-ij t}$.

If $\text{dist}(\alpha, \mathbb{Z}) = \frac{1}{2}$, then all eigenvalues have multiplicity 2.

if $a_0 < (\text{dist}(\alpha, \mathbb{Z}))^2$, $\text{dist}(\alpha, \mathbb{Z}) = \frac{1}{2} \Rightarrow \exists j_0 \in \mathbb{Z}, \beta_1, \beta_2 \in \mathbb{C}$ s.t.

$$\lambda^{-\sqrt{(\alpha - j_0)^2 - a_0}} u(\lambda \cos t, \lambda \sin t) \xrightarrow{\lambda \rightarrow 0^+} \beta_1 e^{-ij_0 t} + \beta_2 e^{-i(2\alpha - j_0)t} \text{ in } C^{1,\tau}(0, 2\pi)$$

Example: Aharonov-Bohm magnetic potentials in dim $N = 2$

$$\mathbf{A}(\cos t, \sin t) = \alpha(-\sin t, \cos t), \alpha \in \mathbb{R} \setminus \mathbb{Z}, \quad a(\cos t, \sin t) = a_0, a_0 \in \mathbb{R}$$

$$\left(-i \nabla + \alpha \left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2} \right) \right)^2 u - \frac{a_0}{|x|^2} u = h u, \quad \text{with } x = (x_1, x_2) \in \Omega \subset \mathbb{R}^2, \\ \Omega \text{ bounded, } 0 \in \Omega, \\ \text{and } h \text{ verifying } (\mathbf{H}_0).$$

$$\text{Eigenvalues: } \{(\alpha - j)^2 - a_0 : j \in \mathbb{Z}\} \Rightarrow \mu_1(\mathbf{A}, a) = (\text{dist}(\alpha, \mathbb{Z}))^2 - a_0$$

If $\text{dist}(\alpha, \mathbb{Z}) \neq \frac{1}{2}$, then all eigenvalues are simple and the eigenspace associated to the first eigenvalue is generated by $\psi(\cos t, \sin t) = e^{-ij t}$.

If $\text{dist}(\alpha, \mathbb{Z}) = \frac{1}{2}$, then all eigenvalues have multiplicity 2.

Furthermore, in view of the Corollary,

if $(\text{dist}(\alpha, \mathbb{Z}))^2 < 1 + a_0 \Rightarrow u \in C_{\text{loc}}^{0,\gamma}(\Omega)$ with $\gamma = \sqrt{(\text{dist}(\alpha, \mathbb{Z}))^2 - a_0}$

if $(\text{dist}(\alpha, \mathbb{Z}))^2 \geq 1 + a_0 \Rightarrow u$ is locally Lipschitz continuous in Ω .

In preparation [F.-Ferrero-Terracini]...

- Quantum many-body problem: $-\Delta - V$ with

$$V(x) = \sum_{\substack{J_1, J_2 \in \mathcal{A}_k \\ J_1 \cap J_2 = \emptyset}} \frac{\alpha_{J_1 J_2}}{|x_{J_1} - x_{J_2}|^2},$$

$3 \leq k \leq N$, $\mathcal{A}_k := \{\text{multi-indices of length } k\}$, $x_J = (x_i)_{i \in J}$.

In this case $V(x) = \frac{a(x/|x|)}{|x|^2}$ with

$$a(\theta) = \sum_{\substack{J_1, J_2 \in \mathcal{A}_k \\ J_1 \cap J_2 = \emptyset}} \frac{\alpha_{J_1 J_2}}{|\theta_{J_1} - \theta_{J_2}|^2} \notin L^\infty(\mathbb{S}^{N-1}), \quad \theta_J = \frac{x_J}{|x|}.$$

In preparation [F.-Ferrero-Terracini]...

- Quantum many-body problem: $-\Delta - V$ with

$$V(x) = \sum_{\substack{J_1, J_2 \in \mathcal{A}_k \\ J_1 \cap J_2 = \emptyset}} \frac{\alpha_{J_1 J_2}}{|x_{J_1} - x_{J_2}|^2},$$

$3 \leq k \leq N$, $\mathcal{A}_k := \{\text{multi-indices of length } k\}$, $x_J = (x_i)_{i \in J}$.

In this case $V(x) = \frac{a(x/|x|)}{|x|^2}$ with

$$a(\theta) = \sum_{\substack{J_1, J_2 \in \mathcal{A}_k \\ J_1 \cap J_2 = \emptyset}} \frac{\alpha_{J_1 J_2}}{|\theta_{J_1} - \theta_{J_2}|^2} \notin L^\infty(\mathbb{S}^{N-1}), \quad \theta_J = \frac{x_J}{|x|}.$$

- Cylindrical singularities: $V(x) = \sum_{J \in \mathcal{A}_k} \frac{\alpha_J}{|x_J|^2}$, $\alpha_J \in \mathbb{R}$.