

CONGRESSO
SGI-SIMP 2014

Milano
10-12 Settembre 2014

THE ITALIAN GEOSCIENCES

THE FUTURE OF



OF THE FUTURE

GIS analysis to apply theoretical Minimal Model on glacier flow line and assess glacier response in climate change scenarios

Moretti M., Mattavelli M., DeAmicis M. & Maggi V.

Department of Earth and Environmental Sciences (DISAT)


University of Milano-Bicocca

Project NextData




UN A/Res/62/196, 2008 Glacier are sentinels of Climate Change.

“Recognizes that mountains provide indications of global climate change through phenomena such as [...] the retreat of mountain glaciers [...]”



How glaciers respond to Climate Change


Glacial dynamics are too complex to be modelled in every aspects. Theoretical Model was implemented to reduce the complex situation and focus to one aspect.



Minimal Glacier Models (J. Oerlemans 2008, 2011)

Minimal Model is based on continuity equation, that is integrated on entire volume of glacier, and on perfect plasticity principle, a first-order estimate of how the thickness of a glacier varies with its horizontal dimension.

The elaboration is based on reconstruction of historical time series, after have obtained **meteorological**, **physical** and **morphological data** to start the model it is possible compare the **flow line length variation**, the model results, with real measured variations.



Minimal Model fundamentals and GIS interaction:

Minimal Model Input: Mass Balance & ELA

Boundary Condition:

B_0 = highest elevation [m] (β)
 H_m = mean thickness [m] (ΔH)
 b_m = mean bed elevation [m]
 L = length of the flow line [m]
 s = mean slope

$$B_S = W(\beta(H_m + b_m - E)L)$$

$$\frac{dL}{dt} = \left(\frac{3\alpha_m}{2(1 + vs)} L^{1/2} - \frac{\alpha_m v}{(1 + vs)^2} L^{3/2} \frac{\partial s_m}{\partial L} \right)^{-1} \frac{B_S}{W}$$

Mass Balance gradient

$$\beta = \frac{db}{dz} = \frac{\dot{b}}{\bar{h} - E}$$

$$\bar{h} = H_m + b_0 - \frac{L \cdot s}{2}$$

Model BC computable by GIS


Data obtainable by DTM analysis, to evaluate the accuracy it is required a multitemporal dataset → Developed of iterative GIS module.

From DETERMINISTIC to SPATIAL approach using GIS




QGIS Algorithms: MMGlacierData(MMGD)

Development of an algorithm to calculate and iterate all the GIS operations to obtain the input for the minimal model. Developed in QGIS using its different available instruments: GRASS module and GDAL/OGR-libraries.



Study Area

Module MMGlacierData and Minimal Model were tested on Careser and Rutor glacier.



Careser is one the most studied glacier. All dataset used derived by UNIPD TESAF work (Carturan et al, 2007, 2012, 2013)



Rutor glacier is the most studied glacier by UNIMIB DISAT and there is a sufficient dataset to start.

DTMs from:
Carturan et. all, 2013

2007

2000

DTM analysis
[1933 – 2007]
Hillshade movie

1990

1959


1980

1969

1933

MMGD Input: FLOWLINE

Flowlines calculated with Grass **r.flow** used in Qgis and corrected by a geomorphological analysis to choose the most probably.



Parameters Log Help

Show advanced parameters

Elevation
dtm_2007_2x2_32N [EPSG:32632]

Aspect
hill_2000 [EPSG:32632]

Barriers
df_1990_2000_poly90 [EPSG:32632]

Number of cells between flowlines
1,000000

Maximum number of segments per flowline
5,000000

Compute upslope flowlines instead of default downhill flowlines
No

3-D lengths instead of 2-D
No

GRASS region extent(xmin, xmax, ymin, ymax)
[Leave blank to use map covering extent]

GRASS region cellsz (leave 0 for default)
0,000000

Output flowline vector layer
[Salva in un file temporaneo]

Open output file after running algorithm

Output flowpath length raster layer
[Salva in un file temporaneo]

Open output file after running algorithm

Output flowline density raster layer
[Salva in un file temporaneo]

Open output file after running algorithm

0%


Run Close

MMGD Input: Polygons


Polygons are used in MMGD as intersect surface to measure the length of the flowline and to obtain the DTM statistics for a single year.

Polygons years:

- 1933
- 1959
- 1969
- 1980
- 1990
- 2000
- 2006
- 2012




From MMGlacierData to Minimal Model



Minimal Model Results:


Minimal model accuracy using input data from MMGD(b) or input data from literature and averages.

Rutor: east flow line - Miminal Model, improving input



Minimal Model Results:


Careser: minimal model - global climate model RCP 4.5 and regional Protheus



Future projection using RCP 4,5 scenario for CMIP5 and CSIRO global model. Comparison with regional climate model PROTHEUS based on SRES.

Future projection using RCP 8,5 scenario for CMIP5 and CSIRO global model. Comparison with regional climate model PROTHEUS based on SRES.

Careser: minimal model - global climate model RCP 8.5 and regional Protheus



Glacier retreat

I want to be a 'Panda' in China...



www.pulci.org



References

- Haeberli W., Hoelzle M. (2012): Application of inventory data for estimating characteristics of and regional climate-change effects on mountain glaciers: a pilot study with the European Alps. *Annals of Glaciol.*, 21, 206–212.
- Hamming R.W. (1986): Numerical Methods for Scientists and Engineers, *Unabridged Dover*, republication of the 2nd edition published by McGraw-Hill 1973
- Linsbauer A., Paul F., Haberli W. (2012): Modeling glacier thickness distribution and bed topography over entire mountain range with GlabTop: Application of fast and robust approach. *Journal of Geo. Res.*, Vol. 117, F03007, 2012
- Oerlemans J. (2008): Minimal Glacier Models. *Igitur, Utrecht University*, 90 pp., 2008
- Paterson, W. (1994), The Physics of Glaciers, *Pergamon*, Tarrytown, N.Y.
- Carturan L. and Seppi R. (2007): Recent mass balance results and morphological evolution of Careser Glacier (Central Alps). *Geogr. Fis. Din. Quat.*, 30(1), 33–42
- Carturan L. and Seppi R. (2009): Comparison of current behaviour of three glaciers in western Trentino (Italian Alps). In *Epitome: Geoitalia 2009, Settimo Forum Italiano di Scienze della Terra, 9–11 September 2009, Rimini, Italy, Vol. 3*. Federazione Italiana di Scienze della Terra, 298
- Carturan L., Dalla Fontana G. and Cazorzi F. (2009a): The mass balance of La Mare Glacier (Ortles-Cevedale, Italian Alps) from 2003 to 2008. In *Epitome: Geoitalia 2009, Settimo Forum Italiano di Scienze della Terra, 9–11 September 2009, Rimini, Italy, Vol. 3*. Federazione Italiana di Scienze della Terra, 298
- Carturan L., Cazorzi F. and Dalla Fontana G. (2012): Distributed mass-balance modelling on two neighbouring glaciers in Ortles-Cevedale, Italy, from 2004 to 2009. *Journal of Glaciol.*, Vol. 58, No. 209, 2012
- Carturan, L., Baroni, C., Becker, M., Bellin, A., Cainelli, O., Carton, A. & Seppi, R. (2013). Decay of a long-term monitored glacier: the Careser glacier (Ortles-Cevedale, European Alps). *The Cryosphere*, 7(6), 1819-1838.
- General Assembly of the United Nation: Sustainable mountain development, *UN A/Res/62/196*, 2008.
- Knutti R., Masson D., and Gettelman A. (2013): Climate model genealogy: Generation CMIP5 and how we got there. *Geophys. Res. Lett.*
- Jeffrey S., Rotstajn L., Collier M., Dravitzki S., Hamalainen C., Moeseneder C., Wong K. and Syktus J. (2012): Australia's CMIP5 submission using the CSIRO-Mk3.6 model. *Austr. Meteo. Ocean. Journ.* V.63 1-13.
- Taylor K. E., Stouffer R. J. and Meehl G.A. (2012): An Overview of CMIP5 and the Experiment Design. *Am. Meteo. Soc.* DOI:10.1175.