CLASSIFICATION OF FINITE GROUPS THAT ADMIT AN ORIENTED REGULAR REPRESENTATION

JOY MORRIS AND PABLO SPIGA

ABSTRACT. This is the third, and last, of a series of papers dealing with oriented regular representations. Here we complete the classification of finite groups that admit an oriented regular representation (or ORR for short), and give a complete answer to a 1980 question of László Babai: "Which [finite] groups admit an oriented graph as a DRR?" It is easy to see and wellunderstood that generalised dihedral groups do not admit ORRs. We prove that, with 11 small exceptions (having orders ranging from 8 to 64), every finite group that is not generalised dihedral has an ORR.

1. INTRODUCTION

All groups and graphs in this paper are finite. Let G be a group and let S be a subset of G. The **Cayley digraph**, denoted by $\operatorname{Cay}(G, S)$, over G with connection set S is the digraph with vertex set G and with (x, y) being an arc if $yx^{-1} \in S$. (An **arc** is an ordered pair of adjacent vertices.) Since the group G acts faithfully as a group of automorphisms of $\operatorname{Cay}(G, S)$ via the right regular representation, Cayley digraphs represent groups geometrically and combinatorially as groups of automorphisms of digraphs. Naively, the closer G is to the full automorphism group of $\operatorname{Cay}(G, S)$, the closer this representation is to a precise graphical encoding of G.

Following this line of thought, it is natural to ask which groups G admit a subset S with G being the automorphism group of Cay(G, S); that is, Aut(Cay(G, S)) = G. We say that G admits a **digraphical regular representation** (or DRR for short) if there exists a subset S of G with Aut(Cay(G, S)) = G. Babai [1, Theorem 2.1] has given a complete classification of the groups admitting a DRR: except for

(1) $Q_8, C_2^2, C_2^3, C_2^4 \text{ and } C_3^2,$

every group admits a DRR.

In light of Babai's result, it is natural to try to combinatorially represent groups as automorphism groups of special classes of Cayley digraphs. Observe that, if Sis inverse-closed (that is, $S = S^{-1} := \{s^{-1} \mid s \in S\}$), then $\operatorname{Cay}(G, S)$ is undirected. Now, we say that G admits a **graphical regular representation** (or GRR for short) if there exists an inverse-closed subset S of G with $\operatorname{Aut}(\operatorname{Cay}(G, S)) = G$. With a considerable amount of work culminating in [6, 8], the groups admitting a GRR have been completely classified. (In fact, this question attracted significant interest well before the DRR problem, although the final solutions to both problems appeared at about the same time.)

We recall that a **tournament** is a digraph $\Gamma := (V, A)$ with vertex set V and arc set A such that, for every two distinct vertices $x, y \in V$, exactly one of (x, y) and

This research was supported in part by the National Science and Engineering Research Council of Canada grant RGPIN-2017-04905.

(y, x) is in A. After the completion of the classification of DRRs and GRRs, Babai and Imrich [2] proved that every group of odd order except for C_3^2 and C_3^3 admits a **tournament regular representation** (or TRR for short). That is, each finite odd-order group G different from C_3^2 and C_3^3 contains a subset S with $\operatorname{Cay}(G, S)$ being a tournament and with $\operatorname{Aut}(\operatorname{Cay}(G, S)) = G$. In terms of the connection set S, the Cayley digraph $\operatorname{Cay}(G, S)$ is a tournament if and only if $S \cap S^{-1} = \emptyset$ and $G \setminus \{1\} = S \cup S^{-1}$. This observation makes it clear that a Cayley digraph on G cannot be a tournament if G contains an element of order 2, so only groups of odd order can admit TRRs.

In [1, Problem 2.7], Babai observed that one class of Cayley digraphs is rather interesting and had not been investigated in the context of regular representations; that is, the class of oriented Cayley digraphs (or as Babai called them, oriented Cayley graphs). An **oriented Cayley digraph** is in some sense a "proper" Cayley digraph. More formally, it is a Cayley digraph Cay(G, S) whose connection set Shas the property that $S \cap S^{-1} = \emptyset$. Equivalently, in graph-theoretic terms, it is a Cayley digraph with no digons.

Definition 1.1. The group G admits an *oriented regular representation* (or ORR for short) if there exists a subset S of G with $S \cap S^{-1} = \emptyset$ and $\operatorname{Aut}(\operatorname{Cay}(G, S)) = G$.

Babai asked in [1] which (finite) groups admit an ORR. Since a TRR is a special type of ORR, and C_3^2 is one of the five groups in Eq. 1 that do not admit a DRR (so cannot admit an ORR), the answer to this question for groups of odd order was already known when Babai published his question.

In this paper, answering the question of Babai and also confirming the conjecture given in [12, Conjecture 1.5], we prove the following result.

Theorem 1.2. Every finite group G admits an ORR, unless one of the following holds:

- (i): G is generalised dihedral with |G| > 2 (see Definition 2.1 for the meaning of generalised dihedral);
- (ii): G is isomorphic to one of the following eleven groups

$$Q_8, C_4 \times C_2, C_4 \times C_2^2, C_4 \times C_2^3, C_4 \times C_2^4, C_3^2, C_3 \times C_2^3, \langle a, b \mid a^4 = b^4 = (ab)^2 = (ab^{-1})^2 = 1 \rangle \text{ (of order 16)}, \langle a, b, c \mid a^4 = b^4 = c^4 = (ba)^2 = (ba^{-1})^2 = (bc)^2 = (bc^{-1})^2 = 1, a^2 = c^2, a^c = a^{-1}, a^2 = b^2 \rangle \text{ (of order 16)}, \langle a, b, c \mid a^4 = b^4 = c^4 = (ab)^2 = (ab^{-1})^2 = 1, \\ \end{cases}$$

$$(ac)^2 = (ac^{-1})^2 = (bc)^2 = (bc^{-1})^2 = a^2b^2c^2 = 1$$
 (of order 32),

 $D_4 \circ D_4$ (the central product of two dihedral groups of order 8,

which is the extraspecial group of order 32 of plus type).

We remark that since this theorem relies on results in the previous two papers [12, 13] and since the results in [12] depend upon the Classification of Finite Simple Groups, this theorem also depends on the Classification.

In our opinion this is not the final word on oriented regular representations of finite groups. In fact, it is still unclear whether ORRs behave asymptotically like DRRs and GRRs. (It is believed that most Cayley digraphs are DRRs and that most Cayley graphs are GRRs. One should be very careful about how to understand "most" in these statements and we refer the reader to the introduction of [5] for two distinct, natural interpretations of "most".)

We conclude this introductory section by observing that regular representations have shown a new vitality lately. For instance, Marston Conder, Mark Watkins and Tom Tucker [4] have been studying finite groups admitting a graphical Frobenius representation and have posed some very intriguing conjectures in this context. All of these conjectures are in line with the classification of groups that admit DRRs, GRRs, TRRs and now ORRs: except for some "low level noise" (yielding a finite number of small exceptions) and any obvious obstructions, regular representations of the desired type will exist. For DRRs, there are no obvious general obstructions; for TRRs, groups of even order are problematic and yield the only general obstruction; for GRRs, groups admitting automorphisms that map each element to itself or to its inverse are problematic and yield the only general obstruction; for ORRs, groups for which every generating set contains at least one involution (that is, generalised dihedral groups) are problematic and (in light of Theorem 1.2) yield the only general obstruction.

Finally, we refer to [5, 11, 14, 15] for some recent work on similar problems.

2. Earlier work and preliminaries

Before moving to the proof of Theorem 1.2 we need to review the main results that have been proved on oriented regular representations. We start with a few definitions.

Babai pointed out in [1] that generalised dihedral groups of order greater than 2 can never admit an ORR. (Given a group element g, we denote by o(g) its **order**.)

Definition 2.1. Let A be an abelian group. The generalised dihedral group over A is the group $\langle \tau, A \rangle$ with $o(\tau) = 2$ and $\tau a \tau = a^{-1}$ for every $a \in A$.

In the special case where A is cyclic, this is the dihedral group over A. Observe that, unless |G| = 2, if Cay(G, S) is an ORR, then Cay(G, S) is connected and hence S is a generating set for G. Now, Babai's observation follows immediately from the fact that if G is the generalised dihedral group over the abelian group A, then every element of $G \setminus A$ has order 2. Thus every generating set S for G must contain an involution, so that $S \cap S^{-1} \neq \emptyset$. This renders understanding generalised dihedral groups very important when we are studying ORRs.

Let G be a finite group. As customary, we denote by d(G) the **minimum** number of generators for G. Following [13, Section 2], we say that a generating set $\{g_1, \ldots, g_d\}$ for G is *irredundant* if, for each $i \in \{1, \ldots, d\}$, the d-1 elements

$$g_1, g_2, \ldots, g_{i-1}, g_{i+1}, \ldots, g_d$$

do not generate G. Observe that each generating set for G of cardinality d(G) is irredundant.

We say that the *d*-tuple (g_1, \ldots, g_d) of elements of *G* is **beautiful** if the following conditions hold:

(i): $\{g_1, \ldots, g_d\}$ is an irredundant generating set for G, (ii): $o(g_i) > 2$ for every $i \in \{1, ..., d\}$, (iii): $o(g_{i+1}g_i^{-1}) > 2$ for every $i \in \{1, ..., d-1\}$.

Observe that being beautiful is a property of ordered tuples and not of sets; that is, it depends upon the ordering of the generating set $\{g_1, \ldots, g_d\}$ for G.

An important connection between beautiful generating tuples and ORRs is given in the next theorem.

Theorem 2.2. Let G be a finite group admitting a beautiful generating tuple. Then G admits an ORR if and only if $G \not\cong Q_8$, $G \not\cong C_3 \times C_2^3$, and $G \not\cong C_3 \times C_3$.

This theorem is implicit in [12] and follows immediately from the theory developed therein. For a proof see [13, Theorem 2.1].

Developing the theory of beautiful generating tuples (and actually something more general, which we called five-product-avoiding generating sets), we have proved in [12] that each non-soluble group admits an ORR. Building on this result, the second author has proved the following result, which (among other things) reduces the classification of groups admitting an ORR to some very specific infinite families of 2-groups.

Theorem 2.3 (([13], Theorem 1.2)). Let G be a finite group. Then one of the following holds:

- (i): G admits an ORR;
- (ii): G has an abelian 2-subgroup A, a normal subgroup N and two elements $g \in G \setminus N$ and $n \in N \setminus A$ with A < N < G, |G:N| = |N:A| = 2, $g^2 = 1$, $n^g = n^{-1}$ and $a^g = a^{-1}$ for each $a \in A$;
- (iii): there exists a normal subgroup N of G, $g \in G$ and $n_0 \in N$ with |G:N| = 2, $G = \langle N, g \rangle$, $g^2 = 1$, N is a 2-group and the action of g by conjugation on N inverts precisely half of the elements of N and $N = H \cup n_0 H$, where $H := \{n \in N \mid n^g = n^{-1}\}$. Moreover, N has no automorphism inverting more than half of its elements. (Every group N that has an automorphism inverting half of its elements and no automorphism that inverts more is classified in [7] by Hegarty and MacHale);
- (iv): G is isomorphic to Q_8 , to $C_3 \times C_3$ or to $C_3 \times C_2^3$;
- (v): G is generalised dihedral.

In view of this theorem, the classification of finite groups admitting an ORR is reduced to the groups in (ii) and (iii). We address these two families in Sections 4 and 5, respectively.

We remark that the "flavour" of many of the ORRs that are produced in this paper is quite different from those produced in [12] and [13]. Those papers focused on the use of beautiful generating tuples, so that for any two consecutive elements g_i and g_{i+1} of the generating tuple, the product $g_{i+1}g_i^{-1}$ has order greater than 2. As we will see, a common situation for a group G in the families we study in this paper is that G contains an elementary abelian subgroup B of high rank and low index. In order to generate such a group G, we require a large number of elements from at least one of the cosets Bg of B. If we place any two elements $g_i := b_1g$ and $g_{i+1} := b_2g$ of Bg (where $b_1, b_2 \in B$) consecutively in the generating set, then $g_{i+1}g_i^{-1} = b_2gg^{-1}b_1^{-1} = b_2b_1^{-1} \in B$, so $o(g_{i+1}g_i^{-1}) \leq 2$. In principle it could be possible to ensure that whenever $g_i \in Bg$ we have $g_{i+1} \notin Bg$, but in practice we may still find that $o(g_{i+1}g_i^{-1}) = 2$. We will therefore take a very different approach that uses a GRR for B as a starting point. In our previous papers, if we look at the induced subdigraph of each of our ORRs on the vertices that lie in the connection set (equivalently, the induced subdigraph on the open neighbourhood

 $\mathbf{4}$

CLASSIFICATION OF FINITE GROUPS THAT ADMIT AN ORIENTED REGULAR REPRESENTATION

of any vertex), that digraph is (weakly) connected and asymmetric. The ORRs we produce in this paper will often include many isolated vertices in that induced subdigraph.

Before moving into our analysis of the groups in (ii) and (iii) (from above), we introduce some other results from the literature that will be important in our proofs.

The following result is found in the proof of the theorem in [9]. Note that the statement in [9] assumes only $k \ge 5$, but there is a mistake in the case k = 5 that has been pointed out by multiple researchers. Although the elementary abelian 2-group of rank 5 also admits a GRR, it requires a different connection set so we omit it from our statement.

Lemma 2.4 ((Imrich [9])). Let G be an elementary abelian 2-group of rank $k \ge 6$, and let $\{x_1, \ldots, x_k\}$ be a generating set for G. Then G has a GRR; furthermore, a GRR is given by the Cayley graph on G whose connection set consists of the 2k + 1elements:

(2) $x_1, \ldots, x_k, x_1x_2, x_2x_3, \ldots, x_{k-1}x_k, x_1x_2x_{k-2}x_{k-1}, x_1x_2x_{k-1}x_k.$

Note that the connection set found in this lemma depends on the order as well as the choice of the elements in the generating set for G. We therefore use the following definition.

Definition 2.5. Let G be an elementary abelian 2-group of rank $k \ge 6$. Given the generating tuple (x_1, \ldots, x_k) for G, we refer to the connection set given in Eq. 2 as the *Imrich generating set for G with respect to* (x_1, \ldots, x_k) .

In their work on the GRR problem, Nowitz and Watkins proved a lemma that is very useful in our context also. (Given a graph Γ and a vertex v of Γ , we denote by Aut(Γ) the automorphism group of Γ and by Aut(Γ)_v the stabiliser of the vertex vin Aut(Γ).)

Lemma 2.6 ((Nowitz and Watkins [10])). Let G be a group, let S be a subset of G, let $\Gamma := \operatorname{Cay}(G, S)$ and let X be a subset of G. If φ fixes X pointwise for every $\varphi \in \operatorname{Aut}(\Gamma)_1$, then φ fixes $\langle X \rangle$ pointwise for every $\varphi \in \operatorname{Aut}(\Gamma)_1$.

Thus, if $\operatorname{Aut}(\Gamma)_1$ fixes every element of a generating set for G or if the subgraph induced by Γ on the neighbourhood $\Gamma(1) = S$ is asymmetric, then $\operatorname{Aut}(\Gamma) = G$. Hence Γ is a DRR for G, and is therefore an ORR if the connection set satisfies $S \cap S^{-1} = \emptyset$. We will use this fact repeatedly when we cite the above lemma. Also, although Nowitz and Watkins did not make this explicit, the same proof applies if we replace both occurrences of the word "pointwise" in the statement of Lemma 2.6 with the word "setwise." We will also use this sometimes when we cite the above lemma.

We include in this section two more lemmas that we will need. Lemma 2.7 follows fairly easily from the work in [12], but we include a complete proof as the precise connection set (and therefore the fact that the induced subgraph on that connection set is weakly connected) is not easy to see from the statements in that paper.

Lemma 2.7. Let A be an abelian 2-group of order 2^k . Assume that A is not elementary abelian and $A \not\cong C_4 \times C_2^{k-2}$. Then there is a generating set S for A with $|S| \geq 2$ such that the induced subgraph of $\Gamma := \operatorname{Cay}(G, S)$ on S is weakly connected with trivial automorphism group, so that Γ is an ORR.

Proof. From the structure of finite abelian groups, we may write $A = A_1 \times \cdots \times A_m$, where $A_i := \langle a_i \rangle$ is a non-trivial cyclic group for each $i \in \{1, \ldots, m\}$. Moreover, we may assume that $o(a_{i+1})$ divides $o(a_i)$ for every $i \in \{1, \ldots, m-1\}$.

Since A is not elementary abelian and $A \not\cong C_4 \times C_2^{k-2}$, we must have either $o(a_1) > 4$, or $m \ge 2$ and $o(a_1) = o(a_2) = 4$. In the first case, set $x_1 := a_1$, $x_2 := a_1^{-1}a_2$ and $x_i := a_1^{(-1)^{i-1}}a_i$ for every $i \in \{3, \ldots, m\}$; in the second case, set $x_1 := a_1, x_2 := a_2, x_{2i+1} := a_1a_{2i+1}$ and $x_{2j} := a_2a_{2j}$ for every $2i + 1, 2j \in \{3, \ldots, m\}$. In both cases, it is easy to verify that the set $X := \{x_1, \ldots, x_m\}$ generates A irredundantly. (In fact, (x_1, \ldots, x_m) is a beautiful generating tuple for A.)

Consider $S := X \cup Y$ where

$$Y := \begin{cases} \{x_1^2\} & \text{if } m = 1, \\ \{x_2 x_1^{-2}\} \cup \{x_2 x_1^{-1}, x_3 x_2^{-1}, \dots, x_m x_{m-1}^{-1}\} & \text{if } m \ge 2 \text{ and } o(a_1) > 4, \\ \{x_1 x_2\} \cup \{x_2 x_1^{-1}, x_3 x_2^{-1}, \dots, x_m x_{m-1}^{-1}\} & \text{if } m \ge 2 \text{ and } o(a_1) = o(a_2) = 4. \end{cases}$$

Let $\Gamma := \operatorname{Cay}(A, S)$ and observe that $S \cap S^{-1} = \emptyset$ so that Γ is an oriented digraph.




FIGURE 1. The oriented graph Δ in the proof of Lemma 2.7: top left when m = 1; top right when $m \ge 2$ and $o(a_1) > 4$; and bottom when $m \ge 2$ and $o(a_1) = o(a_2) = 4$

Let Δ be the subgraph induced by Γ on S. It is clear that (x_{i-1}, x_i) and $(x_i x_{i-1}^{-1}, x_i)$ are arcs of Δ for every $i \in \{2, \ldots, m\}$. Moreover, when m = 1, (x_1, x_1^2) is an arc of Δ ; when $m \geq 2$ and $o(a_1) > 4$, $(x_1, x_2 x_1^{-1})$ and $(x_2 x_1^{-2}, x_2 x_1^{-1})$ are arcs of Δ ; while, when $m \geq 2$ and $o(a_1) = o(a_2) = 4$, $(x_1, x_1 x_2)$ and $(x_2, x_1 x_2)$ are arcs of Δ . (See Figure 1.) This shows that Δ is weakly connected. It is not hard (but rather tedious because it requires some detailed computations) to show that these are the only arcs of Δ . We do not give a complete proof of this easy fact, but we deal with one case to show the type of computations that are required. Suppose that there exists an arc $(x_i x_{i-1}^{-1}, x_j x_{j-1}^{-1})$ between two distinct vertices in Y, for some $i, j \in \{2, \ldots, m\}$. Then $x_j x_{j-1}^{-1} x_{i-1} x_i^{-1} \in S$ and hence either $x_j x_{j-1}^{-1} x_{i-1} x_i^{-1} = x_k$ for some $k \in \{1, \ldots, m\}$, or $x_j x_{j-1}^{-1} x_{i-1} x_i^{-1} = x_k x_{k-1}^{-1}$ for some $k \in \{2, \ldots, m\}$. or $x_j x_{j-1}^{-1} x_{i-1} x_i^{-1} \in \{x_1^2, x_1 x_2, x_2 x_1^{-2}\}$. Each of these relations contradicts the irredundancy of the generating set X.

The structure of the arcs that we have described in Δ (see again Figure 1) shows that Δ has trivial automorphism group. Lemma 2.6 then implies that Γ is an ORR.

We conclude this section with a technical lemma. Although we will only apply this lemma with $|G:N| \leq 8$, we state it in greater generality as the proof is no different and the general result may be useful in future work on ORRs.

Lemma 2.8. Let G be a 2-group and suppose that G is not generalised dihedral. Let N be a normal subgroup of G with G/N elementary abelian. Suppose that there exists $T \subseteq N$ with $|T| \ge 2$ such that Cay(N,T) is an ORR and such that the subgraph induced by Cay(N,T) on the neighbourhood T of the vertex 1 is weakly connected. Let $\kappa := d(G/N)$. Then there is a set of elements $\{a_1, \ldots, a_\kappa\} \subseteq G \setminus N$ such that:

- $G = \langle a_1, \ldots, a_{\kappa}, N \rangle$; and
- $o(a_i) > 2$ for each $i \in \{1, ..., \kappa\}$.

Furthermore, if there exists such a set of elements with the additional property that, for every $i, j \in \{1, ..., \kappa\}$ with $i \neq j$, we have a_i^2 centralises N, and either $a_i^2 \neq a_j^2$ or $a_i a_j$ does not centralise N, then $\operatorname{Cay}(G, T \cup \{a_1, ..., a_\kappa\})$ is an ORR

Proof. Let $T \subseteq N$ satisfying the hypothesis of this lemma; that is: $|T| \geq 2$, $\operatorname{Cay}(N,T)$ is an ORR, and the subgraph induced by $\operatorname{Cay}(N,T)$ on the neighbourhood T of the vertex 1 is weakly connected.

We begin by proving the first conclusion of this lemma. (The argument here is similar to the proof of [12, Lemma 2.6] and uses the ideas therein.) Among all κ -tuples a_1, \ldots, a_{κ} of elements of G such that $\{a_1N, \ldots, a_{\kappa}N\}$ is a generating set for G/N, choose one with as few involutions as possible. If no element in $\{a_1, \ldots, a_{\kappa}\}$ is an involution, then our first conclusion is proved. Hence we suppose that $\{a_1, \ldots, a_{\kappa}\}$ has at least one involution. Relabeling the index set $\{1, \ldots, \kappa\}$ if necessary, we may assume that a_1 is an involution.

Let $n \in N$. Now, $\{a_1n, a_2, \ldots, a_\kappa\}$ is still a generating set for G modulo N of cardinality κ . Since this generating set cannot contain fewer involutions than the original generating set, the element a_1n must be an involution. Thus $1 = (a_1n)^2 = a_1na_1n = n^{a_1}n$, that is, $n^{a_1} = n^{-1}$. Since this argument does not depend upon $n \in N$, we obtain that a_1 acts by conjugation inverting each element of N. In particular, N is abelian.

Let $j \in \{2, \ldots, \ell\}$ and let $n \in N$. Now, $\{a_1a_jn, a_2, a_3, \ldots, a_\kappa\}$ is still a generating set for G modulo N of cardinality κ . Since this generating set cannot contain fewer involutions than the original generating set, the element a_1a_jn must be an involution. Thus

$$1 = (a_1 a_j n)^2 = a_1 a_j n a_1 a_j n = a_1^2 (a_j n)^{a_1} a_j n = a_j^{a_1} n^{a_1} a_j n = a_j^{a_1} n^{-1} a_j n;$$

so $a_j^{a_1} = n^{-1}a_j^{-1}n$. By applying this equality with n := 1, we deduce that conjugation by a_1 inverts a_j . Therefore $a_j^{-1} = a_j^{a_1} = n^{-1}a_j^{-1}n$ for each $n \in N$, that is, a_j commutes with N.

Let $i, j \in \{2, ..., \kappa\}$ with $i \neq j$. Arguing as above, $\{a_1 a_i a_j, a_2, a_3, ..., a_\ell\}$ is still a generating set for G modulo N of cardinality κ . Since this generating set cannot contain fewer involutions than the original generating set, the element $a_1a_ia_j$ must be an involution. Thus

$$1 = (a_1 a_i a_j)^2 = a_1 a_i a_j a_1 a_i a_j = a_1^2 (a_i a_j)^{a_1} a_i a_j = (a_i^{a_1} a_j^{a_1}) a_i a_j = (a_i^{-1} a_j^{-1}) a_i a_j,$$

so $a_i a_j = a_j a_i$; that is, a_i and a_j commute.

This shows that $M := \langle N, a_2, \ldots, a_\ell \rangle$ is an abelian normal subgroup of G. Since $G = \langle M, a_1 \rangle$ and a_1 has order 2, we have |G : M| = 2. Moreover, since the action of a_1 by conjugation inverts a generating set for M, we see that G is a generalised dihedral group over M, contrary to our assumption. Our first conclusion is thus proven.

To complete the proof, we assume from now on that $\{a_1, \ldots, a_\kappa\}$ satisfies the additional hypothesis that, for every $i, j \in \{1, \ldots, \kappa\}$ with $i \neq j$, we have a_i^2 centralises N, and either: $a_i^2 \neq a_j^2$, or $a_i a_j$ does not centralise N.

Consider the set $S := T \cup \{a_1, \ldots, a_\kappa\}$. By construction, $S \cap S^{-1} = \emptyset$ and hence $\Gamma := \operatorname{Cay}(G, S)$ is an oriented digraph. When $\kappa = 0$, that is, G = N, Γ is an ORR as part of our hypothesis. Thus, for the rest of the proof, we assume that $\kappa \ge 1$.

Let φ be an automorphism of Γ with $1^{\varphi} = 1$. Now, φ fixes the neighbourhood $\Gamma(1) = S$ of the vertex 1, that is, φ acts as a group of automorphisms of the subgraph Δ induced by Γ on S. By construction, the vertices a_1, \ldots, a_{κ} are isolated in Δ . As $|T| \geq 2$ and a_1, \ldots, a_{κ} are isolated in Δ , T is the (unique) connected component of Δ of largest possible order. Thus φ fixes T setwise. Therefore, by Lemma 2.6, φ fixes $\langle T \rangle = N$ setwise. Thus φ acts as an automorphism of the graph induced by Γ on N; that is, on the Cayley digraph $\operatorname{Cay}(N, S \cap N) = \operatorname{Cay}(N, T)$. By hypothesis, $\operatorname{Cay}(N, T)$ is an ORR and hence φ fixes N pointwise. Furthermore, this shows that, for every $n \in N$ and every vertex v of Γ , we have

$$(3) (nv)^{\varphi} = nv^{\varphi}.$$

Suppose that, for some $i \in \{1, ..., \kappa\}$, we have $a_i^{\varphi} \neq a_i$. Since $a_i^{\varphi} \in S$, there exists $j \in \{1, ..., \kappa\} \setminus \{i\}$ with $a_i^{\varphi} = a_j$. Now (a_i, a_i^2) is an arc of Γ and hence $(a_i^{\varphi}, (a_i^2)^{\varphi})$ is also an arc of Γ . Since G/N is elementary abelian, we have $a_i^2 \in N$ and, since φ fixes N pointwise, we have $(a_i^2)^{\varphi} = a_i^2$. Therefore, $a_i^2(a_i^{\varphi})^{-1} = a_i^2 a_j^{-1} \in S \setminus N = \{a_1, ..., a_{\kappa}\}$. Since $d(G/N) = \kappa$, this forces $a_i^2 a_j^{-1} = a_j$, so $a_i^2 = a_j^2$.

By Eq. (3), $(na_i)^{\varphi} = na_j$ for every $n \in N$. Fix $n \in N$, and observe that $a_i na_i \in N$ because G/N is elementary abelian and that $(a_i na_i)^{\varphi} = a_i na_i$ because φ fixes N pointwise. Since $(na_i, a_i na_i)$ is an arc of Γ , $((na_i)^{\varphi}, (a_i na_i)^{\varphi}) = (na_j, a_i na_i)$ is also an arc of Γ ; that is,

$$a_i n a_i (n a_j)^{-1} = a_i^2 (a_i^{-1} n a_i) (a_j^{-1} n^{-1} a_j) a_j^{-1} = a_i^2 n^{a_i} (n^{-1})^{a_j} a_j^{-1} \in S.$$

As $a_i^2 n^{a_i} (n^{-1})^{a_j} a_j^{-1}$ lies in the coset Na_j^{-1} and G/N is elementary abelian, we deduce $a_i^2 n^{a_i} (n^{-1})^{a_j} a_j^{-1} = a_j$; that is, $n^{a_i} = n^{a_j}$. Thus $n^{a_i a_j} = (n^{a_i})^{a_j} = n^{a_j^2} = n$ because a_j^2 centralises N by hypothesis. As this argument does not depend upon $n \in N$, $a_i a_j$ centralises N, contradicting our hypothesis.

Since *i* was arbitrary and our only assumption was that $(a_i)^{\varphi} \neq a_i$, we conclude that φ fixes *S* pointwise, so by Lemma 2.6 we have $\varphi = 1$ and Γ is an ORR.

3. Constructing ORRs in groups that have low-index elementary Abelian subgroups

In this section, we prove some results that will be useful for constructing ORRs on many groups that have elementary abelian subgroups of low index. These results make intense use of Imrich generating sets for GRRs on elementary abelian groups of rank at least 6, so they require the elementary abelian subgroup to be of sufficiently high rank.

Given a Cayley digraph $\operatorname{Cay}(G, S)$ and two vertices x and y, we say that x and y are **adjacent via** s if $yx^{-1} = s \in S$. Similarly, we say that x and y are **adjacent via** X if $yx^{-1} \in X \subset S$.

Lemma 3.1. Let B be an elementary abelian 2-group of rank $k \ge 6$, and let T be the Imrich generating set for B (with respect to some generating tuple). Let G be a group with $B \le G$, let Bx be a coset of B in G and let S be any subset of G with $S \cap Bx = Tx \cup \{x\}$. Then in Cay(G, S), x is the only vertex of $S \cap Bx$ that has at least three mutual inneighbours via elements of $S \cap Bx$ with every other vertex of $S \cap Bx$.

Proof. Let the generating tuple be (z_1, \ldots, z_k) . We begin by showing that x has at least three common inneighbours via elements of $S \cap Bx$ with every other vertex of $S \cap Bx$. Let tx be an arbitrary vertex of $(S \cap Bx) \setminus \{x\}$, so $t \in T$. Observe that y is a mutual inneighbour via elements of $S \cap Bx$ of x and tx if and only if there exist $t_1, t_2 \in T \cup \{1\}$ such that $t_1xy = x$ and $t_2xy = tx$. Equivalently, $y = t_1^x$ and $t_1t_2 = t$. This is clearly satisfied with $y = t_1 = 1$ and $t_2 = t$, so that 1 is a mutual inneighbour. It is also always satisfied by taking $y = t^x$, $t_1 = t$, and $t_2 = 1$, so that t^x is a mutual inneighbour. The third mutual inneighbour depends on t.

If $t = z_i$ for some $i \in \{1, \ldots, k-1\}$, then taking $t_1 := z_i z_{i+1}$, $t_2 := z_{i+1}$, and $y := t_1^x$ gives y as a third mutual inneighbour. If $t = z_k$, then taking $t_1 := z_{k-1}z_k$, $t_2 := z_{k-1}$, and $y := t_1^x$ gives y as a third mutual inneighbour. If $t = z_i z_{i+1}$ for some $i \in \{1, \ldots, k-1\}$, then taking $t_1 := z_i$, $t_2 := z_{i+1}$, and $y := t_1^x$ gives y as a third mutual inneighbour. If $t = z_i z_{i+1}$ for some $i \in \{1, \ldots, k-1\}$, then taking $t_1 := z_i$, $t_2 := z_{i+1}$, and $y := t_1^x$ gives y as a third mutual inneighbour. Finally, if $t = z_1 z_2 z_{\ell-1} z_\ell$ for $\ell \in \{k-1, k\}$, then taking $t_1 := z_1 z_2$, $t_2 := z_{\ell-1} z_\ell$, and $y := t_1^x$ gives y as a third mutual inneighbour.

Now we show that for any vertex tx of Tx, there is some vertex t'x of $Tx \setminus \{tx\}$ such that tx and t'x have fewer than three common inneighbours via elements of $S \cap Bx$. Observe that y is a mutual inneighbour via elements of $S \cap Bx$ of tx and t'x if and only if there exist $t_1, t_2 \in T \cup \{1\}$ such that $t_1xy = tx$ and $t_2xy = t'x$. Equivalently, $y = (t_1t)^x$ and $t_1t_2 = tt'$. This is clearly satisfied with $y = 1, t_1 = t$, and $t_2 = t'$. It is also satisfied with $y = (tt')^x$, $t_1 = t'$, and $t_2 = t$. Thus 1 and $(tt')^x$ are mutual inneighbours of tx and t'x.

If $t = z_i$ for some $i \in \{1, \ldots, k\}$, then let $t' := z_j$ for some $j \notin \{i - 2, i - 1, i, i + 1, i + 2\}$ (such a j exists since $k \ge 6$). There is no way to write $z_i z_j$ as a product of two elements $t_1, t_2 \in T \cup \{1\}$ except when $\{t_1, t_2\} = \{z_i, z_j\}$. Thus there is no third mutual inneighbour of $z_i x$ and $z_j x$ via elements of $S \cap Bx$. Similarly, if $t = z_i z_{i+1}$, then let t' be either $z_{i+2} z_{i+3}$ when $i \le k-3$, or $z_{i-2} z_{i-1}$ otherwise. Again, if tx and t'x had a third mutual inneighbour, then there would be some other way of writing some $z_j z_{j+1} z_{j+2} z_{j+3}$ (where $j \in \{i, i-2\}$) as a product of two elements of $T \cup \{1\}$, but this is not possible. Finally, if $t = z_1 z_2 z_{\ell-1} z_{\ell}$ where $\ell \in \{k-1, k\}$, then let $t' := z_3$. There is no other way of writing $z_1 z_2 z_3 z_{\ell-1} z_{\ell}$ as a product of two elements of $T \cup \{1\}$. This completes the proof.

Unfortunately, to make the above lemma directly useful in building ORRs, we would need to know that every automorphism of $\operatorname{Cay}(G, S)$ that fixes 1 also fixes $S \cap Bx = Tx \cup \{x\}$ setwise. This is challenging to prove in general. It turns out to be much easier to prove that if $S \cap Bx = (Bx \setminus Tx) \setminus \{x\}$, then every automorphism of $\operatorname{Cay}(G, S)$ that fixes 1 also fixes $S \cap Bx$. We therefore prove the following result about the case where $S \cap Bx = (Bx \setminus Tx) \setminus \{x\}$, which is a corollary to the above lemma.

Corollary 3.2. Let B be an elementary abelian 2-group of rank $k \ge 6$, and let T be the Imrich generating set for B (with respect to some generating tuple). Let G be a group with $B \le G$, let Bx be a coset of B in G and let S be any subset of G with $S \cap Bx = (Bx \setminus Tx) \setminus \{x\}$. If $Aut(Cay(G, S))_1$ fixes $S \cap Bx$ setwise, then $Aut(Cay(G, S))_1$ fixes x.

Proof. Assume that, for every $\varphi \in \operatorname{Aut}(\operatorname{Cay}(G, S))_1$, $(S \cap Bx)^{\varphi} = S \cap Bx$. Observe first that B^x is uniquely determined setwise as the set of vertices that have an outneighbour in $S \cap Bx$ via some element of $S \cap Bx$. Furthermore, $X := Tx \cup \{x\}$ is uniquely determined as the set of vertices that are not in S, but are outneighbours of some vertex in B^x via some element of $S \cap Bx$. Now Lemma 3.1 (applied with the set S replaced by $G \setminus S$) tells us that x is the only vertex in X having the property that, for every other vertex $y \in X$, there are at least three vertices of B^x that are not inneighbours of either y or x. Thus $x^{\varphi} = x$, for every $\varphi \in \operatorname{Aut}(\operatorname{Cay}(G, S))_1$. \Box

Finally, we can describe a situation that will arise frequently in which $S \cap Bx = (Bx \setminus Tx) \setminus \{x\}$ is fixed setwise by every automorphism φ of Cay(G, S) that fixes 1. We show that in this situation, B and x are fixed pointwise by φ , which is a significant step toward proving that Cay(G, S) is an ORR.

Proposition 3.3. Let G be a group, and suppose that B < G where B is an elementary abelian group of rank $k \ge 6$, and $x \in G$ centralises B. Let T be the Imrich generating set for B with respect to some generating tuple, and let S be a generating set for G such that $S = [(Bx \setminus Tx) \setminus \{x\}] \cup X$, where $X \subseteq G \setminus \langle B, x \rangle$ with $|X| \le 17$. Then $\operatorname{Aut}(\operatorname{Cay}(G, S))_1$ fixes B and x pointwise.

Proof. We claim that a vertex of Cay(G, S) is an outneighbour of at least $2^k - 4k - 4$ vertices of S if and only if it is an element of Bx^2 .

Let bx^2 be an arbitrary element of Bx^2 . Then, for any $c \in (B \setminus T) \setminus \{1\}$, we have $bc \in B$, so $(bcx)(cx) = bx^2$ is an outneighbour of cx unless $bc \in T \cup \{1\}$; that is, unless $c \in b(T \cup \{1\})$. Since there are 2k + 2 elements in $b(T \cup 1)$, there are at most 2k + 2 vertices in $(Bx \setminus Tx) \setminus \{x\}$ that are not inneighbours of bx^2 . This means that there are at least $2^k - 4k - 4$ vertices in S that are inneighbours of bx^2 .

On the other hand, if we take a vertex v that is not in Bx^2 and it is an outneighbour of at least $2^k - 4k - 4$ vertices of S, then since $2^k - 4k - 4 > 17$, v must be an outneighbour of some vertex $bx \in Bx \cap S$. As we observed above, outneighbours of vertices of Bx via elements of $S \cap Bx$ are in Bx^2 , so $v \notin Bx^2$ implies that v is an outneighbour of at most 17 vertices that are in Bx. Therefore, v is an outneighbour of at most 34 vertices of S (at most 17 in Bx and at most 17 in X). Since $2^k - 4k - 4 > 34$, this completes the proof of our claim.

Let $\varphi \in \operatorname{Aut}(\operatorname{Cay}(G,S))_1$. The previous paragraph shows that $(Bx^2)^{\varphi} = Bx^2$. Since Bx is the set of vertices of $\operatorname{Cay}(G,S)$ such that all but at most 17 of their outneighbours lie in Bx^2 , we also have $(Bx)^{\varphi} = Bx$. Thus $(S \cap Bx)^{\varphi} = S \cap Bx = (Bx \setminus Tx) \setminus \{x\}$. Applying Corollary 3.2 to our graph, we see that $x^{\varphi} = x$, and so $(Tx)^{\varphi} = Tx$. Since T is the connection set for a GRR on B and x is fixed, this implies that Bx is fixed pointwise, which means that (using Lemma 2.6) B is fixed pointwise. \Box

We will be using this proposition for most groups G arising from Theorem 2.3 that contain an elementary abelian subgroup of low index. As long as G is sufficiently large, the elementary abelian subgroup will have high enough rank to allow us to apply this result. For small orders of G, we will make use of computations in magma to complete the proof.

4. The groups in Theorem 2.3 (ii)

In this section we deal with the groups arising from Theorem 2.3 (ii). First, we give a result that finds an ORR for all but three specific families of groups.

Proposition 4.1. Let G be a 2-group with an abelian subgroup A, with a normal subgroup N and with two elements $g \in G \setminus N$ and $n \in N \setminus A$ such that

$$A < N < G, |G:N| = |N:A| = 2, g^2 = 1, n^g = n^{-1}, and a^g = a^{-1}, for each a \in A$$

Then one of the following holds:

- (a) G has an ORR;
- (b) l and κ are non-negative integers, V is an elementary abelian 2-group of rank 2l + κ with generating set

$$\{v_1, w_1, \ldots, v_\ell, w_\ell, e_1, \ldots, e_\kappa\},\$$

and $G = \langle V, x \rangle$, where $v_i^x = w_i$, $w_i^x = v_i$ for $i \in \{1, \ldots, \ell\}$ and $e_i^x = e_i$ for $i \in \{1, \ldots, \kappa\}$; and

- (i) $x^2 = 1$; or
- (ii) $\kappa \geq 1$ and $x^2 = e_1$;
- (c) G contains a maximal subgroup D isomorphic to $D_4 \times C_2^{\ell}$ for some $\ell \in \mathbb{N}$ (D_4 is the dihedral group of order 8); or
- (d) G is generalised dihedral.

Proof. We assume that conclusion (d) does not hold, that is, G is not generalised dihedral.

Suppose that A has exponent 2. Then $V := \langle A, g \rangle$ is a maximal subgroup of G and is elementary abelian. Now, V is a $\mathbb{Z}\langle n \rangle$ -module. Therefore, from the structure theorem of finitely generated $\mathbb{Z}\langle n \rangle$ -modules, we may write $V = A_1 \times A_2 \times \cdots \times A_{\ell+\kappa}$, where $A_i := \langle v_i \rangle$ is a non-identity non-trivial cyclic $\mathbb{Z}\langle n \rangle$ -module for each $i \in \{1, \ldots, \ell\}$, and $A_{\ell+i} := \langle e_i \rangle$ is a non-identity trivial (and hence cyclic) $\mathbb{Z}\langle n \rangle$ -module for each $i \in \{1, \ldots, \kappa\}$. For each $i \in \{1, \ldots, \ell\}$, write $w_i := v_i^n$. As $n^2 \in A$ centralises $V, v_1, w_1, v_2, w_2, \ldots, v_\ell, w_\ell, e_1, \ldots, e_\kappa$ is a basis of V (as a vector space over the field of cardinality 2) and $|V| = 2^{2\ell+\kappa}$.

Now the action of n on V is encoded in the pair (ℓ, κ) . Thus, to determine the isomorphism class of G it suffices to determine n^2 . Write $n^2 = v_1^{\varepsilon_1} w_1^{\varepsilon'_1} \cdots v_\ell^{\varepsilon_\ell} w_\ell^{\varepsilon'_\ell} e_1^{\eta_1} \cdots e_{\kappa}^{\eta_{\kappa}}$, for some $\varepsilon_1, \varepsilon'_1, \ldots, \varepsilon_\ell, \varepsilon'_\ell, \eta_1, \ldots, \eta_{\kappa} \in \{0, 1\}$. Since n centralizes n^2 and $v_i^n = w_i$, $w_i^n = v_i$, we deduce that $\varepsilon_i = \varepsilon'_i$ for every $i \in \{1, \ldots, \ell\}$. Let $I := \{i \in \{1, \ldots, \ell\} \mid \varepsilon_i = 1\}$ and write $n' := n \prod_{i \in I} v_i$. Now, n and n' induce the same action by conjugation on V and $(n')^2 = e_1^{\eta_1} \cdots e_{\kappa}^{\eta_{\kappa}}$. In particular, replacing n by n' if necessary, we may assume that $\varepsilon_i = 0$ for each $i \in \{1, \ldots, \ell\}$.

If $\eta_1 = \cdots = \eta_{\kappa} = 0$, then $n^2 = 1$ and G splits over V. Taking x := n, this is conclusion (b)(i).

If $\eta_i \neq 0$ for some $i \in \{1, \ldots, \kappa\}$, then $n^2 \neq 1$ and $e_1, \ldots, e_{i-1}, n^2, e_{i+1}, \ldots, e_{\kappa}$ are linearly independent and span $\langle e_1, \ldots, e_{\kappa} \rangle$. Therefore, up to a change of basis in $\langle e_1, \ldots, e_{\kappa} \rangle$, we may assume that $n^2 = e_1$. This is conclusion (b)(ii).

Suppose that $A \cong C_4 \times C_2^{\ell}$, for some $\ell \in \mathbb{N}$. Now $D := \langle A, g \rangle$ is generalised dihedral isomorphic to $D_4 \times C_2^{\ell}$. This is conclusion (c).

For the rest of the proof, we may assume that A is neither elementary abelian nor isomorphic to $A \cong C_4 \times C_2^{\ell}$, for some $\ell \in \mathbb{N}$. By Lemma 2.7, there exists $T \subseteq A$ with $|T| \ge 2$ such that $\operatorname{Cay}(A, T)$ is an ORR and such that the induced subgraph of $\operatorname{Cay}(A, T)$ on T is weakly connected. As G/A is elementary abelian of order 4 and G is not generalised dihedral, by Lemma 2.8 applied to the normal subgroup A, there exist $a_1, a_2 \in G \setminus N$ with $G = \langle a_1, a_2, N \rangle$, $o(a_1) > 2$ and $o(a_2) > 2$. Since $G \setminus A = nA \cup gA \cup ngA$ and every element of gA is an involution, the set $\{a_1, a_2\}$ consists of one element of nA and one of ngA. Since G/A is elementary abelian and A is abelian, a_1^2 and a_2^2 are in A and both centralise A. Now $a_1a_2 \in gA$ and no element of gA centralises A, because A does not have exponent 2. Thus, the second hypothesis of Lemma 2.8 holds, and we deduce that G admits an ORR; that is, conclusion (a) holds.

We break the rest of this section into subsections, each dealing with one of the three families of groups described in this proposition that are not generalised dihedral, but have not yet been shown to admit ORRs.

4.1. The groups of Proposition 4.1(b)(i). Let G be a group that arises in Proposition 4.1(b)(i). Then we have $G = V \rtimes \langle x \rangle$, where $V = \langle v_1, w_1, \ldots, v_\ell, w_\ell, e_1, \ldots, e_\kappa \rangle$ is an elementary abelian 2-group, $x^2 = 1$, $e_i^x = e_i$ for every $i \in \{1, \ldots, \kappa\}$, and $v_i^x = w_i$ for every $i \in \{1, \ldots, \ell\}$.

We begin by observing that, if $\ell \leq 1$, then G is a generalised dihedral group.

Lemma 4.2. Let G be a group that arises in Proposition 4.1(b)(i). If $\ell \leq 1$, then G is generalised dihedral.

Proof. If $\ell = 0$, then G is elementary abelian, which is a special form of generalised dihedral. If $\ell = 1$, then $V' = \langle v_1 x, e_1, \ldots, e_{\kappa} \rangle$ is an abelian group having index 2 in G (since $(v_1 x)^2 = v_1 w_1$). Also, the action of w_1 by conjugation inverts every element of V'. Thus, G is a generalised dihedral group over V'.

We now show that there is an ORR for any remaining sufficiently large group of this type.

Lemma 4.3. Let G be a group that arises in Proposition 4.1(b)(i). If $\ell \geq 2$ and $2\ell + \kappa \geq 8$, then G has an ORR.

Proof. We will use two different generating sets, according to whether or not $\kappa \geq 2$, or $\kappa \in \{0, 1\}$.

If $\kappa \geq 2$, then let

$$V_1 := \langle v_2, \dots, v_\ell, w_2, \dots, w_\ell, e_1, \dots, e_\kappa \rangle$$

and let T_1 be the Imrich generating set for V_1 with respect to this generating tuple (this exists because the rank of V_1 is $2(\ell-1)+\kappa \ge 6$). If $\kappa \in \{0,1\}$, then $2\ell+\kappa \ge 8$ implies $\ell \geq 4$. In this case, let

 $V_2 := \langle v_2, v_4 w_4, v_3, v_2 w_2, v_4, v_2 w_3, v_5, \dots, v_{\ell}, e_1, \dots, e_{\kappa} \rangle$

and let T_2 be the Imrich generating set for V_2 with respect to this generating tuple (this exists because the rank of V_2 is at least 6 since the first 6 elements are always in the tuple).

Let

$$S := [(V_i \setminus T_i) \setminus \{1\}] v_1 x \cup \{v_2 x\},$$

where i = 1 if $\kappa \ge 2$ and i = 2 if $\kappa \in \{0, 1\}$.

Observe that the product of any two elements of S has the form $vv_1xv'v_1x = vv_1(v')^x w_1 \in V_1v_1 w_1$ (where $v, v' \in V_i$), or $vv_1xv_2x = vv_1w_2 \in V_1v_1$ (where $v \in V_i$), or $v_2xvv_1x = v_2v^xw_1 \in V_1w_1$ (where $v \in V_i$), or $(v_2x)^2 = v_2w_2$. Since none of these can be 1, S is the connection set for an oriented digraph.

CLAIM: $v_2 x$ is the only vertex of S for which all of its outneighbours have no other inneighbour from S.

Proof of the Claim: Observe that the outneighbours of v_2x are all in $V_1v_1 \cup V_1$, while the outneighbours of any vertex of V_iv_1x are in $V_1v_1w_1 \cup V_1w_1$. Thus every outneighbour of v_2x has a unique inneighbour from S. It remains to prove the uniqueness.

Let $vv_1x \in S$ be arbitrary, so that $v \in V_i$ but $v \notin T_i \cup \{1\}$. If i = 1, then let $f_1 := e_1$ and $f_2 := e_2$; if i = 2, then let $f_1 := v_2w_2$ and $f_2 := v_3w_3$. We claim that $v \notin (T_i \cup \{1\})f_1 \cap (T_i \cup \{1\})f_2$. Otherwise, $v = tf_1 = t'f_2$ for some $t, t' \in T_i \cup \{1\}$ implies $tt' = f_1f_2$. Due to the structure of T_2 , this has no solutions if i = 2. When i = 1, the only solutions are $\{t, t'\} = \{1, e_1e_2\}$ or $\{t, t'\} = \{e_1, e_2\}$. Thus $v \in \langle e_1, e_2 \rangle \subset T_1 \cup \{1\}$, a contradiction that completes the proof of the claim.

Let $j \in \{1, 2\}$ be such that $v \notin (T_i \cup \{1\})f_j$. Then $vf_j \notin T_i \cup \{1\}$, so $vf_jv_1x \in S$ and $(vf_jv_1x)^2 = v_1w_1vv^x$ is an outneighbour of vf_jv_1x . It is also an outneighbour of vv_1x since $(vv_1x)^2 = v_1w_1vv^x$.

Let $\varphi \in \operatorname{Aut}(\operatorname{Cay}(G,S))_1$. From the previous claim, φ fixes v_2x . This implies that $V_iv_1x \cap S$ is fixed setwise by φ . Now we can apply Corollary 3.2 to see that φ fixes v_1x . This also implies that $(T_iv_1x)^{\varphi} = T_iv_1x$, and our choice of T_i as the generating set for a GRR on V_i implies that V_iv_1x is fixed pointwise by φ , and therefore so is V_i . Thus φ fixes pointwise the generating set $\{v_2x, v_1x\} \cup V_i$ of G. By Lemma 2.6, $\operatorname{Cay}(G, S)$ is an ORR.

This shows that every group that arises in Proposition 4.1(b)(i) with $\ell \geq 2$ that has order at least 2⁹ has an ORR.

4.2. The groups of Proposition 4.1(b)(ii). Let G be a group that arises in Proposition 4.1(b)(ii). Then we have $G = \langle V, x \rangle$, where $V = \langle v_1, w_1, \ldots, v_\ell, w_\ell, e_1, \ldots, e_\kappa \rangle$ is an elementary abelian 2-group, $x^2 = e_1$, $e_i^x = e_i$ for every $i \in \{1, \ldots, \kappa\}$, and $v_i^x = w_i$ for every $i \in \{1, \ldots, \ell\}$.

Lemma 4.4. Let G be a group that arises in Proposition 4.1(b)(ii). If $2\ell + \kappa \ge 7$, then G has an ORR.

Proof. Let $(v_1, \ldots, v_\ell, w_1, \ldots, w_\ell, e_2, \ldots, e_\kappa)$ be a generating tuple for

 $V' := \langle v_1, \ldots, v_\ell, w_1, \ldots, w_\ell, e_2, \ldots e_\kappa \rangle < V,$

and let T be the Imrich generating set for V' with respect to this generating tuple (this exists because V' has rank $2\ell + \kappa - 1 \ge 6$). Let

$$S := Tx \cup \{x\}.$$

As the product of any two elements of S is not 1 (since it will be in $V'e_1$), Cay(G, S) is an oriented Cayley digraph.

By Lemma 3.1, x is the unique vertex of S that has at least three mutual inneighbours with every other vertex of S. Since any automorphism $\varphi \in \operatorname{Aut}(\operatorname{Cay}(G,S))_1$ fixes S setwise, we must have $x^{\varphi} = x$. Furthermore $(Tx)^{\varphi} = Tx$ and since T is the generating set for a GRR on V', any automorphism that fixes x and fixes Tx setwise must actually fix $\langle T \rangle x = V'x$ pointwise. Since φ fixes every point of $\{x, V'x\}$ and $\langle x, V'x \rangle = G$, by Lemma 2.6 we have $\varphi = 1$ and this completes the proof.

This shows that every group that arises in Proposition 4.1(b)(ii) that has order at least 2^8 has an ORR.

4.3. The groups of Proposition 4.1(c).

Notation 4.5. Let G be a group that arises in Proposition 4.1(c). Then G has a maximal subgroup D isomorphic to $D_4 \times C_2^{\ell}$ for some $\ell \in \mathbb{N}$. Studying the initial description of the groups in Proposition 4.1, we see that we may assume that $G = \langle A, g, n \rangle$, where $A \cong C_4 \times C_2^k$ is generated by a_1, \ldots, a_{k+1} with $o(a_1) = 4$ and $o(a_i) = 2$ for $i \in \{2, \ldots, k+1\}$, and g acts by conjugation on A inverting every $a_i, g^2 = 1, n^g = n^{-1}, n$ normalises A and $n^2 \in A$. Furthermore, if n centralises A, then $N := \langle A, n \rangle$ is abelian and $G = \langle N, g \rangle$ is generalised dihedral. Thus, if we assume that G is not generalised dihedral, n does not centralise A.

Lemma 4.6. Let G be as in Notation 4.5. If G is not generalised dihedral, then there exists $a \in A$ such that o(a) > 2 and a is not centralised by n.

Proof. Assume that G is not generalised dihedral. Therefore n must not centralise A. Also, there is some $a_1 \in A$ with $o(a_1) > 2$. If a_1 is not centralised by n then taking $a := a_1$ we are done, so we may assume that n centralises every element of A of order greater than 2. Since n does not centralise A, there is some $b \in A$ such that b is not centralised by n. Thus o(b) = 2. Let $c := b^n$, so $c \neq b$ has order 2. Now, a_1b has order $o(a_1) > 2$, and $(a_1b)^n = a_1b^n = a_1c \neq a_1b$, so taking $a := a_1b$ satisfies our conclusion.

Lemma 4.7. Let G be as in Notation 4.5. If o(n) = 2 and G is not generalised dihedral, then there is some element $a \in A$ such that o(a) > 2 and a is not inverted by n.

Proof. Assume o(n) = 2. Let a_1 be an element of A with $o(a_1) > 2$. If a_1 is not inverted by n then taking $a := a_1$ we are done, so we may assume that n inverts every element of A of order greater than 2. Consider a_1b for any b of order 2 in A. As $o(a_1b) = o(a_1) > 2$, we have $a_1^{-1}b = (a_1b)^{-1} = (a_1b)^n = a_1^nb^n = a_1^{-1}b^n$, and $b^n = b$. Thus n centralises every element of order 2, and this implies that every element in A is inverted by n. Hence ng centralises A and $\langle A, ng \rangle$ is abelian. Since the action of g by conjugation inverts ng (because $(ng)^g = n^g g = n^{-1}g = ng$) and each element of A, we deduce that G is a generalised dihedral group over $\langle A, ng \rangle$, a contradiction.

We now have the key points for dealing with these groups. Our final lemma in this section shows that all sufficiently large groups that arise in Proposition 4.1(c) are either generalised dihedral or have ORRs.

Lemma 4.8. Let G be as in Notation 4.5, with $k \ge 6$. Then either G is generalised dihedral, or G admits an ORR.

Proof. Suppose that G is not generalised dihedral; let $B := \langle a_2, \ldots, a_{k+1} \rangle$, and let T be the Imrich generating set for B with respect to some generating tuple. By Lemma 4.6, there exists $a \in A$ with o(a) > 2 and $a^n \neq a$. Let

$$S := \begin{cases} [(Ba \setminus Ta) \setminus \{a\}] \cup \{n^{-1}, an^{-1}g, an\} & \text{when } a^n \neq n^{-2}a^{-1} \text{ and } o(n) \neq 2; \\ [(Ba \setminus Ta) \setminus \{a\}] \cup \{an^{-1}g, an\} & \text{when } a^n \neq a^{-1} \text{ and } o(n) = 2; \\ [(Ba \setminus Ta) \setminus \{a\}] \cup \{an^{-1}g, n\} & \text{when } (a')^n = n^{-2}(a')^{-1} \text{ for every } a' \in A \text{ with } o(a') = 4. \end{cases}$$

Observe that the three cases above do not cover all possibilies: in the remaining case every element a' of order 4 in A is either centralised by n, or has $(a')^n = n^{-2}(a')^{-1}$, and they are not all in the second category. In the next paragraph, we study this latter possibility.

So there is $a \in A$ as in Lemma 4.6, and $a^n = n^{-2}a^{-1}$. Notice that $a \neq a^n = n^{-2}a^{-1}$ implies that $n^2 \neq a^2$ in this case.

CLAIM: There exist three elements $b_1, b_2, b_3 \in B$ such that ab_1, ab_1b_2, ab_1b_3 are distinct and centralised by n.

Proof of claim: Since not all elements of order 4 of A are in the second category, there exists an element of order 4 centralised by n. As $A = \langle a \rangle \times B$, we may assume that this element is of the form ab_1 , for some $b_1 \in B \setminus \{1\}$. If $|\mathbf{C}_B(n)| > 2$, then we may choose two distinct elements $b_2, b_3 \in \mathbf{C}_B(n) \setminus \{1\}$, and now n centralises the three distinct elements ab_1, ab_1b_2, ab_1b_3 . Assume $|\mathbf{C}_B(n)| \leq 2$.

Now, $B_0 := \langle a^2, B \rangle$ is characteristic in A because it consists of all elements of order 2 of A, and hence B_0 is *n*-invariant. Clearly, $|\mathbf{C}_{B_0}(n)| \leq 2|\mathbf{C}_B(n)| \leq 4$. As $n^2 \in A$ centralises A, the mapping $f : B_0 \to \mathbf{C}_{B_0}(n)$ defined by $x \mapsto xx^n$ is a homomorphism with kernel $\mathbf{C}_{B_0}(n)$. Therefore, from the first homomorphism theorem, $|B_0/\mathbf{C}_{B_0}(n)| \leq |\mathbf{C}_{B_0}(n)|$ and $|B_0| \leq |\mathbf{C}_{B_0}(n)|^2 = 16$. However, $|B_0| = 2^{k+1} \geq 2^{6+1}$, a contradiction.

Let $b_1, b_2, b_3 \in B$ be as in the previous claim and let

$$S := [(Ba \setminus Ta) \setminus \{a\}] \cup \{an^{-1}g, ab_1n, ab_1b_2n, ab_1b_3n\}.$$

We claim that, in all four cases, $\Gamma := \operatorname{Cay}(G, S)$ is an ORR for G. First we show that S is asymmetric and all of its elements have order greater than 2. Certainly in every case the elements of $S \cap A$ have order 4 and do not contain any inverse-closed pair. Any products that are not in A will clearly not be 1. The remaining pairwise products are $n^{-1}an = a^n \neq 1$; $ann^{-1} = a \neq 1$; n^{-2} or n^2 (notice that, if o(n) = 2, then Lemma 4.7 implies that $n \notin S$); $(an^{-1}g)^2 = a(a^n)^{-1} \neq 1$ by our choice of a; $(an)^2 = aa^n n^2 \neq 1$ by our choice of a in the cases where $an \in S$; and for every $b_i, b_j \in \{1, b_1, b_2, b_3\}$, we have $ab_1b_inab_1b_jn = a^2b_ib_jn^2 \neq 1$ since $n^2 \neq a^2b_ib_j$. Thus Γ is an oriented digraph.

Let $\varphi \in \operatorname{Aut}(\Gamma)_1$. By Proposition 3.3, $a^{\varphi} = a$ and φ fixes every element of B pointwise. Therefore φ fixes every element of $\langle a, B \rangle = A$ pointwise.

Now if S has the first of the four possible forms, then $an^{-1}g$ is the only vertex of $\{an^{-1}g, an, n^{-1}\}$ that has only one outneighbour (via these three elements) in

A, so $(an^{-1}g)^{\varphi} = an^{-1}g$. The outneighbours of n^{-1} via $\{n^{-1}, an\}$ are n^{-2} and a, while the outneighbours of an are a^n and aa^nn^2 . We know that $a^n \neq a$, and $a^n = n^{-2}$ implies $n^{-2} = a$, but then $a^n = a$, a contradiction. Thus an is unique in having a^n as an outneighbour, so $(an)^{\varphi} = an$. This completes the proof since φ fixes every element of the generating set $\{an^{-1}g, an\} \cup A$ of G.

If S has the second of the four possible forms, then $n^2 = 1$ and the outneighbours of $an^{-1}g$ via $an^{-1}g$ and an are $a(a^n)^{-1}$ and $anan^{-1}g = aa^ng$, while the outneighbours of an are aa^n and $an^{-1}gan = a(a^n)^{-1}g$. Thus an is the only one of these vertices that has $a(a^n) \in A$ as an outneighbour, so $(an)^{\varphi} = an$ and $(an^{-1}g)^{\varphi} = an^{-1}g$. This completes the proof since φ fixes every element of the generating set $\{an, an^{-1}g\} \cup A$ of G.

If S has the third of the four possible forms, then $(a')^n = n^{-2}(a')^{-1}$ for every a' of order 4 in A. The pairwise products of $\{an^{-1}g,n\}$ are $a(a^n)^{-1} = a(n^{-2}a^{-1})^{-1} = n^2a^2$, $an^{-2}g$, n^2 , and a^ng . Since $a^2 \neq 1$, $an^{-1}g$ and n do not have the same outneighbours, so $n^{\varphi} = n$ and $(an^{-1}g)^{\varphi} = an^{-1}g$. This completes the proof since φ fixes every element of the generating set $\{n, an^{-1}g\} \cup A$ of G.

Finally, if S has the fourth of the four possible forms, then $an^{-1}g$ is the only vertex of $S \setminus Ba$ that has only one outneighbour in A, so $(an^{-1}g)^{\varphi} = an^{-1}g$. Of the remaining three vertices of $S \setminus Ba$, ab_1n is the only one that has both $a^2b_2n^2$ and $a^2b_3n^2$ as outneighbours, so $(ab_1n)^{\varphi} = ab_1n$. This completes the proof since φ fixes every element of the generating set $\{an^{-1}g, ab_1n\} \cup A$ of G.

This shows that every group that arises in Proposition 4.1(c) that has order at least 2^{10} has an ORR.

5. The groups in Theorem 2.3 (III)

We discuss in some detail the work in [7] and establish some notation that we use for the rest of this section. Let G, N, g and n_0 be as in Theorem 2.3 (iii). Thus |G : N| = 2, $g^2 = 1$, $N = H \cup n_0 H$ where $H := \{n \in N \mid n^g = n^{-1}\}$ (note that H is a set and not necessarily a group), |H| = |N|/2 and N has no automorphisms inverting more than half of its elements (according to Theorem 2.3, N is a group described in the Hegarty and MacHale paper, and hence N admits an automorphism inverting precisely half of its elements, namely conjugation via g, and no automorphisms inverting more than half of its elements). We observe that our assumption about the existence and properties of n_0 is not amongst the assumptions of Hegarty and MacHale, and we will be able to use this assumption to eliminate some of the groups in their classification.

The classification of Hegarty and MacHale is very satisfactory, but not very easy to use in our application. These groups fall into ten isoclinism classes and, for each class, the authors give a very explicit description of a stem group in the class. (We denote by X' the derived subgroup and by $\mathbf{Z}(X)$ the centre of the group X.) We recall that the groups X and Y are isoclinic if there exist two group automorphisms $\varphi: X' \to Y'$ and $\psi: X/\mathbf{Z}(X) \to Y/\mathbf{Z}(Y)$ with

$$[x_1 \mathbf{Z}(X), x_2 \mathbf{Z}(X)]^{\varphi} = [(x_1 \mathbf{Z}(X))^{\psi}, (x_2 \mathbf{Z}(X))^{\psi}], \quad \text{for every } x_1, x_2 \in X.$$

For instance, the dihedral group of order 8 and the quaternion group of order 8 are isoclinic. Being isoclinic is an equivalence relation coarser than the equivalence relation determined by the notion of group isomorphism. This means that finite groups are subdivided into isoclinism classes. Two groups in the same isoclinism

class may not have the same order, for instance, X and $X \times Z$ are isoclinic for every finite group X and for every abelian group Z. It is well-known and also easy to prove that, for every group X, in the isoclinism class of X there exists a group Y with $\mathbf{Z}(Y) \leq Y'$. A group satisfying $\mathbf{Z}(Y) \leq Y'$ is said to be *stem* group. The definition of isoclinism yields that the stem groups are precisely the groups of smallest possible order within their isoclinism class. It is quite unfortunate that two stem groups may be isoclinic but not necessarily isomorphic: consider again the example of the dihedral and the quaternion group of order 8.

We can now explain in some detail the work of Hegarty and MacHale. (Just for this paragraph, we say that X is half-inverting, if X has an automorphism inverting half of its elements and no automorphisms inverting more than half of its elements.) As a by-product of their main theorem, they prove that if X and Y are isoclinic and X is half-inverting, then Y is half-inverting. Moreover, Hegarty and MacHale prove that half-inverting groups fall into 10 distinct isoclinism classes. To describe these 10 isoclinism classes, Hegarty and MacHale exhibit 10 (non-isoclinic) stem groups: two of order 32, six of order 64 and two of order 128. We emphasise once again that this does not mean that there are only two stem groups of order 128 that are half-inverting. (A computation with the computer algebra system magma [3] shows that there are 5, 55 and 251 half-inverting stem groups of order 32, 64 and 128, respectively, up to isomorphism.)

Now that we have explained some details required for understanding the classification of Hegarty and MacHale, we can proceed to deal with the groups that arise in our context.

Proposition 5.1. Let G be a group that arises in Theorem 2.3 (iii). Then G admits an ORR except possibly if N has an abelian subgroup A with |N : A| = 4, and A is either isomorphic to $C_4 \times C_2^{\ell}$ for some $\ell \leq 5$, or A is elementary abelian of rank at most 7.

Proof. By Theorem 2.3 (iii), G is a 2-group with a normal subgroup N, |G:N| = 2, and elements $g \in G \setminus N$ and $n_0 \in N$ with $g^2 = 1$; the action of g by conjugation on N inverts precisely half of the elements of N; and $N = H \cup n_0 H$, where $H := \{n \in N \mid n^g = n^{-1}\}$, and |H| = |N|/2.

From [7, Lemma 1] and its proof (see also the last paragraph of [7, page 132]), N has an abelian subgroup A with |N : A| = 4 and with $a^g = a^{-1}$ for each $a \in A$. Some more information on the interaction between A and g and N is available by reading the proof of Lemma 1 in [7] (again, see also the last paragraph of page 132 in [7]). Let $1, x_2, x_3, x_4$ be left coset representatives for A in N; thus $N = A \cup x_2 A \cup x_3 A \cup x_4 A$. According to Hegarty and MacHale, there are only two cases:

CASE I: g inverts some element in each coset of A in N;

CASE II: g inverts some element in the cosets A, x_3A and x_4A , and g inverts no elements in the coset x_2A .

For the benefit of the reader, we have used the same subdivision into cases here, as the subdivision used in [7]. Replacing x_2, x_3, x_4 by suitable coset representatives we may assume that g inverts x_3 and x_4 and that g also inverts x_2 in CASE I. Let $x \in \{x_2, x_3, x_4\}$ in CASE I and $x \in \{x_3, x_4\}$ in CASE II. Let $y \in H \cap xA$. Then, y = xa for some $a \in A$ and $a^{-1}x^{-1} = (xa)^{-1} = y^{-1} = y^g = (xa)^g = x^g a^g = x^{-1}a^{-1}$.

This shows that xa = ax. Therefore

$$H = \begin{cases} A \cup x_2 \mathbf{C}_A(x_2) \cup x_3 \mathbf{C}_A(x_3) \cup x_4 \mathbf{C}_A(x_4) & \text{in CASE I,} \\ A \cup x_3 \mathbf{C}_A(x_3) \cup x_4 \mathbf{C}_A(x_4) & \text{in CASE II.} \end{cases}$$

Theorem 2.3 (iii) gives us information about the element n_0 in addition to the structure that was studied by Hegarty and MacHale, and we have not used this in our analysis yet. We do so now. Recall that $N = H \cup n_0 H$, and $H \cap n_0 H = \emptyset$ because H has cardinality |N|/2.

Assume that CASE I holds. Now $n_0A \subseteq n_0H$ and hence n_0H contains a whole left coset of A in N. Since H contains elements from each left coset of A in N, we get $n_0H \cap H \neq \emptyset$, a contradiction. Therefore 2-groups in CASE I do not arise in Theorem 2.3 (iii).

For the rest of the proof, we assume that CASE II holds. From [7, line 11 from the bottom of page 134], we have $A \triangleleft N$ and N/A is elementary abelian of order 4; therefore, $A \triangleleft G$ because g also normalises A.

 \mathbf{As}

$$2|A| = |N|/2 = |H| = |A| + |\mathbf{C}_A(x_3)| + |\mathbf{C}_A(x_4)|,$$

we deduce $|A : \mathbf{C}_A(x_3)| = |A : \mathbf{C}_A(x_4)| = 2$. Clearly, $n_0 \in x_2A$, otherwise, arguing as in the previous paragraph, we get $n_0H \cap H \neq \emptyset$. Replacing x_2 if necessary, we may assume that $n_0 = x_2$. Thus

$$m_0H = x_2A \cup x_2x_3\mathbf{C}_A(x_3) \cup x_2x_4\mathbf{C}_A(x_4).$$

It is important to observe that since A is normal in N and N/A is elementary abelian, we have $x_2x_3A = (x_2A)(x_3A) = x_4A$ and $x_2x_4A = (x_2A)(x_4A) = x_3A$; therefore $x_2x_3\mathbf{C}_A(x_3)$ is contained in the coset x_4A and $x_2x_4\mathbf{C}_A(x_4)$ is contained in the coset x_3A . Now, the condition $N = H \cup n_0H$ yields $x_3A = x_3\mathbf{C}_A(x_3) \cup x_2x_4\mathbf{C}_A(x_4)$. This implies that A is the union of a coset of $\mathbf{C}_A(x_4)$ with a coset of $\mathbf{C}_A(x_3)$, and since each of these subgroups of A has cardinality |A|/2, this can happen only when $\mathbf{C}_A(x_3) = \mathbf{C}_A(x_4)$. Since $N = \langle A, x_3, x_4 \rangle$, we have

$$\mathbf{Z}(N) = \mathbf{C}_A(x_3) \cap \mathbf{C}_A(x_4) = \mathbf{C}_A(x_4) < A;$$

hence, $|N : \mathbf{Z}(N)| = 8$. Finally we observe that G/A is elementary abelian of order 8 because g acts by conjugation inverting x_3 and x_4 and hence fixes the cosets x_3A and x_4A .

Suppose now that A is neither elementary abelian nor isomorphic to $C_4 \times C_2^{\ell}$, for some $\ell \in \mathbb{N}$. From Lemma 2.7, there exists a subset T of A of cardinality at least 2 with $\operatorname{Cay}(A, T)$ an ORR and such that the subgraph induced by $\operatorname{Cay}(A, T)$ on T is weakly connected.

We claim that $A = \mathbf{C}_G(A)$, that is, no element of $G \setminus A$ centralises A. Observe that

 $G \setminus A = x_3 A \cup x_4 A \cup x_3 x_4 A \cup g x_3 A \cup g x_4 A \cup g x_3 x_4 A.$

From [7, Lemma 1], N has no abelian subgroup of index less then four. Therefore no element in the cosets x_3A , x_4A and x_3x_4A centralises A and, moreover,

$$\mathbf{Z}(N) = \mathbf{C}_A(x_3) = \mathbf{C}_A(x_4) = \mathbf{C}_A(x_3x_4).$$

Observe now that $\mathbf{Z}(N)$ is not elementary abelian because A has no elementary abelian subgroup of index 2. Since g acts by conjugation inverting each element of $\mathbf{Z}(N)$, we deduce that no element in the cosets gA, gx_3A , gx_4A and gx_3x_4A centralises $\mathbf{Z}(N) \leq A$. Thus our claim is proved.

By Lemma 2.8, since |G:A| = 8, there are generators a_1, a_2, a_3 for G modulo A none of which is an involution. Furthermore, since G/A is elementary abelian and A is abelian, a_i^2 centralises A for every $i \in \{1, 2, 3\}$. From the previous paragraph, $a_i a_j$ does not centralise A, for every $i, j \in \{1, 2, 3\}$ with $i \neq j$. In particular, the second hypothesis of Lemma 2.8 holds, and we deduce that G admits an ORR.

Suppose now that A is either elementary abelian or isomorphic to $C_4 \times C_2^{\ell}$ for some $\ell \in \mathbb{N}$. We deal with the two possible structures for A individually.

Suppose first that A is isomorphic to $C_4 \times C_2^{\ell}$. By ignoring the exceptions from our statement, we may assume that $\ell \geq 6$. Let $a_0 \in A \setminus \mathbf{Z}(N)$ with $o(a_0) = 4$ (observe that this is possible because $\mathbf{Z}(N) < N$ cannot contain all the elements of order 4 of A). Let B be an elementary abelian subgroup of rank ℓ in A that does not include the non-identity square element of A, and let T be the Imrich generating set for B with respect to some generating tuple. Let

 $X := \{gx_3a_0, gx_3a_0b, gx_3a_0b', gx_3a_0b''\},\$

where b, b', and b'' are chosen from B so that $\langle b, b', b'' \rangle$ has order 8 and none of the pairwise products from $\{1, b, b', b''\}$ is $(a_0^{-1})^{x_3}a_0$: the rank of B is easily large enough that such choices are possible. Let

 $Y := \{gx_4a_0, gx_4a_0c, gx_4a_0c'\},\$

where c and c' are chosen from B so that $\langle c, c' \rangle$ has order 4 and none of the pairwise products from $\{1, c, c'\}$ is $(a_0^{-1})^{x_4}a_0$ (again, this is possible because B has rank at least 6). Let

 $S := [(Ba_0 \setminus Ta_0) \setminus \{a_0\}] \cup X \cup Y \cup \{gx_3x_4\}.$

The elements of $S \cap Ba_0$ have order 4 and none is the inverse of another since $a_0^2 \notin B$. The pairwise products of the elements of S that are not in A cannot yield the identity. Furthermore, observe that, for every $z, z' \in \{1, b, b', b''\}$, we have

$$(gx_3a_0z)(gx_3a_0z') = (a_0^{-1})^{x_3}a_0zz' \neq 1$$

by our choice of b, b' and b'' and the fact that $a_0 \notin \mathbf{Z}(N) = \mathbf{C}_A(x_3)$. Similarly, for every $z, z' \in \{1, c, c'\}$, we have

$$(gx_4a_0z)(gx_4a_0z') = (a_0^{-1})^{x_4}a_0zz' \neq 1$$

by our choice of c and c' and the fact that $a_0 \notin \mathbf{Z}(N) = \mathbf{C}_A(x_4)$. Finally,

$$(gx_3x_4)^2 = gx_3x_4gx_3x_4 = x_3^g x_4^g x_3x_4 = x_3^{-1} x_4^{-1} x_3x_4 \neq 1,$$

where in the third equality we have used $x_3, x_4 \in H = \{n \in N \mid n^g = n^{-1}\}$, and in the last inequality we used the fact that x_3 and x_4 do not commute because $x_3x_4 \notin H$ and

$$x_4^{-1}x_3^{-1} = (x_3x_4)^{-1} \neq (x_3x_4)^g = x_3^g x_4^g = x_3^{-1}x_4^{-1}.$$

This proves that $S \cap S^{-1} = \emptyset$ and $\operatorname{Cay}(G, S)$ is an oriented graph.

By Proposition 3.3, any automorphism $\varphi \in \operatorname{Aut}(\operatorname{Cay}(G,S))_1$ fixes B and a_0 pointwise, so fixes $\langle a_0, B \rangle = A$ pointwise.

Observe that, for any of the elements of X, its outneighbours via elements of $X \cup Y \cup \{gx_3x_4\}$ have four outneighbours in A and four not in A; the elements of Y each have three outneighbours via $X \cup Y \cup \{gx_3x_4\}$ in A and five not in A; and gx_3x_4 has only one outneighbour via $X \cup Y \cup \{gx_3x_4\}$ that is in A. Thus $X^{\varphi} = X, Y^{\varphi} = Y$, and $(gx_3x_4)^{\varphi} = gx_3x_4$, for every $\varphi \in \text{Aut}(\text{Cay}(G, S))_1$. Furthermore, in X, since $\langle b, b', b'' \rangle$ has order 8, gx_3a_0 is the only vertex that does not have either $(a_0^{-1})^{x_3}a_0bb'$

or $(a_0^{-1})_3^x a_0 bb''$ as an outneighbour. Therefore gx_3a_0 is fixed by $\operatorname{Aut}(\operatorname{Cay}(G,S))_1$. Similarly, in Y, since $\langle c, c' \rangle$ has order 4, gx_4a_0 is the only vertex that does not have $(a_0^{-1})^{x_4}a_0cc'$ as an outneighbour. Therefore gx_4a_0 is fixed by $\operatorname{Aut}(\operatorname{Cay}(G,S))_1$. Thus $\operatorname{Aut}(\operatorname{Cay}(G,S))_1$ fixes the generating set $\{gx_3x_4, gx_3a_0, gx_4a_0\} \cup A$ for G pointwise, so by Lemma 2.6, $\operatorname{Aut}(\operatorname{Cay}(G,S))_1 = 1$ and $\operatorname{Cay}(G,S)$ is an ORR.

Suppose now that A is elementary abelian, that is, $A \cong C_2^{\ell}$ for some $\ell \in \mathbb{N}$. By ignoring the exceptions from our statement, we may assume that $\ell \geq 8$. As x_3 and x_4 do not commute, the commutator $d := x_4^{-1}x_3^{-1}x_4x_3 \neq 1$. Recall that $N/\mathbb{Z}(N)$ is elementary abelian and hence the commutator subgroup N' is contained in $\mathbb{Z}(N)$. Thus $d \in \mathbb{Z}(N)$. Let $d, b_1, \ldots, b_{\ell-2}$ be an irredundant generating set for $\mathbb{Z}(N) \cong C_2^{\ell-1}$, and let $B := \langle b_1, \ldots, b_{\ell-2} \rangle$. So B is elementary abelian of rank $\ell-2 \geq 6$. Let T be the Imrich generating set for B with respect to some generating tuple.

Let $a_0 \in A \setminus \mathbf{Z}(N)$ and observe that $\langle \mathbf{Z}(N), g, x_3 a_0, x_4 a_0 \rangle$ is a subgroup of G having index 2. Since G is not generalised dihedral, there exists $v \in G \setminus \langle \mathbf{Z}(N), g, x_3 a_0, x_4 a_0 \rangle$ with o(v) > 2. Let

 $X := \{gx_3a_0, gx_3a_0b, gx_3a_0b', gx_3a_0b''\},\$

where b, b', and b'' are chosen from B so that $\langle b, b', b'' \rangle$ has order 8 and none of the pairwise products from $\{1, b, b', b''\}$ is $(a_0^{-1})^{x_3}a_0$: the rank of B is easily large enough that such choices are possible. Let

$$Y := \{gx_4a_0, gx_4a_0c, gx_4a_0c'\},\$$

where c and c' are chosen from B so that $\langle c, c' \rangle$ has order 4 and none of the pairwise products from $\{1, c, c'\}$ is $(a_0^{-1})^{x_4}a_0$ (again, this is possible because B has rank at least 6). Let

$$S := gx_3x_4[(B \setminus T) \setminus \{1\}] \cup X \cup Y \cup \{v\} \cup \{gx_3x_4d\}.$$

For any $z, z' \in B$, we have

$$(gx_3x_4z)(gx_3x_4z') = x_3^{-1}x_4^{-1}x_3x_4zz' = dzz' \neq 1,$$

because $d \notin B$. The argument that no two elements of X or Y are inverses of one another is exactly as in the previous case. Notice also that v is not in $\langle S \setminus \{v\} \rangle$, so o(v) > 2 implies that $v^{-1} \notin S$. Now, observe that, for any $b \in B \setminus \{1\}$, we have

$$gx_3x_4bgx_3x_4d = d^{x_4}bd = dbd = b \neq 1$$

since $d \in \mathbf{Z}(N)$ which is elementary abelian. Finally, $(gx_3x_4d)^2 = d \neq 1$. Thus $S \cap S^{-1} = \emptyset$ and $\operatorname{Cay}(G, S)$ is an oriented Cayley graph.

Observe that $(gx_3x_4)^2 = d$ implies $o(gx_3x_4) = 4$, and since $\mathbb{Z}(N)$ is elementary abelian we have $B < \mathbb{Z}(N) = \mathbb{Z}(G)$, so gx_3x_4 centralises B. Thus Proposition 3.3 applies to show that $\operatorname{Aut}(\operatorname{Cay}(G,S))_1$ fixes B and gx_3x_4 pointwise. Thus the cosets of B are blocks of imprimitivity for $\operatorname{Aut}(\operatorname{Cay}(G,S))_1$. Since the four vertices of Xlie in one coset of B, the three vertices of Y lie in a different coset, and the vertices v and gx_3x_4d each lie in a different coset of B, we have $X^{\varphi} = X$ and $Y^{\varphi} = Y$, for every $\varphi \in \operatorname{Aut}(\operatorname{Cay}(G,S))_1$. The same argument as in the previous case again shows that gx_3a_0 and gx_4a_0 are fixed by $\operatorname{Aut}(\operatorname{Cay}(G,S))_1$.

Let $\varphi \in \operatorname{Aut}(\operatorname{Cay}(G,S))_1$. Since $d \in \mathbb{Z}(N)$ has order 2, we see that gx_3x_4d has $2^{\ell} - 2\ell - 2$ outneighbours in B, but v has at most one outneighbour in B. Thus $v^{\varphi} = v$ and $(gx_3x_4d)^{\varphi} = gx_3x_4d$. We now see that φ fixes every point of the

generating set $\{gx_3a_0, gx_4a_0, v, gx_3x_4, gx_3x_4d\} \cup B$ for G. Since φ was arbitrary, Lemma 2.6 gives that Cay(G, S) is an ORR.

6. Proof of Theorem 1.2

We are now ready to prove our main theorem, Theorem 1.2.

Proof of Theorem 1.2. If |G| = 2, then $Cay(G, \emptyset)$ is an ORR for G. We henceforth assume that G is not generalised dihedral.

By Theorem 2.3, G admits an ORR unless G is as in Theorem 2.3 (ii), Theorem 2.3 (iii), or $G \cong Q_8, C_3^2$, or $C_3 \times C_2^3$. The final three possibilities are in our list of exceptions in the statement of Theorem 1.2. Suppose now that G is as in Theorem 2.3 (ii). By Proposition 4.1, either G admits an ORR, or one of the following possibilities holds:

(1) ℓ and κ are non-negative integers, V is an elementary abelian 2-group of rank $2\ell + \kappa$ with generating set $\{v_1, w_1, \ldots, v_\ell, w_\ell, e_1, \ldots, e_\kappa\}$, and $G = \langle V, x \rangle$, where $v_i^x = w_i$, $w_i^x = v_i$ for $i \in \{1, \ldots, \ell\}$ and $e_i^x = e_i$ for $i \in \{1, \ldots, \kappa\}$; and (i) $x^2 = 1$; or

(ii)
$$\kappa \ge 1$$
 and $x^2 = e_1$; or

(2) $\langle A, g \rangle$ is isomorphic to $D_4 \times C_2^{\ell}$ for some $\ell \in \mathbb{N}$.

If (1)(i) holds, then by Lemma 4.2 we have $\ell \geq 2$. Now by Lemma 4.3, G admits an ORR as long as $2\ell + \kappa \geq 8$; equivalently, as long as $|G| \geq 2^9$. If (1)(ii) holds, then by Lemma 4.4, G admits an ORR as long as $2\ell + \kappa \geq 7$; equivalently, as long as $|G| \geq 2^8$. If (2) holds, then by Lemma 4.8, G admits an ORR as long as $\ell \geq 6$; equivalently, as long as $|G| \geq 2^{10}$. On the other hand, if G is as in Theorem 2.3 (iii), then by Proposition 5.1, G admits an ORR when $|G| \geq 2^{11}$.

It remains to check 2-groups of order at most 2^9 that are not generalised dihedral and that satisfy Theorem 2.3 (ii) and the 2-groups of order at most 2^{10} that satisfy Proposition 5.1. These were all checked with the aid of magma, and the only groups that do not admit an ORR are those listed.

References

- BibliographyL. Babai, 'Finite digraphs with given regular automorphism groups', Periodica Mathematica Hungarica 11 (1980), 257–270.
- BibliographyL. Babai and W. Imrich, 'Tournaments with given regular group', Aequationes Mathematicae 19 (1979), 232–244.
- [3] BibliographyW. Bosma, J. Cannon and C. Playoust, 'The Magma algebra system. I. The user language', J. Symbolic Comput. 24 (1997), 235–265.
- [4] BibliographyM. Conder, T. Tucker, and M. Watkins, 'Graphical Frobenius Representations with even complements', https://www.math.auckland.ac.nz/ dleemans/SCDO2016/SCDO-TALKS/Monday/Tucker.pdf
- [5] BibliographyE. Dobson, P. Spiga, and G. Verret, 'Cayley graphs on abelian groups', Combinatorica 36 (2016), 371–393.
- [6] BibliographyC. D. Godsil, 'GRRs for nonsolvable groups', Algebraic Methods in Graph Theory, (Szeged, 1978), 221–239, Colloq. Math. Soc. János Bolyai 25, North-Holland, Amsterdam-New York, 1981.
- [7] BibliographyP. Hegarty and D. MacHale, 'Two-groups in which an automorphism inverts precisely half of the elements', Bull. London Math. Soc. 30 (1998), 129–135.
- [8] BibliographyD. Hetzel, 'Über reguläre graphische Darstellung von auflösbaren Gruppen', Technische Universität, Berlin, 1976.

- BibliographyW. Imrich, 'Graphs with transitive abelian automorphism group', Combinat. Theory (Proc. Collog., Balatonfűred, 1969), Budapest, 1970, 651–656.
- [10] BibliographyL. A. Nowitz and M. E. Watkins, 'Graphical regular representations of nonabelian groups. I', Canad. J. Math. 24 (1972), 993–1008.
- [11] BibliographyJ. Morris, P. Spiga and G. Verret, 'Automorphisms of Cayley graphs on generalised dicyclic groups', European Journal of Combinatorics 43 (2015), 68–81.
- [12] BibliographyJ. Morris and P. Spiga, 'Every Finite Non-Solvable Group admits an Oriented Regular Representation', Journal of Combinatorial Theory Series B, to appear.
- [13] BibliographyP. Spiga, 'Finite groups admitting an oriented regular representation', *Journal of Combinatorial Theory Series A*, to appear.
- [14] BibliographyP. Spiga, 'Cubic graphical regular representations of finite non-abelian simple groups', *submitted*.
- [15] BibliographyS. J. Xu, X. G. Fang, J. Wang and M. Xu, 'On cubic s-arc transitive Cayley graphs of finite simple groups', *European J. Combin.* 26 (2005), 133–143.

JOY MORRIS, DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF LETHBRIDGE, LETHBRIDGE, AB. T1K 3M4. CANADA

 $E\text{-}mail\ address: \texttt{joy.morris@uleth.ca}$

PABLO SPIGA, DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, UNIVERSITY OF MILANO-BICOCCA, VIA COZZI 55, 20125 MILANO, ITALY

E-mail address: pablo.spiga@unimib.it