Microstates of accelerating and supersymmetric

 AdS_4 black holes from the spindle index

Edoardo Colombo,^{1,2} Seyed Morteza Hosseini,³ Dario Martelli,^{1,2} Antonio Pittelli,^{1,2} and Alberto Zaffaroni^{4,5}

¹Dipartimento di Matematica, Università di Torino, Via Carlo Alberto 10, 10123 Torino, Italy

²INFN, Sezione di Torino, Via Pietro Giuria 1, 10125 Torino, Italy

³Department of Physics, Imperial College London, London, SW7 2AZ, UK

⁴Dipartimento di Fisica, Università di Milano-Bicocca, Piazza della Scienza 3, 20126 Milano, Italy

⁵INFN, Sezione di Milano-Bicocca, Piazza della Scienza 3, 20126 Milano, Italy

We provide a first principles derivation of the microscopic entropy of a very general class of supersymmetric, rotating and accelerating black holes in AdS_4 . This is achieved by analysing the large-Nlimit of the spindle index and completes the construction of the first example of a holographic duality involving supersymmetric field theories defined on orbifolds with conical singularities.

Introduction

The explanation of the microscopic origin of the entropy of supersymmetric black holes in anti de Sitter (AdS) is one of the most spectacular successes of the holographic duality. This was first accomplished in [1] for a class of AdS_4 black holes through the study of the large-N limit of the topologically twisted index [2]. The landscape of supersymmetric black holes was significantly broadened in [3], which constructed a supersymmetric, rotating and accelerating black hole with *spindle* horizon, displaying a number of remarkable features. Most strikingly, in this solution supersymmetry is preserved via a novel mechanism, referred to as anti-twist. It is was later noted that supersymmetry on the spindle may be preserved by means of a more standard topological twist [4, 5]. Utilising the insight of [6], it was shown in [7] that the on-shell action of a supersymmetric and complex deformation of the black hole of [3] takes the form of an *entropy func*tion, whose extremization yields the Bekenstein-Hawking entropy. A generalization of such entropy function was conjectured in [8], where it was proposed that it can be expressed in terms of gravitational blocks [9], as in all previous examples of black holes. The block decomposition of the gravitational entropy function was proved in [10] using the formalism of [11] and then in [12] employing equivariant localization in supergravity.

Motivated by these developments, [13, 14] computed the localized partition function of $\mathcal{N} = 2$ Chern-Simonsmatter theories defined on $\Sigma \times S^1$, where $\Sigma = \mathbb{WCP}_{[n_+,n_-]}^1$ is the spindle, with either twist or anti-twist for the Rsymmetry connection A:

$$\int_{\Sigma} \frac{\mathrm{d}A}{2\pi} = \frac{1}{2} \left(\frac{1}{n_{-}} + \frac{\sigma}{n_{+}} \right) \equiv \frac{\chi_{\sigma}}{2} , \qquad (1)$$

with $\sigma = \pm 1$. The result can be expressed by a single formula, dubbed *spindle index* [13], which can be defined [7] as a flavoured Witten index

$$Z_{\Sigma \times S^1} = \operatorname{Tr}_{\mathscr{H}[\Sigma]} \left[e^{-i\sum_{\alpha=1}^{\mathfrak{d}}\varphi_{\alpha}Q_{\alpha} + i\epsilon J} \right], \qquad (2)$$

where Q_{α} are the generators of global symmetries of rank \mathfrak{d} , J generates angular momentum on \mathbb{Z} , $\mathscr{H}[\mathbb{Z}]$ is the Hilbert space of BPS states on the spindle and the complex chemical potentials are related by the constraint

$$\sum_{\alpha=1}^{\mathfrak{d}} \varphi_{\alpha} + \frac{\chi_{-\sigma}}{2} \epsilon = 2\pi n , \qquad n \in \mathbb{Z} .$$
 (3)

In this letter we will demonstrate that the large-N limit of the spindle index reproduces the entropy functions associated to the supersymmetric and accelerating AdS₄ black holes. Explicitly, the entropy of a black hole with electric charges Q_{α} and angular momentum J is obtained by extremizing with respect to the variables φ_{α} and ϵ the entropy function

$$\mathcal{S} \equiv \log Z_{\Sigma \times S^1} + \mathrm{i} \sum_{\alpha=1}^{\mathfrak{d}} \varphi_{\alpha} Q_{\alpha} - \mathrm{i} \epsilon J \,, \tag{4}$$

under the constraint (3), setting n = 1 and requiring that J, Q_{α} and S are real. In the case $n_{+} = n_{-} = 1$ our result encompasses the large-N limit of both the topologically twisted index and the generalized superconformal index. More details and generalizations will be discussed in [15].

The spindle index matrix model

We consider $\mathcal{N} = 2$ Chern-Simons-matter quiver gauge theories with gauge group $\mathcal{G} = \prod_{a=1}^{|\mathcal{G}|} U(N)_a$ and chiral multiplets transforming in either bi-fundamental or adjoint representations of the gauge group factors. The index has been derived using supersymmetric localization in [13, 14] and it is written as the matrix model

$$Z_{\Sigma \times S^1}(\varphi, \mathfrak{n}, \epsilon) = \sum_{\mathfrak{m} \in \Gamma_{\mathfrak{h}}} \oint_{\mathcal{C}} \frac{\mathrm{d}u}{|W_{\mathcal{G}}|} \widehat{Z}(u, \mathfrak{m} | \varphi, \mathfrak{n}, \epsilon) , \quad (5)$$

where \mathfrak{h} , $\Gamma_{\mathfrak{h}}$ and $W_{\mathcal{G}}$ denote the Cartan algebra, the coroot lattice and the Weyl group of the gauge group \mathcal{G} , respectively; while \mathcal{C} is a suitable integration contour for u. Here we have collectively expressed by $u \in \mathfrak{h}$ and $\mathfrak{m} \in \Gamma_{\mathfrak{h}}$ the gauge holonomies on S^1 and fluxes through \mathbb{Z} , respectively. Similarly, φ and \mathfrak{n} are flavour/topological charges and fluxes, with (5) implicitly depending on the spindle data n_+, n_- and the twist parameter σ .

We focus on theories whose gauge group and matter content can be represented by a quiver diagram with $|\mathcal{G}|$ nodes, where an arrow from node a to node b corresponds to a bifundamental field in the representation $\mathbf{N}_a \otimes \overline{\mathbf{N}}_b$, and a = b indicates the adjoint representation. For each $\mathrm{U}(N)_a$ factor there are N holonomies and fluxes, $(u_i^a, \mathfrak{m}_i^a)_{i=0}^{N-1}$; for each arrow we assign flavour charges and fluxes $(\varphi_I, \mathfrak{n}_I)$, where the index I runs over *all* the $|\mathcal{R}|$ chiral multiplets of the theory. If the corresponding arrow stretches from a node a to a node b, we write $I \in (a, b)$. Moreover, for each node we assign charges/fluxes $(\varphi_m^a, \mathfrak{n}_m^a)$ for the topological symmetries. As in [13, 14] we consider a choice of R-symmetry that assigns even charges to the chiral multiplets: $r_I \in 2\mathbb{Z}$. For a chiral multiplet the corresponding chemical potential φ_I is related to the flavour holonomy u_I^F via

$$\varphi_I = 2\pi u_I^F + \left(\pi n - \frac{\epsilon}{4}\chi_{-\sigma}\right)r_I , \qquad (6)$$

where, for each monomial term W in the superpotential,

$$\sum_{I \in W} u_I^F = \sum_{I \in W} \mathfrak{n}_I = 0, \qquad \sum_{I \in W} r_I = 2, \qquad (7)$$

so that

$$\sum_{I \in W} \varphi_I + \frac{\chi_{-\sigma}}{2} \epsilon = 2\pi n .$$
(8)

Notice that the index I runs over the fields belonging to a superpotential term, while in (3) the index α labels the generators of the global symmetries of the theory. For the ABJM model, that is the main focus in this letter, these two sets coincide. More general quivers will be discussed in [15].

The integrand of (5) is the product of a classical part and the 1-loop determinants of chiral and vector multiplets. In order to write it explicitly we need to introduce some further notation [13]: first, we define the symbols $\sigma_{+} = \sigma$ and $\sigma_{-} = -1$. Then, we set

$$\mathfrak{b}_{ij}^{I} = 1 - \frac{\mathfrak{m}_{i}^{a} - \mathfrak{m}_{j}^{b}}{n_{+}n_{-}} - \frac{\mathfrak{n}_{I}}{n_{+}n_{-}} - \frac{r_{I}}{2}\chi_{\sigma} - \mathcal{A}_{I;\,ij}^{-} - \sigma \mathcal{A}_{I;\,ij}^{+} ,$$

$$\mathfrak{c}_{ij}^{I} = \mathcal{A}_{I;\,ij}^{-} - \sigma \mathcal{A}_{I;\,ij}^{+} , \qquad (9)$$

for each arrow $I \in (a, b)$, with

$$\mathfrak{l}_{a;i}^{\pm} = n_{\pm} \left\{ \frac{\sigma_{\pm} a_{\pm} \mathfrak{m}_{i}^{a}}{n_{\pm}} \right\}, \\
\mathcal{A}_{I;ij}^{\pm} = \left\{ \frac{\mathfrak{l}_{a;i}^{\pm} - \mathfrak{l}_{b;j}^{\pm} + \sigma_{\pm} a_{\pm} \mathfrak{n}_{I} - r_{I}/2}{n_{\pm}} \right\}, \quad (10)$$

and $a_{\pm} \in \mathbb{Z}$ such that $n_{+}a_{-} - n_{-}a_{+} = 1$. Moreover $\{x\} \equiv x - \lfloor x \rfloor$. Notice that $\mathfrak{b}_{ij}^{I} \in \mathbb{Z}$, while $\mathfrak{l}_{a;i}^{\pm}, n_{\pm}\mathcal{A}_{I;ij}^{\pm} \in \mathbb{Z}_{n_{\pm}}$. Denoting by

$$y_{ij}^I = e^{-i\varphi_I - 2\pi i (u_i^a - u_j^b)} \cdot q^{\frac{1}{2}\mathfrak{c}_{ij}^I} , \qquad q = e^{i\epsilon} , \qquad (11)$$

the gauge holonomies, the 1-loop determinant contribution of the chiral multiplets can be written as [13, 14]

$$Z_{1-\mathrm{L}}^{\mathrm{CM}} = \prod_{I=1}^{|\mathcal{R}|} \prod_{i,j=0}^{N-1} \zeta_q^{\sigma}(y_{ij}^I, \mathfrak{b}_{ij}^I) , \qquad (12)$$

in terms of the function

$$\zeta_{q}^{\sigma}(y,\mathfrak{b}) \equiv (-y)^{\frac{1-\sigma-2\mathfrak{b}}{4}} q^{\frac{(1-\sigma)(\mathfrak{b}-1)}{8}} \frac{(q^{\frac{1+\mathfrak{b}}{2}}y^{-1};q)_{\infty}}{(q^{\frac{1-\mathfrak{b}}{2\sigma}}y^{-\sigma};q)_{\infty}}, \quad (13)$$

where $(z;q)_{\infty}$ is the *q*-Pochhammer symbol, $y,q \in \mathbb{C}$, $\mathfrak{b} \in \mathbb{Z}$ and $\sigma = \pm 1$. This is the 1-loop determinant of a single chiral multiplet in an Abelian theory, satisfying

$$\zeta_q^{\sigma}(y,\mathfrak{b}) = \zeta_q^{\sigma}(y^{-\sigma}, 1 - \sigma - \mathfrak{b})^{-\sigma}.$$
(14)

The 1-loop determinant of all vector multiplets reads

$$Z_{1-L}^{VM} = \prod_{a=1}^{|\mathcal{G}|} \prod_{i,j=0}^{N-1} \zeta_q^{\sigma}(y_{ij}^a, \mathfrak{b}_{ij}^a) , \qquad (15)$$

where y_{ij}^a , \mathfrak{b}_{ij}^a , \mathfrak{c}_{ij}^a , and $\mathcal{A}_{a;ij}^{\pm}$ are defined as in (9), (10), and (11), with all the instances of I and b replaced by a, and with the following identifications: $r_a \equiv 2$, $\mathfrak{n}_a \equiv 0$, $\varphi_a \equiv 2\pi n - \frac{\epsilon}{2}\chi_{-\sigma}$.

The classical part receives contributions from the Chern-Simons terms, which can be written as [15]

$$Z_{\text{eff}}^{\text{CS}} = \prod_{a=1}^{|\mathcal{G}|} \prod_{i=0}^{N-1} \left(-y_i^a \right)^{\mathbf{k}_a (\mathfrak{b}_i^a - 1)}, \qquad (16)$$

where we defined

$$y_{i}^{a} = e^{-2\pi i u_{i}^{a}} \cdot q^{\frac{l_{a;i}}{2n_{-}} - \sigma \frac{l_{a;i}}{2n_{+}}},$$

$$\mathfrak{b}_{i}^{a} = 1 - \frac{\mathfrak{m}_{i}^{a}}{n_{+}n_{-}} - \frac{\mathfrak{l}_{a;i}}{n_{-}} - \sigma \frac{\mathfrak{l}_{a;i}^{+}}{n_{+}}.$$
 (17)

In this paper we restrict to the case where $\sum_a k_a = 0$, corresponding to $\mathcal{N} = 2$ Chern-Simons-matter quiver gauge theories with an M theory dual $\mathrm{AdS}_4 \times M_7$. The topological symmetries also contribute to the classical part but the explicit expression will not be needed in this letter.

Holomorphic block factorization

The spindle index factorizes into the product of dual holomorphic blocks [16]. It is convenient to use a choice

of factorization that breaks the Weyl-symmetry of the gauge group, generalizing the one introduced in [17] for the superconformal index. Starting from (12), we split the product over i, j into a product over i < j and one over i > j, ignoring the diagonal terms that are subleading at large N; then we apply (14) to the i > j terms and we find

$$Z_{1-\mathrm{L}}^{\mathrm{CM}} = \prod_{I=1}^{|\mathcal{R}|} \Psi_I \cdot \mathcal{B}_I^+ \cdot \mathcal{B}_I^-, \qquad (18)$$

where for $I \in (a, b)$ we defined

$$\Psi_{I} = \prod_{i < j} (y_{ij}^{I})^{\frac{1 - \sigma - 2\mathfrak{b}_{ij}^{I}}{4}} (y_{ji}^{I})^{-\frac{1 - \sigma - 2\mathfrak{b}_{ji}^{I}}{4}} \cdot q^{\frac{(1 - \sigma)(\mathfrak{b}_{ij}^{I} - \mathfrak{b}_{ji}^{I})}{8}},$$
$$\mathcal{B}_{I}^{\pm} = \prod_{\substack{\mathcal{B}^{-}: i < j \\ \mathcal{B}^{+}: i > j}} \frac{\left(\left(\frac{z_{a;i}^{\pm}}{z_{b;j}^{\pm}} \right)^{\pm \sigma_{\pm}} \mathrm{e}^{-\mathrm{i}\sigma_{\pm}\Delta_{I}^{\pm}} q^{1 - \mathcal{A}_{I;ij}^{\pm}; q} \right)_{\infty}}{\left(\left(\frac{z_{a;i}^{\pm}}{z_{b;i}^{\pm}} \right)^{\mp \sigma_{\pm}} \mathrm{e}^{\mathrm{i}\sigma_{\pm}\Delta_{I}^{\pm}} q^{\mathcal{A}_{I;ji}^{\pm}; i}; q \right)_{\infty}}.$$
 (19)

Notice that here we have swapped the role of i, j in the blocks for convenience. The Ψ_I will turn out to be subleading after the cancellation of the long-range forces. The blocks \mathcal{B}_I^{\pm} depend on the combinations

$$\Delta_I^{\pm} = \varphi_I \pm \frac{\epsilon}{2} \left(\frac{\mathfrak{n}_I}{n_+ n_-} + \frac{\chi_\sigma}{2} r_I \right) ,$$

$$z_{a;i}^{\pm} = e^{\pm 2\pi i u_i^a} q^{-\frac{\mathfrak{m}_i^a}{2n_+ n_-}} .$$
(20)

Notice that the variables Δ_I^{\pm} satisfy the constraints

$$\sum_{I \in W} \Delta_I^{\pm} = 2\pi n + \frac{\sigma_{\pm} \epsilon}{n_{\pm}}.$$
 (21)

We derive the vector-multiplet counterparts of (18) and (19) by replacing the indices I and b with a and applying standard identifications.

Strategy for the large-N limit

We will implement the large-N limit of the spindle index by relying on its factorization into holomorphic blocks, generalizing the approach of [17–19]. For the partition functions on $S^2 \times S^1$ the sum over all the possible values of the gauge fluxes $\mathfrak{m} \in \Gamma_{\mathfrak{h}} \equiv \mathbb{Z}^{|\mathcal{G}|N}$ is usually approximated at large N by promoting the fluxes \mathfrak{m} to continuous variables. However, for $\mathbb{Z} \times S^1$ this approximation is hindered by the presence of fractional parts in (10). To take care of this, we split each gauge flux as $\mathfrak{m}_i^a \equiv n_+n_-(\mathfrak{m}')_i^a + \mathfrak{r}_i^a$, with $(\mathfrak{m}')_i^a \in \mathbb{Z}$ and $\mathfrak{r}_i^a \in \mathbb{Z}_{n+n_-}$. We then observe that there is a one-to-one correspondence between the possible values of $\mathfrak{l}_{a;i}^{\pm}$ and \mathfrak{r}_i^a :

$$\frac{\mathfrak{l}_{a;i}^{\pm}}{n_{\pm}} = \left\{ \frac{\sigma_{\pm} a_{\pm} \mathfrak{r}_{i}^{a}}{n_{\pm}} \right\} , \quad \frac{\mathfrak{r}_{i}^{a}}{n_{+} n_{-}} = \left\{ -\frac{\mathfrak{l}_{a;i}^{-}}{n_{-}} - \sigma \frac{\mathfrak{l}_{a;i}^{+}}{n_{+}} \right\} . \quad (22)$$

We can therefore split the sum over \mathfrak{m}_i^a as

$$\sum_{\mathfrak{m}_{i}^{a} \in \mathbb{Z}} = \sum_{\mathfrak{l}_{a;i}^{-}=0}^{n_{-}-1} \sum_{\mathfrak{l}_{a;i}^{+}=0}^{n_{+}-1} \sum_{(\mathfrak{m}')_{i}^{a} \in \mathbb{Z}} , \qquad (23)$$

and in the large-N limit we may promote the $(\mathfrak{m}')_i^a$ to be continuous variables while keeping the $\mathfrak{l}_{a;i}^{\pm}$ discrete. Thus, we approximate the integration measure of (5) by

$$\sum_{\mathfrak{m}\in\Gamma_{\mathfrak{h}}} \oint_{\mathcal{C}} \frac{\mathrm{d}u}{|W_{\mathcal{G}}|} \longrightarrow \sum_{\mathfrak{l}^{\pm}\in(\mathbb{Z}_{n_{\pm}})^{|\mathcal{G}|N}} \int_{\mathcal{C}^{+}} \mathrm{d}z^{+} \int_{\mathcal{C}^{-}} \mathrm{d}z^{-}, \quad (24)$$

at large N, where the variables z^{\pm} were defined in (20), \mathcal{C}^{\pm} are appropriate middle-dimensional contours in $\mathbb{C}^{|\mathcal{G}|N}$, and the order of the Weyl group can be ignored since $\log |W_{\mathcal{G}}| = \mathcal{O}(N \log N)$.

Since the \mathcal{B}_{I}^{\pm} blocks depend separately on z^{\pm} , we will be able to perform the saddle point approximation in z^{-} and z^+ independently of one another. However the right hand side of (24) also features a sum over the vectors of integers l^{\pm} , which can take a total of $(n_{+}n_{-})^{|\mathcal{G}|N}$ possible values, exponentially growing with N. At large Nonly one value of the l^{\pm} is expected to dominate at any given region of the parameter space: in particular, there will be one such value associated to the saddle point that reproduces the accelerating AdS_4 black holes. Two observations are in order to find the correct ansatz for l^{\pm} : first, we need to restrict our attention to the set of possible choices of l^{\pm} that, up to an appropriate permutation of the index i, are periodic under shifts $i \to i+T$ for some $T \ll N$. This assumption is necessary in order to be able to take (partially) the continuum limit: splitting the index *i* as $i = Ti' + \tilde{i}$ with $\tilde{i} \in \{0, \dots, T-1\}$, makes the fluxes $\mathfrak{l}_{a;i}^{\pm}$ only depend on the index $\tilde{i}, \mathfrak{l}_{a;i}^{\pm} \equiv \mathfrak{l}_{a;\tilde{i}}^{\pm}$. Hence, at large N the index i' can be replaced with a continuous variable t. Second, all the known methods for computing 3d partition functions at large N [1, 20, 21] require that terms with $i \sim j$ dominate over the terms with $|i-j| \gg 1$. The latter are called "long-range forces" and with the appropriate assumptions they cancel out at leading order, at least for a class of quiver theories that we shall discuss momentarily. The cancellation of long-range forces constrains the possible choices of l^{\pm} , although in general the constraint is complicated and it involves the value of z^{\pm} as well. Remarkably, the special value

$$\mathfrak{l}_{a;i}^{\pm} = i \mod n_{\pm} \tag{25}$$

makes the long-range forces vanish for any z^{\pm} . We also anticipate that (25), along with a simple ansatz for z^{\pm} , reproduces the entropy of accelerating AdS₄ black holes. Curiously, (25) exhibits a strong similarity to the ansatz reproducing the entropy of AdS₅ black holes with arbitrary momenta, as discussed in [22, 23].

Long-range forces cancellation

In (19) the prefactors Ψ_I encode long-range forces among the variables z^{\pm} that could spoil the large-N limit. As in previous work on 3d theories, we cancel the long-range forces by restricting to "non-chiral" quivers, where for any bi-fundamental connecting the nodes a and b there is a bi-fundamental connecting b and a and

$$\sum_{I \in (a)} \mathfrak{n}_I = \sum_{I \in (a)} u_I^F = \sum_{I \in (a)} (r_I - 1) + 2 = 0, \qquad (26)$$

at each node a, where the sum is taken over all the arrows in the quiver with an endpoint at the node a, with adjoint chirals counting twice. In a four-dimensional quiver this condition would be equivalent to the absence of ABJ anomalies for any symmetry. The conditions (26) also imply that Tr Q = 0 for any global or R-symmetry symmetry with generator Q, where the trace is taken over all the fermions in the theory.

Using the periodicity relation $\mathfrak{l}_{a;i}^{\pm} = \mathfrak{l}_{a;i+T}^{\pm}$ that we have assumed, for the non-chiral quivers satisfying (26) the product of all the prefactor terms (19) at large N can be simplified down to

$$\prod_{I=1}^{|\mathcal{R}|} \Psi_I \cdot \prod_{a=1}^{|\mathcal{G}|} \Psi_a \longrightarrow \prod_{I=1}^{|\mathcal{R}|} \widetilde{\Psi}_I \cdot \prod_{a=1}^{|\mathcal{G}|} \widetilde{\Psi}_a , \qquad (27)$$

where for $I \in (a, b)$

$$\widetilde{\Psi}_{I} = \prod_{s=\pm} \prod_{i,j=0}^{N-1} \left(\frac{z_{a;i}^{s}}{z_{b;j}^{s}} \right)^{\frac{\sigma_{s}}{4} \left(1 - \frac{1}{n_{s}} - 2\mathcal{A}_{I;ij}^{s} \right) \cdot \operatorname{sign}(i-j)}$$
(28)

and a similar definition holds for $\widetilde{\Psi}_a$. Requiring the right hand side of (27) to vanish yields a mixed constraint on l^{\pm} and z^{\pm} . Crucially, the ansatz (25) is the only one that satisfies the property

$$\frac{1}{n_{\pm}} \sum_{j=j_0}^{j_0+n_{\pm}-1} \mathcal{A}_{I;\,ij}^{\pm} = \frac{1}{2} \left(1 - \frac{1}{n_{\pm}} \right) \tag{29}$$

(and a similar relation with i, j inverted) ensuring that the long-range forces coming from (27) vanish for any z^{\pm} . Thanks to the Weyl-symmetry breaking factorization that we have used, the blocks \mathcal{B}_{I}^{\pm} will not produce any long-range term at leading order, as will now show.

Holomorphic blocks at large N

In order to compute the large-N limit of the blocks \mathcal{B}_{I}^{\pm} , we will first consider the usual ansatz for the saddle point distribution of z^{\pm} [1, 20, 21],

$$\log z_{a;i}^{\pm} = -\sigma_{\pm} N^{\alpha} t_i \mp \mathrm{i} y_a^{\pm}(t_i) , \qquad (30)$$

4

where t_i , $y_a(t_i)$ are real and are assumed to be ordered so that $t_i \leq t_j$ for i < j. The power of N must be set to $\alpha = \frac{1}{2}$, otherwise the 1-loop contributions and the Chern-Simons terms would grow with a different power law at large N and it would not be possible to find non-trivial critical points. When we take the continuum limit we split the index $i \equiv Ti' + \tilde{i}$: assuming that the eigenvalues t_i conform to a single continuous distribution at large N allows to make the replacements $t_i \equiv t_{i'} \equiv t$ and define the eigenvalue density $\rho^{\pm}(t)$ such that

$$\frac{1}{N}\sum_{i=0}^{N-1} \bullet \longrightarrow \frac{1}{T}\sum_{\tilde{i}=0}^{T-1} \int \mathrm{d}t\rho^{\pm}(t)\bullet, \quad \int \mathrm{d}t\rho^{\pm}(t) = 1.$$
(31)

Expanding the *q*-Pochhammer symbols in terms of polylogarithms at all orders in ϵ and taking the large-N limit of each term as in [1, 20, 21] yields

$$\log \mathcal{B}_{I}^{\pm} = N^{\frac{3}{2}} \sum_{k=0}^{2} \epsilon^{k-1} \frac{B_{k}}{k!} \frac{1}{T^{2}} \sum_{\tilde{\imath}, \tilde{\jmath}=0}^{T-1} \int \mathrm{d}t \rho^{\pm}(t)^{2} \cdot \qquad (32)$$
$$\cdot g_{3-k}(-\sigma_{\pm} \delta y_{ab}^{\pm}(t) - \sigma_{\pm} \Delta_{I}^{\pm} - \epsilon \mathcal{A}_{I;\tilde{\imath}\tilde{\jmath}}^{\pm}) + o(N^{\frac{3}{2}}) ,$$

with $B_k = B_k(1) = \{1, \frac{1}{2}, \frac{1}{6}, \ldots\}$ and

$$g_k(x) = \frac{(2\pi)^k}{k!} B_k\left(\frac{x}{2\pi} + \nu\right),$$
 (33)

where $B_k(w)$ are the Bernoulli polynomials and the integer ν in (33) must be chosen so that $\operatorname{Im}(\frac{1}{\epsilon}) < 1$ $\operatorname{Im}\left(\frac{1}{\epsilon}\left(\frac{x}{2\pi}+\nu\right)\right) < 0. \text{ We are using the notation } \delta y_{ab}^{\pm\epsilon'}(t) \equiv y_{a}^{\pm}(t) - y_{b}^{\pm}(t).$

The Large-N limit of the spindle index

We assume that the index is dominated by the configuration (25), which leads to a consistent large-N limit. The large-N limit of the classical Chern-Simons terms simplifies to

$$\log Z_{\text{eff}}^{\text{CS}} = N^{3/2} \sum_{a} \sum_{s=\pm} \frac{\sigma_s}{\epsilon} k_a \int \mathrm{d}t \, t \rho^s(t) y_a^s(t) \,. \tag{34}$$

Consistently with fact that for saddles with gravity duals flux quantization implies $N/(n_+n_-) \in \mathbb{N}$ [3], we can take $T = n_{+}n_{-}$. Moreover, in order to compare with the black hole solutions, we need to take n = 1 [7]. Finally, we also need to choose a determination: we assume that $\operatorname{Im}\left(\frac{1}{\epsilon}\right) < \operatorname{Im}\left(\frac{1}{2\pi\epsilon}(y_{ab}^{\pm}(t) + \varphi_I)\right) < 0.$ After some algebra, the explicit expression for $\mathfrak{l}_{a;i}^{\pm}$ and the conditions (26) vield

$$\log Z_{\mathbb{Z} \times S^1} = -\sum_{s=\pm} \sigma_s \frac{F(\rho^s, \delta y^s_{ab}, \Delta^s_I)}{\epsilon}$$
(35)

with

$$\frac{F(\rho^{\pm}, \delta y_{ab}^{\pm}, \Delta_I^{\pm})}{N^{3/2}} = -\sum_a k_a \int dt \, t \rho^{\pm}(t) y_a^{\pm}(t) + \sum_{I \in (a,b)} \int dt \rho^{\pm}(t)^2 G_3^{\pm}(\delta y_{ab}^{\pm}(t) + \Delta_I^{\pm}) \,, \qquad (36)$$

where $G_3^{\pm}(x) = \frac{1}{6}x(x - \sum_{I \in W} \Delta_I^{\pm}/2)(x - \sum_{I \in W} \Delta_I^{\pm})$. The functions $G_3^{\pm}(x)$ are obtained by $g_3(x)$ in the range Re $x \in [0, 2\pi]$ by replacing all occurrences of π with $\sum_{I \in W} \Delta_I^{\pm}/2$. The two terms in (35) depend on different variables and they can be extremized independently.

For example, for the ABJM theory dual to $\operatorname{AdS}_4 \times S^7$, with $|\mathcal{G}| = 2$, Chern-Simons level $k_1 = 1$ and $k_2 = -1$ and four bi-fundamental fields transforming as $\mathbf{N}_1 \otimes \overline{\mathbf{N}}_2$ for I = 1, 2 and as $\mathbf{N}_2 \otimes \overline{\mathbf{N}}_1$ for I = 3, 4, we find

$$(36) = \int dt t \rho^{\pm} \delta y_{21}^{\pm} - \frac{1}{2} \int dt (\rho^{\pm})^2 \Big(\sum_I \Delta_I^{\pm} (\delta y_{21}^{\pm})^2 - 2(\Delta_1^{\pm} \Delta_2^{\pm} - \Delta_3^{\pm} \Delta_4^{\pm}) \delta y_{21}^{\pm} - \sum_{I < J < K} \Delta_I^{\pm} \Delta_J^{\pm} \Delta_K^{\pm} \Big).$$
(37)

This functional coincides with the large-N limit of the effective twisted superpotential for the ABJM theory derived in [1], expressed in terms of \pm quantities, and its extremization is straightforward [24]. The explicit expressions for ρ^{\pm} and δy_{21}^{\pm} can be found for example in [1, (2.70)-(2.75)]. The critical value is

$$F(\rho^{\pm}, \delta y_{21}^{\pm}, \Delta_I^{\pm})\Big|_{\text{crit}} = \frac{2}{3}N^{3/2}\sqrt{2\Delta_1^{\pm}\Delta_2^{\pm}\Delta_3^{\pm}\Delta_4^{\pm}} \,. \tag{38}$$

Using (35) we recover the gravitational block form [8] [25] of the entropy function obtained in [7] and more generally conjectured in [5]. The density of eigenvalues ρ^{\pm} also agrees with the gravitational analysis performed in [26].

We can extend the result to more general quivers: indeed, the term of order zero of F in the ϵ expansion coincides with the large-N limit of the effective twisted superpotential of the $\mathcal{N} = 2$ theory [27]

$$i\frac{\mathcal{W}(\rho,\delta y_{ab},\Delta_I)}{N^{3/2}} = -\sum_a k_a \int dt \, t\rho(t) y_a(t) + \sum_{I \in (a,b)} \int dt \rho(t)^2 g_3(\delta y_{ab}(t) + \Delta_I) \,, \tag{39}$$

where $\Delta_I = \Delta_I^{\pm}|_{\epsilon=0} = 2\pi u_I^F + \pi r_I$ and we are ignoring topological symmetries for simplicity. This agrees with well-known asymptotic behaviour of the holomorphic blocks [16]: $\log(\text{block}) = i\frac{\mathcal{W}}{\epsilon} + O(\epsilon)$. We then observe that (36) is a homogeneous form of the large-Nlimit of the effective twisted superpotential \mathcal{W} obtained by replacing Δ_I with Δ_I^{\pm} and all occurrences of π with $\sum_{I \in \mathcal{W}} \Delta_I^{\pm}/2$. The extremization of (36) is then equivalent to the extremization of \mathcal{W} with the constraint $\sum_{I \in W} \Delta_I = 2\pi$. One concludes that the entropy function has always a block form

$$\log Z_{\Sigma \times S^1} = \frac{F_{\text{crit}}(\Delta_I^-)}{\epsilon} - \sigma \frac{F_{\text{crit}}(\Delta_I^+)}{\epsilon} \,. \tag{40}$$

We also see that the block function $F_{\text{crit}}(\Delta_I)$, up to factors, is the homogeneous form of the large-N on-shell value of the effective twisted superpotential \mathcal{W} . This has been computed for many examples in [28]. At large N \mathcal{W} coincides with the S^3 partition function of the $\mathcal{N} = 2$ theory [27] and, for theories with an $\text{AdS}_4 \times M_7$ dual, the latter is in turn related [20, 21] to the Sasakian volume [29, 30] of M_7 . Using this chain of equalities, one provides a field theory derivation of the gravitational block decomposition obtained in [10, 12, 31] for configurations with a "mesonic" (or, "flavour") twist [32]. More details about topological symmetries and issues with baryonic symmetries will be discussed in [15].

Discussion

In this letter we solved the fundamental problem of elucidating the microscopic origin of the Bekenstein-Hawking entropy of the most general class of rotating BPS black holes currently known in four dimensions. Specifically, our findings demonstrate that the microstates contributing to the entropy of accelerating black holes in fourdimensional anti de Sitter space-time are precisely mirrored by the physical degrees of freedom characterizing three-dimensional gauge theories quantized on a spindle. To successfully solve this problem we developed a novel approach tailored to deal with the degrees of freedom of gauge theories on orbifolds. This technique holds vast potential impact as it applies to supersymmetric systems in any number of dimensions, including e.g. three-dimensional orbifold partition functions [14] and four-dimensional orbifold indices [33]. Our results complete the construction of the first duality between a gravitational theory and a quantum field theory defined on an orbifold, paving the way for a re-energized research program in holography.

Ackowledgments

The work of EC and DM is supported in part by a grant Trapezio (2023) of the Fondazione Compagnia di San Paolo. AZ is partially supported by the MUR-PRIN grant No. 2022NY2MXY. EC, DM, AP and AZ acknowledge partial support by the INFN. SMH is supported in part by the STFC Consolidated Grants ST/T000791/1 and ST/X000575/1.

- F. Benini, K. Hristov, and A. Zaffaroni, JHEP 05, 054 (2016), arXiv:1511.04085 [hep-th].
- [2] F. Benini and A. Zaffaroni, JHEP 07, 127 (2015), arXiv:1504.03698 [hep-th].
- [3] P. Ferrero, J. P. Gauntlett, J. M. P. Ipiña, D. Martelli, and J. Sparks, Phys. Rev. D 104, 046007 (2021), arXiv:2012.08530 [hep-th].
- [4] P. Ferrero, J. P. Gauntlett, and J. Sparks, JHEP 01, 102 (2022), arXiv:2112.01543 [hep-th].
- [5] P. Ferrero, M. Inglese, D. Martelli, and J. Sparks, Phys. Rev. D 105, 126001 (2022), arXiv:2109.14625 [hep-th].
- [6] A. Cabo-Bizet, D. Cassani, D. Martelli, and S. Murthy, JHEP 10, 062 (2019), arXiv:1810.11442 [hep-th].
- [7] D. Cassani, J. P. Gauntlett, D. Martelli, and J. Sparks, Phys. Rev. D 104, 086005 (2021), arXiv:2106.05571 [hep-th].
- [8] F. Faedo and D. Martelli, JHEP 02, 101 (2022), arXiv:2111.13660 [hep-th].
- [9] S. M. Hosseini, K. Hristov, and A. Zaffaroni, JHEP 12, 168 (2019), arXiv:1909.10550 [hep-th].
- [10] A. Boido, J. P. Gauntlett, D. Martelli, and J. Sparks, Commun. Math. Phys. 403, 917 (2023), arXiv:2211.02662 [hep-th].
- [11] C. Couzens, J. P. Gauntlett, D. Martelli, and J. Sparks, JHEP 01, 212 (2019), arXiv:1810.11026 [hep-th].
- [12] P. Benetti Genolini, J. P. Gauntlett, Y. Jiao, A. Lüscher, and J. Sparks, (2024), arXiv:2401.10977 [hep-th].
- [13] M. Inglese, D. Martelli, and A. Pittelli, J. Phys. A 57, 085401 (2024), arXiv:2303.14199 [hep-th].
- [14] M. Inglese, D. Martelli, and A. Pittelli, (2023), arXiv:2312.17086 [hep-th].
- [15] E. Colombo, S. M. Hosseini, D. Martelli, A. Pittelli, and A. Zaffaroni, (2024).
- [16] C. Beem, T. Dimofte, and S. Pasquetti, JHEP 12, 177 (2014), arXiv:1211.1986 [hep-th].
- [17] S. Choi, C. Hwang, and S. Kim, (2019), arXiv:1908.02470 [hep-th].
- [18] S. Choi and C. Hwang, JHEP 03, 068 (2020), arXiv:1911.01448 [hep-th].

- [19] S. M. Hosseini and A. Zaffaroni, JHEP 12, 025 (2022), arXiv:2209.09274 [hep-th].
- [20] C. P. Herzog, I. R. Klebanov, S. S. Pufu, and T. Tesileanu, Phys. Rev. D83, 046001 (2011), arXiv:1011.5487 [hep-th].
- [21] D. Martelli and J. Sparks, Phys. Rev. D84, 046008 (2011), arXiv:1102.5289 [hep-th].
- [22] F. Benini, E. Colombo, S. Soltani, A. Zaffaroni, and Z. Zhang, Class. Quant. Grav. **37**, 215021 (2020), arXiv:2005.12308 [hep-th].
- [23] E. Colombo, JHEP 12, 013 (2022), arXiv:2110.01911 [hep-th].
- [24] The functional \mathcal{W} written in [1] has a critical point under the condition $\sum_{I} \Delta_{I} = 2\pi$. We are using a homogeneous form of \mathcal{W} where this condition has been used to eliminate all occurrences of π . The variables Δ_{I} in [1] correspond to our $\Delta_{I}^{\pm}|_{\epsilon=0}$.
- [25] To compare with the formulas in [8] we set n = 1 and identify the variables as follows: $\pi \Delta_i^{\pm}|_{\text{there}} = \Delta_I^{\mp}|_{\text{here}}$, $-2\pi\epsilon|_{\text{there}} = \epsilon|_{\text{here}}, \pi\varphi_i|_{\text{there}} = \varphi_I|_{\text{here}}, r_i|_{\text{there}} = r_I|_{\text{here}},$ $\mathfrak{n}_i|_{\text{there}} = \mathfrak{n}_I|_{\text{here}}/(n_+n_-) + r_I\chi_{\sigma}/2.$
- [26] A. Boido, A. Lüscher, and J. Sparks, JHEP 05, 226 (2024), arXiv:2312.11640 [hep-th].
- [27] S. M. Hosseini and A. Zaffaroni, JHEP 08, 064 (2016), arXiv:1604.03122 [hep-th].
- [28] S. M. Hosseini and N. Mekareeya, JHEP 08, 089 (2016), arXiv:1604.03397 [hep-th].
- [29] D. Martelli, J. Sparks, and S.-T. Yau, Commun. Math. Phys. 268, 39 (2006), arXiv:hep-th/0503183 [hep-th].
- [30] D. Martelli, J. Sparks, and S.-T. Yau, Commun. Math. Phys. 280, 611 (2008), arXiv:hep-th/0603021 [hep-th].
- [31] A. Boido, J. P. Gauntlett, D. Martelli, and J. Sparks, Phys. Rev. Lett. **130**, 091603 (2023), arXiv:2210.16069 [hep-th].
- [32] S. M. Hosseini and A. Zaffaroni, JHEP 07, 174 (2019), arXiv:1904.04269 [hep-th].
- [33] A. Pittelli, (2024), arXiv:2403.12318 [hep-th].