

Available online on 15.03.2023 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

Copyright © 2023 The Author(s): This is an open-access article distributed under the terms of the CC BY-NC 4.0 which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited



Open Access Full Text Article



Check for updates

Letter to Editor

All About Lercanidipine: Ten Questions and Answers

Francesco Fici ^{1,2} , Gokhan Faikoglu ^{3*} , Guido Grassi ^{4,5} , Nicolas Roberto Robles ^{1,2,6} , Kubra Saygisever-Faikoglu ³

1. *Catedra de Riesgo Cardiovascular, Universidad de Salamanca, Salamanca, Spain*

2. *Milano-Bicocca, University, Milan, Italy*

3. *Department of Pharmacology, Cerrahpasa Faculty of Medicine, Istanbul University, Cerrahpasa, Istanbul, Turkey*

4. *Department of Medicine and Surgery, University of Milano-Bicocca, Milan, Italy*

5. *IRCCS Multimedica, Sesto San Giovanni, Milan, Italy*

6. *Hospital Universitario de Badajoz. Badajoz. Spain*

Article Info:



Article History:

Received 03 Feb 2023
Reviewed 24 Feb 2023
Accepted 01 March 2023
Published 15 March 2023

Cite this article as:

Fici F, Faikoglu G, Grassi G, Robles NR, Faikoglu KS, All About Lercanidipine: Ten Questions and Answers, Journal of Drug Delivery and Therapeutics. 2023; 13(3)P1-P3 DOI: <http://dx.doi.org/10.22270/jddt.v13i3.5769>

*Address for Correspondence:

Gokhan Faikoglu, Department of Pharmacology, Cerrahpasa Faculty of Medicine, Istanbul University, Cerrahpasa, Istanbul, Turkey

INTRODUCTION

Dihydropyridine calcium channel blockers (DHP-CCBs) are first-line drugs in antihypertensive therapy, either as monotherapy, or in combination with other antihypertensive drugs, as recommended by ESC/ESH ¹. Lercanidipine, a third-generation calcium channel blocker, is characterized by high vascular selectivity, high lipophilic properties, slow onset and long duration of pharmacological effect. Therefore, lercanidipine is different from a number of other CCBs ².

1. What are the most important pharmacological properties?

Dihydropyridine-CCBs reversibly inhibit voltage-activated L-type Ca⁺⁺ channels, located in the smooth muscle cells of the arteries ^{3,4}. The reduction of intracellular calcium level leads to vasodilation and consequently to blood pressure (BP) reduction. Differently from other dihydropyridine CCBs, lercanidipine inhibits both L and T calcium channels ⁵. There is evidence that T channels are particularly expressed in the renal efferent arterioles, while L channels are particularly located in the afferent arterioles ^{6,7}.

Therefore, lercanidipine blocking both L and T channels, dilates afferent and efferent renal arterioles and thereby prevents the increase of glomerular pressure. The result is a decrease of filtration fraction and, therefore, a kidney function protective effect ⁸.

2. What is the evidence of antihypertensive efficacy?

Double-blind, randomized, comparative trials and large open, observational studies have shown that, lercanidipine (5-20mg once daily) significantly decreases systolic and diastolic blood

pressure (SBP/DBP), assessed either as office and home measurement or as 24-hours monitoring (ABPM). Responder rate (SBP/DBP reduction greater than 20 and 10 mmHg respectively) is achieved by 62%-72% of patients ^{9,10,11}. The antihypertensive efficacy of lercanidipine has also been successfully reported in patients with isolated systolic hypertension, ^{12,13} with diabetes, ¹⁴ and with cardiovascular disease ¹⁵. The therapeutic activity does not differ statistically from that of amlodipine, felodipine, nifedipine GITS, lacidipine, manidipine, ¹⁶ losartan ¹⁷ and candesartan ¹⁸.

3. What about the duration of the antihypertensive effect?

Blood pressure reduction during lercanidipine treatment is sustained throughout 24-hour, with a significant reduction of morning BP rise and BP variability ^{10,11,19,20}.

4. Is the antihypertensive effect different according to the age and gender of patients?

The antihypertensive effect of lercanidipine is not different between young and elderly patients, as well between women and men ^{10,21}.

5. What is the relationship between lercanidipine and endothelial dysfunction?

Essential hypertension is associated with impaired endothelium-mediated nitric oxide (NO) release, induced by oxidative stress. Therefore, the vascular tone shifts to vasoconstriction, resulting in higher peripheral vascular resistance ²². In patients with essential hypertension, lercanidipine significantly increases endothelium-mediated vasodilation, through the release of NO. Additionally the drug shows, antioxidant activity, lowering some markers of

oxidative stress, as lipoperoxides, isoprostanes and malondialdehyde and asymmetric dimethylarginine^{20,23,24,25}.

6. What is the effect on Augmentation Index/Central aortic SBP?

Lercanidipine significantly reduces the Augmentation index, as well the aortic SBP and pulse pressure^{13,20}.

The increase in central aortic pressure is a marker of arterial stiffness and is involved in the development of cardiovascular events^{26,27,28}. Lercanidipine, improving arterial stiffness and decreasing aortic pressure, shows a cardiovascular protective effect in hypertensive patients.

7. What about the effect on sympathetic system?

Differently from felodipine and nifedipine, chronic administration of lercanidipine, does not induce sympathetic activation, and does not increase plasma norepinephrine level. This aspect has an important clinical relevance considering that, in hypertensive patients, sympathetic overdrive is associated with tachycardia and development of cardiovascular events^{29,30}.

8. What is the evidence of renal protection?

Hypertension remains a major risk factor for kidney disease³¹. Lercanidipine blocking both L and T channels, dilates afferent and efferent renal arteries, thus decreases intraglomerular pressure. The result shows that lercanidipine reduces chronic kidney disease progression³². This is evident considering that lercanidipine lowers microalbuminuria in patients with type 2 diabetes similarly to ramipril, as reported in the DIAL study³³. The improvement of renal function was also obtained in patients with chronic renal failure³⁴ and in subjects after renal artery intervention for atherosclerotic lesions³⁵ in whom lercanidipine significantly increased glomerular filtration rate, after 6 months of treatment and decreased proteinuria.

9. What are the pleiotropic effects?

Lercanidipine improving endothelial function and increasing NO bioavailability shows an atheroprotective effects. NO decreases oxidative stress, reduces vascular intimal and smooth muscle cell proliferation, decreases the plasma levels of E-selectin, P-selectin, adhesion molecules, inhibits cholesterol accumulation, LDL oxidation and platelet aggregation to the endothelium. Through these effects, lercanidipine decreases the risk of atherothrombotic events^{11,36,37}. Moreover, in hypertensive patients lercanidipine treatment is associated with a regression of microvascular structural changes, evaluated as wall-to-lumen ratio²⁰.

10. What about the tolerability?

Lercanidipine is well tolerated, with a very low rate of adverse events, such as dizziness, headache, flushing, palpitations, and vertigo.^{38, 39}. Particularly, compared with other CCBs (amlodipine, nifedipine, felodipine) lercanidipine, decreases ankle edema, by 56%^{16,40}. Chronic treatment does not change laboratory parameters^{4,41} and is associated with very low (2.1%–<1%) withdrawal rate^{10,42}.

Conflicts of interest

Gokhan Faikoglu and Kubra Saygisever-Faikoglu are employees of Recordati.

References

- Williams B, Mancia G, Spierin W et al 2018 ESC/ESH Guidelines for the management of arterial hypertension Eur Heart J 2018; 39:3021-3104 <https://doi.org/10.1093/eurheartj/ehy439>
- Meredith PA. Lercanidipine: a novel lipophilic dihydropyridine calcium antagonist with long duration of action and high vascular selectivity. Expert Opin Investig Drugs. 1999; 8:1043-1062 <https://doi.org/10.1517/13543784.8.7.1043>
- Bang LM, Chapman TM, Goa KL. Lercanidipine: a review of its efficacy in the management of hypertension. Drugs. 2003; 63:2449-2472 <https://doi.org/10.2165/00003495-200363220-00013>
- McClellan KJ, Jarvis B. Lercanidipine: a review of its use in hypertension. Drugs. 2000; 60:1123-1140 <https://doi.org/10.2165/00003495-200060050-00009>
- Cerbai E, Mugelli U Lercanidipine and T-type calcium current Eur Rev Med Pharmacol Sci 2018; 22:4025-4031
- Abe M, Okada K, Soma M. T-type Ca channel blockers in patients with chronic kidney disease in clinical practice Curr Hypertens Rev 2013; 9:202-9 <https://doi.org/10.2174/1573402110666140131155028>
- Rosenthal T, Rosenmann E, Tomassoni D et al Effect of Lercanidipine on Kidney Microanatomy in Cohen-Rosenthal Diabetic Hypertensive Rats Journal of Cardiovascular Pharmacology 2007; 12:145-152 <https://doi.org/10.1177/1074248407300621>
- Robles NR, Fici F, Grassi G Dihydropyridine calcium channel blockers and renal disease Hypertension Res 2017; 40:21-28 <https://doi.org/10.1038/hr.2016.85>
- Mancia G, Coca A, Chazova I et al Effects on office and home blood pressure of the lercanidipine-enalapril combination in patients with Stage 2 hypertension: a European randomized, controlled clinical trial Journal of Hypertension 2014, 32:1700-1707 <https://doi.org/10.1097/HJH.0000000000000239>
- Burnier M, Gasser UE. Efficacy and tolerability of lercanidipine in patients with hypertension: Results of a phase IV study in general practice. Expert Opin Pharmacother 2007; 8:2215-23 <https://doi.org/10.1517/14656566.8.14.2215>
- Grassi G, Robles N R, Seravalle G et al Lercanidipine in the Management of Hypertension: An Update J Pharmacol Pharmacother 2017; 8:155-65 https://doi.org/10.4103/jpp.JPP_34_17
- Millar-Craig M, Shaffu B, Greenough A Lercanidipine vs lacidipine in isolated systolic hypertension J Hum Hypertens 2003; 17:799-806 <https://doi.org/10.1038/sj.jhh.1001614>
- Mackenzie IS, McEnery CM, Dhakam Z, et al. Comparison of the effects of antihypertensive agents on central blood pressure and arterial stiffness in isolated systolic hypertension. Hypertension 2009; 54:409-13 <https://doi.org/10.1161/HYPERTENSIONAHA.109.133801>
- Viviani GL. Lercanidipine in type II diabetic patients with mild to moderate arterial hypertension. J Cardiovasc Pharmacol. 2002; 40:133-9 <https://doi.org/10.1097/00005344-200207000-00016>
- Barrios V, Escobar C, Navarro A et al Lercanidipine is an effective and well tolerated antihypertensive drug regardless the cardiovascular risk profile: the LAURA Study Int J Clin Pract 2006; 60:1364-70 <https://doi.org/10.1111/j.1742-1241.2006.01176.x>
- Makarounas-Kirchmann K, Glover-Koudounas S, Ferrari P. Results of a meta-analysis comparing the tolerability of lercanidipine with the 1st and 2nd generation dihydropyridine calcium channel blockers. Clin Ther. 2009; 31:1652-63 <https://doi.org/10.1016/j.clinthera.2009.08.010>
- James IG, Jones A, Davies P. A randomised, double-blind, double-dummy comparison of the efficacy and tolerability of lercanidipine tablets and losartan tablets in patients with mild to moderate essential hypertension. J Hum Hypertens. 2002; 16:605-10 <https://doi.org/10.1038/sj.jhh.1001430>
- Aranda P, Aranda FJ, Bianchi JL et al. Therapeutic efficacy and tolerability of lercanidipine versus candesartan, alone or in combination, in mild-moderate essential hypertensives. J Hypertens. 2000; 18-Suppl 2:S152 <https://doi.org/10.1097/00004872-200006001-00518>
- Omboni S, Zanchetti A. Antihypertensive efficacy of lercanidipine at 2.5, 5 and 10 mg in mild to moderate essential hypertensives assessed by clinic and ambulatory blood pressure measurements.

- Multicenter study investigators. *J Hypertens* 1998; 16:1831-8
<https://doi.org/10.1097/00004872-199816120-00017>
- 20 De Ciuceis C, Salvetti M, Rossini C et al Effect of antihypertensive treatment on microvascular structure, central blood pressure and oxidative stress in patients with mild essential hypertension. *J Hypertens* 2014; 32:565-574
<https://doi.org/10.1097/HJH.0000000000000067>
- 21 Poncelet P, Ribstein J, Goullard L, Bassous M, et al. Efficacy and acceptability of lercanidipine are not age dependent in patients with essential hypertension: The AGATE study. *Ann Cardiol Angeiol (Paris)* 2004; 53:123-30
<https://doi.org/10.1016/j.ancard.2004.03.004>
- 22 Panza JA, Casino PR, Kilcoyne CM et al Role of endothelium-derived nitric oxide in the abnormal endothelium-dependent vascular relaxation of patients with essential hypertension. *Circulation*. 1993; 87:1468-1474
<https://doi.org/10.1161/01.CIR.87.5.1468>
- 23 Taddei S, Virdis A, Ghiadoni L et al Calcium Antagonist Treatment by Lercanidipine Prevents Hyperpolarization in Essential Hypertension *Hypertension* 2003; 41:950-955
<https://doi.org/10.1161/01.HYP.0000063361.70525.3C>
- 24 Incandela L, Belcaro G M R Cesarone M R et al Oxygen-free radical decrease in hypertensive patients treated with lercanidipine *International Angiology* 2001;20: 136-140
- 25 Martinez M LL, Lopes L F, Coelho E B, et al. Lercanidipine reduces matrix metalloproteinase-9 activity in patients with hypertension *J Cardiovasc Pharmacol* 2006; 47:117-122
<https://doi.org/10.1097/01.fjc.0000196241.96759.71>
- 26 Chirinos JA, Segers P, Hughes T et al Large-Artery Stiffness in health and disease JACC State-of-the-Art Review *J Am. Coll. Cardiol* 2019; 74:1237-1263 <https://doi.org/10.1016/j.jacc.2019.07.012>
- 27 Roman MJ, Devereux RB, Kizer JR, et al. Central pressure more strongly relates to vascular disease and outcome than does brachial pressure: the Strong Heart Study. *Hypertension*. 2007; 50:197-203
<https://doi.org/10.1161/HYPERTENSIONAHA.107.089078>
- 28 Weber T, Auer J, O'Rourke MF, et al. Arterial stiffness, wave reflections, and the risk of coronary artery disease. *Circulation*. 2004; 109(2):184-189
<https://doi.org/10.1161/01.CIR.0000105767.94169.E3>
- 29 Grassi G, Seravalle G, Turri C, et al Short-versus long-term effects of different dihydropyridines on sympathetic and baroreflex function in hypertension. *Hypertension* 2003; 41:558-62
<https://doi.org/10.1161/01.HYP.0000058003.27729.5A>
- 30 Fogari R, Mugellini A, Zoppi A et al Differential effects of lercanidipine and nifedipine GITS on plasma norepinephrine in chronic treatment of hypertension *Am J Hypertens* 2003; 16:596-9
[https://doi.org/10.1016/S0895-7061\(03\)00901-4](https://doi.org/10.1016/S0895-7061(03)00901-4)
- 31 Robles NR, Fici F, Ari Bakir E, et al Does established vascular kidney disease exist? *J Clin Hypertens (Greenwich)*. 2020; 22:296-298
<https://doi.org/10.1111/jch.13818>
- 32 Robles NR, Fici F, Grassi G Dihydropyridine calcium channel blockers and renal Disease *Hypertension Res* 2017; 40:21-28
<https://doi.org/10.1038/hr.2016.85>
- 33 Dalla Vestra M, Pozza G, Mosca A, Grazioli V et al. Effect of lercanidipine compared with ramipril on albumin excretion rate in hypertensive type II diabetic patients with microalbuminuria: DIAL Study (diabete, ipertensione, albuminuria, lercanidipina). *Diab Nutr Metab* 2004; 17:259-266
- 34 Robles NR, Ocon J, Gomez CF et al Lercanidipine in patients with chronic renal failure: the ZAFRA study. *Ren Fail*. 2005; 27:73-80
<https://doi.org/10.1081/JDI-42801>
- 35 Peng M, Jiang XJ, Dong H, et al. Can lercanidipine improve renal function in patients with atherosclerotic renal artery stenosis undergoing renal artery intervention? *Curr Med Res Opin* 2015; 31:177-82 <https://doi.org/10.1185/03007995.2014.960071>
- 36 Harrison DG. Cellular and molecular mechanisms of endothelial cell dysfunction. *J Clin Invest*. 1997; 100:2153-2157
<https://doi.org/10.1172/JCI119751>
- 37 Rachmani R, Levi Z, Zadok BS, Ravid M. Losartan and lercanidipine attenuate low-density lipoprotein oxidation in patients with hypertension and type 2 diabetes mellitus: A randomized, prospective crossover study. *Clin Pharmacol Ther* 2002; 72:302-7
<https://doi.org/10.1067/mcp.2002.127110>
- 38 Barrios V, Navarro A, Esteras A, et al. Antihypertensive efficacy and tolerability of lercanidipine in daily clinical practice. The ELYPSE study. Eficacia de lercanidipino y su perfil de seguridad. *Blood Press* 2002; 11:95-100
<https://doi.org/10.1080/08037050211265>
- 39 Barrios V, Escobar C, de la Figuera M, et al. High doses of lercanidipine are better tolerated than other dihydropyridines in hypertensive patients with metabolic syndrome: Results from the TOLERANCE study. *Int J Clin Pract* 2008; 62:723-8
<https://doi.org/10.1111/j.1742-1241.2008.01736.x>
- 40 Makani H, Bangalore S, Romero J, et al Peripheral edema associated with calcium channel blockers: Incidence and withdrawal rate - A meta-analysis of randomized trials. *J Hypertens* 2011;29:1270-80
<https://doi.org/10.1097/HJH.0b013e3283472643>
- 41 Barrios V, Calderon A, Navarro A et al Lercanidipine effectiveness and tolerability profile is not influenced by overweight or body fat increase. The LERZAMIG study. *J Hypertens* 2004; 22 Suppl 2:S258-9 <https://doi.org/10.1097/00004872-200406002-00908>
- 42 Barrios V, Navarro A, Esteras A, Luque M, et al. Antihypertensive efficacy and tolerability of lercanidipine in daily clinical practice. The ELYPSE study. Eficacia de lercanidipino y su perfil de seguridad. *Blood Press* 2002; 11:95-100
<https://doi.org/10.1080/08037050211265>