

RIGIDITY AT INFINITY FOR LATTICES IN RANK-ONE LIE GROUPS

ALESSIO SAVINI

ABSTRACT. Let Γ be a non-uniform lattice in $PU(p, 1)$ without torsion and with $p \geq 2$. By following the approach developed in [FK06], we introduce the notion of volume for a representation $\rho : \Gamma \rightarrow PU(m, 1)$ where $m \geq p$. We use this notion to generalize the Mostow–Prasad rigidity theorem. More precisely, we show that given a sequence of representations $\rho_n : \Gamma \rightarrow PU(m, 1)$ such that $\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(M)$, then there must exist a sequence of elements $g_n \in PU(m, 1)$ such that the representations $g_n \circ \rho_n \circ g_n^{-1}$ converge to a reducible representation ρ_∞ which preserves a totally geodesic copy of $\mathbb{H}_{\mathbb{C}}^p$ and whose $\mathbb{H}_{\mathbb{C}}^p$ -component is conjugated to the standard lattice embedding $i : \Gamma \rightarrow PU(p, 1) < PU(m, 1)$. Additionally, we show that the same definitions and results can be adapted when Γ is a non-uniform lattice in $PSp(p, 1)$ without torsion and for representations $\rho : \Gamma \rightarrow Sp(m, 1)$, still maintaining the hypothesis $m \geq p \geq 2$.

1. INTRODUCTION

One of the most celebrated theorem about rigidity of lattices in semisimple Lie groups is the Mostow rigidity theorem which states that real hyperbolic lattices are strongly rigid. More precisely, let Γ_1 and Γ_2 in $PO^\circ(p, 1) = \text{Isom}^+(\mathbb{H}_{\mathbb{R}}^p)$ be two non-uniform lattices without torsion. If Γ_1 is isomorphic to Γ_2 , there exists $g \in PO(p, 1)$ such that $\Gamma_2 = g\Gamma_1g^{-1}$. Equivalently the p -dimensional complete hyperbolic manifolds $M_1 = \Gamma_1 \backslash \mathbb{H}_{\mathbb{R}}^p$ and $M_2 = \Gamma_2 \backslash \mathbb{H}_{\mathbb{R}}^p$ are isometric. Here $\mathbb{H}_{\mathbb{R}}^p$ denotes the real hyperbolic space of dimension p .

If we assume $\Gamma < PO^\circ(p, 1)$ and we look at representations $\rho : \Gamma \rightarrow PO(m, 1)$, the result above may be strengthened by introducing the notion of volume for representations $\rho : \Gamma \rightarrow PO(m, 1)$, where $m \geq p \geq 3$. For $m = p$ a way to define the volume of a representation ρ is based on the properties of the bounded cohomology of the group $PO^\circ(p, 1)$. In [BBI13] the authors prove that the volume class ω_p is a generator for the cohomology group $H_{cb}^p(PO^\circ(p, 1))$, hence, starting from it, we can construct a class in $H_b^p(\Gamma)$ by pulling back ω_p along ρ_b^* and then evaluate this class with a relative fundamental class $[N, \partial N] \in H^3(N, \partial N)$ via the Kronecker pairing. Here N is any compact core of $M = \Gamma \backslash \mathbb{H}_{\mathbb{R}}^p$.

Generalizing [BBI13], in [FK06] the authors define the volume for representations $\rho : \Gamma \rightarrow PO(m, 1)$ when $m \geq p$ by considering the infimum all over the volumes $\text{Vol}(f)$, where $f : \mathbb{H}_{\mathbb{R}}^p \rightarrow \mathbb{H}_{\mathbb{R}}^m$ is a properly ending smooth ρ -equivariant map. Intuitively a map f properly ends if it respects the action of the peripheral subgroups. The existence of such a map is proved in [FK06].

For any representation ρ the volume is invariant by conjugation and hence we have a well-defined function on the character variety $X(\Gamma, PO(m, 1))$ which is continuous with respect to the topology of pointwise convergence. This function is rigid, that is it satisfies $\text{Vol}(\rho) \leq \text{Vol}(M)$ and, if equality holds, ρ preserves a totally geodesic copy of $\mathbb{H}_{\mathbb{R}}^p$ and its $\mathbb{H}_{\mathbb{R}}^p$ -component is conjugated to the standard lattice embedding $i : \Gamma \rightarrow PO(p, 1) < PO(m, 1)$.

By generalizing both [CS83] and [MS84], in [Mor86] J. Morgan proposed a compactification of the variety $X(\Gamma, PO(m, 1))$ whose ideal points can be interpreted as projective length functions of isometric Γ -actions on real trees. We call this compactification the Morgan–Shalen compactification of $X(\Gamma, PO(m, 1))$. It is natural to ask if there exists a way to extend continuously the volume function to this compactification and which are the possible values attained at any ideal point. For instance, one could ask if it is possible to extend the rigidity of the volume function also at ideal points. In [FS] the authors answered to this question by proving a rigidity result which we may call strong rigidity at infinity. Let Γ be as above and let $\rho_n : \Gamma \rightarrow PO(m, 1)$ be a sequence of representations such that $\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(M)$. There must exist a sequence of elements $g_n \in PO(m, 1)$ such that the sequence $g_n \circ \rho_n \circ g_n^{-1}$ converges to a representation ρ_∞ which preserves a totally geodesic copy of $\mathbb{H}_{\mathbb{R}}^p$ and whose $\mathbb{H}_{\mathbb{R}}^p$ -component is conjugated to the standard lattice embedding $i : \Gamma \rightarrow PO(p, 1) < PO(m, 1)$. As a consequence the sequence of representations cannot diverge to an ideal point of the character variety $X(\Gamma, PO(m, 1))$, or equivalently if $\rho_n : \Gamma \rightarrow PO(m, 1)$ is a sequence of representations converging to any ideal point of the Morgan–Shalen compactification of $X(\Gamma, PO(m, 1))$, then the sequence of volumes $\text{Vol}(\rho_n)$ must be bounded from above by $\text{Vol}(M) - \varepsilon$ with a suitable $\varepsilon > 0$.

The proof of this theorem is based essentially on the properties of the so-called BCG–natural map associated to a non-elementary representation $\rho : \Gamma \rightarrow PO(m, 1)$, described in [BCG95], [BCG96] and [BCG99]. Given such a representation there exists a map $F : \mathbb{H}_{\mathbb{R}}^p \rightarrow \mathbb{H}_{\mathbb{R}}^m$ which is equivariant with respect to ρ , smooth and satisfies $Jac_p(F) \leq 1$ for every $x \in \mathbb{H}_{\mathbb{R}}^p$, where $Jac_p(F)$ is the p -Jacobian of the map F . Moreover, the equality holds if and only if $D_x F$ is an isometry.

However the construction of the BCG–natural map is much more general. Let Γ be a non-uniform lattice in G_p without torsion, with either $G_p = PU(p, 1)$ or $G_p = PSp(p, 1)$. We say that the lattice

Γ is complex in the former case, quaternionic in the latter. Given a representation of $\rho : \Gamma \rightarrow G_m$, where $G_m = PU(m, 1)$ if Γ is complex or $G_m = PSp(m, 1)$ if Γ is quaternionic, we can adapt the procedure described by both [BCG99] and [Fra09] to obtain a natural map which satisfies the same properties listed previously.

To prove the strong rigidity at infinity in both the complex and the quaternionic case we introduce the notion of volume for representations $\rho : \Gamma \rightarrow G_m$, with $m \geq p$. For uniform complex lattices the definition of volume for representations $\rho : \Gamma \rightarrow PU(m, 1)$ is given both by [BCG99] and by [BCG07], whereas for non-uniform complex lattices we refer to [BI01] and to [KM08]. We find another interesting approach in [KK12], where the authors use the pairing between bounded cohomology and l^1 -Lipschitz homology to define the volume of a representation. However, here we give a different version of volume to adapt this notion to the non compact case, also for quaternionic lattices. By following the strategy of the proof of [FS, Theorem 1] we get

Theorem 1.1. *Denote by G_p either $PU(p, 1)$ or $PSp(p, 1)$ and let X^p the globally symmetric space associated to G_p . Let Γ be a non-uniform lattice in G_p without torsion. Assume $p \geq 2$. Let $\rho_n : \Gamma \rightarrow G_m$ be a sequence of representations with $m \geq p$. If $\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(\Gamma \backslash X^p)$, then there must exist a sequence of elements $g_n \in G_m$ such that the sequence $g_n \circ \rho_n \circ g_n^{-1}$ converges to a representation ρ_∞ which preserves a totally geodesic copy of X^p and whose X^p -component is conjugated to the standard lattice embedding $i : \Gamma \rightarrow G_p < G_m$.*

We want to stress that the quaternionic case could be obtained as a direct consequence of [Cor92], but we reported it here for sake of completeness and to show how we can get it using natural maps.

The first section is dedicated to preliminary definitions. Let G_p and X^p be as in the theorem above. We briefly recall the notion of barycentre of a positive Borel measure on $\partial_\infty X^p$ and the definition of natural map F associated to a non-elementary representation $\rho : \Gamma \rightarrow G_m$. We introduce the definition of volume for representations $\rho : \Gamma \rightarrow G_m$ and we compare it with the volume of the ε -natural maps F^ε . These maps are smooth, ρ -equivariant and converge to F with respect to the C^1 -topology, as in [FK06]. The second section is devoted to the proof of the main theorem. We conclude with some comments and remarks about the main results.

Acknowledgements: I would like to thank Marc Burger, Alessandra Iozzi and Michelle Bucher for the precious help and the enlightening conversations. I am also very grateful to Stefano Francaviglia for having introduced me to the study of BCG natural maps.

I finally thank the referee for all the comments and suggestions.

2. PRELIMINARY DEFINITIONS

2.1. **Barycentre of a measure.** We start by fixing some notation. Let G_p be either $PU(p, 1)$ or $PSp(p, 1)$. Denote by $\mathfrak{g}_p = T_e G_p$ the tangent space to G_p at the neutral element. If we endow \mathfrak{g}_p with its natural structure of Lie algebra, we recall that \mathfrak{g}_p admits an involution $\Theta : \mathfrak{g}_p \rightarrow \mathfrak{g}_p$ which allows us to decompose $\mathfrak{g}_p = \mathfrak{l} \oplus \mathfrak{p}$, where \mathfrak{l} and \mathfrak{p} are the eigenspaces with respect to 1 and -1 of the involution Θ . Moreover \mathfrak{p} is naturally identified with the tangent space at the point fixed by $\exp(\mathfrak{l})$ of the symmetric space X^p associated to G_p and the restriction of the Killing form to \mathfrak{p} is positive definite. Via the group action we can transport this scalar product to any other tangent space of X^p and hence we get in a canonical way a Riemannian metric on X^p . If $G_p = PU(p, 1)$ then the associated symmetric space X^p is the complex hyperbolic space of rank p , namely $\mathbb{H}_{\mathbb{C}}^p$, whereas if $G_p = PSp(p, 1)$ the symmetric space X^p coincides with the quaternionic hyperbolic space of rank p , that is $\mathbb{H}_{\mathbb{H}}^p$. In both cases we recall that the sectional curvature of these spaces lies between -4 and -1 . In particular, since X^p is negatively curved, we can talk about the visual boundary of X^p and we denote it by $\partial_{\infty} X^p$.

From now until the end of the paper we are going to fix a point in X^p as basepoint and we are going to denote it by O . Moreover, we will use the same letter O to denote basepoints in symmetric spaces of different dimension. Let $B_O(x, \theta)$ be the Busemann function of X^p normalized at O , that means for every $\theta \in \partial_{\infty} X^p$ we set

$$B_O(x, \theta) = \lim_{t \rightarrow \infty} d(x, c(t)) - t,$$

where c is the geodesic ray starting at $O = c(0)$ and ending at $\theta = c(\infty)$.

Let β be a positive probability measure on $\partial_{\infty} X^p$. Thanks to the convexity of Busemann functions the map

$$\varphi_{\beta} : X^p \rightarrow \mathbb{R}, \quad \varphi_{\beta}(y) := \int_{\partial_{\infty} X^p} B_O(y, \theta) d\beta(\theta)$$

is strictly convex, if we assume that β is not the sum of two Dirac measures. Additionally, if the measure β does not contain any atom of mass greater or equal than $1/2$, the following condition holds

$$\lim_{y \rightarrow \partial_{\infty} X^p} \varphi_{\beta}(y) = \infty.$$

and this implies that φ_{β} admits a unique minimum in X^p (see [BCG95, Appendix A]). On the other hand, if β contains an atom of mass at least $1/2$, then it is easy to check that the minimum of φ_{β} is $-\infty$ and it is attained when y coincides with the atom.

Definition 2.1. Let β be any positive probability measure on the visual boundary $\partial_{\infty} X^p$ which is not the sum of two Dirac masses with

the same weight. If β contains an atom x of mass greater or equal than $1/2$ then we define its *barycentre* as

$$\text{bar}_{\mathcal{B}}(\beta) = x,$$

otherwise we define it as the point

$$\text{bar}_{\mathcal{B}}(\beta) = \text{argmin}(\varphi_{\beta}).$$

The letter \mathcal{B} emphasizes the dependence of the construction on the Busemann functions. The barycentre of β will be a point in $\overline{X^p}$ which satisfies the following properties:

- it is continuous with respect to the weak-* topology on the set of probability measures on $\partial_{\infty}X^p$, that is if $\beta_n \rightarrow \beta$ in the weak-* topology (and no measure is the sum of two atoms with equal weight) it holds

$$\lim_{n \rightarrow \infty} \text{bar}_{\mathcal{B}}(\beta_n) = \text{bar}_{\mathcal{B}}(\beta)$$

- it is G_p -equivariant, indeed for every $g \in G_p$ (if β is not the sum of two equal atoms) we have

$$\text{bar}_{\mathcal{B}}(g_*\beta) = g(\text{bar}_{\mathcal{B}}(\beta)),$$

- when β does not contain any atom of weight greater or equal than $1/2$, it is characterized by the following equation

$$(1) \quad \int_{\partial_{\infty}X^p} dB_O|_{(\text{bar}_{\mathcal{B}}(\beta), y)}(\cdot) d\beta(y) = 0.$$

We refer to [BCG95, Appendix A] for the equation above.

2.2. The Patterson-Sullivan family of measures and the BCG–natural map. For more details about the following definitions and constructions we recommend the reader see [BCG95], [BCG99] and [Fra09]. Even if the last one refers only to real hyperbolic lattices, every definition that appears can be easily adapted in our context, as we will see. Before starting, fix $k = 2p$ if $G_p = PU(p, 1)$ and $k = 4p$ if $G_p = PSp(p, 1)$. The value k is simply the real dimension of the symmetric space X^p associated to G_p .

Let Γ be a non-uniform lattice in G_p without torsion, that is a discrete subgroup of G_p which can be thought of as the fundamental group of a complete manifold $M = X^p/\Gamma$ with finite volume and which is not compact. We say that Γ is a complex lattice if $\Gamma < PU(p, 1)$, whereas we call it quaternionic if $\Gamma < PSp(p, 1)$.

Definition 2.2. The *critical exponent* associated to the lattice Γ is defined as

$$\delta(\Gamma) = \inf\{s \in [0, \infty] \mid \sum_{\gamma \in \Gamma} e^{-sd(x, \gamma x)} < \infty\}$$

for any point $x \in X^p$. The definition above does not depend on the choice of the basepoint $x \in X^p$ used to compute the series.

When Γ is a non-uniform lattice in G_p , the critical exponent is always finite and by [Alb97, Theorem 2] we have that $\delta(\Gamma) = k + d - 2$. The number d is the real dimension of the algebra on which the hyperbolic space X^p is defined. Hence we have either $d = 2$ if Γ is complex or $d = 4$ if Γ is quaternionic. Moreover, we remind the reader that for $s = \delta(\Gamma)$ the series diverges by [Alb99, Proposition D], that is

$$\sum_{\gamma \in \Gamma} e^{-\delta(\Gamma)d(x, \gamma x)} = +\infty.$$

and for this reason we may refer to Γ as a lattice of divergence type.

Definition 2.3. Let $\mathcal{M}^1(Y)$ be the set of positive probability measures on a space Y . The *family of Patterson-Sullivan measures associated to a non-uniform lattice* Γ is a family of measures $\{\mu_x\} \in \mathcal{M}^1(\partial_\infty X^p)$, where $x \in X^p$, which satisfies the following properties

- the family is Γ -equivariant, that is $\mu_{\gamma x} = \gamma_*(\mu_x)$ for every $\gamma \in \Gamma$ and every $x \in X^p$,
- For every $x, y \in X^p$ it holds

$$d\mu_x(\theta) = e^{-\delta(\Gamma)B_y(x, \theta)} d\mu_y(\theta)$$

where $B_y(x, \theta)$ is the Busemann function normalized at y .

Remark 2.4. The construction of the family of Patterson-Sullivan measures has been generalized by [Alb97, Alb99] to any lattice in a Lie group G of non-compact type. The support of the measures μ_x is the Furstenberg boundary $\partial_{\mathcal{F}} X$ of X , which can be thought of as the G -orbit of a regular point $\xi \in \partial_\infty X$. Since we are considering rank one Lie group and

$$\text{codim}_{\partial_\infty X} \partial_{\mathcal{F}} X = \text{rank}(X) - 1$$

we have that $\partial_\infty X = \partial_{\mathcal{F}} X$ in our context.

Let Γ be as above and let $\{\mu_x\}$ be the family of Patterson-Sullivan measures associated to Γ . We set $\mu = \mu_O$.

Let $\rho : \Gamma \rightarrow G_m$ be a non-elementary representation. We are assuming that either $G_m = PU(m, 1)$ if Γ is complex or $G_m = PSp(m, 1)$ if Γ is quaternionic. Recall that the action of Γ on $(\partial_\infty X^p \times \partial_\infty X^p, \mu \times \mu)$ is ergodic by [Nic89, Yue96, BM96, Rob00], for instance. Hence by [BM96, Corollary 3.2] there exists a ρ -equivariant measurable map

$$D : \partial_\infty X^p \rightarrow \partial_\infty X^m$$

and two different maps of this type must agree on a full μ -measure set.

We define

$$\nu_x := D_*(\mu_x).$$

Clearly the measure ν_x lives in $\mathcal{M}^1(\partial_\infty X^m)$ for every x .

Since we have a non-elementary representation, ν_x does not contain any atom of mass greater or equal than $1/2$. Indeed the following holds

Lemma 2.5. *Let $\rho : \Gamma \rightarrow G_m$ be a non-elementary representation and let $D : \partial_\infty X^p \rightarrow \partial_\infty X^m$ be a ρ -equivariant measurable map. Then $D(x) \neq D(y)$ for almost every $(x, y) \in \partial_\infty X^p \times \partial_\infty X^p$.*

Proof. Define the set $A := \{(x, y) \in \partial_\infty X^p \times \partial_\infty X^p \mid D(x) = D(y)\}$. Since the map D is ρ -equivariant, A is a Γ -invariant measurable subset of $\partial_\infty X^p \times \partial_\infty X^p$. Recall that Γ acts ergodically on $\partial_\infty X^p \times \partial_\infty X^p$ with respect to the measure $\mu \times \mu$. In particular, the set A must have either null measure or full measure. By contradiction, suppose that A has full measure. This implies that for almost all x , the slice $A(x) := \{y \in \partial_\infty X^p \mid D(x) = D(y)\}$ has full measure in $\partial_\infty X^p$. The G_m -action preserves the class of μ , in particular, for any $\gamma \in \Gamma$, if $A(x)$ has full measure then so does $\gamma A(x)$. Since Γ is countable, this implies that for almost all x , the set $A_\Gamma(x) := \bigcap_{\gamma \in \Gamma} \gamma^{-1} A(x)$ has full measure in $\partial_\infty X^p$. Fix now a point $y \in A_\Gamma(x)$. For any $\gamma \in \Gamma$ we have $(x, \gamma y) \in A$. In particular

$$D(y) = D(x) = D(\gamma y) = \rho(\gamma)D(y)$$

for every $\gamma \in \Gamma$. We use $\gamma = id$ in the first equality and the last follows by equivariance of D .

The computation above would imply that ρ is elementary, which is a contradiction. \square

By the previous lemma, for all $x \in X^p$, we can define

$$F(x) := \text{bar}_{\mathcal{B}}(\nu_x)$$

and this point will lie in X^m . In this way we get a map $F : X^p \rightarrow X^m$.

Definition 2.6. The map $F : X^p \rightarrow X^m$ is called *natural map* for the representation $\rho : \Gamma \rightarrow G_m$.

Equation (1) becomes

$$(2) \quad \int_{\partial_\infty X^m} dB_O|_{(F(x), y)}(\cdot) d\nu_x(y) = 0.$$

and since $\nu_x = D_*(\mu_x)$, it can be rewritten as

$$(3) \quad \int_{\partial_\infty X^p} dB_O|_{(F(x), D(z))}(\cdot) d\mu_x(z) = 0.$$

The natural map is smooth and satisfies the following properties:

- Recall that k is the real dimension of the symmetric space X^p . Define the k -Jacobian of F as

$$Jac_k(F)(x) := \max_{u_1, \dots, u_k \in T_x X^p} \|D_x F(u_1) \wedge \dots \wedge D_x F(u_k)\|_{X^m}$$

where $\{u_1, \dots, u_k\}$ is an orthonormal frame of the tangent space $T_x X^p$ with respect to the standard metric induced by g_{X^p} and the norm $\|\cdot\|_{X^m}$ is the norm on $T_{F(x)} X^m$ induced by g_{X^m} . We have $Jac_k(F) \leq 1$ and the equality holds at x if and only if $D_x F : T_x X^p \rightarrow T_{F(x)} X^m$ is an isometry (see for instance [BCG99, Theorem 1.10]).

- The map F is ρ -equivariant, that is $F(\gamma x) = \rho(\gamma)F(x)$.
- By differentiating (3), one gets that for all $x \in X^p$, $u \in T_x X^p$, $v \in T_{F(x)} X^m$ it holds that

$$(4) \quad \int_{\partial_\infty X^p} \nabla dB_O|_{(F(x), D(z))}(D_x F(u), v) d\mu_x(z) = \delta(\Gamma) \int_{\partial_\infty X^p} dB_O|_{(F(x), D(z))}(v) dB_O|_{(x, z)}(u) d\mu_x(z)$$

where ∇ is the Levi-Civita connection on X^m .

2.3. Volume of representations and ε -natural maps. Let Γ be a non-uniform lattice in G_p without torsion. If we denote by $M = X^p/\Gamma$ we obtain a complete manifold of finite volume which is locally symmetric X^p and not compact. Moreover, as a consequence of the Margulis lemma, it admits a decomposition

$$M := N \cup \bigcup_{i=1}^h C_i$$

where N is a compact core of finite volume and each C_i is a cuspidal neighborhood which can be seen as $N_i \times (0, \infty)$ where $\pi_1(N_i)$ is a discrete virtually nilpotent parabolic subgroup of G_p (see [BGS85] or [Bow95]).

As before, we set $k = 2p$ if $G_p = PU(p, 1)$ or $k = 4p$ if $G_p = PSp(p, 1)$. Let $\rho : \Gamma \rightarrow G_m$ be a representation, with $m \geq p$, and let $f : X^p \rightarrow X^m$ be a smooth ρ -equivariant map. By following the definition of [FK06] we want to define the volume $\text{Vol}(f)$. Let g_{X^m} be the natural metric on X^m . The pullback of g_{X^m} along f defines in a natural way a pseudo-metric on X^p , which can be possibly degenerate, and hence it defines a natural k -form given by $\tilde{\omega}_D = \sqrt{|\det f^* g_{X^m}|}$. The equivariance of f with respect to ρ implies that the form $\tilde{\omega}_f$ is Γ -invariant and hence it determines a k -form on M . Denote this form by ω_f .

Definition 2.7. Let $\rho : \Gamma \rightarrow G_m$ be a representation and let $f : X^p \rightarrow X^m$ be any smooth ρ -equivariant map. The *volume* of f is defined as

$$\text{Vol}(f) := \int_M \omega_f$$

We keep denoting by $f : X^p \rightarrow X^m$ a generic smooth ρ -equivariant map. For each cuspidal neighborhood $C_i = N_i \times (0, \infty)$, we know that $\pi_1(N_i)$ is parabolic, so it fixes a unique point in $\partial_\infty X^p$. Suppose $c_i = \text{Fix}(\pi_1 N_i)$ and let $r(t)$ be a geodesic ray ending at c_i . We say that f is a *properly ending map* if all the limit points of $f(r(t))$ lie either in $\text{Fix}(\rho(\pi_1 N_i))$ or in a finite union of $\rho(\pi_1 N_i)$ -invariant geodesics.

Definition 2.8. Given a representation $\rho : \Gamma \rightarrow G_m$, we define its *volume* as

$$\text{Vol}(\rho) := \inf\{\text{Vol}(f) \mid f \text{ is smooth, } \rho\text{-equivariant and properly ending}\}.$$

When ρ is non-elementary, a priori the BCG-natural map $F : X^p \rightarrow X^m$ associated to ρ is not a properly ending map, hence we cannot compare its volume with the volume of representation ρ . However, by adapting the proofs contained in [FK06], for any $\varepsilon > 0$ it is possible to construct a family of smooth functions $F^\varepsilon : X^p \rightarrow X^m$ that C^1 -converges to F as $\varepsilon \rightarrow 0$ and such that F^ε is a properly ending map for every $\varepsilon > 0$.

Definition 2.9. For any $\varepsilon > 0$ there exists a map $F^\varepsilon : X^p \rightarrow X^m$ called ε -*natural map* associated to ρ which satisfies the following properties

- F^ε is smooth and ρ -equivariant,
- at every point of X^p we have $\text{Jac}_k(F^\varepsilon) \leq 1 + \varepsilon$,
- for every $x \in X^p$ it holds $\lim_{\varepsilon \rightarrow 0} F^\varepsilon(x) = F(x)$ and $\lim_{\varepsilon \rightarrow 0} D_x F^\varepsilon = D_x F$,
- F^ε is a properly ending map.

The existence of ε -natural maps for real hyperbolic lattices is proved in [FK06, Lemma 4.5], but the proof can be suitably adapted both to the complex and the quaternionic case.

Remark 2.10. The properly ending property of F^ε is guaranteed by the fact that $\pi_1(N_i)$ is parabolic and stabilizes each horosphere through the fixed point c_i . In particular, since F^ε is a properly ending map, it holds by definition that

$$\text{Vol}(\rho) \leq \int_M \sqrt{|\det((F^\varepsilon)^* g_{X^m})|}.$$

We are going to use the previous estimate later.

The rigidity of volume holds also in this context. Indeed the same proof of [FK06] leads us to

Theorem 2.11. *Denote by G_p either $PU(p, 1)$ or $PSp(p, 1)$ and let X^p be the globally symmetric space associated to G_p . Let Γ be a non-uniform lattice in G_p without torsion and assume $p \geq 2$. Let $\rho : \Gamma \rightarrow G_m$ be a representation, where $m \geq p$. Then $\text{Vol}(\rho) \leq \text{Vol}(M)$ and equality holds if and only if the representation ρ is a discrete and faithful representation of Γ into the isometry group of a totally geodesic copy of X^p contained in X^m .*

For complex lattices, this result is exactly the one obtained in [BI01] or in [KK12]. The statement regarding the quaternionic case is compatible with the result obtained in [Cor92].

3. PROOF OF THE MAIN THEOREM

We start by fixing the following setting.

- A lattice $\Gamma < G_p$ where $G_p = PU(p, 1)$ or $G_p = PSp(p, 1)$ so that $M := X^p/\Gamma$ is a (non-compact) complete manifold of finite volume. Recall that X^p is the Riemannian symmetric space associated to G_p . Assume $p \geq 2$.
- A base-point $O \in X^p$ used to normalize the Busemann function $B_O(x, \theta)$, with $x \in X^p$ and $\theta \in \partial_\infty X^p$.
- The family $\{\mu_x\}$ of Patterson-Sullivan probability measures associated to Γ . Set $\mu = \mu_O$.
- A sequence of representations $\rho_n : \Gamma \rightarrow G_m$ such that $\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(M)$.

We start observing that the condition $\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(M)$ implies that, up to passing to a subsequence, we can suppose that no ρ_n is elementary. Indeed elementary representations have zero volume and $\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(M)$, which is strictly positive. With an abuse of notation we still denote the subsequence of non-elementary representations by ρ_n .

Since no ρ_n is elementary, we can consider the sequence of ρ_n -equivariant measurable maps $D_n : \partial_\infty X^p \rightarrow \partial_\infty X^m$ and the corresponding sequence of BCG-natural maps $F_n : X^p \rightarrow X^m$. Since conjugating ρ_n by $g \in G_m$ reflects in post-composing F_n with g , up to conjugating ρ_n by a suitable element $g_n \in G_m$, we can suppose $F_n(O) = O$ for every $n \in \mathbb{N}$.

The choice to fix the origin of X^m as the image of $F_n(O)$ is made to avoid pathological behaviour. For instance consider a sequence of hyperbolic elements $g_n \in G_m$ which is divergent and define the representations $\rho_n := g_n \circ i \circ g_n^{-1}$, where $i : \Gamma \rightarrow G_p < G_m$ is the standard lattice embedding. Clearly this sequence of representations satisfies $\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(M)$, since for every $n \in \mathbb{N}$ we have $\text{Vol}(\rho_n) = \text{Vol}(M)$. However, there does not exist any subsequence of ρ_n converging to the holonomy of the manifold M .

Definition 3.1. For any $n \in \mathbb{N}$ and every $x \in X^p$ we can define the following quadratic forms on $T_{F_n(x)}X^m$:

$$k_n|_{F_n(x)}(u, u) := \langle K_n|_{F_n(x)}u, u \rangle = \int_{\partial_\infty X^p} \nabla dB_O|_{(F_n(x), D_n(\theta))}(u, u) d\mu_x(\theta)$$

$$h_n|_{F_n(x)}(u, u) := \langle H_n|_{F_n(x)}u, u \rangle = \int_{\partial_\infty X^p} (dB_O|_{(F_n(x), D_n(\theta))}(u))^2 d\mu_x(\theta)$$

for any $u \in T_{F_n(x)}X^m$. The notation $\langle \cdot, \cdot \rangle$ stands for the scalar product on $T_{F_n(x)}X^m$ induced by the natural metric on X^m . Since the rank m is bigger than p , we will need to define another quadratic form, this time on $T_x X^p$. For any $v \in T_x X^p$, we define

$$h'_n|_x(v, v) = \langle H'_n|_x v, v \rangle = \int_{\partial_\infty X^p} (dB_O|_{(x, \theta)}(v))^2 d\mu_x(\theta).$$

For any quadratic form we are going to drop the subscript which refers to the tangent space on which the form is defined. Since

$$Jac_k(F_n)(x) := \max_{u_1, \dots, u_k \in T_x X^p} \|D_x F_n(u_1) \wedge \dots \wedge D_x F_n(u_k)\|_{X^m},$$

let $\{u_1, \dots, u_k\}$ be any frame which realizes the maximum and denote by U_x the subspace $U_x := \text{span}_{\mathbb{R}}\{u_1, \dots, u_k\}$ of $T_x X^p$ (since we are working with k -tuples, the subspace U_x coincides exactly with $T_x X^p$, but we prefer to maintain the same notation of [BCG99]). Set $V_{F_n(x)} := D_x F_n(U_x)$. We denote by $K_n^V(x)$, $H_n^V(x)$ and $H_n'^U(x)$ the restrictions of the form $K_n|_{F_n(x)}$, $H_n|_{F_n(x)}$ and $H'_n|_x$ to the subspace $V_{F_n(x)}$, $V_{F_n(x)}$ and U_x , respectively. As consequence of the Cauchy–Schwarz inequality, as in [BCG99, Section 2] it follows that

$$\begin{aligned} & \det(K_n(x)^V) Jac_k(F_n)(x) \\ & \leq (k + d - 2)^k (\det(H_n^V(x)))^{\frac{1}{2}} (\det(H_n'^U(x)))^{\frac{1}{2}} \\ & \leq (k + d - 2)^k (\det(H_n^V(x)))^{\frac{1}{2}} (\text{Tr}(H_n'^U(x))/k)^{\frac{1}{2}} \\ & \leq k^{-\frac{k}{2}} (k + d - 2)^k (\det(H_n^V(x)))^{\frac{1}{2}} \end{aligned}$$

Lemma 3.2. *Suppose $\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(M)$. Then we have that $Jac_k(F_n)$ converges to 1 almost everywhere in X^p with respect to the measure induced by the standard metric.*

Proof. Denote by $F_n^\varepsilon : X^p \rightarrow X^m$ the ε -natural maps introduced in Section 2.3. Recall that we have the following estimate

$$\text{Vol}(\rho_n) \leq \text{Vol}(F_n^\varepsilon) = \int_M Jac_k(F_n^\varepsilon) d\text{vol}_{X^p}(x)$$

and since $Jac_k(F_n) \leq 1 + \varepsilon$ and $\lim_{\varepsilon \rightarrow 0} D_x F_n^\varepsilon = D_x F_n$, by the dominated convergence theorem we get

$$\text{Vol}(\rho_n) \leq \int_M Jac_k(F_n) d\text{vol}_{X^p}(x) \leq \text{Vol}(M)$$

from which it follows the statement. \square

By the previous lemma we have $\lim_{n \rightarrow \infty} Jac_k(F_n) = 1$ almost-everywhere on X^p with respect to the measure induced by the standard volume form. If \mathcal{N} is the set of zero measure outside of which $Jac_k(F_n)$ converges, for every $x \in X^p \setminus \mathcal{N}$ and fixed $\varepsilon > 0$ there must exist $n_0 = n_0(\varepsilon, x)$ such that $Jac_k(F_n) \geq 1 - \varepsilon$ for every $n > n_0$. Thus it holds

$$\left(\frac{(k+d-2)^2}{k} \right)^{\frac{k}{2}} \frac{\det(H_n^V)^{\frac{1}{2}}}{\det(K_n^V)} > 1 - \varepsilon$$

from which we can deduce

$$\frac{\det(H_n^V)}{(\det(K_n^V))^2} > \left(\frac{k}{(k+d-2)^2} \right)^k (1 - \varepsilon)^2 > \left(\frac{k}{(k+d-2)^2} \right)^k (1 - 2\varepsilon).$$

Moreover, since X^p has sectional curvature which varies between -4 and -1 , we can write $K_n^V = I - H_n^V - \sum_{i=1}^{d-1} J_i H_n^V J_i$, where $J_i(x)$ are orthogonal endomorphisms used to define the complex or the quaternionic structure on $T_{F_n(x)} X^m$ (see [BCG95, Section 5.b]). Recall that $J_i^2 = -I$ at every point. Here I stands for the identity on $T_{F_n(x)} X^m$. Hence, by substituting the expression of K_n in the previous inequality, we get

$$(5) \quad \frac{\det(H_n^V)}{(\det(I - H_n^V - \sum_{i=1}^{d-1} J_i H_n^V J_i))^2} > \left(\frac{k}{(k+d-2)^2} \right)^k (1 - 2\varepsilon).$$

Consider now the set of positive definite symmetric matrices of dimension k with real entries and trace equal to 1, namely

$$\text{Sym}_1^+(k, \mathbb{R}) := \{H \in \text{Sym}(k, \mathbb{R}) \mid H > 0, \text{Tr}(H) = 1\}.$$

Once we have fixed a basis of $V_{F_n(x)}$, we can identify H_n^V , K_n^V and J_i with the matrices representing these endomorphisms with respect to the fixed basis. Under this assumption, recall that $H_n \in \text{Sym}_1^+(k, \mathbb{R})$ for every $n \in \mathbb{N}$, as shown in [BCG95, Proposition B.1]. If we define

$$\varphi : \text{Sym}_1^+(k, \mathbb{R}) \rightarrow \mathbb{R}, \quad \varphi(H) := \frac{\det(H)}{(\det(I - H - \sum_{i=1}^{d-1} J_i H J_i))^2},$$

we know that

$$\varphi(H) \leq \left(\frac{k}{(k+d-2)^2} \right)^k$$

and equality holds if and only if $H = I/k$ (see [BCG95, Appendix B]). It is worth noticing that the space $Sym_1^+(k, \mathbb{R})$ is not compact and a priori there could exist a divergent sequence of elements $H_n \in Sym_1^+(k, \mathbb{R})$ such that

$$\lim_{n \rightarrow \infty} \varphi(H_n) = \left(\frac{k}{(k+d-2)^2} \right)^k.$$

We are going to show that this is impossible.

Proposition 3.3. *Suppose $H_n \in Sym_1^+(k, \mathbb{R})$ is sequence such that*

$$\lim_{n \rightarrow \infty} \varphi(H_n) = \left(\frac{k}{(k+d-2)^2} \right)^k.$$

Then the sequence H_n must converge to I/k .

Proof. We are not going to work directly on the function φ but we will use the auxiliary function

$$\psi(H) := \frac{(k-1)^{\frac{2k(k-1)}{k+d-2}} \det(H)^{\frac{k-d}{k+d-2}}}{(k+d-2)^{2k} \det(I-H)^{\frac{2(k-1)}{k+d-2}}}.$$

By [BCG95, Lemme B.3], for every $H \in Sym_1^+(k, \mathbb{R})$ we have that $\varphi(H) \leq \psi(H)$. Moreover both functions attain the same maximum value

$$\max_{H \in Sym_1^+(k, \mathbb{R})} \varphi = \max_{H \in Sym_1^+(k, \mathbb{R})} \psi = \left(\frac{k}{(k+d-2)^2} \right)^k$$

at $H = I/k$.

We are going to study the properties of the function ψ . We start by observing that the function ψ is invariant by conjugation by an element $g \in GL(k, \mathbb{R})$. Indeed, $\psi(H)$ can be expressed as

$$\psi(H) = \frac{(k-1)^{\frac{2k(k-1)}{k+d-2}} p_H(0)^{\frac{k-d}{k+d-2}}}{(k+d-2)^{2k} p_H(1)^{\frac{2(k-1)}{k+d-2}}},$$

where p_H is the characteristic polynomial of H . Hence the claim follows. In particular, we have an induced function

$$\tilde{\psi} : O(k, \mathbb{R}) \backslash Sym_1^+(k, \mathbb{R}) \rightarrow \mathbb{R}, \quad \tilde{\psi}(\bar{H}) = \psi(H),$$

where \bar{H} denotes the equivalence class of the matrix H and the orthogonal group $O(k, \mathbb{R})$ acts on $Sym_1^+(k, \mathbb{R})$ by conjugation. We can think of the space $O(k, \mathbb{R}) \backslash Sym_1^+(k, \mathbb{R})$ as the interior $\bar{\Delta}_{k-1}$ of the standard $(k-1)$ -simplex quotiented by the action of the symmetric group \mathfrak{S}_k

which permutes the coordinates of an element $(a_1, \dots, a_k) \in \mathring{\Delta}_{k-1}$. An explicit homeomorphism between the two spaces is given by

$\Lambda : O(k, \mathbb{R}) \backslash Sym_1^+(k, \mathbb{R}) \rightarrow \mathfrak{S}_k \backslash \mathring{\Delta}_{k-1}$, $\Lambda(\bar{H}) := [a_1(H), \dots, a_k(H)]$, where $a_i(H)$ for $i = 1, \dots, k$ are the eigenvalues of H . By defining $\Psi = \psi \circ \Lambda^{-1}$, we can express this function as

$$\Psi : \mathfrak{S}_k \backslash \mathring{\Delta}_{k-1} \rightarrow \mathbb{R}, \quad \Psi([a_1, \dots, a_k]) = \frac{(k-1)^{\frac{2k(k-1)}{k+d-2}}}{(k+d-2)^{2k}} \prod_{i=1}^k \frac{(a_i)^{\frac{k-d}{k+d-2}}}{(1-a_i)^{\frac{2(k-1)}{k+d-2}}}.$$

We are going to think of Ψ as defined on $\mathring{\Delta}_{k-1}$ and we are going to estimate this function on the boundary of Δ_{k-1} . Since $\sum_{i=1}^k a_i = 1$, we can define the function

$$\hat{\Psi}(a_1, \dots, a_{k-1}) = \frac{(k-1)^{\frac{2k(k-1)}{k+d-2}}}{(k+d-2)^{2k}} \frac{(a_1 \dots a_{k-1} (1 - \sum_{i=1}^{k-1} a_i))^{\frac{k-d}{k+d-2}}}{((1-a_1) \dots (1-a_{k-1}) (\sum_{i=1}^{k-1} a_i))^{\frac{2(k-1)}{k+d-2}}},$$

which is the composition of Ψ with identification of $\mathring{\Delta}_{k-1}$ with the interior of the simplex τ in \mathbb{R}^{k-1} whose vertices are the origin $(0, 0, \dots, 0)$ and the vectors $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ of the canonical basis, for $i = 1, \dots, k-1$. If a sequence of points is converging to a boundary point of Δ_{k-1} , then we have a sequence $\{(a_1^{(n)}, \dots, a_{k-1}^{(n)})\}_{n \in \mathbb{N}}$ of points in τ converging to a boundary point. If the limit point is not a vertex of τ then $\lim_{n \rightarrow \infty} \hat{\Psi}(a_1^{(n)}, \dots, a_{k-1}^{(n)}) = 0$, because the numerator is converging to 0 whereas the denominator converges to a non-zero number.

The delicate points are given by the vertices of τ . On these points the function $\hat{\Psi}$ a priori cannot be continuously extended. Suppose we have a sequence $\{(a_1^{(n)}, \dots, a_{k-1}^{(n)})\}_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow \infty} (a_1^{(n)}, \dots, a_{k-1}^{(n)}) = (0, 0, \dots, 0)$. We have

$$\hat{\Psi}(a_1^{(n)}, \dots, a_{k-1}^{(n)}) = \frac{(k-1)^{\frac{2k(k-1)}{k+d-2}}}{(k+d-2)^{2k}} \frac{(a_1^{(n)} \dots a_{k-1}^{(n)} (1 - \sum_{i=1}^{k-1} a_i^{(n)}))^{\frac{k-d}{k+d-2}}}{((1-a_1^{(n)}) \dots (1-a_{k-1}^{(n)}) (\sum_{i=1}^{k-1} a_i^{(n)}))^{\frac{2(k-1)}{k+d-2}}},$$

and since we are in a neighborhood of $(0, 0, \dots, 0)$ the sequence $\hat{\Psi}(a_1^{(n)}, \dots, a_{k-1}^{(n)})$ will have the same behaviour of the following sequence

$$\hat{\Psi}(a_1^{(n)}, \dots, a_{k-1}^{(n)}) \sim \frac{(k-1)^{\frac{2k(k-1)}{k+d-2}}}{(k+d-2)^{2k}} \frac{(a_1^{(n)} \dots a_{k-1}^{(n)})^{\frac{k-d}{k+d-2}}}{(\sum_{i=1}^{k-1} a_i^{(n)})^{\frac{2(k-1)}{k+d-2}}}.$$

By looking carefully at the right-hand side of the estimate above, we can apply the inequality relating the geometric mean and arithmetic mean to get

$$\begin{aligned}
& \frac{(k-1)^{\frac{2k(k-1)}{k+d-2}} (a_1^{(n)} \dots a_{k-1}^{(n)})^{\frac{k-d}{k+d-2}}}{(k+d-2)^{2k} \left(\sum_{i=1}^{k-1} a_i^{(n)}\right)^{\frac{2(k-1)}{k+d-2}}} \\
& \leq \frac{(k-1)^{\frac{2k(k-1)}{k+d-2}}}{(k+d-2)^{2k}} \frac{1}{\left(\sum_{i=1}^{k-1} a_i^{(n)}\right)^{\frac{2(k-1)}{k+d-2}}} \left(\frac{\left(\sum_{i=1}^{k-1} a_i^{(n)}\right)^{k-1}}{(k-1)^{k-1}}\right)^{\frac{k-d}{k+d-2}} \\
& = \frac{(k-1)^{\frac{(k-1)(k+d)}{k+d-2}}}{(k+d-2)^{2k}} \left(\sum_{i=1}^{k-1} a_i^{(n)}\right)^{\frac{(k-1)(k-d-2)}{k+d-2}}.
\end{aligned}$$

The last term which appears in the inequality above depends on the exponent $k-d-2$. More precisely, by the assumption $p \geq 2$ we already know that $k \geq d+2$, but we need to distinguish the case $k = d+2$ from the case $k > d+2$. Since we assumed either $G_p = PU(p, 1)$ or $G_p = PSp(p, 1)$, we can have either $d = 2$ or $d = 4$. Thus, if $k = d+2$, we should have $k = 4$ or $k = 6$. The case $k = 6$ is not possible because the dimension of the tangent space of a quaternionic hyperbolic space is a multiple of 4, so we are going to analyze only the case $k = 4$. When $k = 4$, the space X^p becomes the complex hyperbolic space $\mathbb{H}_{\mathbb{C}}^2$ and we get the estimate

$$\hat{\Psi}(0, \dots, 0) \leq \frac{3^{\frac{9}{2}}}{4^8}$$

which is strictly less than the maximum of $\hat{\Psi}$. When $k > d+2$ the right-hand side of the inequality becomes a function which is continuous at $(0, \dots, 0)$ and it converges to 0. Moreover, since $\Psi(0, 0, \dots, 0, 1) = \hat{\Psi}(0, \dots, 0)$ and Ψ is a function which is invariant under the action of the group \mathfrak{S}_k , we get that

$$\Psi(1, 0, 0, \dots, 0) = \Psi(0, 1, 0, \dots, 0) = \dots = \Psi(0, 0, \dots, 0, 1)$$

are all bounded away from the maximum value of Ψ . Hence, in all the possible cases, we can bound $\Psi(a_1, \dots, a_k)$ away from its maximum in a suitable neighborhood of the boundary $\partial\Delta_{k-1}$. The claim follows because $\varphi(H) \leq \psi(H)$ for every $H \in Sym_1^+(k, \mathbb{R})$. \square

By the Estimate (5), we know that in our context we have

$$\left(\frac{k}{(k+d-2)^2}\right)^k (1-2\varepsilon) \leq \varphi(H_n^V) \leq \left(\frac{k}{(k+d-2)^2}\right)^k$$

for $n \geq n_0$. As a consequence of Proposition 3.3, the sequence H_n^V must converge to I/k . Hence H_n^V converges to I/k almost-everywhere on X^p . By following the same proof of [FS, Proposition 3.8] we immediately get

Proposition 3.4. *Suppose the sequence H_n^V converges almost everywhere to I/k . Let K be any compact set of X^p . Then we have that*

$$\lim_{n \rightarrow \infty} \|H_n^V - \frac{I}{k}\| = 0$$

uniformly for every $x \in K$. The norm which appears above is the one obtained by thinking of H_n^V as an operator between Euclidean vector spaces.

As a consequence of the Cauchy–Schwarz inequality applied to Equation (4), we can write

$$(6) \quad |k_n(v, D_x F(u))| \leq (k + d - 2)h_n(v, v)^{\frac{1}{2}}h'_n(u, u)^{\frac{1}{2}},$$

for every $u \in T_x X^p$ and $v \in T_{F_n(x)} X^m$. We are going to use the previous inequality, combined with Proposition 3.4, to prove the following

Lemma 3.5. *Let $r > 0$ and let $\overline{B_r(O)}$ be the closed ball of radius r around O in X^p . Then, for n large enough, there exists a real constant $C > 0$ such that*

$$\|D_x F_n\| < C$$

for every x uniformly on $\overline{B_r(O)}$.

Proof. By Proposition 3.4, we have that $\lim_{n \rightarrow \infty} H_n^V(x) = I/k$ for every x uniformly on $\overline{B_r(O)}$. This implies that

$$\lim_{n \rightarrow \infty} K_n^V(x) = \frac{k + d - 2}{k}I, \quad \lim_{n \rightarrow \infty} H_n'^U(x) = \frac{1}{k}I.$$

for every x uniformly on $\overline{B_r(O)}$. Let $\delta > 0$ and let $u \in U_x$ and $v = D_x F_n(u)$. By taking $n > n_1$, we get

$$\begin{aligned} (k + d - 2)/k \|D_x F_n(u)\|_{X^m}^2 - \delta &\leq k_n(D_x F_n(u), D_x F_n(u)) \\ h_n(D_x F_n(u), D_x F_n(u))^{\frac{1}{2}} &\leq \|D_x F_n(u)\|_{X^m} / \sqrt{k} + \delta \\ h'_n(u, u)^{\frac{1}{2}} &\leq \|u\|_{X^p} / \sqrt{k} + \delta. \end{aligned}$$

Hence, as a consequence of Inequality (6), it hold that

$$(k+d-2)/k \|D_x F_n(u)\|_{X^m}^2 - \delta \leq (k+d-2)(\|D_x F_n(u)\|_{X^m} / \sqrt{k} + \delta)(\|u\|_{X^p} / \sqrt{k} + \delta).$$

By considering on both sides the supremum on all the vectors u of norm equal to 1 we get

$$\|D_x F_n\|^2 < k(\|D_x F_n\| / \sqrt{k} + \delta)(1 / \sqrt{k} + \delta) + k\delta,$$

hence $\|D_x F_n\|$ is uniformly bounded on $\overline{B_r(O)}$ for any $n > n_1$ and for any choice of $r > 0$. \square

We are now ready to prove Theorem 1.1.

Proof. Since we know that $\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(M)$, by Lemma 3.5 we have that $\|D_x F_n\|$ must be eventually uniformly bounded $\overline{B_r(O)}$, for every $r > 0$. Let $x \in X^p$ be any point and let $\gamma \in \Gamma$. Let c be the geodesic joining x to γx . Denote by $L = d(x, \gamma x)$ so that the interval $[0, L]$ parametrizes the curve c . Consider a closed ball $\overline{B_r(O)}$ sufficiently large to contain in its interior both x and γx . On this ball there must exist a constant C such that $\|D_x F_n\| < C$ for n bigger than a suitable value n_0 . Thus, it holds that

$$d(F_n(x), F_n(\gamma x)) \leq \int_0^L \|D_{c(t)} F_n(\dot{c}(t))\| dt \leq \int_0^L \|D_{c(t)} F_n\| dt \leq C d(x, \gamma x).$$

Recall that given an element $g \in G_m$ its translation length is defined as $\ell_{X^m}(g) := \inf_{y \in X^m} d(gy, y)$. The previous estimate implies that the translation length of the element $\rho_n(\gamma)$ can be bounded by

$$\ell_{X^m}(\rho_n(\gamma)) \leq d(\rho_n(\gamma)F_n(x), F_n(x)) \leq C d(\gamma x, x).$$

We claim that it is sufficient to bound the translation length of $\rho_n(\gamma)$ for each $\gamma \in \Gamma$ to show that the sequence ρ_n is bounded in the character variety $X(\Gamma, G_m)$. Let $\gamma \in \Gamma$. If $\ell_{X^m}(\rho_n(\gamma)) = 0$ there is nothing to prove. Assume $\ell_{X^m}(\rho_n(\gamma)) > 0$. Then, up to conjugation, the element $\rho_n(\gamma)$ can be put in the form

$$\begin{pmatrix} \alpha \cosh(\ell_{X^m}(\rho_n(\gamma))/2) & \alpha \sinh(\ell_{X^m}(\rho_n(\gamma))/2) & 0 \\ \alpha \sinh(\ell_{X^m}(\rho_n(\gamma))/2) & \alpha \cosh(\ell_{X^m}(\rho_n(\gamma))/2) & 0 \\ 0 & 0 & M \end{pmatrix}$$

where $M = (m_{ij})$ is a matrix which lies in either in $PU(p-1)$ or in $PSp(p-1)$ and $|\alpha| = 1$. Then

$$|\text{Tr}(\rho_n(\gamma))| \leq 2 \cosh(\ell_{X^m}(\rho_n(\gamma))/2) + |\text{Tr}(M)| \leq C d(\gamma x, x) + C_0$$

where $|\text{Tr}(M)| < C_0$ is obtained by a compactness argument. Hence $|\text{Tr}(\rho_n(\gamma))|$ is bounded, as claimed.

Moreover the choice made before to fix $F_n(O) = O$ guarantees that the sequence ρ_n must converge to a representation ρ_∞ . Since the volume function is continuous with respect to the pointwise convergence, we get

$$\lim_{n \rightarrow \infty} \text{Vol}(\rho_n) = \text{Vol}(\rho_\infty) = \text{Vol}(M)$$

and hence we conclude by the rigidity of the volume function. \square

We want to conclude with some comments about the proof. A key point to show the rigidity at infinity for both complex and quaternionic lattices is given by the estimate on the Jacobian $Jac_k(F)$ of the natural

map. This estimate is sharp and this fact guarantees the rigidity at infinity of the volume function.

In both [CF03a] and [CF03b] the authors generalize the construction of natural maps for lattices in Lie groups of any rank by obtaining a similar estimate on the Jacobian. The estimate is sharp for lattices in products of rank one Lie groups, but this does not hold any longer for Lie groups which are not products of this type. However, the sharpness of the estimate for lattices in products of rank one Lie groups suggests us that it should be possible to extend the strong rigidity at infinity at least in this more general context.

REFERENCES

- [Alb97] P. Albuquerque, *Patterson-Sullivan measures in higher rank symmetric spaces*, C. R. Acad. Sci. Paris Sr. I Math. **324** (1997), no. 4, 427–432.
- [Alb99] P. Albuquerque, *Patterson-Sullivan theory in higher rank symmetric spaces*, Geom. Funct. Anal. **9** (1999), no. 1, 1–28
- [BGS85] W. Ballman, M. Gromov, V. Schroeder, *Manifolds of Nonpositive Curvature*, Birkhäuser, Boston, 1985.
- [BCG95] G. Besson, G. Courtois, S. Gallot, *Entropies et rigidités des espaces localement symétriques de courbure strictement négative*, Geom. Funct. Anal. **5** (1995), no. 5, 731–799.
- [BCG96] G. Besson, G. Courtois, S. Gallot, *Minimal entropy and Mostow’s rigidity theorems*, Ergodic Theory Dynam. Systems **16** (1996), no. 4, 623–649.
- [BCG99] G. Besson, G. Courtois, S. Gallot, *A real Schwarz lemma and some applications*, Rend. Mat. Appl. (7) **18** (1998), no. 2, 381–410.
- [BCG07] G. Besson, G. Courtois, S. Gallot, *Inégalités de Milnor–Wood géométriques*, Comment. Math. Helv. **82** (2007), no. 4, 753–803.
- [Bow95] B. Bowditch, *Geometrical finiteness with variable negative curvature*, Duke J. Math., **77** (1995).
- [BBI13] M. Bucher, M. Burger, A. Iozzi, *A dual interpretation of the Gromov-Thurston proof of Mostow rigidity and volume rigidity for representations of hyperbolic lattices*, Trends in harmonic analysis, 47–76, Springer, Milan, 2013.
- [BI01] M. Burger, A. Iozzi, *Bounded Cohomology and representation varieties of lattices in $PSU(1, n)$* , preprint, 2001.
- [BM96] M. Burger, S. Mozes, *$CAT(-1)$ -spaces, divergence groups and their commensurators*, J. Amer. Math. Soc. **9** (1996), no. 1, 57–93.
- [CF03a] C. Connell, B. Farb, *Minimal entropy rigidity for lattices in products of rank one symmetric spaces*, Comm. Anal. Geom. **11** (2003), no. 5, 1001–1026.
- [CF03b] C. Connell, B. Farb, *The degree theorem in higher rank*, J. Differential Geom. **65** (2003), no. 1, 19–59.
- [Cor92] K. Corlette, *Archimedean superrigidity and hyperbolic geometry*, Ann. of Math. (2) **135** (1992), no. 1, 165–182.
- [CS83] M. Culler, P. B. Shalen, *Varieties of group representations and splitting of 3-manifolds*, Ann. of Math. **117** (1983), 109–146.
- [Fra04] S. Francaviglia, *Hyperbolic volume of representations of fundamental groups of cusped 3-manifolds*, Int. Math. Res. Not. 2004, no. 9, 425–459.
- [Fra09] S. Francaviglia, *Constructing equivariant maps for representations*, Ann. Inst. Fourier (Grenoble) **59** (2009), no. 1, 393–428.

- [FK06] S. Francaviglia, B. Klaff, *Maximal volume representations are Fuchsian*, *Geom. Dedicata* **117** (2006), 111–124.
- [FS] S. Francaviglia, A. Savini, *Volume rigidity at ideal points of the character variety of hyperbolic 3-manifolds*, <https://arxiv.org/abs/1706.07347>
- [KK12] S. Kim, I. Kim, *Volume invariant and maximal representations of discrete subgroups of Lie groups*, *Math. Z.* **276** (2014), no. 3–4, 1189–1213.
- [KM08] V. Koziarz, J. Maubon, *Harmonic maps and representations of non-uniform lattices of $PU(m, 1)$* , *Ann. Inst. Fourier (Grenoble)* **58** (2008), no. 2, 507–558.
- [Mor86] J. W. Morgan, *Group actions on trees and the compactification of the space of classes of $SO(n, 1)$ -representations*, *Topology* **25** (1986), 1–33.
- [MS84] J. W. Morgan, P. B. Shalen, *Valuations, trees and degeneration of hyperbolic structures I*, *Ann. of Math.*, **120**, 1984, 401–476
- [Mos73] G. D. Mostow, *Strong rigidity of locally symmetric spaces*, *Ann. of Math. Studies*, **78**, Princeton University Press (1973).
- [Nic89] P.J. Nicholls, *The ergodic theory of discrete groups*, London Mathematical Society Lecture Note Series, 143, Cambridge University Press (1989).
- [Rob00] T. Roblin, Thomas, *Sur l'ergodicité rationnelle et les propriétés ergodiques du flot géodésique dans les variétés hyperboliques*, *Ergodic Theory Dynam. Systems* **20** (2000), no. 6, 1785–1819.
- [Yue96] C. Yue, *The ergodic theory of discrete isometry groups on manifolds of variable negative curvature*, *Trans. Amer. Math. Soc.* (1996), **12**, 4965–5005.

Alessio Savini
Department of Mathematics,
University of Bologna,
Piazza di Porta San Donato 5,
40126 Bologna,
Italy
alessio.savini5@unibo.it