

Merit functions: a bridge between optimization and equilibria*

Massimo Pappalardo[†] Giandomenico Mastroeni[†] Mauro Passacantando[†]

Abstract. In the last decades, many problems involving equilibria, arising from engineering, physics and economics, have been formulated as variational mathematical models. In turn, these models can be reformulated as optimization problems through merit functions. This paper aims at reviewing the literature about merit functions for variational inequalities, quasi-variational inequalities and abstract equilibrium problems. Smoothness and convexity properties of merit functions and solution methods based on them will be presented.

Keywords. Merit functions, gap functions, variational inequalities, equilibrium problems, descent methods.

1 Introduction

Optimization is a widespread mathematical technique in many engineering and economic applications. However, in many real-world problems, an objective function to be optimized is missing and the concept of equilibrium becomes crucial. Roughly speaking, if optimization takes care of the system utility function, the equilibrium takes into account the mutual interaction between users. In recent years, the interest in equilibrium problems has widely grown. The main applications are concerned with traffic over telecommunication networks or over public roads, oligopolistic and spatial price markets, financial markets, risk management, climate competition, migration problems, power allocation in radio systems, internet advertising, cloud computing (see, e.g., Altman and Wynter (2004); Ardagna et al (2011, 2013); Bigi et al (2009); Bigi and Passacantando (2012); Dafermos (1980); Drouet et al (2008); Ferris and Pang (1997); Forgó et al (2005); Konnov (2007, 2008a,b); Liu and Nagurney (2007); Miller and Ruszczyński (2008); Mordukhovich et al (2007); Nagurney (1993, 2010); Patriksson (1994); Pang et al (2010); Scutari et al (2010); Wardrop (1952); Zhao and Nagurney (2008) and references therein).

All these problems have been formulated in the literature through variational mathematical models as complementarity problems, variational inequalities, quasi-variational inequalities, and Nash equilibrium problems among others. Variational inequalities (VIs) are one of the most known variational models. They were introduced by Hartman and Stampacchia (1966) as a tool for studying partial differential equations in infinite dimensional spaces arising from mechanics (free-obstacle problem, friction problem, etc.). Later, their applications to contact problems in mechanical structures provided a vast source of finite dimensional problems.

A finite-dimensional VI is defined as follows:

$$\text{find } x^* \in C \text{ such that } \langle F(x^*), y - x^* \rangle \geq 0, \text{ for all } y \in C, \quad (\text{VI})$$

*This paper has been published in Annals of Operations Research, vol. 240 (1), pp. 271-299, see: <https://link.springer.com/article/10.1007/s10479-015-2013-3>.

[†]Dipartimento di Informatica, Università di Pisa, Largo B. Pontecorvo 3, 56127 Pisa, Italy.

where $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$, C is a closed and convex subset of \mathbb{R}^n and $\langle \cdot, \cdot \rangle$ is the scalar product in \mathbb{R}^n .

Several kinds of numerical methods to solve VIs have been proposed (see, e.g., Facchinei and Pang (2003); Harker and Pang (1990) and references therein). One popular approach is based on the reformulation of (VI) as an optimization problem through suitable merit functions.

A function $p : \mathbb{R}^n \rightarrow \mathbb{R}$ is called *merit function* for (VI) if there exists a set $\Omega \subseteq \mathbb{R}^n$ such that:

- p is nonnegative on Ω ,
- x^* is a solution to (VI) if and only if $x^* \in \Omega$ and $p(x^*) = 0$.

If the set Ω coincides with the feasible set C of (VI), a merit function is also known in the literature as a *gap function*. Hence, if (VI) has at least a solution, then it is equivalent to the optimization problem

$$\min_{x \in \Omega} p(x).$$

Therefore, merit functions are the key concept to build a bridge between VIs and optimization.

In this paper we aim at reviewing the state of the art concerning the merit function approach for VIs and two interesting generalization of VIs: quasi-variational inequalities and abstract equilibrium problems. This is an updated version of the previous review from Pappalardo et al (2014).

The rest of the paper is organized as follows: Section 2 is devoted to the preliminary concepts that will be used in the paper. Section 3 deals with both constrained and unconstrained optimization reformulations of (VI). In particular, we will describe continuity and differentiability properties of merit functions, conditions under which merit functions are convex or their stationary points solve (VI) and error bound results, i.e., how the distance between an arbitrary point x and the solution set of (VI) can be estimated in terms of the merit function value at x . Furthermore, *ad-hoc* descent methods for minimizing merit functions will be shown. Sections 4 and 5 are devoted to the results about merit functions for quasi-variational inequalities and abstract equilibrium problems, respectively. Examples of applications of the presented models are provided in Sections 3, 4 and 5. Some concluding remarks and suggestions for future research are collected in Section 6. We hope that this paper may stimulate further interest in merit functions and may be the basis to obtain new results.

2 Preliminaries

In this section, we show two classic particular cases of (VI), and we recall the main definitions and preliminary results that will be used throughout the paper. We make the blanket assumptions that the feasible set C of (VI) is closed and convex and the operator F is continuous on C .

Optimality conditions. As first particular case, let us consider the problem of finding a local minimum x^* of a differentiable function $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$ over the set C . The classic first order necessary optimality condition states that the directional derivative of ψ at x^* in any feasible direction is nonnegative, i.e.

$$\langle \nabla \psi(x^*), y - x^* \rangle \geq 0, \quad \forall y \in C.$$

This condition is a particular case of (VI) where $F(x) = \nabla \psi(x)$.

Complementarity problems. Another example of (VI) is provided by a complementarity problem described as follows: given a closed convex cone $C \subseteq \mathbb{R}^n$ and a mapping $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$, the complementarity problem asks to determine a point $x^* \in C$ such that

$$\langle F(x^*), x^* \rangle = 0 \quad \text{and} \quad F(x^*) \in C^*,$$

where C^* denotes the dual cone of C , i.e.

$$C^* := \{d \in \mathbb{R}^n : \langle d, y \rangle \geq 0 \text{ for all } y \in C\}.$$

Solving the complementarity problem amounts to solving (VI). In fact, if x^* solves the complementarity problem, then for any $y \in C$ we have

$$\langle F(x^*), y - x^* \rangle = \langle F(x^*), y \rangle \geq 0,$$

hence x^* solves (VI); vice versa, if x^* solves (VI), then setting $y = 0$ and $y = 2x^*$ (which belong to C because C is a cone) we obtain $\langle F(x^*), x^* \rangle = 0$ and hence $F(x^*) \in C^*$, that is x^* is a solution to the complementarity problem. Note that if we define

$$p(x) := \langle F(x), x \rangle, \quad \Omega := \{x \in C : F(x) \in C^*\}, \tag{1}$$

then $p(x) \geq 0$ for any $x \in \Omega$ and x^* solves the complementarity problem if and only if $x^* \in \Omega$ and $p(x^*) = 0$, i.e., p is a merit function for the complementarity problem.

Monotonicity definitions. Monotonicity is a key assumption to establish existence of solutions, convergence results for algorithms and to provide error bounds for (VI). We now recall the main monotonicity properties that will be exploited in the paper. F is said *monotone* on C if

$$\langle F(x) - F(y), x - y \rangle \geq 0, \quad \forall x, y \in C;$$

the corresponding concept of strict monotonicity is analogously defined just requiring strict inequality to hold for any $x, y \in C$ with $x \neq y$; F is said *strongly monotone* on C with modulus μ if

$$\langle F(x) - F(y), x - y \rangle \geq \mu \|x - y\|^2, \quad \forall x, y \in C,$$

for some $\mu > 0$; F is said *pseudomonotone* on C if for any $x, y \in C$ one has

$$\langle F(y), x - y \rangle \geq 0 \quad \implies \quad \langle F(x), x - y \rangle \geq 0.$$

In the particular case where $F(x) = \nabla\psi(x)$, monotonicity and strong monotonicity of F on C are equivalent to convexity and strong convexity of ψ on C , respectively.

Existence results. We now recall two basic results concerning the existence of a solution to (VI). For the sake of simplicity, we will not consider the sharpest possible assumptions. The solution set of (VI) is nonempty if either the feasible set C is bounded (Hartman and Stampacchia, 1966) or the following coercivity condition holds: there exists a point $y \in C$ such that

$$\lim_{\|x\| \rightarrow \infty, x \in C} \langle F(x), y - x \rangle = -\infty. \tag{2}$$

In particular, condition (2) holds if F is strongly monotone on C . Moreover, the strong monotonicity of F ensures that (VI) has a unique solution.

Fixed point problem reformulation. If we denote by π_C the Euclidean projection operator on C , i.e.,

$$\pi_C(x) := \arg \min_{y \in C} \|y - x\|,$$

then it is well-known that (VI) is equivalent to finding a fixed point of the operator $x \mapsto \pi_C(x - F(x))$ (see, e.g., Facchinei and Pang (2003)).

Complementarity problem reformulation. Assuming

$$C := \{x \in \mathbb{R}^n : g_i(x) \leq 0, \quad i = 1, \dots, m\}, \quad (3)$$

where the functions g_i are differentiable and convex for all $i = 1, \dots, m$, it is possible to derive the Karush-Kuhn-Tucker (KKT) conditions for (VI). In fact, x^* is a solution to (VI) if and only if it is a global minimum of the following convex optimization problem:

$$\min_{y \in C} \langle F(x^*), y \rangle.$$

Under some constraint qualification, the following KKT conditions

$$\begin{cases} F(x^*) + \sum_{i=1}^m \lambda_i^* \nabla g_i(x^*) = 0, \\ \lambda_i^* g_i(x^*) = 0, & i = 1, \dots, m, \\ \lambda_i^* \geq 0, \quad g_i(x^*) \leq 0, & i = 1, \dots, m, \end{cases} \quad (4)$$

are necessary and sufficient for optimality and, in turn, for the existence of solutions to (VI). It is well-known that KKT system (4) is equivalent to a complementarity problem.

3 Merit functions for variational inequalities

In this section, we summarize several approaches in order to express (VI) as a constrained or unconstrained optimization problem by means of different merit functions.

A first merit function can be defined by exploiting the fixed point reformulation stated in Section 2. In fact, the function $x \mapsto \|x - \pi_C(x - F(x))\|$ is a gap function for (VI) (see Eaves (1971)).

A further merit function for (VI) can be obtained by means of the complementarity reformulation (4), which in turn can be associated with the merit function (1). Further examples of merit functions associated with a complementarity problem can be found in Fischer and Jiang (2000).

3.1 Constrained optimization reformulations

This section is devoted to gap functions for (VI).

3.1.1 Auslender gap function

A first example of gap function was given in Auslender (1976), where the following function was introduced:

$$p(x) := \sup_{y \in C} \langle F(x), x - y \rangle. \quad (5)$$

It is trivial to prove that p is a gap function for (VI). Since the supremum in (5) can be infinite or not attained in a unique point, this function is in general neither finite, nor differentiable, nor convex. However, when C is bounded and F is continuously differentiable, it is finite and admits directional derivatives $p'(x; d)$ at any point $x \in C$ in any direction d . Moreover, if F is monotone, any stationary point x^* of p on C , i.e.

$$p'(x^*; y - x^*) \geq 0, \quad \forall y \in C,$$

is a solution to (VI). In the particular case of monotone affine VIs, i.e., if $F(x) = Ax + b$ where A is a positive semidefinite matrix, p also turns out to be convex (Marcotte, 1985). If F is strongly monotone with modulus μ and x^* is the unique solution to (VI), then p provides the following error bound:

$$\|x - x^*\| \leq \sqrt{p(x)/\mu}, \quad \forall x \in C.$$

A descent method based on the function p has been proposed in Marcotte and Dussault (1989) in the case where C is a bounded polyhedron. At each iteration, the descent direction is obtained by minimizing a linearization of a further gap function. If F is monotone, then the algorithm is globally convergent¹ to a solution to (VI). Moreover, the convergence is quadratic² if F is strongly monotone and the termination is achieved in a finite number of iterations if F is affine. Under a so-called “geometric stability condition”, it is shown that p also provides an error bound for (VI).

3.1.2 Regularized gap functions

Many efforts of the research have been directed to the study of differentiable gap functions in order to simplify the computational aspects of the problem. Important results in this sense have been obtained in Fukushima (1992); Larsson and Patriksson (1994); Wu et al (1993); Zhu and Marcotte (1994).

First, Auchmuty (1989) proposed a scheme in order to define a general class of gap functions:

$$p^A(x) := \sup_{y \in C} [\langle F(x), x - y \rangle + f(x) - f(y) - \langle \nabla f(x), x - y \rangle], \quad (6)$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex and continuously differentiable. It was proved that if (x^*, y^*) is a saddle point of the function

$$L(x, y) := \langle F(x), x - y \rangle + f(x) - f(y) - \langle \nabla f(x), x - y \rangle$$

on $C \times C$, i.e.,

$$L(x^*, y) \leq L(x^*, y^*) \leq L(x, y^*), \quad \forall (x, y) \in C \times C,$$

then x^* is a solution to (VI) and p^A is a gap function. We observe that if f is strongly convex on C and F is differentiable, then the function p^A is finite and differentiable.

¹The convergence is said global if it does not depend on the choice of the starting point.

²A sequence $\{x^k\}$ is said to be convergent to \bar{x} with rate of convergence equal to r if

$$\limsup_{k \rightarrow +\infty} \frac{\|x^{k+1} - \bar{x}\|}{\|x^k - \bar{x}\|^r} = \gamma \in (0, +\infty).$$

If $r = 1$ and $\gamma \in (0, 1)$, then the convergence is said to be linear, if $r > 1$, then the convergence is said to be superlinear, and, in particular, if $r = 2$, the convergence is said to be quadratic.

Later, Fukushima (1992) introduced a gap function which is a special case of (6), setting $f(x) = \langle x, Mx \rangle / 2$, where M is a symmetric and positive definite matrix. It is defined by

$$p^F(x) := \max_{y \in C} \left[\langle F(x), x - y \rangle - \frac{1}{2} \langle x - y, M(x - y) \rangle \right]. \quad (7)$$

Note that the maximum in (7) is always attained in a unique point $y(x)$ since the objective function is strongly concave with respect to the variable y , hence p^F is always finite. If F is continuously differentiable, then also p^F is so (Danskin, 1966) and

$$\nabla p^F(x) = F(x) - [(\nabla F(x))^T - M](y(x) - x).$$

Moreover, if x^* is a stationary point of p^F on C , i.e.

$$\langle \nabla p^F(x^*), y - x^* \rangle \geq 0, \quad \forall y \in C,$$

and the Jacobian matrix $\nabla F(x^*)$ is positive definite, then x^* is a solution to (VI). In the special case of strongly monotone affine VIs, i.e., $F(x) = Ax + b$ with A positive definite, p^F turns out to be convex (strongly convex) provided that the matrix $A + A^T - M$ is positive semidefinite (positive definite) (see Larsson and Patriksson (1994)).

A descent algorithm for minimizing p^F has been proposed in Fukushima (1992): given any starting point $x^0 \in C$, the sequence $\{x^k\}$ is generated by the iterations

$$x^{k+1} = x^k + t_k d^k, \quad (8)$$

where the search direction $d^k = y(x^k) - x^k$ and the stepsize $t_k \in (0, 1]$ is such that

$$p^F(x^k + t_k d^k) = \min_{t \in (0, 1]} p^F(x^k + t d^k). \quad (9)$$

Under the assumptions that C is bounded, F is continuously differentiable on C and $\nabla F(x)$ is positive definite for all $x \in C$, the sequence $\{x^k\}$ belongs to C and converges to the unique solution to (VI). This algorithm converges also employing an inexact line search rule, provided that F is strongly monotone on C and ∇F is Lipschitz continuous on C .

A variant of the above method which does not require the strong monotonicity of F has been proposed in Zhu and Marcotte (1993), setting the matrix $M = \alpha I$, where I is the identity matrix and $\alpha > 0$. In fact, the monotonicity of F paired with the boundedness of C guarantees that at any point $x \in C$ the vector $y(x) - x$, which depends on the matrix M and hence depends on α , is a descent direction for p^F , provided that α is small enough. The method performs a line search at the current iterate x^k if $y(x^k) - x^k$ is a descent direction for p^F ; otherwise the value of α is decreased. The method is globally convergent if C is bounded and F is continuously differentiable and monotone on C . This algorithm has been extended to nonsmooth VIs in Panicucci et al (2009).

Marcotte and Dussault (1987) and Taji et al (1993) propose a modified Newton method: at each iteration it finds the solution to the linearized (VI) at x , i.e.,

$$\text{find } z(x) \in C \text{ s.t. } \langle F(x) + \nabla F(x)[z(x) - x], y - z(x) \rangle \geq 0, \quad \forall y \in C. \quad (10)$$

In the hypothesis of strong monotonicity of F , problem (10) admits a unique solution $z(x)$ such that $d = z(x) - x$ is a descent direction for the gap functions p^A and p^F : the function p^A has

been considered in Marcotte and Dussault (1987), while p^F in Taji et al (1993). By employing a line search strategy, the method is shown to be quadratically convergent under suitable additional assumptions.

When the feasible set C , defined as in (3), is not a polyhedron, the evaluation of the regularized gap function p^F at a given point x could be computationally expensive. In order to overcome this drawback, the following gap function has been proposed in Taji and Fukushima (1996):

$$p^{TF}(x) := \max_{y \in T(x)} \left[\langle F(x), x - y \rangle - \frac{1}{2} \langle x - y, M(x - y) \rangle \right], \quad (11)$$

where

$$T(x) := \{y \in \mathbb{R}^n : g_i(x) + \langle \nabla g_i(x), y - x \rangle \leq 0, \quad i = 1, \dots, m\} \quad (12)$$

is an outer polyhedral approximation of C at x . If F is continuously differentiable, g_i 's are continuously differentiable and a constraint qualification holds, then p^{TF} is directionally differentiable. Furthermore, if ∇F is positive definite and g_i 's are twice continuously differentiable, then any stationary point of p^{TF} on C is a solution to (VI). A successive quadratic programming algorithm based on the minimization of an exact penalty function associated with p^{TF} has been proposed in Taji and Fukushima (1996).

A generalization of the gap function introduced by Fukushima has been proposed in Wu et al (1993); Zhu and Marcotte (1994) by replacing in (7) the regularizing term $\langle x - y, M(x - y) \rangle / 2$ with a general bifunction $G : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that:

$$\begin{aligned} G(x, y) &\geq 0 \text{ for all } (x, y) \in \mathbb{R}^n \times \mathbb{R}^n, \\ G &\text{ is continuously differentiable,} \\ G(x, \cdot) &\text{ is strongly convex on } C \text{ for all } x \in C, \\ G(x, x) &= \nabla_y G(x, x) = 0 \text{ for all } x \in C. \end{aligned} \quad (13)$$

For any $\alpha > 0$, the function

$$p_\alpha(x) := \max_{y \in C} [\langle F(x), x - y \rangle - \alpha G(x, y)] \quad (14)$$

turns out to be a gap function, which is continuously differentiable, if F is so, with

$$\nabla p_\alpha(x) = F(x) - (\nabla F(x))^T (y_\alpha(x) - x) - \alpha \nabla_x G(x, y_\alpha(x)), \quad (15)$$

where $y_\alpha(x)$ is the unique solution to problem (14). Note that when F is only locally Lipschitz continuous, the function p_α is also locally Lipschitz and its Clarke generalized gradient satisfies a formula similar to (15) (see Ng and Tan (2007b)). Moreover, if x^* is a stationary point of p_α on C and $\nabla F(x^*)$ is positive definite, then x^* solves (VI). A further important feature of p_α is that, under the assumption that F is strongly monotone and $\nabla_y G(x, \cdot)$ is Lipschitz continuous on C , for every $x \in C$, it provides an error bound (see Zhu and Marcotte (1994)), i.e., there exists $M > 0$ such that

$$\|x - x^*\| \leq M \sqrt{p_\alpha(x)}, \quad \forall x \in C, \quad (16)$$

where x^* is the unique solution to (VI). In particular, (16) implies the boundedness of the sublevel sets of p_α , which is of crucial importance in the convergence of the minimization algorithms.

The descent methods developed in Fukushima (1992) have been generalized to a more general framework exploiting several classes of gap functions defined by (14) (see Zhu and Marcotte (1994)).

Given a continuous mapping $\Gamma : C \times C \rightarrow \mathbb{R}^n$, such that $\Gamma(x, \cdot)$ is strongly monotone on C for any $x \in C$, the following auxiliary variational inequality is considered at a given point $x \in C$: find $y^* \in C$ such that

$$\langle \Gamma(x, y^*) - \Gamma(x, x) + F(x), y - y^* \rangle \geq 0, \quad \forall y \in C. \quad (\text{AVI}(x))$$

Having denoted by $w(x)$ the unique solution to $(\text{AVI}(x))$, one can prove that the mapping $w : C \rightarrow C$ is continuous and x^* is a solution to (VI) if and only if $x^* = w(x^*)$. In view of this result the following iterative method is proposed. Given $x^k \in C$, compute $w(x^k)$: if $w(x^k) = x^k$, then x^k is a solution to (VI), otherwise find x^{k+1} performing an Armijo³ inexact line search for the gap function p_α along the direction $d^k := w(x^k) - x^k$.

Each combination of G and Γ generates a different descent algorithm, which is globally convergent to the unique solution to (VI) under the assumption of continuous differentiability and strong monotonicity of F and suitable additional assumptions on G and Γ (Zhu and Marcotte, 1994). Note that the algorithm (8)–(9) previously described is recovered by setting $\Gamma(x, y) = M(y - x)$.

The analysis of the convergence properties of the descent methods based on the regularized gap functions p^F and p_α in the case where the operator F is nondifferentiable has been considered in Ng and Tan (2007a); Tan (2007).

3.1.3 Minty (dual) gap functions

The *Minty (or dual) variational inequality* was introduced in Minty (1967) and consists in finding $x^* \in C$ such that

$$\langle F(y), y - x^* \rangle \geq 0, \quad \forall y \in C. \quad (\text{MVI})$$

Its relevance to applications was pointed out in Giannessi (1998); Pappalardo and Passacantando (2002, 2004). In particular, Minty states the equivalence between (VI) and (MVI) when F is pseudomonotone on C (Minty, 1967).

In parallel with the Auslender gap function, it can be shown that

$$p^M(x) := \sup_{y \in C} \langle F(y), x - y \rangle \quad (17)$$

is a gap function for (MVI) and hence it is a gap function for (VI) provided that F is pseudomonotone on C .

The most important feature of this function, known in the literature as *Minty (or dual) gap function*, is its convexity. However, it is in general nondifferentiable; subdifferentiability and related properties have been analysed in Marcotte and Dussault (1987); Marcotte and Zhu (1999); Nguyen and Dupuis (1984); Yamashita and Fukushima (1997); Zhang et al (2003). Furthermore, it can be difficult to evaluate p^M since the optimization problem in (17) is generally not convex.

A cutting plane method for minimizing p^M has been proposed in Nguyen and Dupuis (1984): at each iteration it solves a linear programming problem, provided that C is a polyhedron, and it converges to a solution to (VI) if F is strictly monotone. Later, this method has been combined with the Tikhonov regularization technique in order to deal with monotone VIs (Bigi and Panicucci, 2010).

³The Armijo inexact line search along the direction d^k consists in finding the smallest non negative integer m such that

$$p_\alpha(x^k + \beta^m d^k) \leq p_\alpha(x^k) - \sigma \beta^m \|d^k\|^2,$$

where $\beta, \sigma \in (0, 1)$ are parameters, and then setting $x^{k+1} := x^k + \beta^m d^k$.

Following the scheme described before for (VI), it is possible to regularize the function p^M exploiting a bifunction G which satisfies conditions (13). In fact, the function

$$p_G^M(x) := \sup_{y \in C} [\langle F(y), x - y \rangle - G(x, y)] \quad (18)$$

is a gap function for (MVI) (see Mastroeni (1999)). Moreover, if the optimization problem in (18) has a unique solution $y(x)$, then p_G^M is continuously differentiable and its gradient is given by

$$\nabla p_G^M(x) = F(y(x)) - \nabla_x G(x, y(x)).$$

In parallel with the analysis developed for (VI), a descent method for the function p_G^M has been proposed in Mastroeni (1999). Given any starting point $x^0 \in C$, any sequence $\{x^k\}$ generated by an exact line search algorithm with descent direction given by $y(x) - x$ converges to the unique solution to (VI), provided that C is compact, F is continuously differentiable, ∇F is positive definite on C and

$$\nabla_x G(x, y) + \nabla_y G(x, y) = 0, \quad \forall x, y \in C.$$

Observe that the latter condition is fulfilled, for instance, by

$$G(x, y) = \frac{1}{2} \langle x - y, M(x - y) \rangle.$$

This algorithm has been extended in Mastroeni (2005) employing an inexact line search rule and replacing the assumption that $\nabla F(x)$ is positive definite for $x \in C$ with the strong monotonicity of F on C .

A different regularization of the gap function p^M has been proposed in Yamashita and Fukushima (1997), where the following function is considered:

$$p_\beta^M(x) := \sup_{y \in C} [\langle F(y), x - y \rangle + \beta \|x - y\|^2], \quad (19)$$

where β is a positive parameter. This function is convex and lower semicontinuous as the original p^M . It is continuously differentiable provided that F is so and the supremum in (19) is attained in a unique point. Moreover, if F is strongly monotone on C with modulus μ , and $\beta \in (0, \mu]$, it is a gap function for (VI) and provides an error bound, i.e.,

$$\|x - x^*\| \leq \sqrt{p_\beta^M(x)/\beta}, \quad \forall x \in C, \quad (20)$$

where x^* is the unique solution to (VI).

3.1.4 Gap functions based on conjugate duality

Given a convex function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and a concave function $g : \mathbb{R}^n \rightarrow \mathbb{R}$, we recall that

$$f^*(y) := \sup_{x \in \mathbb{R}^n} [\langle y, x \rangle - f(x)], \quad g_*(y) := \inf_{x \in \mathbb{R}^n} [\langle y, x \rangle - g(x)]$$

are the Fenchel conjugate, in convex and concave sense respectively, of f and g (see, e.g., Rockafellar (1970)). Moreover, the Fenchel dual of the problem

$$\inf_{x \in \mathbb{R}^n} [f(x) - g(x)]$$

is defined as

$$\sup_{y \in \mathbb{R}^n} [g_*(y) - f^*(y)].$$

Note that the Fenchel dual of the constrained problem

$$\inf_{x \in C} f(x)$$

can be obtained defining $g(x) = -\delta_C(x)$, where δ_C is the indicator⁴ function of the set C , so that

$$\inf_{x \in C} f(x) = \inf_{x \in \mathbb{R}^n} [f(x) - (-\delta_C(x))],$$

and the associated Fenchel dual turns out to be

$$\sup_{y \in \mathbb{R}^n} [-f^*(y) + \inf_{x \in C} \langle y, x \rangle].$$

When the feasible set C is explicitly defined by convex constraints as in (3), the value of the Auslender gap function p at a given point x coincides (see Altangerel et al (2007); Larsson and Patriksson (1994)) with the opposite of the optimal value of the Fenchel dual of the problem

$$\inf_{y \in C} \langle F(x), y - x \rangle. \quad (P(x))$$

Moreover, the opposite of the optimal value of the so called Lagrangian dual and the Fenchel-Lagrange dual associated with $P(x)$ leads to define a further gap function that coincides with

$$p^L(x) := \inf_{\lambda \geq 0} \sup_{y \in \mathbb{R}^n} [\langle F(x), x - y \rangle - \langle \lambda, g(y) \rangle], \quad (21)$$

which has been proposed in Giannessi (1995).

Similarly, considering the opposite of the optimal values of the Lagrange and of the Fenchel dual associated with the problem

$$\inf_{y \in C} \langle F(y), y - x \rangle,$$

which is equivalent to the one which appears in the right-hand side of (17), the following gap functions for (MVI) are defined (Altangerel et al, 2007):

$$p^{ML}(x) := \inf_{\lambda \geq 0} \sup_{y \in \mathbb{R}^n} [\langle F(y), x - y \rangle - \langle \lambda, g(y) \rangle], \quad (22)$$

$$p^{MF}(x) := \inf_{p \in \mathbb{R}^n} \left\{ \sup_{y \in \mathbb{R}^n} [\langle F(y), x - y \rangle + \langle p, y \rangle] + \delta_C^*(-p) \right\}, \quad (23)$$

where $\delta_C^*(x) = \sup_{y \in C} \langle x, y \rangle$ is the support function to the set C .

The proposed gap functions are all convex if F is an affine monotone map.

3.2 Unconstrained optimization reformulations

In this section, we review merit functions which allow to reformulate (VI) as an unconstrained optimization problem.

⁴ δ_C is defined as follows: $\delta_C(x) = 0$ if $x \in C$ and $\delta_C(x) = +\infty$ otherwise.

3.2.1 D-gap functions

The difference of two regularized gap functions

$$p_{\alpha\beta}(x) := p_{\alpha}(x) - p_{\beta}(x), \quad (24)$$

where p_{α} and p_{β} are defined by (14) with $0 < \alpha < \beta$, is called D-gap function (where D stands for “difference”). This function is nonnegative on the whole space \mathbb{R}^n and $p_{\alpha\beta}(x^*) = 0$ if and only if x^* is a solution to (VI) (Yamashita et al, 1997). Therefore, solving (VI) is equivalent to finding the optimal solutions of the problem

$$\min_{x \in \mathbb{R}^n} p_{\alpha\beta}(x). \quad (25)$$

When (VI) is a nonlinear complementarity problem, the D-gap function with $\beta = 1/\alpha$ and $\alpha \in (0, 1)$ coincides with the implicit Lagrangian proposed and studied in Mangasarian and Solodov (1993); Peng (1997); Peng and Yuan (1997).

Clearly, the D-gap function inherits the differentiability properties of p_{α} and p_{β} , i.e., if F is differentiable, the function $p_{\alpha\beta}$ is also differentiable and

$$\nabla p_{\alpha\beta}(x) = (\nabla F(x))^T [y_{\beta}(x) - y_{\alpha}(x)] + \beta \nabla_x G(x, y_{\beta}(x)) - \alpha \nabla_x G(x, y_{\alpha}(x)), \quad (26)$$

where $y_{\alpha}(x)$ and $y_{\beta}(x)$ are the solutions of the optimization problem (14) with α and β respectively. When the mapping F is locally Lipschitz continuous, the D-gap function is also locally Lipschitz and the Clarke generalized gradient of $p_{\alpha\beta}$ satisfies a formula similar to (26) (see Ng and Tan (2007a)).

The D-gap function is not convex in general and the stationary points of (25) may not be global minima. However, if x^* is a stationary point, i.e., $\nabla p_{\alpha\beta}(x^*) = 0$, and the Jacobian matrix $\nabla F(x^*)$ is positive definite, then x^* is a solution to (VI) (see Yamashita et al (1997)). Notice that the positive definiteness of $\nabla F(x^*)$ can not be replaced by the strict monotonicity assumption on F (see the counterexample in Yamashita et al (1997)). When the feasible set C is a box, it is sufficient to assume that $\nabla F(x^*)$ is a P -matrix (i.e. its principal minors are all positive) to obtain the same conclusion (Kanzow and Fukushima, 1998b). In the special case of strongly monotone affine VIs, the D-gap function is convex (strongly convex) provided that the parameters α and β are chosen so that the matrix

$$A + A^T - \alpha I - \beta^{-1} A^T A \quad (27)$$

is positive semidefinite (positive definite) (Peng and Fukushima, 1999).

Since the D-gap functions allow to reformulate (VI) as an unconstrained problem, the boundedness of their level sets is an important issue in order to develop minimization algorithms for solving problem (25). The level sets of the D-gap function, denoted by

$$L(c) := \{x \in \mathbb{R}^n : p_{\alpha\beta}(x) \leq c\},$$

are bounded for all $c \geq 0$ if either C is bounded (Kanzow and Fukushima, 1998b) or F is strongly monotone (Peng and Fukushima, 1999; Qu et al, 2003). Recently, it has been proved in Li and Ng (2009) that the strong monotonicity assumption on F can be replaced by a coercivity condition stronger than (2).

If F is strongly monotone on \mathbb{R}^n and either F is Lipschitz continuous on \mathbb{R}^n or C is bounded, then $\sqrt{p_{\alpha\beta}}$ provides an error bound for (VI), i.e. there exists a constant $M > 0$ such that

$$\|x - x^*\| \leq M \sqrt{p_{\alpha\beta}(x)}, \quad \forall x \in \mathbb{R}^n,$$

(see Yamashita et al (1997)). Notice that when C is unbounded, the strong monotonicity on F , without the Lipschitz continuity assumption, is not sufficient to guarantee the same result (see the counterexample in Huang and Ng (2005)). However, when F is strongly monotone, (VI) has a unique solution and hence it is possible to reformulate the problem by replacing the set C by its intersection with a sphere large enough to contain the solution. In the special case where C is a box, the strong monotonicity of F can be replaced by the assumption that F is a uniform P -function (Kanzow and Fukushima, 1998b). Recently, new global error bounds have been proposed in Li et al (2010).

On the other hand, the strong monotonicity on F only guarantees a local error bound on the level sets of the D-gap function (Qu et al, 2003), that is for any $c \geq 0$ there exists $M > 0$ such that

$$\|x - x^*\| \leq M \sqrt{p_{\alpha\beta}(x)}, \quad \forall x \in L(c).$$

Recently, this result has been extended to locally ξ -monotone and coercive mappings (Li et al, 2010) and to general nonmonotone mappings (Li and Ng, 2009).

There are several solution methods for VIs based on the minimization of D-gap functions. A descent method with Armijo-type line search has been proposed in Yamashita et al (1997): at each iteration it exploits the search direction $d = r(x) + \rho s(x)$, where $r(x) = y_\alpha(x) - y_\beta(x)$, $s(x) = \alpha \nabla_x G(x, y_\alpha(x)) - \beta \nabla_x G(x, y_\beta(x))$ and $\rho > 0$ is a sufficiently small constant. This method converges to the solution to (VI) if F is strongly monotone on \mathbb{R}^n and either F is Lipschitz continuous on \mathbb{R}^n or C is bounded. Another descent method was developed in Solodov and Tseng (2000) for solving monotone VIs with bounded feasible set. It is similar to the method proposed in Zhu and Marcotte (1993) based on the Fukushima's regularized gap function: at each iteration it uses $d = y_\alpha(x) - y_\beta(x)$ as search direction along with a suitable update of the parameters α and β . A descent method for solving nonmonotone VIs, which is based on the minimization of the function $\sqrt{p_{\alpha\beta}}$, has been presented recently in Li and Ng (2009).

A hybrid Newton method has been proposed in Peng and Fukushima (1999): at each iteration, it finds the solution $z(x)$ to the linearized VI (10) at x and it uses the direction $d = z(x) - x$ whenever it provides a sufficient decrease in the D-gap function $p_{\alpha\beta}$; otherwise the direction $d = -\nabla p_{\alpha\beta}(x)$ is used. Then, an inexact line search is performed to get the next iterate. The generated sequence converges superlinearly to the unique solution x^* to (VI) if F is continuously differentiable and strongly monotone on \mathbb{R}^n . Furthermore, the convergence is quadratic if ∇F is Lipschitz continuous around x^* . A variant of this method has been proposed in Peng et al (1999) for box constrained VIs.

In Kanzow and Fukushima (1998b) a nonsmooth Gauss-Newton type method for solving box constrained VIs has been presented. At each iteration, it solves a linear system of equations involving the generalized Hessian of the D-gap function and uses this vector as search direction if a descent condition is satisfied; otherwise the direction $d = -\nabla p_{\alpha\beta}(x)$ is used. Then an inexact line search is performed. The algorithm is globally and superlinearly convergent under suitable assumptions. A similar Gauss-Newton strategy has also been adopted in a trust region method for minimizing the D-gap function (Sun et al, 1997).

Another Newton type method for the solution of box constrained VIs is based on the reformulation of (VI) as a system of nonsmooth and nonlinear equations involving the natural residual. This method, based on the minimization of the D-gap function, is globally and superlinearly convergent (Kanzow and Fukushima, 1998a).

3.2.2 Merit functions via the Moreau-Yosida regularization

Another approach to get unconstrained optimization reformulations of (VI) is based on the Moreau-Yosida regularization of some gap functions (Yamashita and Fukushima, 1997).

The function

$$p_{\alpha\lambda}^{MY}(x) := \inf_{z \in C} \left\{ \sup_{y \in C} [\langle F(z), z - y \rangle - \alpha \|y - z\|^2] + \lambda \|z - x\|^2 \right\}, \quad (28)$$

with $\alpha \geq 0$ and $\lambda > 0$, is derived from the Moreau-Yosida regularization of the regularized gap function p^F , with $M = \alpha I$. It is nonnegative on the whole space \mathbb{R}^n and $p_{\alpha\lambda}^{MY}(x^*) = 0$ if and only if x^* solves (VI).

Notice that this merit function may not be easy to evaluate in practice unless (VI) has a certain special structure (e.g., F is affine and C is a polyhedron (see Yamashita and Fukushima (1997))). However, it enjoys some nice theoretical properties that other merit functions do not have. For instance, if the infimum in (28) is uniquely attained in $z_{\alpha\lambda}(x)$ for each $x \in \mathbb{R}^n$, then $p_{\alpha\lambda}^{MY}$ is differentiable on \mathbb{R}^n and

$$\nabla p_{\alpha\lambda}^{MY}(x) = 2\lambda [x - z_{\alpha\lambda}(x)],$$

even if F is not differentiable.

In general, the function $p_{\alpha\lambda}^{MY}$ is not convex. However, if the gap function p is convex, then $p_{0\lambda}^{MY}$ is differentiable and convex on \mathbb{R}^n for any $\lambda > 0$; while if the regularized gap function p^F , with $M = \alpha I$, is convex then $p_{\alpha\lambda}^{MY}$ is differentiable and convex on \mathbb{R}^n for any $\lambda > 0$.

The function $p_{\alpha\lambda}^{MY}$ provides also a global error bound under the strong monotonicity of F (without assuming Lipschitz continuity as is the case for the D-gap functions). In fact, if F is strongly monotone on C with modulus μ , $\alpha \in [0, \mu]$ and $\lambda > 0$, then

$$\frac{1}{2} \min\{\mu - \alpha, \lambda\} \|x - x^*\|^2 \leq p_{\alpha\lambda}^{MY}(x) \leq \lambda \|x - x^*\|^2, \quad \forall x \in \mathbb{R}^n,$$

i.e., the growth rate of $p_{\alpha\lambda}^{MY}$ is in the order of the squared distance from the unique solution x^* to (VI).

When F satisfies suitable monotonicity assumptions, further merit functions for (VI) can be obtained by the Moreau-Yosida regularization of Minty gap functions. The function

$$p_{\beta\lambda}^M(x) := \inf_{z \in C} \left\{ \sup_{y \in C} [\langle F(y), z - y \rangle + \beta \|y - z\|^2] + \lambda \|z - x\|^2 \right\}, \quad (29)$$

with $\beta \geq 0$ and $\lambda > 0$, is the Moreau-Yosida regularization of the Minty gap function p^M (when $\beta = 0$) and the regularized Minty gap function p_{β}^M (when $\beta > 0$). If F is pseudomonotone on C , then $p_{0\lambda}^M$ turns out to be a merit function for (VI) for any $\lambda > 0$, because it is nonnegative on \mathbb{R}^n and $p_{0\lambda}^M(x^*) = 0$ if and only if x^* solves (VI). Furthermore, if F is strongly monotone with modulus μ , then the same happens for $p_{\beta\lambda}^M$ provided that $\beta \in [0, \mu]$.

Note that also $p_{\beta\lambda}^M$ may not be easy to evaluate in practice, but some nice theoretical properties hold. In fact, it is differentiable and convex on \mathbb{R}^n for any $\beta \geq 0$ and $\lambda > 0$, without making any additional assumption on F . Moreover, if F is strongly monotone on C with modulus μ , $\beta \in (0, \mu]$ and $\lambda > 0$, then the quadratic growth rate of $p_{\beta\lambda}^M$ is ensured, i.e.,

$$\frac{1}{2} \min\{\beta, \lambda\} \|x - x^*\|^2 \leq p_{\beta\lambda}^M(x) \leq \lambda \|x - x^*\|^2, \quad \forall x \in \mathbb{R}^n,$$

where x^* is the unique solution to (VI).

3.3 An application to traffic network equilibrium problems

A traffic network consists of a set of nodes \mathcal{N} , a set of arcs $\mathcal{A} \subseteq \mathcal{N} \times \mathcal{N}$ and a set of origin/destination pairs $\mathcal{W} \subseteq \mathcal{N} \times \mathcal{N}$. For each O/D pair w , a traffic demand d_w has to be distributed among the paths connecting w . We denote \mathcal{P}_w the set of all paths connecting w , x_p the flow on path p and $x = (x_p)_{p \in \mathcal{P}_w, w \in \mathcal{W}}$ the vector of all path flows. The set of feasible path flows is given by

$$X = \left\{ x \geq 0 : \sum_{p \in \mathcal{P}_w} x_p = d_w, \quad \forall w \in \mathcal{W} \right\}.$$

The flow f_a on each arc a is the sum of all flows on paths to which the arc belongs, hence the arc flow vector $f = (f_a)_{a \in \mathcal{A}}$ can be written as $f = \Delta x$, where Δ is the arc-path incidence matrix:

$$\Delta_{a,p} = \begin{cases} 1 & \text{if } a \in p, \\ 0 & \text{otherwise.} \end{cases}$$

For each arc a , there is a nonnegative cost function $t_a(f)$, which represents the travel time associated with arc a and depends on the arc flow vector f . The corresponding path cost function is assumed to be additive, i.e., the travel time $T_p(x)$ on path p is the sum of the travel times of the arcs belonging to p :

$$T_p(x) = \sum_{a \in p} t_a(\Delta x).$$

According to the Wardrop equilibrium principle (Wardrop, 1952), a path flow $x^* \in X$ is called a *network equilibrium* if it is positive only on minimum cost paths, i.e., the following implication

$$x_p^* > 0 \implies T_p(x^*) = \min_{q \in \mathcal{P}_w} T_q(x^*)$$

holds for any O/D pair $w \in \mathcal{W}$ and path $p \in \mathcal{P}_w$.

It is well-known (Dafermos, 1980) that the problem of finding network equilibria is equivalent to solving the following variational inequality:

$$\text{find } x^* \in X \text{ such that } \langle T(x^*), y - x^* \rangle \geq 0, \text{ for all } y \in X. \quad (30)$$

Next example shows how the merit function approach for VIs can be applied to network equilibria.

Example 1. Consider the network in Fig. 1 with two O/D pairs: $w_1 = (1, 4)$ with demand $d_1 = 4$ and $w_2 = (1, 5)$ with $d_2 = 6$. Each O/D pair is connected by two paths: $\mathcal{P}_{w_1} = \{(1, 2), (2, 4); (1, 3), (3, 4)\}$ and $\mathcal{P}_{w_2} = \{(1, 2), (2, 5); (1, 3), (3, 5)\}$. We denote the flow on paths as x_1, \dots, x_4 , respectively. Hence the set of feasible path flows is given by

$$X = \{x \in \mathbb{R}_+^4 : x_1 + x_2 = 4, \quad x_3 + x_4 = 6\}.$$




Figure 1: Traffic network in Example 1.

Assume that the arc cost functions are defined as follows:

$$\begin{cases} t_{12} := f_{12} + 1 = x_1 + x_3 + 1, \\ t_{13} := 3 f_{13} + 2 = 3(x_2 + x_4) + 2, \\ t_{24} := 2 f_{24} + f_{34} + 1 = 2x_1 + x_2 + 1, \\ t_{25} := 2 f_{25} + f_{35} + 3 = 2x_3 + x_4 + 3, \\ t_{34} := f_{34} + 2 = x_2 + 2, \\ t_{35} := 4 f_{35} + 1 = 4x_4 + 1, \end{cases}$$

thus the corresponding path costs are

$$\begin{cases} T_1 = t_{12} + t_{24} = 3x_1 + x_2 + x_3 + 2, \\ T_2 = t_{13} + t_{34} = 4x_2 + 3x_4 + 4, \\ T_3 = t_{12} + t_{25} = x_1 + 3x_3 + x_4 + 4, \\ T_4 = t_{13} + t_{35} = 3x_2 + 7x_4 + 3, \end{cases}$$

i.e., the operator of VI (30) is $T(x) = Ax + b$, with

$$A = \begin{pmatrix} 3 & 1 & 1 & 0 \\ 0 & 4 & 0 & 3 \\ 1 & 0 & 3 & 1 \\ 0 & 3 & 0 & 7 \end{pmatrix}, \quad b = \begin{pmatrix} 2 \\ 4 \\ 4 \\ 3 \end{pmatrix}.$$

Note that the matrix A is positive definite, thus the mapping T is strongly monotone and there exists a unique solution of VI (30), i.e., a unique network equilibrium.

We now consider the Fukushima's regularized gap function (7) with $M = I$, i.e.,

$$p^F(x) = \max_{y \in X} \left[\langle T(x), x - y \rangle - \frac{1}{2} \|x - y\|^2 \right].$$

This function is continuously differentiable and strongly convex since the matrix $A + A^T - I$ is positive definite. In Fig. 2 we show the graph of p^F defined on the 2-dimensional space (x_1, x_3) , with $x_1 \in [0, 4]$ and $x_3 \in [0, 6]$ (the demand constraints allow to express variables x_2 and x_4 as function of x_1 and x_3 , respectively).




Figure 2: The regularized gap function p^F with $M = I$ in Example 1.

Iteration	x_1	x_2	x_3	x_4	$p^F(x)$
0	4.000000	0	6.000000	0	1.5000 e+02
1	2.679622	1.320378	4.019434	1.980566	8.8840 e-03
2	2.631471	1.368529	4.052452	1.947548	1.5178 e-06
3	2.631587	1.368413	4.052626	1.947374	2.5929 e-10
4	2.631579	1.368421	4.052632	1.947368	2.8747 e-14

Table 1: Numerical results of the descent algorithm (8)-(9) applied to Example 1.

The descent algorithm (8)-(9) applied to p^F can be exploited to compute the network equilibrium. Table 1 reports the first four iterations of the algorithm (implemented in MATLAB) starting from the feasible flow (4, 0, 6, 0).

Note that the path costs corresponding to the equilibrium solution

$$x^* = (2.6316, 1.3684, 4.0526, 1.9474)$$

are

$$T(x^*) = (15.3158, 15.3158, 20.7368, 20.7368),$$

i.e., the two paths connecting each O/D pair have the same cost. Furthermore, the Lagrange multipliers λ^* associated with x^* in the KKT conditions (4) coincide with the equilibrium costs, i.e., $\lambda^* = (15.3158, 20.7368)$.

Figure 3 shows the graph of the D-gap function $p_{\alpha\beta}$, with $\alpha = 1$, $\beta = 2$ and $G(x, y) = \|x - y\|^2/2$, defined on the space (x_1, x_3) . Note that this function is always nonnegative (even in unfeasible points) and its global minimum is x^* .




Figure 3: The D-gap function $p_{\alpha\beta}$ with $\alpha = 1$, $\beta = 2$ and $G(x, y) = \|x - y\|^2/2$ in Example 1.

4 Merit functions for quasi-variational inequalities

In this section we consider the merit function approach for quasi-variational inequalities (QVIs), i.e., VIs in which the feasible region depends on the variable x . Given a vector-valued mapping

$F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and a set-valued mapping $C : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$, such that $C(x)$ are closed and convex sets for any $x \in \mathbb{R}^n$, the QVI is defined as follows:

$$\text{find } x^* \in C(x^*) \text{ such that } \langle F(x^*), y - x^* \rangle \geq 0, \text{ for all } y \in C(x^*). \quad (\text{QVI})$$

When all the sets $C(x)$ coincide with the same set C , (QVI) collapses to (VI). The set of fixed points of the mapping C , i.e.

$$X := \{x \in \mathbb{R}^n : x \in C(x)\},$$

is the feasible region of (QVI). In the following we suppose that sets $C(x)$ are defined by constraints, i.e.,

$$C(x) := \{y \in \mathbb{R}^n : g_i(x, y) \leq 0, \quad i = 1, \dots, m\},$$

where the functions $g_i : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ are assumed to be continuous and $g_i(x, \cdot)$ convex for any fixed $x \in \mathbb{R}^n$. Furthermore, in order to guarantee the convexity of the set X , we assume that the functions $x \mapsto g_i(x, x)$ are convex for all $i = 1, \dots, m$.

QVIs were introduced in Bensoussan et al (1973); Bensoussan and Lions (1973) and subsequently exploited to model several finite and infinite-dimensional problems (see Baiocchi and Capelo (1984); Chan and Pang (1982); Facchinei et al (2014, 2013) and references therein).

Some merit functions have been proposed in the literature extending to QVIs similar ideas developed for VIs. Similarly to VIs, the reformulation of (QVI) as a fixed point problem leads to define a merit function. In fact, it follows from the definition that x solves (QVI) if and only if $x = \pi_{C(x)}(x - F(x))$, thus $\|x - \pi_{C(x)}(x - F(x))\|$ is a merit function for (QVI). Another approach is based on reformulating (QVI), under suitable constraint qualifications, as a complementarity problem via the following KKT conditions:

$$\begin{cases} F(x) + \sum_{i=1}^m \lambda_i \nabla_y g_i(x, x) = 0, \\ \lambda_i g_i(x, x) = 0, & i = 1, \dots, m, \\ \lambda_i \geq 0, \quad g_i(x, x) \leq 0, & i = 1, \dots, m. \end{cases}$$

Recently, a solution method based on these conditions has been proposed in Facchinei et al (2014).

A straightforward extension of the gap function (5) to QVIs is defined as follows (Giannessi, 1995):

$$\mathfrak{p}(x) := \sup_{y \in C(x)} \langle F(x), x - y \rangle. \quad (31)$$

This function is nonnegative on the set X and x^* solves (QVI) if and only if $x^* \in X$ and $\mathfrak{p}(x^*) = 0$. However, the gap function \mathfrak{p} is nondifferentiable and it may occur that $\mathfrak{p}(x) = +\infty$ for some point in X .

The regularized gap function (7) has been extended to QVIs in Fukushima (2007) and is defined by

$$\mathfrak{p}_\alpha(x) := \max_{y \in C(x)} \left[\langle F(x), x - y \rangle - \frac{\alpha}{2} \|x - y\|^2 \right]. \quad (32)$$

This function is a gap function, it is finite and the maximum in (32) is attained in a unique point $y_\alpha(x)$, provided that the set $C(x)$ is nonempty. Actually, it is possible to define \mathfrak{p}_α replacing the regularization term $\alpha \|x - y\|^2 / 2$ and the set $C(x)$ with more general expressions satisfying suitable conditions (Fukushima, 2007; Taji, 2008).

In contrast to VIs, this function is nondifferentiable even if F is so (see examples in Harms et al (2014b)). If F and g_i are continuously differentiable and a constraint qualification holds, then \mathbf{p}_α is directionally differentiable everywhere and its directional derivative at x along direction d is given by

$$\mathbf{p}'_\alpha(x; d) = \min_{\lambda \in \Lambda_\alpha(x)} \langle F(x) - [(\nabla F(x))^T - \alpha I][y_\alpha(x) - x] - \sum_{i=1}^m \lambda_i \nabla_x g_i(x, y_\alpha(x)), d \rangle,$$

where $\Lambda_\alpha(x)$ is the set of Lagrange multipliers associated with $y_\alpha(x)$, i.e.,

$$\Lambda_\alpha(x) = \{ \lambda \in \mathbb{R}_+^m : F(x) + \alpha[y_\alpha(x) - x] + \sum_{i=1}^m \lambda_i \nabla_y g_i(x, y_\alpha(x)) = 0, \\ \lambda_i g_i(x, y_\alpha(x)) = 0, \quad i = 1, \dots, m \},$$

(see Fukushima (2007)). Furthermore, \mathbf{p}_α turns out to be continuously differentiable in the special case of QVI with ‘moving sets’, i.e., when $C(x) = Q + c(x)$, where Q is a closed and convex set and $c : \mathbb{R}^n \rightarrow \mathbb{R}^n$, provided that mappings F and c are continuously differentiable (Dietrich, 2001). Recently, this latter result has been extended to QVIs with generalized moving sets (Harms et al, 2014b).

Similarly to VIs, the regularized gap function \mathbf{p}_α is nonconvex in general. In Taji (2008) it is proved that, whenever \mathbf{p}_α is directionally differentiable, a stationary point x^* of \mathbf{p}_α on X , i.e.,

$$\mathbf{p}'_\alpha(x^*; y - x^*) \geq 0, \quad \forall y \in X,$$

is a solution to (QVI) provided that the matrix $\nabla F(x^*)$ is positive definite and

$$\lambda_i \langle \nabla_x g_i(x^*, y_\alpha(x^*)), y_\alpha(x^*) - x^* \rangle \geq 0, \quad \text{for all } i = 1, \dots, m \text{ and } \lambda \in \Lambda_\alpha(x).$$

Notice that in Taji (2008) the key assumption $y_\alpha(x^*) \in X$ is not explicitly stated in the statement, but it is exploited in the proof and must be therefore considered as hypothesis.

An unconstrained minimization reformulation of (QVI) can be obtained via the D-gap functions, i.e., the difference of two regularized gap functions. In fact, given $0 < \alpha < \beta$, the function

$$\mathbf{p}_{\alpha\beta}(x) := \mathbf{p}_\alpha(x) - \mathbf{p}_\beta(x)$$

is nonnegative on \mathbb{R}^n and $\mathbf{p}_{\alpha\beta}(x^*) = 0$ if and only if x^* solves (QVI). The directional differentiability of $\mathbf{p}_{\alpha\beta}$ directly follows from that of \mathbf{p}_α and \mathbf{p}_β . Recently, extending an idea of Dietrich (see Dietrich (1999)), another unconstrained optimization reformulation has been obtained in Harms et al (2014a) by making use of Toland’s and Singer’s duality theory.

The functions $\sqrt{\mathbf{p}_\alpha}$ and $\sqrt{\mathbf{p}_{\alpha\beta}}$ provide error bound results for (QVI) provided that F is strongly monotone and Lipschitz continuous on \mathbb{R}^n and an additional technical assumption on the Euclidean projection on the sets $C(x)$ is fulfilled (Gupta and Mehra, 2012). Another error bound result based on the function $\sqrt{\mathbf{p}_\alpha}$ has been proved in Aussel et al (2011).

4.1 Application to generalized Nash equilibrium problems

Let us consider a noncooperative game with N players, in which each player i controls a set of variables $x_i \in \mathbb{R}^{n_i}$. The vector of all players strategies is denoted by $x = (x_1, \dots, x_N) \in \mathbb{R}^n$, with $n = n_1 + \dots, n_N$; the vector x is also denoted by $x = (x_i, x_{-i})$, where x_{-i} denotes the strategy vector of all the players different from player i . Each player i has a cost function $\theta_i : \mathbb{R}^n \rightarrow \mathbb{R}$, which

possibly depends on all players strategies x , and a feasible set $X_i(x_{-i}) \subseteq \mathbb{R}^{n_i}$, possibly depending on the rival players' strategies x_{-i} .

A generalized Nash equilibrium (GNE) of the game is a vector

$$x^* = (x_1^*, \dots, x_N^*) \in X_1(x_{-1}^*) \times \dots \times X_N(x_{-N}^*)$$

such that, for any $i = 1, \dots, N$, x_i^* is an optimal solution of the following optimization problem:

$$\min_{x_i} \theta_i(x_i, x_{-i}^*) \quad \text{subject to} \quad x_i \in X_i(x_{-i}^*).$$

In other words, x^* is a GNE if no player can improve its own cost function by unilaterally changing its strategy.

It is well-known (see, e.g., Facchinei and Kanzow (2010)) that under the following assumptions:

- θ_i is continuously differentiable for any $i = 1, \dots, N$,
- $\theta_i(\cdot, x_{-i})$ is convex for any x_{-i} and $i = 1, \dots, N$,
- the feasible sets $X_i(x_{-i})$ are closed and convex for all $x \in \mathbb{R}^n$ and $i = 1, \dots, N$,

the problem of finding GNE is equivalent to solving the QVI with operator

$$F(x) = (\nabla_{x_1} \theta_1(x), \dots, \nabla_{x_N} \theta_N(x))$$

and set-valued mapping

$$C(x) = X_1(x_{-1}) \times \dots \times X_N(x_{-N}).$$

Example 2. (Cavazzuti et al, 2002) Consider a two-person noncooperative game, in which player i selects the coordinate $x_i \in \mathbb{R}$ subject to a individual constraint $x_i \leq 0$ and a shared constraint $x_1 + x_2 \leq -1$. The aim of player i is to minimize the (squared) distance between (x_1, x_2) and his favourite goal $P_i \in \mathbb{R}^2$, with $P_1 = (1, 0)$ and $P_2 = (0, 1)$. Thus the optimization problems of the two players are defined as follows:

$$\text{Player 1: } \begin{cases} \min_{x_1} (x_1 - 1)^2 + x_2^2 \\ x_1 \leq 0 \\ x_1 + x_2 \leq -1 \end{cases} \quad \text{Player 2: } \begin{cases} \min_{x_2} x_1^2 + (x_2 - 1)^2 \\ x_2 \leq 0 \\ x_1 + x_2 \leq -1 \end{cases}$$

The set of GNE of the game coincides with the solution set of the QVI given by $F(x) = (2x_1 - 2, 2x_2 - 2)$ and

$$C(x) = (-\infty, \min\{0, -1 - x_2\}] \times (-\infty, \min\{0, -1 - x_1\}].$$

The feasible region of the QVI, i.e. the set of fixed point of the set-valued mapping C , is

$$X = \{x \in \mathbb{R}^2 : x_1 \leq 0, x_2 \leq 0, x_1 + x_2 \leq -1\}.$$

It is easy to check that the solution set of the QVI is the segment connecting $(-1, 0)$ and $(0, -1)$.

The value of the gap function (31) can be explicitly computed:

$$\begin{aligned}
\mathfrak{p}(x) &= \sup_{y \in C(x)} \langle F(x), x - y \rangle \\
&= \sup_{y \in C(x)} [2(x_1 - 1)(x_1 - y_1) + 2(x_2 - 1)(x_2 - y_2)] \\
&= 2x_1(x_1 - 1) + 2x_2(x_2 - 1) + \sup_{y_1 \leq \min\{0, -1 - x_2\}} 2(1 - x_1)y_1 \\
&\quad + \sup_{y_2 \leq \min\{0, -1 - x_1\}} 2(1 - x_2)y_2 \\
&= \begin{cases} 2x_1(x_1 - 1) + 2x_2(x_2 - 1) + \\ \quad + 2(1 - x_1) \min\{0, -1 - x_2\} + \\ \quad + 2(1 - x_2) \min\{0, -1 - x_1\}, & \text{if } x_1 \leq 1 \text{ and } x_2 \leq 1, \\ +\infty, & \text{otherwise.} \end{cases}
\end{aligned}$$

This function is equal to zero on the solution set, but is not finite everywhere on \mathbb{R}^2 and it is not differentiable on the half-lines $\{-1\} \times (-\infty, 1]$ and $(-\infty, 1] \times \{-1\}$.

Figures 4 and 5 show the graphs of the regularized gap function \mathfrak{p}_α , with $\alpha = 5$, and the D-gap function $\mathfrak{p}_{\alpha\beta}$, with $\alpha = 5$ and $\beta = 10$, respectively. Note that both functions are finite on \mathbb{R}^2 and equal to zero in the solution set; \mathfrak{p}_α is negative in points not belonging to X , while $\mathfrak{p}_{\alpha\beta}$ is nonnegative on the whole space \mathbb{R}^2 .




Figure 4: The regularized gap function \mathfrak{p}_α with $\alpha = 5$ in Example 2.




Figure 5: The D-gap function $\mathbf{p}_{\alpha\beta}$ with $\alpha = 5$ and $\beta = 10$ in Example 2.

5 Merit functions for abstract equilibrium problems

The abstract equilibrium problem is a general mathematical model which includes optimization, multi-objective optimization, variational inequalities, fixed point and complementarity problems, Nash equilibria in noncooperative games and inverse optimization as special cases (see Bigi et al (2013); Blum and Oettli (1994)). It is defined as follows:

$$\text{find } x^* \in C \text{ such that } f(x^*, y) \geq 0, \text{ for all } y \in C, \quad (\text{EP})$$

where C is a closed and convex subset of \mathbb{R}^n and $f : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is a bifunction such that $f(x, \cdot)$ is convex and satisfies $f(x, x) = 0$ for all $x \in C$. Setting $f(x, y) = \langle F(x), y - x \rangle$ we obtain (VI).

In the last decade, several merit functions for (EP) have been introduced in the literature. These functions often extend to (EP) those originally conceived for VIs. For instance, a direct extension of the gap function (5) from VIs to (EP) is defined as follows (Mastroeni, 2003):

$$\mathbf{p}(x) := \sup_{y \in C} [-f(x, y)].$$

This function is nonnegative on C and x^* solves (EP) if and only if $x^* \in C$ and $\mathbf{p}(x^*) = 0$. However, \mathbf{p} has the same disadvantages of function (5), i.e., it is in general neither finite, nor differentiable nor convex. For these reasons, the regularized gap function has been proposed (Mastroeni, 2003):

$$\mathbf{p}_\alpha(x) := \max_{y \in C} \left[-f(x, y) - \frac{\alpha}{2} \|x - y\|^2 \right]. \quad (33)$$

It allows to reformulate (EP) as the problem of minimizing \mathbf{p}_α on C , it is continuously differentiable, if the bifunction f is so, and

$$\nabla \mathbf{p}_\alpha(x) = -\nabla_x f(x, y_\alpha(x)) - \alpha[x - y_\alpha(x)],$$

where $y_\alpha(x)$ is the unique maximizer of problem (33). Note that the regularization term $\|y - x\|^2$ can be replaced by a more general bifunction G satisfying condition (13) (see Mastroeni (2003)). Similarly to VIs, the regularized gap function is nonconvex in general. However, if x^* is a stationary point of \mathbf{p}_α on C , i.e.,

$$\langle \nabla \mathbf{p}_\alpha(x^*), y - x^* \rangle \geq 0, \quad \forall y \in C,$$

and f is strictly ∇ -monotone on C , i.e.,

$$\langle \nabla_x f(x, y) + \nabla_y f(x, y), y - x \rangle > 0, \quad \forall x, y \in C \text{ with } x \neq y,$$

then x^* is a solution to (EP). The strict ∇ -monotonicity of f plays a role similar to those of positive definiteness of ∇F for VIs. In fact, it guarantees, in addition to the above “stationarity” property, that $y_\alpha(x) - x$ is a descent direction for \mathbf{p}_α at any non-stationary point x . Solution methods based on the minimization of \mathbf{p}_α along this direction have been developed in Chadli et al (2004); Mastroeni (2003). An inexact version of these methods has been proposed in Di Lorenzo et al (2014).

A descent method which does not require the strict ∇ -monotonicity of f has been introduced in Bigi et al (2009). It is similar to that developed in Zhu and Marcotte (1993) for VIs: at any iteration it performs a line search if $y_\alpha(x) - x$ is a descent direction for \mathbf{p}_α at x , otherwise the value of α is reduced. Convergence is guaranteed provided that C is bounded and f is c -monotone on C , i.e.,

$$f(x, y) + \langle \nabla_x f(x, y), y - x \rangle \geq 0, \quad \forall x, y \in C. \quad (34)$$

The latter condition is neither stronger nor weaker than strict ∇ -monotonicity and it is satisfied if $f(\cdot, y)$ is concave for all $y \in C$ (see Bigi et al (2009); Bigi and Passacantando (2015c)).

Similarly to VIs, function $\sqrt{\mathbf{p}_\alpha}$ provides error bound results under suitable monotonicity assumptions on f (see Chadli et al (2004); Konnov and Pinyagina (2003b); Mastroeni (2003)).

Since the evaluation of the regularized gap function \mathbf{p}_α could be computationally expensive if C is defined as in (3) by nonlinear constraints, a variant of function \mathbf{p}_α can be exploited as in the case of VIs. In Bigi and Passacantando (2012) the following function has been introduced:

$$\mathbf{p}_\alpha^{\text{BP}}(x) := \max_{y \in T(x)} \left[-f(x, y) - \frac{\alpha}{2} \|x - y\|^2 \right],$$

where $T(x)$ is the outer polyhedral approximation of C at x defined as in (12). This function turns out to be a locally Lipschitz gap function for (EP). Furthermore, if g_i 's are continuously differentiable and a constraint qualification holds, then $\mathbf{p}_\alpha^{\text{BP}}$ is directionally differentiable. Solution methods for (EP) exploiting this merit function have been proposed in Bigi and Passacantando (2012, 2015b).

D-gap functions have been extended from VIs to (EP) as well. Indeed, the difference of two regularized gap functions

$$\mathbf{p}_{\alpha\beta}(x) := \mathbf{p}_\alpha(x) - \mathbf{p}_\beta(x), \quad (35)$$

with $0 < \alpha < \beta$, is nonnegative on \mathbb{R}^n and $\mathbf{p}_{\alpha\beta}(x^*) = 0$ if and only if x^* solves (EP). Thus, the global minima of $\mathbf{p}_{\alpha\beta}$ on \mathbb{R}^n coincide with the solutions to (EP) (see Konnov and Pinyagina (2003a); Zhang and Han (2009)). The D-gap function inherits the differentiability properties of \mathbf{p}_α and \mathbf{p}_β but in general is not convex. Stationary points of $\mathbf{p}_{\alpha\beta}$ coincide with the solutions to (EP) if the mappings $\nabla_x f(x, \cdot) + \nabla_y f(x, \cdot)$ are strictly monotone on \mathbb{R}^n for any $x \in \mathbb{R}^n$ (Zhang and Han, 2009). Similarly to VIs, function $\sqrt{\mathbf{p}_{\alpha\beta}}$ provides error bound results under suitable monotonicity assumptions on f (Cherugondi, 2013; Zhang and Wu, 2009).

Several solution methods for (EP) are based on D-gap functions. Descent methods exploiting the direction $d = r(x) + \rho s(x)$, where $r(x) = y_\alpha(x) - y_\beta(x)$, $s(x) = \alpha[x - y_\alpha(x)] - \beta[x - y_\beta(x)]$ and $\rho > 0$ is small enough, have been introduced in Cherugondi (2013); Konnov and Pinyagina (2003a). A descent method, which is similar to that proposed in Solodov and Tseng (2000) for VIs, is based on direction $d = y_\alpha(x) - y_\beta(x)$ and suitable updates of parameters α and β (Bigi and Passacantando, 2015a). Another descent method relies on the same direction $d = y_\alpha(x) - x$ which is exploited by the solution methods for \mathbf{p}_α (Zhang and Wu, 2009).

The regularized Minty gap function (19) has been extended to (EP) in Quoc and Muu (2012) and it has been used to develop an iterative method for solving strongly monotone equilibrium problems, while gap functions based on conjugate duality have been extended to (EP) in Altangerel et al (2006).

5.1 Application to a class of Nash-Cournot equilibrium problems

We now describe a problem of production competition over a network between several firms which produce the same commodity. We consider a modification of the oligopolistic model originally proposed in Marcotte (1987). Given a transportation network $(\mathcal{N}, \mathcal{A})$, where \mathcal{N} is the set of nodes and \mathcal{A} the set of arcs, the firms and the markets are located at some subsets of nodes I and J , respectively. Each firm $i \in I$ chooses the quantity x_j^i to supply to each market $j \in J$ and the quantities v_a^i to be sent on each arc $a \in \mathcal{A}$. These variables are subject to flow-conservation constraints, i.e., for any $i \in I$ and $k \in \mathcal{N}$ we have

$$(E v^i)_k = \begin{cases} -\sum_{j \in J} x_j^i & \text{if } k = i, \\ 0 & \text{if } k \notin J, \\ x_k^i & \text{if } k \in J, \end{cases} \quad (36)$$

where E is the node-arc incidence matrix of the network and $v^i = (v_a^i)_{a \in \mathcal{A}}$. Moreover, q^i denotes the maximum quantity that firm i may produce, i.e.,

$$\sum_{j \in J} x_j^i \leq q^i. \quad (37)$$

The goal of the firm i is to maximize its profit given by

$$\sum_{j \in J} x_j^i p_j \left(\sum_{\ell \in I} x_j^\ell \right) - \sum_{a \in \mathcal{A}} s_a v_a^i - \pi_i \left(\sum_{j \in J} x_j^i \right), \quad (38)$$

where $p_j : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is the inverse demand function for market j , that is $p_j(z)$ denotes the unitary price at which the market j requires a total quantity z , s_a is the unitary transportation cost on arc a and $\pi_i : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is the production cost function of firm i . Note that the first term of (38) depends on the quantities x_j^ℓ chosen by all the firms $\ell \in I$.

We say that an equilibrium state is reached when the flows and the quantities produced by the firms are such that no firm would increase its profit by changing its own choices while the other firms keep their own. This equilibrium definition coincides with the concept of Nash equilibrium in a noncooperative game where firms are the players and (38) are their payoff functions. Setting $x = (x_j^i)_{i \in I, j \in J}$, $v = (v_a^i)_{i \in I}$ and analogously y and w , Nash equilibria of this game are the solutions

of the abstract equilibrium problem (EP), where the bifunction f is the Nikaidô-Isoda function associated with the game (Nikaidô and Isoda, 1955), that is:

$$f((x, v), (y, w)) = \sum_{i \in I} \left[\sum_{j \in J} x_j^i p_j \left(\sum_{\ell \in I} x_j^\ell \right) - \sum_{j \in J} y_j^i p_j \left(y_j^i + \sum_{\ell \in I, \ell \neq i} x_j^\ell \right) + \sum_{a \in \mathcal{A}} s_a (w_a^i - v_a^i) + \pi_i \left(\sum_{j \in J} y_j^i \right) - \pi_i \left(\sum_{j \in J} x_j^i \right) \right]$$

and the feasible set C is defined by constraints (36) and (37).

Example 3. Let us consider the transportation network in Fig. 6, where $I = \{1, 2\}$, $J = \{6, 7\}$ and the number associated with each arc a denotes the unitary transportation cost s_a .




Figure 6: Transportation network in Example 3.

We assume that the production bounds are $q_1 = 70$ and $q_2 = 40$; that both markets have the same inverse demand function:

$$p_j(z) = p(z) := \rho^{1/\tau} (z + \sigma)^{-1/\tau}, \quad j \in J,$$

with $\rho = 5000$, $\tau = 1.1$ and $\sigma = 0.01$ (see, e.g., Murphy et al (1982)), and that the production cost functions have the form

$$\pi_i(z) := \gamma_i z + (1 + \delta_i)^{-1} K_i^{-\delta_i} z^{1+\delta_i}, \quad i \in I,$$

where parameters γ_i , δ_i and K_i are reported in Table 2.

i	γ_i	δ_i	K_i
1	10	5/6	5
2	6	1	5

Table 2: Parameters of cost functions π_i in Example 3.

Since the functions p and π_i are convex and differentiable, the function $z \mapsto z p(z)$ is concave and the bifunction $f(\cdot, (y, w))$ is concave for any (y, w) . Therefore, condition (34) is fulfilled and

the convergence of the modified descent algorithm proposed in Bigi et al (2009) is guaranteed. We implemented this algorithm in MATLAB exploiting the built-in function `fmincon` from the Optimization Toolbox to evaluate the regularized gap function \mathbf{p}_α and to compute the search direction $y_\alpha(x) - x$. Table 3 reports the quantities supplied by each firm to each market in the first 10 iterations of the algorithm starting from a zero total production.

Iteration	x_6^1	x_7^1	x_6^2	x_7^2	$\mathbf{p}_\alpha(x)$
0	0.0000	0.0000	0.0000	0.0000	1.0553e+04
1	11.5969	10.1858	11.3801	13.1382	1.1763e+03
2	29.5006	27.9329	19.5714	20.4286	8.7228e+00
3	32.6601	29.7497	18.8909	21.1091	8.6960e-02
4	32.5906	29.8771	18.6625	21.3375	2.5378e-03
5	32.5766	29.8855	18.6660	21.3340	1.5252e-03
6	32.5660	29.8917	18.6692	21.3308	9.1849e-04
7	32.5605	29.8950	18.6712	21.3288	6.5303e-04
8	32.5560	29.8977	18.6730	21.3270	4.6451e-04
9	32.5534	29.8992	18.6742	21.3258	3.6820e-04
10	32.5511	29.9006	18.6753	21.3247	2.9189e-04

Table 3: Numerical results of the modified descent algorithm proposed in Bigi et al (2009) applied to Example 3.

6 Concluding remarks

Merit functions have been introduced for a number of variational mathematical models. In this paper, we focused on three of the most important ones: variational inequalities, quasi-variational inequalities and abstract equilibrium problems. Among other relevant models we recall set-valued variational inequalities, vector variational inequalities and generalized Nash equilibrium problems.

The merit function approach has been extensively developed for VIs in the last two decades, while it is still at a quite early stage for more general problems. We believe that this is partially due to the complexity of these problems, but above all because many real-world applications of these problems have arisen recently. Therefore, there are still many challenging open problems regarding merit functions which are worthy of being investigated.

Merit functions for QVIs need to be further investigated: for instance, the general condition under which the stationary points of the regularized gap function are solutions to (QVI) should be deepened for different classes of problems according to the set-valued mapping defining the feasible region. Furthermore, to the best of our knowledge, no ad-hoc descent method based on merit functions has been developed so far.

Regarding abstract equilibrium problems, the convergence of descent methods is usually based on differentiability assumptions. We think that some efforts should be devoted to develop algorithms for nonsmooth problems, which include nonsmooth Nash equilibrium problems as special cases. Moreover, it would be interesting to extend the Moreau-Yosida regularization to merit functions for abstract equilibrium problems. Finally, we believe that new merit functions could be developed

without assuming the convexity of $f(x, \cdot)$: this might allow to extend the merit function approach to nonconvex Nash equilibrium problems.

References

- Altangerel L, Boş RI, Wanka G (2006) On gap functions for equilibrium problems via Fenchel duality. *Pac J Optim* 2(3):667–678
- Altangerel L, Boş RI, Wanka G (2007) On the construction of gap functions for variational inequalities via conjugate duality. *Asia-Pac J Oper Res* 24(3):353–371
- Altman E, Wynter L (2004) Equilibrium, games, and pricing in transportation and telecommunication networks. *Networks and Spatial Economics* 4(1):7–21
- Ardagna D, Panicucci B, Passacantando M (2011) A game theoretic formulation of the service provisioning problem in cloud systems. In: *Proceedings of the 20th International Conference on World Wide Web, WWW '11*, ACM, New York, NY, USA, pp 177–186
- Ardagna D, Panicucci B, Passacantando M (2013) Generalized nash equilibria for the service provisioning problem in cloud systems. *IEEE Transactions on Services Computing* 6(4):429–442
- Auchmuty G (1989) Variational principles for variational inequalities. *Numer Funct Anal Optim* 10(9-10):863–874
- Auslender A (1976) *Optimisation. Méthodes numériques*. Masson, Paris
- Aussel D, Correa R, Marechal M (2011) Gap functions for quasivariational inequalities and generalized Nash equilibrium problems. *J Optim Theory Appl* 151(3):474–488
- Baiocchi C, Capelo A (1984) *Variational and quasivariational inequalities*. John Wiley & Sons, Inc., New York
- Bensoussan A, Lions JL (1973) Nouvelle formulation de problèmes de contrôle impulsionnel et applications. *C R Acad Sci Paris Sér A-B* 276:A1189–A1192
- Bensoussan A, Goursat M, Lions JL (1973) Contrôle impulsionnel et inéquations quasi-variationnelles stationnaires. *C R Acad Sci Paris Sér A-B* 276:A1279–A1284
- Bigi G, Panicucci B (2010) A successive linear programming algorithm for nonsmooth monotone variational inequalities. *Optim Methods Softw* 25(1):29–35
- Bigi G, Passacantando M (2012) Gap functions and penalization for solving equilibrium problems with nonlinear constraints. *Comput Optim Appl* 53(2):323–346
- Bigi G, Passacantando M (2015a) D-gap functions and descent techniques for solving equilibrium problems. *Journal of Global Optimization* 62(1):183–203
- Bigi G, Passacantando M (2015b) Descent and Penalization Techniques for Equilibrium Problems with Nonlinear Constraints. *J Optim Theory Appl* 164(3):804–818

- Bigi G, Passacantando M (2015c) Twelve monotonicity conditions arising from algorithms for equilibrium problems. *Optim Methods Softw* 30(2):323–337
- Bigi G, Castellani M, Pappalardo M (2009) A new solution method for equilibrium problems. *Optim Methods Softw* 24(6):895–911
- Bigi G, Castellani M, Pappalardo M, Passacantando M (2013) Existence and solution methods for equilibria. *European J Oper Res* 227(1):1–11
- Blum E, Oettli W (1994) From optimization and variational inequalities to equilibrium problems. *Math Student* 63(1-4):123–145
- Cavazzuti E, Pappalardo M, Passacantando M (2002) Nash equilibria, variational inequalities, and dynamical systems. *J Optim Theory Appl* 114(3):491–506
- Chadli O, Konnov IV, Yao JC (2004) Descent methods for equilibrium problems in a Banach space. *Comput Math Appl* 48(3-4):609–616
- Chan D, Pang JS (1982) The generalized quasivariational inequality problem. *Math Oper Res* 7(2):211–222
- Cherugondi C (2013) A note on D-gap functions for equilibrium problems. *Optimization* 62(2):211–226
- Dafermos S (1980) Traffic equilibrium and variational inequalities. *Transportation Sci* 14(1):42–54
- Danskin JM (1966) The theory of max – min, with applications. *SIAM J Appl Math* 14:641–664
- Di Lorenzo D, Passacantando M, Sciandrone M (2014) A convergent inexact solution method for equilibrium problems. *Optim Methods Softw* 29(5):979–991
- Dietrich H (1999) A smooth dual gap function solution to a class of quasivariational inequalities. *J Math Anal Appl* 235(1):380–393
- Dietrich H (2001) Optimal control problems for certain quasivariational inequalities. *Optimization* 49(1-2):67–93
- Drouet L, Haurie A, Moresino F, Vial JP, Vielle M, Viguier L (2008) An oracle based method to compute a coupled equilibrium in a model of international climate policy. *Computational Management Science* 5(1-2):119–140
- Eaves BC (1971) On the basic theorem of complementarity. *Math Programming* 1(1):68–75
- Facchinei F, Kanzow C (2010) Generalized Nash equilibrium problems. *Ann Oper Res* 175:177–211
- Facchinei F, Pang JS (2003) Finite-dimensional variational inequalities and complementarity problems. *Springer Series in Operations Research*, Springer-Verlag, New York
- Facchinei F, Kanzow C, Sagratella S (2013) QVILIB: a library of quasi-variational inequality test problems. *Pac J Optim* 9(2):225–250

- Facchinei F, Kanzow C, Sagratella S (2014) Solving quasi-variational inequalities via their KKT conditions. *Math Program* 144(1-2, Ser. A):369–412
- Ferris MC, Pang JS (1997) Engineering and economic applications of complementarity problems. *SIAM Rev* 39(4):669–713
- Fischer A, Jiang H (2000) Merit functions for complementarity and related problems: a survey. *Comput Optim Appl* 17(2-3):159–182
- Forgó F, Fülöp J, Prill M (2005) Game theoretic models for climate change negotiations. *European J Oper Res* 160(1):252–267
- Fukushima M (1992) Equivalent differentiable optimization problems and descent methods for asymmetric variational inequality problems. *Math Programming* 53(1, Ser. A):99–110
- Fukushima M (2007) A class of gap functions for quasi-variational inequality problems. *J Ind Manag Optim* 3(2):165–171
- Giannessi F (1995) Separation of sets and gap functions for quasi-variational inequalities. In: *Variational inequalities and network equilibrium problems* (Erice, 1994), Plenum, New York, pp 101–121
- Giannessi F (1998) On Minty variational principle. In: *New trends in mathematical programming, Appl. Optim.*, vol 13, Kluwer Acad. Publ., Boston, MA, pp 93–99
- Gupta R, Mehra A (2012) Gap functions and error bounds for quasi variational inequalities. *J Global Optim* 53(4):737–748
- Harker PT, Pang JS (1990) Finite-dimensional variational inequality and nonlinear complementarity problems: a survey of theory, algorithms and applications. *Math Programming* 48(2, (Ser. B)):161–220
- Harms N, Hoheisel T, Kanzow C (2014a) On a smooth dual gap function for a class of quasi-variational inequalities. *J Optim Theory Appl* 163(2):413–438
- Harms N, Kanzow C, Stein O (2014b) Smoothness properties of a regularized gap function for quasi-variational inequalities. *Optim Methods Softw* 29(4):720–750
- Hartman P, Stampacchia G (1966) On some non-linear elliptic differential-functional equations. *Acta Math* 115:271–310
- Huang LR, Ng KF (2005) Equivalent optimization formulations and error bounds for variational inequality problems. *J Optim Theory Appl* 125(2):299–314
- Kanzow C, Fukushima M (1998a) Solving box constrained variational inequalities by using the natural residual with D-gap function globalization. *Oper Res Lett* 23(1-2):45–51
- Kanzow C, Fukushima M (1998b) Theoretical and numerical investigation of the D-gap function for box constrained variational inequalities. *Math Programming* 83(1, Ser. A):55–87
- Konnov IV (2007) On variational inequalities for auction market problems. *Optim Lett* 1(2):155–162

- Konnov IV (2008a) Spatial equilibrium problems for auction-type systems. *Russian Mathematics* 52(1):30–44
- Konnov IV (2008b) Variational inequalities for modeling auction markets with price mappings. *Open Oper Res J* 2:29–37
- Konnov IV, Pinyagina OV (2003a) D-gap functions for a class of equilibrium problems in Banach spaces. *Comput Methods Appl Math* 3(2):274–286
- Konnov IV, Pinyagina OV (2003b) Descent method with respect to the gap function for nonsmooth equilibrium problems. *Russian Mathematics* 47(12):67–73
- Larsson T, Patriksson M (1994) A class of gap functions for variational inequalities. *Math Programming* 64(1, Ser. A):53–79
- Li G, Ng KF (2009) Error bounds of generalized D-gap functions for nonsmooth and nonmonotone variational inequality problems. *SIAM J Optim* 20(2):667–690
- Li G, Tang C, Wei Z (2010) Error bound results for generalized D-gap functions of nonsmooth variational inequality problems. *J Comput Appl Math* 233(11):2795–2806
- Liu Z, Nagurney A (2007) Financial networks with intermediation and transportation network equilibria: a supernet network equivalence and reinterpretation of the equilibrium conditions with computations. *Comput Manag Sci* 4(3):243–281
- Mangasarian OL, Solodov MV (1993) Nonlinear complementarity as unconstrained and constrained minimization. *Math Programming* 62(2, Ser. B):277–297
- Marcotte P (1985) A new algorithm for solving variational inequalities with application to the traffic assignment problem. *Math Programming* 33(3):339–351
- Marcotte P (1987) Algorithms for the network oligopoly problem. *Journal of the Operational Research Society* 38(11):1051–1065
- Marcotte P, Dussault JP (1987) A note on a globally convergent Newton method for solving monotone variational inequalities. *Oper Res Lett* 6(1):35–42
- Marcotte P, Dussault JP (1989) A sequential linear programming algorithm for solving monotone variational inequalities. *SIAM J Control Optim* 27(6):1260–1278
- Marcotte P, Zhu D (1999) Weak sharp solutions of variational inequalities. *SIAM J Optim* 9(1):179–189
- Mastroeni G (1999) Minimax and extremum problems associated to a variational inequality. In: *Proceedings of the Workshop “Numerical Methods in Optimization” (Cortona, 1997)*, 58, pp 185–196
- Mastroeni G (2003) Gap functions for equilibrium problems. *J Global Optim* 27(4):411–426
- Mastroeni G (2005) Gap functions and descent methods for Minty variational inequality. In: *Optimization and control with applications, Appl. Optim.*, vol 96, Springer, New York, pp 529–547

- Miller N, Ruszczyński A (2008) Risk-adjusted probability measures in portfolio optimization with coherent measures of risk. *European J Oper Res* 191(1):193–206
- Minty GJ (1967) On the generalization of a direct method of the calculus of variations. *Bull Amer Math Soc* 73:315–321
- Mordukhovich BS, Outrata JV, Červinka M (2007) Equilibrium problems with complementarity constraints: case study with applications to oligopolistic markets. *Optimization* 56(4):479–494
- Murphy FH, Sherali HD, Soyster AL (1982) A mathematical programming approach for determining oligopolistic market equilibrium. *Math Programming* 24(1):92–106
- Nagurney A (1993) *Network economics: a variational inequality approach*. Kluwer Academic Publishers Group, Dordrecht
- Nagurney A (2010) Formulation and analysis of horizontal mergers among oligopolistic firms with insights into the merger paradox: a supply chain network perspective. *Comput Manag Sci* 7(4):377–406
- Ng KF, Tan LL (2007a) D-gap functions for nonsmooth variational inequality problems. *J Optim Theory Appl* 133(1):77–97
- Ng KF, Tan LL (2007b) Error bounds of regularized gap functions for nonsmooth variational inequality problems. *Math Program* 110(2, Ser. A):405–429
- Nguyen S, Dupuis C (1984) Efficient method for computing traffic equilibria in networks with asymmetric transportation costs. *Transportation Science* 18(2):185–202
- Nikaidô H, Isoda K (1955) Note on non-cooperative convex games. *Pacific J Math* 5:807–815
- Pang JS, Scutari G, Palomar D, Facchinei F (2010) Design of cognitive radio systems under temperature-interference constraints: A variational inequality approach. *IEEE Transactions on Signal Processing* 58(6):3251–3271
- Panicucci B, Pappalardo M, Passacantando M (2009) A globally convergent descent method for nonsmooth variational inequalities. *Comput Optim Appl* 43(2):197–211
- Pappalardo M, Passacantando M (2002) Stability for equilibrium problems: from variational inequalities to dynamical systems. *J Optim Theory Appl* 113(3):567–582
- Pappalardo M, Passacantando M (2004) Gap functions and Lyapunov functions. *J Global Optim* 28(3-4):379–385
- Pappalardo M, Mastroeni G, Passacantando M (2014) Merit functions: a bridge between optimization and equilibria. *4OR* 12(1):1–33
- Patriksson M (1994) *The Traffic Assignment problem*. VSP BV
- Peng JM (1997) Equivalence of variational inequality problems to unconstrained minimization. *Math Programming* 78(3, Ser. A):347–355

- Peng JM, Fukushima M (1999) A hybrid Newton method for solving the variational inequality problem via the D-gap function. *Math Program* 86(2, Ser. A):367–386
- Peng JM, Yuan Y (1997) Unconstrained methods for generalized complementarity problems. *J Comput Math* 15(3):253–264
- Peng JM, Kanzow C, Fukushima M (1999) A hybrid Josephy-Newton method for solving box constrained variational inequality problems via the D-gap function. *Optim Methods Softw* 10(5):687–710
- Qu B, Wang CY, Zhang JZ (2003) Convergence and error bound of a method for solving variational inequality problems via the generalized D-gap function. *J Optim Theory Appl* 119(3):535–552
- Quoc TD, Muu LD (2012) Iterative methods for solving monotone equilibrium problems via dual gap functions. *Comput Optim Appl* 51(2):709–728
- Rockafellar RT (1970) *Convex analysis*. Princeton University Press, Princeton, N.J.
- Scutari G, Palomar DP, Barbarossa S (2010) Competitive optimization of cognitive radio MIMO systems via game theory. In: *Convex optimization in signal processing and communications*, Cambridge Univ. Press, Cambridge, pp 387–442
- Solodov MV, Tseng P (2000) Some methods based on the D-gap function for solving monotone variational inequalities. *Comput Optim Appl* 17(2-3):255–277, nonsmooth and smoothing methods (Hong Kong, 1998)
- Sun D, Fukushima M, Qi L (1997) A computable generalized Hessian of the D-gap function and Newton-type methods for variational inequality problems. In: *Complementarity and variational problems* (Baltimore, MD, 1995), SIAM, Philadelphia, PA, pp 452–473
- Taji K (2008) On gap functions for quasi-variational inequalities. *Abstr Appl Anal* pp Art. ID 531,361, 7
- Taji K, Fukushima M (1996) A new merit function and a successive quadratic programming algorithm for variational inequality problems. *SIAM J Optim* 6(3):704–713
- Taji K, Fukushima M, Ibaraki T (1993) A globally convergent Newton method for solving strong monotone variational inequalities. *Math Programming* 58(3, Ser. A):369–383
- Tan LL (2007) Regularized gap functions for nonsmooth variational inequality problems. *J Math Anal Appl* 334(2):1022–1038
- Wardrop JG (1952) Road paper. some theoretical aspects of road traffic research. In: *ICE Proceedings: Engineering Divisions*, Thomas Telford, vol 1, pp 325–362
- Wu JH, Florian M, Marcotte P (1993) A general descent framework for the monotone variational inequality problem. *Math Programming* 61(3, Ser. A):281–300
- Yamashita N, Fukushima M (1997) Equivalent unconstrained minimization and global error bounds for variational inequality problems. *SIAM J Control Optim* 35(1):273–284

- Yamashita N, Taji K, Fukushima M (1997) Unconstrained optimization reformulations of variational inequality problems. *J Optim Theory Appl* 92(3):439–456
- Zhang J, Wan C, Xiu N (2003) The dual gap function for variational inequalities. *Appl Math Optim* 48(2):129–148
- Zhang L, Wu SY (2009) An algorithm based on the generalized D-gap function for equilibrium problems. *J Comput Appl Math* 231(1):403–411
- Zhang LP, Han JY (2009) Unconstrained optimization reformulations of equilibrium problems. *Acta Math Sin (Engl Ser)* 25(3):343–354
- Zhao L, Nagurney A (2008) A network equilibrium framework for internet advertising: models, qualitative analysis, and algorithms. *European J Oper Res* 187(2):456–472
- Zhu DL, Marcotte P (1993) Modified descent methods for solving the monotone variational inequality problem. *Oper Res Lett* 14(2):111–120
- Zhu DL, Marcotte P (1994) An extended descent framework for variational inequalities. *J Optim Theory Appl* 80(2):349–366