

# Investigating Roadside Soil Microbiomes: Metagenomic Analyses to Identify Taxa Driving Rubber Degradation


Alessia Dicesare<sup>1,2</sup>, Daniela Bucchieri<sup>1,2</sup>, Marcella de Divitiis<sup>1,2</sup>, Immacolata Serra<sup>1</sup>, Luca Giannini<sup>3</sup>, Silvia Guerra<sup>3</sup>, Federica Magaletti<sup>3</sup>, Paola Branduardi<sup>1</sup>

<sup>1</sup> Department of Biotechnology and Biosciences, University of Milano-Bicocca, Milano, Italy,  
<sup>2</sup> Department of Material Science and Nanotechnology, University of Milano-Bicocca, Milano, Italy,  
<sup>3</sup> Pirelli Tyre S.p.a., Viale Piero e Alberto Pirelli, 25, Milano, Italy



## Background & Rationale

Tire and road wear particles (TRWPs) represent a significant, recalcitrant environmental pollutant, yet their biodegradation remains significantly under-researched compared to plastics. Existing literature predominantly focuses on isolated microbial strains, overlooking the complex synergistic interactions within consortia required to break down the resilient rubber matrix. This study employs a dual metagenomic approach combining 16S rRNA gene sequencing and Shotgun metagenomics with selective enrichment from high-traffic roadside soils to map the rubber-degradative potential of both culturable and non-culturable taxa. By shifting the focus from single strains to functional microbial communities, we provide novel insights into the taxonomic shifts driven by rubber as a sole carbon source, identifying key players for future bioremediation strategies.



## 16S rRNA gene sequencing










Figure 1: Taxonomic profiles of the soil and liquid samples at phylum level. The profiles include samples collected far from the street (C\_STR), near the street (S\_STR), and those derived from the soil enrichment experiment. The last differentiate between attached (ATT) and planktonic (LIQ) microorganisms grown on either GUM or BT (Tread Compound) rubber films (GUM\_ATT, GUM\_LIQ, BT\_ATT, BT\_LIQ).

Figure 2: Krona charts of single metagenomes show each metagenome represented as concentric rings in a circular form, with each ring corresponding to a taxonomic rank. The white rings in each figure represent the bacteria which in this study is 100% of the population. The area of each ring reflects its abundance in the sample. Krona chart of S\_STR (A) and BT\_ATT (B & C). The taxonomic classification goes from phylum (A & B) to genus level (C).

## Shotgun metagenomics



## Conclusions & Perspectives

Comparative metagenomic analyses reveal significant community shifts driven by selective pressure, providing novel insights into how environments respond to rubber pollutants. These results characterize microbial communities potential for breakdown, establishing a foundational framework for future TRWPs bioremediation, sustainable waste management and a taxonomic identification pipeline.

### CONTACT ME !

Dipartimento di Scienza dei Materiali & Dipartimento di Biotecnologie e Bioscienze  
 Università degli Studi di Milano-Bicocca, P.zza della Scienza, 4 - 20126 Milano, Italy  
 a.dicesare3@campus.unimib.it  
 Tel: 0264483439

