

Determination of the spin and parity of all-charm tetraquarks

<https://doi.org/10.1038/s41586-025-09711-7>

The CMS Collaboration*✉

Received: 9 June 2025

Accepted: 6 October 2025

Published online: 3 December 2025

Open access

 Check for updates

The traditional quark model^{1,2} accounts for the existence of baryons, such as protons and neutrons, which consist of three quarks, as well as mesons, composed of a quark–antiquark pair. Only recently has substantial evidence started to accumulate for exotic states composed of four or five quarks and antiquarks³. The exact nature of their internal structure remains uncertain^{4–29}. Here we report the first measurement of quantum numbers of the recently discovered family of three all-charm tetraquarks^{30–32}, using data collected by the CMS experiment at the Large Hadron Collider from 2016 to 2018 (refs. 33,34). The angular analysis techniques developed for the discovery and characterization of the Higgs boson^{35–37} have been applied to the new exotic states. Here we show that the quantum numbers for parity P and charge conjugation C symmetries are found to be $+1$. The spin J of these exotic states is determined to be consistent with $2\hbar$, while $0\hbar$ and $1\hbar$ are excluded at 95% and 99% confidence levels, respectively. The $J^{PC} = 2^{++}$ assignment implies particular configurations of constituent spins and orbital angular momenta, which constrain the possible internal structure of these tetraquarks.

In 1964, Gell-Mann¹ and Zweig² independently proposed that hadrons, such as protons and neutrons, are made up of elementary particles called quarks (q). They suggested that mesons and baryons consisted of quark–antiquark pairs ($q\bar{q}$) and quark triplets (qqq), respectively. At the time, only three quark types were proposed: the up (u) and down (d), which form protons (uud) and neutrons (udd) and the strange quarks. The discovery of the J/Ψ meson in 1974 (refs. 38,39) was soon recognized as evidence of a fourth quark type, charm (c), bound with its corresponding antiquark. Gell-Mann and Zweig also pointed out that the symmetry principles underlying their model allowed for ‘exotic’ configurations, such as tetraquark and pentaquark systems. Despite numerous searches, the existence of these particles remained uncertain by the end of the last century⁴⁰.

The study of exotic hadrons advanced when access to large samples of hadrons containing heavy c and b quarks became available, highlighted by the 2003 discovery of the new particle $X(3872)$ in $J/\Psi \pi^+ \pi^-$ decays⁴¹. This particle has been interpreted as a tetraquark candidate with content $uc\bar{u}\bar{c}$ and recently classified as $\chi_{c1}(3872)$ (ref. 3), where the number denotes its mass in MeV (megaelectronvolts) and we adopt natural units by setting $c = 1$ and $\hbar = 1$. The discovery of several other candidates for tetraquarks and pentaquarks followed, including a $uc\bar{d}\bar{c}$ candidate $Z_c(3900)^+ \rightarrow J/\Psi \pi^+$ (refs. 42,43), also classified as $T_{cc1}(3900)^+$ (ref. 3). However, the true nature of their structure continues to be widely debated^{4–29}. Most interpretations fall into two categories: tightly bound states of two quarks and two antiquarks, similar to how quarks are bound within protons and neutrons, or loosely bound molecules composed of two mesons, similar to how protons and neutrons compose an atomic nucleus.

The next breakthrough in understanding exotic hadrons may come from studying those composed entirely of heavy quarks. The LHCb, ATLAS and CMS Collaborations observed a state $X(6900)$ (refs. 30–32),

also referred to as $T_{ccc}(6900)^0$ (ref. 3), in the $J/\Psi J/\Psi$ final state. The CMS Collaboration also reported two additional states, $X(6600)$ and $X(7100)$. The triple-peaking structure in the di- J/Ψ invariant mass distribution observed by CMS³² is shown in Fig. 1 and is best described by a quantum-mechanical interference of three amplitudes representing $cc\bar{c}\bar{c}$ states. The interference pattern among the three resonances, along with the mass spacings that follow a radial Regge trajectory²⁹, suggests that these three X particles form a family of states with identical properties, differing only in the radial excitations of their wavefunctions. These properties, which include the spin and symmetry quantum numbers, provide insights into the internal structure of these exotic hadrons and are the primary focus of the experimental study reported here. The corresponding tabulated results are available in the HEPData record for this analysis⁴⁴.

The spin and symmetry properties

Spin is a type of intrinsic angular momentum carried by particles. For composite particles, the total spin J represents the total angular momentum of the system, determined by the combination of the spins S and orbital angular momenta L of the elementary constituent particles. The quantum number for parity (P) symmetry describes how a system behaves under a spatial inversion or, equivalently, a mirror reflection. The charge conjugation (C) symmetry describes how a system transforms when every particle is replaced by its corresponding antiparticle. The $P = \pm 1$ and $C = \pm 1$ values indicate whether the wavefunction of the state changes sign under the respective transformation.

The quarks and antiquarks have a spin of $S = 1/2$, the fundamental quantum of spin, and they serve as the building blocks of the tetraquark state shown in Fig. 2. The quarks inside a hadron can carry three different colour charges, associated with the strong interaction,

*A list of authors and their affiliations appears online. ✉e-mail: cms-publication-committee-chair@cern.ch




Fig. 1 | Candidates for all-charm tetraquarks. The $J/\psi J/\psi \rightarrow \mu^+ \mu^- \mu^+ \mu^-$ invariant mass $m_{4\mu}$ spectrum shows the three exotic states, X(6600), X(6900) and X(7100). Parameterizations of these states are shown both individually and as a combined signal that includes quantum-mechanical interference (denoted by Total signal). The full model³² incorporates both signal and background components, with the background originating from di- J/ψ production, including contributions from nonresonant production and an enhancement near the kinematic threshold of 6.2 GeV.

conventionally referred to as blue, green and red. Antiquarks carry corresponding anticoulour charges (antiblue, antigreen and antired), which are shown as yellow, magenta and cyan, respectively, in Fig. 2. Quarks and antiquarks are held together through the exchange of gluons, which are the mediators of the strong interaction. The hadron as a whole is colour-charge-neutral.

A tightly bound tetraquark state X is shown in Fig. 2 (top). In the ground state with zero orbital angular momentum, two identical charm quarks form an antisymmetric colour state for attraction, leading to symmetric spatial and spin states with total spin 1 (ref. 25). The two charm antiquarks do the same. Each pair carries a colour charge as well, and the two pairs attract each other, forming a strongly bound tetraquark state that is colour-charge-neutral, similar to a quark-antiquark bound state in a meson. The orbital angular momentum L between the quark pair and the antiquark pair can take non-negative integer values. The corresponding parity of the system is then given by $P = (-1)^L$, which arises from the behaviour of the spherical harmonics under spatial inversion.

The lowest and most probable energy state with $L = 0$ is spatially symmetric, $P = +1$. The spins of the two systems combine in a symmetric configuration to yield a total spin $J = 0$ or 2 with $C = +1$, or in an antisymmetric configuration to give $J = 1$ with $C = -1$. For the cc and $\bar{c}\bar{c}$ system, charge conjugation is equivalent to exchanging the pairs, resulting in the associated symmetry. As we will demonstrate later, the $C = -1$ configuration is ruled out and will, therefore, not be considered further. For an antisymmetric spatial state with $L = 1$ and $P = -1$, the spins combine to 1 in an antisymmetric configuration and with $C = +1$, resulting in possible total spins $J = 0, 1$ or 2. States with $L = 2$ are also possible, resulting in $P = +1$ and allowing J values up to 4 when the spins combine to 0 or 2. However, the high orbital angular momentum requires additional energy, making these states less likely.

An alternative model, shown in Fig. 2 (bottom), is a loosely bound molecule of two $c\bar{c}$ mesons. The lowest-energy configuration corresponds to an orbital angular momentum $L = 0$ between the two mesons, resulting in $P = +1$. A key distinction is that, unlike in a tightly bound tetraquark, the two constituent $c\bar{c}$ mesons are not restricted to form spin-1 states. Consequently, lower total spin values such as $J = 0$ or 1 are more likely, although higher spin states cannot be excluded. Another difference is the weaker interaction between the $c\bar{c}$ mesons. Similar to




Fig. 2 | Internal structure models for the particle X. The particle X, composed of $cc\bar{c}\bar{c}$, is shown at rest. Two models of the internal structure of X are presented: a tightly bound tetraquark (top) and a loosely bound molecule of two mesons (bottom). The colours assigned to individual quarks or quark pairs denote possible colour charge assignments in strong interactions, in which attractive forces are mediated by gluon exchange (shown as wavy lines) and meson exchange (shown as a solid pair of arrows). The X decays into two J/ψ mesons with spin projections λ_i along their respective directions of motion; each meson then decays into a $\mu^+ \mu^-$ pair. The polar and azimuthal angles $\Omega_i = (\theta_i, \phi_i)$ describe the direction of the μ^- relative to the z_i -axis, which is defined to point opposite to the X-direction in the centre-of-mass frame of the corresponding J/ψ meson, for $i = 1$ and 2.

how a deuteron is a bound state of a proton and a neutron, the two colour-charge-neutral systems are bound through the exchange of a meson by the Yukawa interaction⁴⁵. However, unlike the deuteron, in an all-charm tetraquark molecule, the exchanged meson must contain charm quarks. A heavier exchange meson substantially suppresses the Yukawa interaction, making the formation of bound states less likely. However, alternative empirical models for these interactions have also been explored^{9–12}.

The three X states under investigation have invariant masses ranging between 6.2 GeV and 8.0 GeV (gigaelectronvolts), as shown in Fig. 1, mean lifetimes between 10^{-24} s and 10^{-23} s (ref. 32), and they decay into either two J/ψ mesons, or potentially several other, yet unobserved, final states. The J/ψ meson has a mass of 3.1 GeV, spin 1, a mean lifetime of approximately 7×10^{-21} s, and in 6% of the cases, the $c\bar{c}$ pair in the J/ψ annihilates into a $\mu^+ \mu^-$ pair³, which is ideal for the detection of the final states and for performing an angular analysis.

Angular distributions

The X quantum numbers can be inferred from the polarizations of the J/ψ mesons, which, in turn, result in specific angular distributions of the decay muons. Figure 2 defines λ_1 and λ_2 as the spin projections of the two J/ψ mesons along their respective directions of motion. Both λ_1 and λ_2 can take values of $-1, 0$ or $+1$. Let $A_{\lambda_1 \lambda_2}$ be the quantum-mechanical amplitude that describes the decay of X into two mesons with spin projections λ_1 and λ_2 . The nine values of $|A_{\lambda_1 \lambda_2}|^2$ can be interpreted as relative probabilities for specific polarization states of the two J/ψ mesons.

The angular distributions in the decay $X \rightarrow J/\psi J/\psi \rightarrow (\mu^+ \mu^-)(\mu^+ \mu^-)$ can be expressed as a function of three angular observables θ_1, θ_2 and $\Phi = (\Phi_1 + \Phi_2)$ as shown in Fig. 2 and further explained in the section ‘Angular observables’. Here, θ_1 and θ_2 are the helicity angles of the two J/ψ mesons, and Φ represents the angle between their decay planes. The nine amplitudes $A_{\lambda_1 \lambda_2}$ appear in the coefficients of the expression. The analytical and numerical calculations of the angular distributions are detailed in refs. 35,46.

We base the prediction of $A_{\lambda_1 \lambda_2}$ on symmetry considerations⁴⁶. Angular momentum conservation implies that $|\lambda_1 - \lambda_2| \leq J$. For two identical J/ψ particles, the relation $A_{\lambda_1 \lambda_2} = (-1)^J A_{\lambda_2 \lambda_1}$ must hold. Finally, assuming that the X has definite P and C quantum numbers, which are

Table 1 | Quantum numbers

J^{PC}	Models J_i^P	Contributing amplitudes
0^{-+}	0^{-}	$A_{++} = -A_{--}$
0^{++}	0_m^+	A_{00} and $A_{++} = A_{--}$
	0_h^+	$A_{++} = A_{--}$ and A_{00}
1^{-+}	1^{-}	$A_{+0} = -A_{0+} = A_{-0} = -A_{0-}$
1^{++}	1^{+}	$A_{+0} = -A_{0+} = -A_{-0} = A_{0-}$
2^{-+}	2_m^{-}	$A_{++} = -A_{--}$
	2_h^{-}	$A_{+0} = A_{0+} = -A_{-0} = -A_{0-}$
2^{++}	2_m^{+}	$A_{+0} = A_{0+} = A_{-0} = A_{0-}$, $A_{+-} = A_{-+}$, A_{00} and $A_{++} = A_{--}$

The possible assignments of quantum numbers J^{PC} , the J_i^P models considered, and the contributing amplitudes in the decay $X \rightarrow J/\psi J/\psi$ are presented.

conserved in the strong decays, we conclude that $A_{\lambda_1\lambda_2} = P(-1)^J A_{(-\lambda_2)(-\lambda_1)}$ and the charge conjugation $C = +1$. The latter is due to the $C = +1$ of the $J/\psi J/\psi$ final state. This results in six possible assignments for the J^{PC} quantum numbers with the corresponding contributions from the amplitudes shown in Table 1. States J^{PC} with $J \geq 3$ would exhibit amplitudes and angular distributions similar to those of 2^{PC} , as inferred from ref. 46.

As shown in Table 1, in the scenarios of 0^{-+} , 1^{-+} and 1^{++} , the relative contributions of all amplitudes $A_{\lambda_1\lambda_2}$ are predetermined. In the scenarios of 0^{++} and 2^{-+} , the relative contributions of the two different sets of amplitudes, associated with different models J_i^P listed in Table 1, are not known a priori. The 2^{++} scenario has four contributions with an undetermined relationship. When pure symmetry alone does not provide a determination, the dynamical properties of the interactions provide additional constraints that relate the contributions. Assuming the resonances differ only by radial excitations of their wavefunctions, we expect all three to exhibit the same structure in their decay interactions. This structure and the corresponding models are defined and motivated in the section ‘Modelling of hadron properties’.

The 0_m^+ model describes the simplest possible structure for the interaction of a spin-0 particle decaying into two spin-1 particles, with the subscript ‘m’ indicating its minimal nature. By contrast, the 0_h^+ model corresponds to a more complex interaction structure, as denoted by the subscript ‘h’ for higher complexity³⁵. The 2_m^{-} and 2_h^{-} models correspond to $A_{++} = -A_{--}$ and $A_{+0} = A_{0+} = -A_{-0} = -A_{0-}$, and their decay angular distributions are indistinguishable from those of 0^{-} and 1^{-} models, respectively. Quantum-mechanical interference between models 0_m^+ and 0_h^+ , or 2_m^{-} and 2_h^{-} , is also taken into account in the data analysis. The 2_m^+ model, as defined in ref. 35, corresponds to the minimal structure of interactions of a spin-2 particle in its decay. It is used to represent the 2^{++} quantum numbers with a distinct contribution from $A_{+-} = A_{-+}$, which is unique to 2^{++} . Moreover, seven other amplitudes contribute simultaneously, another feature unique to 2^{++} . Other possible 2^{++} models, beyond the minimal structure, may exhibit angular distributions similar to those of the 0^{++} or 1^{++} states and are not tested separately.

The reasoning outlined above leads to the predicted probability distributions $\mathcal{P}_i(\theta_1, \theta_2, \Phi, m_{4\mu})$ for the eight individual J_i^P models listed in Table 1, as well as their possible mixtures, at any mass value $m_{4\mu}$ for the particle decay $X \rightarrow J/\psi J/\psi$. Our analysis aims to identify the model that best matches the experimental data.

Experimental data

To create new particles, proton beams are accelerated to a combined energy of 13 TeV (tera-electronvolts) in opposite directions within the Large Hadron Collider (LHC)⁴⁷, an international project hosted by CERN, and collided head-on. Protons are baryons with a mass of 0.94 GeV, composed of three valence quarks. However, approximately half of their momentum is carried by gluons, which bind the quarks together.

Furthermore, sea quark–antiquark pairs emerge and carry roughly one-fifth of the momentum of the proton. As a result, new particles can be created through collisions between gluons, quarks and antiquarks (collectively referred to as partons) within the protons. The data presented in this paper were collected from 2016 to 2018, with a total integrated luminosity of 135 fb^{-1} (inverse femtobarns), which represents approximately 1.5×10^{16} proton–proton (pp) collisions.

Once produced, the X particles immediately decay within the CMS detector^{33,34}, a large general-purpose international experiment at the LHC. Its essential component is a superconducting solenoid with an internal diameter of 6 m, which can generate a magnetic field of 3.8 T (tesla). Within the volume of the solenoid, there are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter and a brass and scintillator hadron calorimeter, each consisting of a barrel and two endcap sections. Muons are identified using gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid. In this analysis, the primary measurements of muon momentum vectors are obtained from the curvature and direction of their paths in the silicon tracker. A fast, real-time data recording decision is made using a two-level trigger system⁴⁸, which reduces the recorded data rate to approximately 1 kHz (kilohertz).

To select events of interest, the trigger criteria include having three muons each with a minimum muon momentum transverse to the beam (p_T) of 3 GeV. The procedure for the offline selection of X candidates is described in ref. 32: there must be at least four muons, each with $p_T > 2 \text{ GeV}$; $\mu^+ \mu^-$ pairs must originate from a common vertex, have a mass consistent with a J/ψ , and the muons from the highest- $p_T J/\psi$ satisfy $p_T > 3.5 \text{ GeV}$. A fit is applied to muon pairs, using the J/ψ meson mass and the common production vertex as constraints, to improve the $m_{4\mu}$ resolution. A total of 8,651 candidate events with two J/ψ mesons have been recorded with a combined mass in the range of 6.2–9.0 GeV, as shown in Fig. 1, with approximately a quarter expected to correspond to the three X states, and the remainder mostly originating from background J/ψ pair production. These candidates are the same as those in ref. 32. For each X candidate, we record the momentum vectors of all four muons for the analysis that follows.

Angular analysis

The objective of the analysis is to compare the angular distributions in the decay of X particles observed in the experiment with the predicted distributions for various J^{PC} models and select the model that is consistent with the data. To prevent a potential bias, the observed distributions were hidden until all aspects of the analysis were finalized. The predicted distributions $\mathcal{P}_i(\theta_1, \theta_2, \Phi, m_{4\mu})$ are precisely known for the muons originating from the decay, but the observed distributions are modified by detector effects. This is because the detection probabilities of the reconstructed muons depend on their momenta and directions, which have complex correlations with the angles. Moreover, the momenta and directions themselves are subject to uncertainties. To account for these effects and to accurately model the background, a comprehensive computer simulation of pp collisions is performed using Monte Carlo (MC) techniques. The simulation proceeds in four stages.

In the first stage, the JHUGEN program, v.7.5.7 (refs. 35,46), originally designed to model variations in the properties of the Higgs boson, is used to simulate the process $pp \rightarrow X \rightarrow J/\psi J/\psi \rightarrow \mu^+ \mu^- \mu^+ \mu^-$. Two models of X production are explored^{9–12}: direct production through parton annihilation within protons and fragmentation of a secondary gluon or quark generated in collisions. In the case of $J \geq 1$, the first model can lead to a preferred spin polarization of X along the beam axis, whereas the second model may result in polarization along the direction of motion of the X particle. JHUGEN enables the modelling of either polarization or the simulation of an unpolarized state through the application of event weights. Polarization variations are used to estimate uncertainties, with the unpolarized case assumed by default.




Fig. 3 | Analysis of angular distributions. **a**, Distributions of $D_{2_m^+ 0^-}$ for the 0^- , 2_m^- , and 2_m^+ models in the range $6.2 < m_{4\mu} < 8.0$ GeV. Distributions for signal only (dashed) and for signal plus background (solid and dash-dot-dotted) models are compared with the experimental data points with error bars, with uncertainty bands representing post-fit model uncertainties, which are

The three stages of the simulation that follow are shared with thousands of other processes analysed by the CMS Collaboration and have, therefore, been extensively tuned to ensure a good agreement with the experimental data. The simulation of the remaining particles appearing in proton collisions and surrounding the four muons is performed by the PYTHIA program, v.8.240 (ref. 49). The simulation takes into account the effects of extra pp collisions that happen at the same time or close in time to the main collision. The generated events are further processed through a dedicated CMS detector simulation, based on the GEANT4 program⁵⁰, which models the detector response. Finally, the simulated events undergo the complete detector reconstruction process, using software and algorithms identical to those used to analyse the experimental data.

The nonresonant J/ψ J/ψ background is simulated with PYTHIA in the first stage and covers cases in which the two mesons are produced either in the same or in different parton collisions in a single pp interaction. The simulated angular distributions were found to be compatible with data in the mass sideband above 8 GeV. The momentum of the J/ψ J/ψ pair in all three directions was tuned for both X and nonresonant production using PYTHIA settings in the second stage to match the data observed in both the sideband and X signal regions.

We conduct pairwise comparisons between different models characterized by distinct J^{PC} quantum numbers. Instead of analysing the multidimensional space of angular observables (θ_1 , θ_2 , Φ) directly, we define a single observable, D_{ij} , constructed from the likelihood ratio between two models labelled i and j . It relies on quantum-mechanical calculations, as detailed in the section ‘Matrix element approach’ and ref. 35. This discriminant D_{ij} depends on the angular observables and is optimal for discriminating between the two hypotheses i and j . The final statistical analysis of the data is carried out using two-dimensional distributions of the events, $\mathcal{P}_{ij}(m_{4\mu}, D_{ij})$. An example of the predicted distributions for the $D_{2_m^+ 0^-}$ observable is presented in Fig. 3a, in which the 0^- , 2_m^- , and 2_m^+ models are shown, along with the background and experimental data. Projecting the two-dimensional distribution ($m_{4\mu}, D_{ij}$) onto the single observable D_{ij} results in a partial loss of statistical power.

Quantum number determination

To distinguish alternative models, the test statistic $q = -2\ln(\mathcal{L}_{J_i^P}/\mathcal{L}_{J_j^P})$ is defined using the likelihood ratio of signal plus background likelihoods for the two signal hypotheses J_i^P and J_j^P , based on the two

partially correlated with the data. The 0^- and 2_m^- distributions are identical apart from systematic uncertainties arising from polarization effects. **b**, Normalized distributions of the test statistic $q = -2\ln(\mathcal{L}_{0^-}/\mathcal{L}_{2_m^+})$ from pseudo-experiments generated under the 2_m^+ (blue, right) and 0^- (orange, left) hypotheses, with the arrow indicating the observed value q_{obs} .

observables ($m_{4\mu}, D_{ij}$), as outlined in the section ‘Statistical analysis’. In a maximum likelihood fit, the likelihood function \mathcal{L} is maximized with respect to the nuisance parameters, which include the yields of signal and background processes, as well as constrained parameters that account for systematic uncertainties. These uncertainties encompass variations in mass shapes and in the discriminants used to model both signal and background components, as detailed in the section ‘Systematic uncertainties’.

An example of the observed value of q , along with the expected distributions for the 2_m^+ and 0^- models, is shown in Fig. 3b. The expectations for each model are derived from a large ensemble of pseudo-experiments, with systematic uncertainties from the parameterization included in the fit procedure. These pseudo-experiments are generated according to the observed data shown in Fig. 1, combined with the kinematic distributions specific to each model. Given the observed value of q , the 2_m^+ model is preferred over 0^- . All the steps outlined above for testing the spin-parity properties of the X states closely follow the methods and tools developed for the discovery and characterization of the Higgs boson in its four-lepton decay in 2012 (refs. 35–37).

Following this methodology, all pairs of the eight J_i^P models listed in Table 1 are tested, and in each case involving the 2_m^+ model, the latter is favoured. The corresponding results are shown in Fig. 4 and Table 2, in which the probability, P -value, and the associated Z -score, expressed as the number of standard deviations derived from the one-sided Gaussian tail integral, are shown for an alternative model J_i^P tested against the 2_m^+ model.

Building on this observation, a mixture of the 0_m^+ and 0_h^+ structures of interactions, as well as that of 2_m^- and 2_h^- , is tested in 10 equal fractional increments against the 2_m^+ model. The 2_m^+ model is again preferred. In each case, one of the mixed models shown in Fig. 4, denoted as 0_{mix}^+ or 2_{mix}^- , represents the mixed scenario with the least separation from 2_m^+ and is listed in Table 2. A mixture of 2_m^- and 2_h^- contributions does not produce interference because of their differing spin projections along the decay axis, as reflected in the amplitude composition presented in Table 1. We examine both constructive and destructive interference between the 0_m^+ and 0_h^+ models, by considering both positive and negative relative signs between their contributing amplitudes. Constructive interference results in the smallest deviation from the 2_m^+ model, except in the first 0_{mix}^+ step shown in Fig. 4, in which the sign-induced model differences are minimal and destructive interference yields a slightly smaller separation.




Fig. 4 | Summary of statistical tests. Distributions of the test statistic q for various J_i^P hypotheses tested against the 2_m^+ model. The observed q_{obs} values are indicated by the black dots. The expected median and the 68.3%, 95.4% and 99.7% confidence level regions for the 2_m^+ model (blue, left) and for each of the

alternative J_i^P hypotheses (orange, right) are shown. The first entry corresponding to 0^- reflects the information shown in Fig. 3b. For 0^- and 2^- models, 11 points correspond to varying fractions in the mixture of the two structures of interaction.

Based on the results presented in Fig. 4 and Table 2, the tests for the $J^{PC} = 0^-$ and 1^+ scenarios reject these hypotheses with a significance level exceeding 5 standard deviations when compared with a 2_m^+ model. The 2^- scenario, along with higher spin values that have the same P and C quantum numbers, is excluded at a significance level of 3 standard deviations. This establishes the quantum numbers $P = +1$ and $C = +1$, as shown by the decay final-state particles and their distributions.

The 1^+ scenario is excluded at more than a 99% confidence level when compared with the 2_m^+ model. The $J^{PC} = 0^+$ scenario, when considering a combination of possible amplitudes, is excluded at more than a 95% confidence level when compared with the 2_m^+ model. It is important to emphasize that the selected 2_m^+ model represents just one possible realization of the $J^{PC} = 2^{++}$ scenario, and an admixture of other amplitudes could lead to angular distributions resembling those of the 0^+ or 1^+ scenarios. The $J \geq 3$ quantum numbers are still possible, but $J = 2$ is more likely, due to the additional energy needed to achieve a higher angular excitation of the hadronic states with $L \geq 2$. This makes the $J^{PC} = 2^{++}$ interpretation preferred for the fully charmed tetraquark states $X(6600)$, $X(6900)$ and $X(7100)$.

Discussion

The study of tetraquark states has attracted considerable interest because of its potential to provide insight into the structure of the

hadronic matter that makes up the world around us^{4–8,8–29}. In this study, we have presented the first measurements of the quantum numbers for a recently discovered family of three all-charm tetraquarks, based on data collected by the CMS experiment at the LHC. Our results, summarized in Table 2, favour a $J^{PC} = 2^{++}$ assignment.

For a system of two constituents, as represented by either model in Fig. 2, an orbital angular momentum of $L = 1$ is excluded by the requirement $P = +1$, whereas higher orbital excitations with $L \geq 2$ are energetically disfavoured. This makes the S -wave ($L = 0$) configuration the most likely. In a molecular scenario, the quark–antiquark pairs are not required to be spin-1 mesons, making a $J = 2$ configuration less likely. This picture agrees with the existing data for all other well-established tetraquark candidates with measured spin, such as the $X(3872)$ and $Z_c(3900)^+$, all of which have $J < 2$ (ref. 3). By contrast, a tightly bound $cc\bar{c}\bar{c}$ tetraquark with a diquark–antidiquark configuration requires both diquarks to be in spin-1 states, which restricts the total spin to $J = 0$ or $J = 2$, making $J = 2$ a natural choice. This spin-1 diquark requirement, however, does not apply to tetraquark candidates with mixed-flavour quark content, a consideration relevant to all previously observed candidates, which contained both heavy and light quarks³. Other theoretical considerations also favour $J = 2$ for the tightly bound tetraquark states^{18,28}.

This advancement in understanding exotic hadrons was enabled by the study of all-heavy tetraquarks, and it brings us closer to uncovering their internal structure. Although our findings do not definitively distinguish between tightly bound tetraquark and meson–meson molecular models, they provide constraints on the possible internal structures and favour the tightly bound scenario.

Table 2 | Summary of statistical tests

J_i^P	<i>P</i> -value	Z-score reject J_i^P
0^-	2.7×10^{-13}	7.2
0_m^+	4.3×10^{-5}	3.9
0_{mix}^+	1.4×10^{-2}	2.2
0_h^+	3.1×10^{-9}	5.8
1^-	8.0×10^{-8}	5.2
1^+	4.7×10^{-3}	2.6
2_m^-	4.1×10^{-12}	6.8
2_{mix}^-	6.5×10^{-4}	3.2
2_h^-	2.2×10^{-8}	5.5

The *P*-value and the associated Z-score are shown for alternative models J_i^P , tested against the 2_m^+ model. A higher Z-score implies that the model is less compatible with the observation.

Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41586-025-09711-7>.

- Gell-Mann, M. A schematic model of baryons and mesons. *Phys. Lett.* **8**, 214–215 (1964).
- Zweig, G. in *Developments in the Quark Theory of Hadrons* Vol. 1 1964–1978 (eds Lichtenberg, D. B. & Rosen, S. P.) 22–101 (Hadronic Press, 1980).
- Particle Data Group. Review of particle physics. *Phys. Rev. D* **110**, 030001 (2024).
- Berezhnoy, A. V., Luchinsky, A. V. & Novoselov, A. A. Heavy tetraquarks production at the LHC. *Phys. Rev. D* **86**, 034004 (2012).

5. Wu, J., Liu, Y.-R., Chen, K., Liu, X. & Zhu, S.-L. Heavy-flavored tetraquark states with the $QQ\bar{Q}\bar{Q}$ configuration. *Phys. Rev. D* **97**, 094015 (2018).
6. Karlöner, M., Nussinov, S. & Rosner, J. L. $QQ\bar{Q}\bar{Q}$ states: masses, production, and decays. *Phys. Rev. D* **95**, 034011 (2017).
7. Richard, J.-M., Valcarce, A. & Vijande, J. String dynamics and metastability of all-heavy tetraquarks. *Phys. Rev. D* **95**, 054019 (2017).
8. Anwar, M. N., Ferretti, J., Guo, F.-K., Santopinto, E. & Zou, B.-S. Spectroscopy and decays of the fully-heavy tetraquarks. *Eur. Phys. J. C* **78**, 647 (2018).
9. Nielsen, M. et al. Supersymmetry in the double-heavy hadronic spectrum. *Phys. Rev. D* **98**, 034002 (2018).
10. Nielsen, M. & Brodsky, S. J. Hadronic superpartners from a superconformal and supersymmetric algebra. *Phys. Rev. D* **97**, 114001 (2018).
11. Ali, A., Maiani, L. & Polosa, A. D. *Multiquark Hadrons* (Cambridge Univ. Press, 2019).
12. Brambilla, N. et al. The XYZ states: experimental and theoretical status and perspectives. *Phys. Rep.* **873**, 1–154 (2020).
13. Liu, M.-S., Lü, Q.-F., Zhong, X.-H. & Zhao, Q. All-heavy tetraquarks. *Phys. Rev. D* **100**, 016006 (2019).
14. Wang, G.-J., Meng, L. & Zhu, S.-L. Spectrum of the fully-heavy tetraquark state $QQ\bar{Q}\bar{Q}'$. *Phys. Rev. D* **100**, 096013 (2019).
15. Bedolla, M. A., Ferretti, J., Roberts, C. D. & Santopinto, E. Spectrum of fully-heavy tetraquarks from a diquark+antidiquark perspective. *Eur. Phys. J. C* **80**, 1004 (2020).
16. Liu, M.-S., Liu, F.-X., Zhong, X.-H. & Zhao, Q. Fully heavy tetraquark states and their evidences in LHC observations. *Phys. Rev. D* **109**, 076017 (2024).
17. Jin, X., Xue, Y., Huang, H. & Ping, J. Full-heavy tetraquarks in constituent quark models. *Eur. Phys. J. C* **80**, 1083 (2020).
18. Becchi, C., Ferretti, J., Giachino, A., Maiani, L. & Santopinto, E. A study of $cc\bar{c}\bar{c}$ tetraquark decays in 4 muons and in $D^{(*)}\bar{D}^{(*)}$ at LHC. *Phys. Lett. B* **811**, 135952 (2020).
19. Chen, H.-X., Chen, W., Liu, X. & Zhu, S.-L. Strong decays of fully-charm tetraquarks into di-charmonia. *Sci. Bull.* **65**, 1994–2000 (2020).
20. Wang, J.-Z., Chen, D.-Y., Liu, X. & Matsuki, T. Producing fully charm structures in the J/ψ -pair invariant mass spectrum. *Phys. Rev. D* **103**, 071503 (2021).
21. Dong, X.-K. et al. Coupled-channel interpretation of the LHCb double- J/ψ spectrum and hints of a new state near the $J/\psi/J/\psi$ threshold. *Phys. Rev. Lett.* **126**, 132001 (2021); erratum **127**, 119901 (2021).
22. Zhang, H.-F., Ma, Y.-Q. & Sang, W.-L. Perturbative QCD evidence for spin-2 particles in the di- J/ψ resonances. *Sci. Bull.* **70**, 1915–1917 (2025).
23. Zhu, R. Fully-heavy tetraquark spectra and production at hadron colliders. *Nucl. Phys. B* **966**, 115393 (2021).
24. Liu, F.-X., Liu, M.-S., Zhong, X.-H. & Zhao, Q. Higher mass spectra of the fully-charmed and fully-bottom tetraquarks. *Phys. Rev. D* **104**, 116029 (2021).
25. Faustov, R. N., Galkin, V. O. & Savchenko, E. M. Fully heavy tetraquark spectroscopy in the relativistic quark model. *Symmetry* **14**, 2504 (2022).
26. Feng, F. et al. Inclusive production of fully charmed tetraquarks at the LHC. *Phys. Rev. D* **108**, L051501 (2023).
27. Celiberto, F. G., Gatto, G. & Papa, A. Fully charmed tetraquarks from LHC to FCC: natural stability from fragmentation. *Eur. Phys. J. C* **84**, 1071 (2024).
28. Belov, I., Giachino, A. & Santopinto, E. Fully charmed tetraquark production at the LHC experiments. *J. High Energy Phys.* **1**, 093 (2025).
29. Zhu, F., Bauer, G. & Yi, K. Experimental road to a charming family of tetraquarks ... and beyond. *Chin. Phys. Lett.* **41**, 111201 (2024).
30. LHCb Collaboration. Observation of structure in the J/ψ -pair mass spectrum. *Sci. Bull.* **65**, 1983–1993 (2020).
31. ATLAS Collaboration. Observation of an excess of dicharmonium events in the four-muon final state with the ATLAS detector. *Phys. Rev. Lett.* **131**, 151902 (2023).
32. CMS Collaboration. New structures in the $J/\psi/J/\psi$ mass spectrum in proton-proton collisions at $\sqrt{s} = 13$ TeV. *Phys. Rev. Lett.* **132**, 111901 (2024).
33. CMS Collaboration. The CMS experiment at the CERN LHC. *J. Instrum.* **3**, S08004 (2008).
34. CMS Collaboration. Development of the CMS detector for the CERN LHC Run 3. *J. Instrum.* **19**, P05064 (2024).
35. Bolognesi, S. et al. On the spin and parity of a single-produced resonance at the LHC. *Phys. Rev. D* **86**, 095031 (2012).
36. CMS Collaboration. Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett. B* **716**, 30–61 (2012).
37. CMS Collaboration. Study of the mass and spin-parity of the Higgs boson candidate via its decays to Z boson pairs. *Phys. Rev. Lett.* **110**, 081803 (2013).
38. E598 Collaboration. Experimental observation of a heavy particle. *J. Phys. Rev. Lett.* **33**, 1404–1406 (1974).
39. SLAC-SP-017 Collaboration. Discovery of a narrow resonance in e^+e^- annihilation. *Phys. Rev. Lett.* **33**, 1406–1408 (1974).
40. Landsberg, L. G. The search for exotic hadrons. *Phys. Usp.* **42**, 871–886 (1999).
41. Belle Collaboration. Observation of a narrow charmoniumlike state in exclusive $B^+ \rightarrow B^0 \pi^+ \pi^- J/\psi$ decays. *Phys. Rev. Lett.* **91**, 262001 (2003).
42. BESIII Collaboration. Observation of a charged charmoniumlike structure in $e^+e^- \rightarrow \pi^+ \pi^- J/\psi$ at $\sqrt{s} = 4.26$ GeV. *Phys. Rev. Lett.* **110**, 252001 (2013).
43. Belle Collaboration. Study of $e^+e^- \rightarrow \pi^+ \pi^- J/\psi$ and observation of a charged charmoniumlike state at Belle. *Phys. Rev. Lett.* **110**, 252002 (2013).
44. The CMS Collaboration. Determination of the spin and parity of all-charm tetraquarks. HEPData <https://doi.org/10.17182/hepdata.158584> (2025).
45. Yukawa, H. On the interaction of elementary particles. I. *Proc. Phys. Math. Soc. Jpn* **17**, 48–57 (1935).
46. Gao, Y. et al. Spin determination of single-produced resonances at hadron colliders. *Phys. Rev. D* **81**, 075022 (2010).
47. Evans, L., & Bryant, P. LHC machine. *J. Instrum.* **3**, S08001 (2008).
48. Khachatryan, V. et al. The CMS trigger system. *J. Instrum.* **12**, P01020 (2017).
49. Sjöstrand, T. et al. An introduction to PYTHIA 8.2. *Comput. Phys. Commun.* **191**, 159–177 (2015).
50. GEANT4 Collaboration. GEANT4—a simulation toolkit. *Nucl. Instrum. Methods Phys. Res. A* **506**, 250–303 (2003).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025, modified publication 2026

Article

The CMS Collaboration

A. Hayrapetyan¹, V. Makarenko¹, A. Tumasyan¹, W. Adam², J. W. Andrejkovic², L. Benato², T. Bergauer², M. Dragicic², C. Giordano², P. S. Hussain², M. Jeitler², N. Krammer², A. L. Li², D. Liko², M. Matthewman², I. Mikulec², J. Schieck², R. Schöfbeck², D. Schwarz², M. Shoshartar², M. Sonawane², W. Waltenberger², C.-E. Wulz², T. Janssen², H. Kwon³, D. Ocampo Henao³, T. Van Laer³, P. Van Mechelen³, J. Bierkens⁴, N. Breugelmans⁴, J. D'Hondt⁴, S. Dansana⁴, A. De Moor⁴, M. Delcourt⁴, F. Heyen⁴, Y. Hong⁴, P. Kashko⁴, S. Lowette⁴, I. Makarenko⁴, D. Müller⁴, J. Song⁴, S. Tavernier⁴, M. Tytgat^{4,5}, G. P. Van Onsem⁴, S. Van Putte⁴, D. Vannerom⁴, B. Bilin⁵, B. Clerbaux⁵, A. K. Das⁵, I. De Bruyn⁵, G. De Lentdecker⁵, H. Evard⁶, P. Favart⁶, P. Gianneos⁶, A. Khalilzadeh⁶, F. A. Khan⁶, A. Malara⁶, M. A. Shahzad⁶, L. Thomas⁶, M. Vanden Bemden⁶, C. Vander Velde⁶, P. Vanlaer⁶, F. Zhang⁶, M. De Coen⁵, D. Dobur⁵, G. Gokbulut⁵, J. Knolle⁵, L. Lambrecht⁵, D. Marckx⁵, K. Skovpen⁵, N. Van Den Bossche⁵, J. van der Linden⁵, J. Vandenbroeck⁵, L. Werenbeek⁵, S. Bein⁷, A. Benecke⁷, A. Bethani⁷, G. Bruno⁷, A. Cappati⁷, J. De Favereau De Jeneret⁷, C. Delaere⁷, A. Giammanco⁷, A. O. Guzel⁷, V. Lemaître⁷, J. Lidrych⁷, P. Malek⁷, P. Mastrapasqua⁷, S. Turkcapar⁷, G. A. Alves⁸, M. Barroso Ferreira Filho⁸, E. Coelho⁸, C. Hensel⁸, T. Menezes De Oliveira⁸, C. Mora Herrera^{8,9}, P. Rebelo Teles⁸, M. Soeiro⁸, E. J. Tonelli Manganote⁸, A. Vilela Pereira^{8,9}, W. L. Aldá Júnior⁹, H. Brandao Malbouisson⁹, W. Carvalho⁹, J. Chinellato⁹, M. Costa Reis⁹, E. M. Da Costa⁹, G. G. Da Silva⁹, D. De Jesus Damiao⁹, S. Fonseca De Souza⁹, R. Gomes De Souza⁹, S. S. Jesus⁹, T. Laux Kuhn⁹, M. Macedo⁹, K. Mota Amarilo⁹, L. Mundim⁹, H. Nogima⁹, J. P. Pinheiro⁹, A. Santoro⁹, A. Sznajder⁹, M. Thiel⁹, F. Torres Da Silva De Araujo⁹, C. A. Bernardes¹⁰, T. R. Fernandez Perez Tomei¹⁰, E. M. Gregores¹⁰, B. Lopes Da Costa¹⁰, I. Maiteti Silverio¹⁰, P. G. Mercadante¹⁰, S. F. Novaes¹⁰, B. Orzari¹⁰, Sandra S. Padula¹⁰, V. Scheurer¹⁰, A. Aleksandrov¹¹, G. Antchev¹¹, P. Danev¹¹, R. Hadjiiska¹¹, P. Iaydjiev¹¹, M. Misheva¹¹, M. Shopova¹¹, G. Sultanov¹¹, A. Dimitrov¹², L. Litov¹², B. Pavlov¹², P. Laykova¹², A. Petrov¹², S. Keshri¹³, D. Laroze¹³, S. Thakur¹³, W. Brooks¹⁴, T. Cheng¹⁵, T. Javaid¹⁵, L. Wang¹⁵, L. Yuan¹⁵, Z. Hu¹⁶, Z. Liang¹⁶, J. Liu¹⁶, X. Wang¹⁶, G. M. Chen¹⁷, H. S. Chen¹⁷, M. Chen¹⁷, Y. Chen¹⁷, Q. Hou¹⁷, X. Hou¹⁷, F. Iemmi¹⁷, C. H. Jiang¹⁷, A. Kapoor¹⁷, H. Liao¹⁷, G. Liu¹⁷, Z.-A. Liu¹⁷, J. N. Song¹⁷, S. Song¹⁷, J. Tao¹⁷, C. Wang¹⁷, J. Wang¹⁷, H. Zhang¹⁷, J. Zhao¹⁷, A. Agapitos¹⁸, Y. Ban¹⁸, A. Carvalho Antunes De Oliveira¹⁸, S. Deng¹⁸, B. Guo¹⁸, Q. Guo¹⁸, C. Jiang¹⁸, A. Levin¹⁸, C. Li¹⁸, Q. Li¹⁸, Y. Mao¹⁸, S. Qian¹⁸, S. J. Qian¹⁸, X. Qin¹⁸, X. Sun¹⁸, D. Wang¹⁸, J. Wang¹⁸, H. Yang¹⁸, M. Zhang¹⁸, Y. Zhao¹⁸, S. Yang¹⁹, Z. You²⁰, K. Jaffel²¹, N. Lu²¹, G. Bauer²², Z. Cui²², B. Li²², H. Wang²², K. Yi^{22,23}, J. Zhang²², Y. Li²⁴, Z. Lin²⁵, C. Lu²⁵, M. Xiao²⁵, C. Avila²⁶, D. A. Barbosa Trujillo²⁶, A. Cabrera²⁶, C. Florez²⁶, J. Fraga²⁶, J. A. Reyes Vega²⁶, C. Rendón²⁷, M. Rodriguez²⁷, A. A. Ruales Barbosa²⁷, J. D. Ruiz Alvarez²⁷, N. Godinovic²⁸, D. Delas²⁸, A. Sculac²⁸, M. Kovac²⁹, A. Petkovic²⁹, T. Sculac²⁹, P. Bargaşa³⁰, V. Brigljevic³⁰, B. K. Chitro³⁰, D. Ferencak³⁰, K. Jakovcic³⁰, A. Starodumov³⁰, T. Susa³⁰, A. Attkis³¹, K. Christoforou³¹, A. Hadjiagapiou³¹, C. Leonidou³¹, C. Nicolaou³¹, L. Paizanos³¹, F. Ptochos³¹, P. A. Raziş³¹, H. Rykaczewski³¹, H. Saka³¹, A. Stepennov³¹, M. Finger³², M. Finger Jr.³², E. Ayala³³, E. Carrera Jarrin³⁴, Y. Assran³⁵, B. El-mahdy³⁵, M. Abdullah Al-Mashhad³⁶, A. Hussein³⁶, H. Mohammed³⁶, K. Ehtahtaj³⁷, M. Kadastik³⁷, T. Lange³⁷, C. Nielsen³⁷, J. Pata³⁷, M. Raidal³⁷, N. Seeb³⁷, L. Tani³⁷, A. Milicevic³⁸, K. Osterberg³⁸, M. Voutilainen³⁸, N. Bin Norjoharudeen³⁹, E. Brücken³⁹, F. Garcia³⁹, P. Inkaev³⁹, K. T. S. Kallonen³⁹, R. Kumar Verma³⁹, T. Lampén³⁹, K. Lassila-Perini³⁹, B. Lehtela³⁹, S. Lehti³⁹, T. Lindén³⁹, N. R. Mancilla Xinto³⁹, M. Myllymäki³⁹, M. M. Rantanen³⁹, S. Saarikari³⁹, N. T. Toikka³⁹, J. Tuominiemi³⁹, H. Kirschenmann⁴⁰, P. Luukka⁴⁰, H. Petrow⁴⁰, M. Beanic⁴¹, F. Couderc⁴¹, M. Dejardin⁴¹, D. Denegri⁴¹, P. Devouge⁴¹, J. L. Faure⁴¹, F. Ferri⁴¹, P. Gaigne⁴¹, S. Ganjour⁴¹, P. Gras⁴¹, G. Hamel de Monchenault⁴¹, M. Kumar⁴¹, V. Lohezic⁴¹, J. Malcles⁴¹, F. Orlandi⁴¹, L. Portales⁴¹, S. Ronchi⁴¹, M. Ö. Sahin⁴¹, A. Savoy-Navarro^{41,42}, P. Simkina⁴¹, M. Titov⁴¹, M. Törnqvist⁴¹, F. Beaudette⁴³, G. Boldrin⁴³, P. Bussón⁴³, C. Charlot⁴³, M. Chiusi⁴³, T. D. Cuisset⁴³, F. Damas⁴³, O. Davignon⁴³, A. De Wit⁴³, T. Debnath⁴³, I. T. Ehle⁴³, B. A. Fontana Santos Alves⁴³, S. Ghosh⁴³, A. Gilbert⁴³, R. Granier de Cassagnac⁴³, L. Kalpitoli⁴³, M. Manoni⁴³, M. Nguyen⁴³, S. Obraztsov⁴³, C. Ochando⁴³, R. Salerno⁴³, J. B. Sauvan⁴³, Y. Sirois⁴³, G. Sokmer⁴³, L. Urda Gómez⁴³, A. Zabi⁴³, A. Zghiche⁴³, J.-L. Agram⁴⁴, J. Andrea⁴⁴, D. Bloch⁴⁴, J.-M. Brom⁴⁴, E. C. Chabert⁴⁴, C. Collard⁴⁴, G. Coulon⁴⁴, S. Falke⁴⁴, U. Goerlach⁴⁴, R. Haeblerle⁴⁴, A. C. Le Bihan⁴⁴, M. Meena⁴⁴, O. Poncet⁴⁴, G. Saha⁴⁴, P. Vaucluse⁴⁴, A. Di Florio⁴⁵, D. Amram⁴⁶, S. Beauceron⁴⁶, B. Blancon⁴⁶, G. Boudout⁴⁶, N. Chanon⁴⁶, D. Contardo⁴⁶, P. Depasse⁴⁶, C. Dozen⁴⁶, H. El Mamouni⁴⁶, J. Fay⁴⁶, S. Gascon⁴⁶, M. Gouzevitch⁴⁶, C. Greenberg⁴⁶, G. Grenier⁴⁶, B. Ille⁴⁶, E. Jourdhuy⁴⁶, I. B. Laktineh⁴⁶, M. Lethuillier⁴⁶, B. MassotEAU⁴⁶, L. Mirabito⁴⁶, A. Purohit⁴⁶, M. Vander Donckt⁴⁶, J. Xiao⁴⁶, I. Lomidze⁴⁷, T. Toriashvili⁴⁷, Z. Tsamalaidze⁴⁷, V. Botta⁴⁸, S. Consuegra Rodriguez⁴⁸, L. Feld⁴⁸, K. Klein⁴⁸, M. Lipinski⁴⁸, D. Meuse⁴⁸, P. Natland⁴⁸, V. Oppenländer⁴⁸, A. Pauls⁴⁸, D. Pérez Adán⁴⁸, N. Röwert⁴⁸, M. Teroerde⁴⁸, C. Daumann⁴⁹, S. Diekmann⁴⁹, A. Dodonova⁴⁹, N. Eich⁴⁹, D. Eliseev⁴⁹, F. Engelke⁴⁹, J. Erdmann⁴⁹, M. Erdmann⁴⁹, B. Fischer⁴⁹, T. Hebbeker⁴⁹, K. Hoepfner⁴⁹, F. Ivone⁴⁹, A. Jung⁴⁹, N. Kumar⁴⁹, M. Y. Lee⁴⁹, F. Mausolf⁴⁹, M. Merschmeyer⁴⁹, A. Meyer⁴⁹, F. Nowotny⁴⁹, A. Pozdnyakov⁴⁹, W. Redjeb⁴⁹, H. Reithler⁴⁹, U. Sarkar⁴⁹, V. Sarkisov⁴⁹, A. Schmidt⁴⁹, C. Seth⁴⁹, A. Sharma⁴⁹, J. L. Spahr⁴⁹, V. Vaulin⁴⁹, S. Zaleski⁴⁹, M. R. Bechers⁵⁰, C. Dziwok⁵⁰, G. Flügge⁵⁰, N. Hoeflich⁵⁰, T. Kress⁵⁰, A. Nowack⁵⁰, O. Pooth⁵⁰, A. Stahl⁵⁰, A. Zolt⁵⁰, H. Aarup Petersen⁵¹, A. Abel⁵¹, M. Aldaya Martin⁵¹, J. Alimena⁵¹, S. Amoroso⁵¹, Y. An⁵¹, I. Andreev⁵¹, J. Bach⁵¹, S. Baxter⁵¹, M. Bayatmakou⁵¹, H. Becerril Gonzalez⁵¹, O. Behnke⁵¹, A. Belvedere⁵¹, F. Blekman^{51,52}, K. Borras^{49,51}, A. Campbell⁵¹, S. Chatterjee⁵¹, L. X. Coll Saravia⁵¹, G. Eckerlin⁵¹, D. Eckstein⁵¹, E. Gallo^{51,52}, A. Geiser⁵¹, V. Guglielmi⁵¹, M. Guffioni⁵¹, A. Hinzmann⁵¹, L. Jepe⁵¹, M. Kasemann⁵¹, C. Reikhsnors⁵¹, R. Kogler⁵¹, M. Kott⁵¹, D. Krücker⁵¹, W. Lange⁵¹, D. Leyva Pernia⁵¹, K.-Y. Lin⁵¹, K. Lipka⁵¹, W. Lohmann⁵¹, J. Malvaso⁵¹, R. Mankel⁵¹, I.-A. Melzer-Pellmann⁵¹, M. Mendizabal Morentin⁵¹, A. B. Meyer⁵¹, G. Miella⁵¹, K. Moral Figueroa⁵¹, A. Mussgiller⁵¹, L. P. Nair⁵¹, J. Niedziela⁵¹, A. Nürnberg⁵¹, J. Park⁵¹, S. E. Ranken⁵¹, A. Raspereza⁵¹, D. Rastorguev⁵¹, L. Rygaard⁵¹, M. Scham^{49,51}, S. Schnake^{49,51}, P. Schütze⁵¹, C. Schwanenberger^{51,52}, D. Selivanova⁵¹, K. Sharko⁵¹, M. Shchedrovskiy⁵¹, D. Stafford⁵¹, M. Torkian⁵¹, F. Vazzoler⁵¹, A. Ventura⁵¹, R. Walsh⁵¹, D. Wang⁵¹, Q. Wang⁵¹, K. Wichmann⁵¹, L. Wiens^{49,51}, C. Wissing⁵¹, Y. Yang⁵¹, S. Zakharov⁵¹, A. Zimmermann Castro Santos⁵¹, A. Albrecht⁵², A. R. Alves Andrade⁵², M. Antonello⁵², S. Bollweg⁵², M. Bonanomi⁵², K. El Morabit⁵², Y. Fischer⁵², M. Frahm⁵², E. Garutti⁵², A. Grohsjean⁵², A. A. Guvenli⁵², J. Haller⁵², D. Hundhausen⁵², G. Kasieczka⁵², P. Keicher⁵², R. Klanner⁵², W. Korcar⁵², T. Kramer⁵², C. C. Kuo⁵², F. Labe⁵², J. Lange⁵², A. Lobanov⁵², L. Moureaux⁵², M. Mrowietz⁵², A. Nigamova⁵², K. Nikolopoulos⁵², Y. Nissan⁵², A. Paasch⁵², K. J. Pena Rodriguez⁵², N. Prouvost⁵²,

T. Quadfasel⁵², B. Raciti⁵², M. Rieger⁵², D. Savoiti⁵², P. Schlexer⁵², M. Schröder⁵², J. Schwandt⁵², M. Sommerhalder⁵², H. Stadié⁵², G. Steinbrück⁵², A. Tews⁵², R. Ward⁵², B. Wiederspan⁵², M. Wolf⁵², S. Brommer⁵³, E. Butz⁵³, Y. M. Chen⁵³, T. Chwalek⁵³, A. Dierlamm⁵³, G. G. Dincer⁵³, U. Elicabul⁵³, N. Faltermann⁵³, M. Giffels⁵³, A. Gottmann⁵³, F. Hartmann^{53,54}, R. Hofsaess⁵³, M. Horzela⁵³, U. Husemann⁵³, J. Kieseler⁵³, M. Klute⁵³, R. Kunlani Muhammed Rafeek⁵³, O. Lavryk⁵³, J. J. M. Lawhorn⁵³, A. Lintuluoto⁵³, S. Maier⁵³, M. Mommile⁵³, Th. Müller⁵³, E. Pfeffer⁵³, M. Presilla⁵³, G. Quast⁵³, K. Rabbertz⁵³, B. Regner^{53,54}, R. Schmieder⁵³, N. Shadskiy⁵³, I. Shvetsov⁵³, H. J. Simonis⁵³, L. Sowa⁵³, L. Stockmeier⁵³, K. Taucje⁵³, M. Toms⁵³, B. Topko⁵³, N. Trevisan⁵³, C. Verstege⁵³, T. Voigtländer⁵³, R. F. Von Cube⁵³, J. Von Den Driesch⁵³, M. Wassmer⁵³, R. Wolf⁵³, W. D. Zeuner⁵³, X. Zuo⁵³, G. Agnastou⁵⁴, G. Daskalakis⁵⁴, A. Kyriakis⁵⁴, G. Melachroinos⁵⁴, Z. Painesis⁵⁴, I. Paraskevas⁵⁴, N. Saoulidou⁵⁴, K. Theofilatos⁵⁴, E. Tziaferi⁵⁴, K. Vellidis⁵⁴, I. Zisopoulos⁵⁴, T. Chatzistavrou⁵⁷, G. Karapostoli⁵⁷, K. Kousouris⁵⁷, E. Siamarkou⁵⁷, G. Tsipolitis⁵⁷, I. Bestintanos⁵⁸, I. Evangelou⁵⁸, C. Foudas⁵⁸, P. Katsoulis⁵⁸, P. Kokkas⁵⁸, P. G. Kosmoglou Kioseoglou⁵⁹, N. Manthos⁵⁹, I. Papadopoulos⁵⁹, J. Strogas⁵⁹, D. Druzhkin⁵⁹, C. Hajdu⁵⁹, D. Horvath^{59,60}, K. Márton⁵⁹, A. J. Rádl^{59,61}, F. Sikler⁵⁹, V. Veszpremi⁵⁹, M. Csanád⁶¹, K. Farkas⁶¹, A. Fehérkúti^{59,61}, M. M. A. Gadallah⁶¹, A. Kadlecik⁶¹, M. León Coelho⁶¹, G. Pásztor⁶¹, G. I. Veres⁶¹, B. Ujvari⁶¹, G. Ziliz⁶², G. Benzecou⁶², S. Czellar⁶², J. Molnar⁶⁰, Z. Szilasi⁶⁰, T. Csorgo^{59,63}, F. Nemes^{59,63}, T. Novak⁶³, I. Szanyi^{63,64}, S. Bansa⁶⁵, S. B. Beri⁶⁵, V. Bhatnagar⁶⁵, G. Chaudhary⁶⁵, S. Chauhan⁶⁵, N. Dhingra⁶⁵, A. Kaur⁶⁵, A. Kaur⁶⁵, H. Kaur⁶⁵, M. Kaur⁶⁵, S. Kumar⁶⁵, T. Shekand⁶⁵, J. B. Singh⁶⁵, A. Singla⁶⁵, A. Bhardwaj⁶⁵, A. Chhetri⁶⁶, B. C. Choudhary⁶⁶, A. Kumar⁶⁶, A. Kumar⁶⁶, M. Naimuddin⁶⁶, S. Pho⁶⁶, K. Ranjan⁶⁶, M. K. Saini⁶⁶, S. Acharya⁶⁷, B. Gomber⁶⁷, B. Sahu⁶⁷, S. Mukherjee⁶⁸, S. Baradia⁶⁹, S. Bhattacharya⁶⁹, S. Das Gupta⁶⁹, S. Dutta⁶⁹, S. Dutta⁶⁹, S. Sarkar⁶⁹, M. M. Ameen⁷⁰, P. K. Behera⁷⁰, S. Chatterjee⁷⁰, G. Dash⁷⁰, A. Dattamuni⁷⁰, P. Jana⁷⁰, P. Kalbhor⁷⁰, S. Kamble⁷⁰, J. R. Komaragiri⁷⁰, T. Mishra⁷⁰, P. R. Pujahari⁷⁰, A. K. Sikdar⁷⁰, R. K. Singh⁷⁰, P. Verma⁷⁰, S. Verma⁷⁰, A. Vijay⁷⁰, B. K. Sivasva⁷⁰, L. Bhatt⁷², S. Dugad⁷², G. B. Mohanty⁷², M. Shelake⁷², P. Suryadevara⁷², A. Bala⁷³, S. Banerjee⁷³, S. Barman⁷³, R. M. Chatterjee⁷³, M. Guchait⁷³, Sh. Jain⁷³, A. Jaiswal⁷³, B. M. Joshi⁷³, S. Kumar⁷³, M. Maity⁷³, G. Majumder⁷³, K. Mazumdar⁷³, S. Parolia⁷³, R. Saxena⁷³, A. Thachayath⁷³, S. Bahinipati⁷⁴, D. Maity⁷⁴, P. Mal⁷⁴, K. Naskar⁷⁴, A. Nayak⁷⁴, S. Nayak⁷⁴, K. Pal⁷⁴, R. Raturi⁷⁴, P. Sadangi⁷⁴, S. K. Swain⁷⁴, S. Varghese⁷⁴, D. Vats⁷⁴, A. Alpna⁷⁵, S. Dube⁷⁵, P. Hazarika⁷⁵, B. Kansal⁷⁵, A. Laha⁷⁵, R. Sharma⁷⁵, S. Sharma⁷⁵, K. Y. Vaish⁷⁵, S. Ghosh⁷⁶, H. Bakhshiansohi⁷⁶, A. Jafari⁷⁷, V. Sedighzadeh Dalavi⁷⁷, M. Zeinali⁷⁷, S. Bashiri⁷⁸, S. Chenarani⁷⁸, S. M. Etesami⁷⁸, Y. Hosseini⁷⁸, M. Khakzad⁷⁸, E. Khaizia⁷⁸, M. Mohammadi Najafabadi⁷⁸, S. Tizchang⁷⁸, M. Felcini⁷⁹, M. Grunewald⁷⁹, M. Abbrescia⁸⁰, M. Barbieri^{80,81}, M. Buonsante^{80,81}, A. Colale^{80,81}, D. Creanza^{80,82}, B. D'Anzi^{80,81}, N. De Filippis^{80,82}, M. De Palma^{80,81}, W. Elmetenawee^{80,81}, N. Ferrara^{80,82}, L. Fiore⁸⁰, L. Longo⁸⁰, M. Louka^{80,81}, G. Maggi^{80,82}, M. Maggi⁸⁰, I. Margjeka⁸⁰, V. Mastrapasqua^{80,81}, S. My^{80,81}, F. Nenna^{80,81}, S. Nuzzo^{80,81}, A. Pellecchia^{80,81}, A. Pompili^{80,81}, G. Pugliese^{80,82}, R. Radogna^{80,81}, D. Ramos⁸⁰, A. Ranieri⁸⁰, L. Silvestris⁸⁰, F. M. Simone^{80,82}, Ü. Stüben⁸⁰, A. Stamerra^{80,81}, D. Troiano^{80,81}, R. Venditti^{80,81}, P. Verwilligen⁸⁰, A. Zaza^{80,81}, G. Abbiendi⁸³, C. Battilana^{83,84}, D. Bonacors^{83,84}, P. Capiluppi^{83,84}, F. R. Cavallo⁸³, G. M. Dallavalle⁸³, T. Diotallevi^{83,84}, F. Fabbri⁸³, A. Fanfani^{83,84}, D. Fasanella⁸³, P. Giacomelli⁸³, C. Grandi⁸³, L. Guiducci^{83,84}, S. Lo Meo⁸³, M. Lorusso^{83,84}, L. Lunerti⁸³, S. Marcellini⁸³, G. Masetti⁸³, F. L. Navarria^{83,84}, G. Paggi^{83,84}, A. Perrotta⁸³, F. Primavera^{83,84}, A. M. Rossi^{83,84}, S. Rossi Tisber^{83,84}, T. Rovelli^{83,84}, G. P. Siroli^{83,84}, S. Costa^{85,86}, A. Di Mattia⁸⁵, A. Lapertosa⁸⁵, R. Potenza^{85,86}, A. Tricom^{85,86}, J. Altkor^{87,88}, P. Assiouras⁸⁷, G. Baglioni⁸⁷, G. Bardelli⁸⁷, M. Bartolini^{87,88}, A. Calandri^{87,88}, B. Camaiani^{87,88}, A. Cassese⁸⁷, R. Ceccarelli⁸⁷, V. Ciulli^{87,88}, C. Cividini⁸⁷, R. D'Alessandro^{87,88}, L. Damenti^{87,88}, E. Focardi^{87,88}, T. Kello⁸⁷, G. Latino^{87,88}, P. Lenzi^{87,88}, M. Lizzo⁸⁷, M. Meschini⁸⁷, S. Paoletti⁸⁷, A. Papanastassiou^{87,88}, G. Sguazzoni⁸⁷, L. Viliani⁸⁷, L. Benussi⁸⁹, S. Colafranceschi⁸⁹, S. Meola⁸⁹, D. Piccolo⁸⁹, M. Alves Gallo Pereira⁹⁰, F. Ferraro⁹⁰, E. Robutti⁹⁰, S. Tosi^{90,91}, A. Benaglia⁹², F. Brivio⁹², V. Camagni^{92,93}, F. Ceterelli^{92,93}, F. De Guio^{92,93}, M. E. Dinardo^{92,93}, P. Dini⁹², S. Genna⁹², R. Gerosa^{92,93}, A. Ghezzi^{92,93}, P. Govoni^{92,93}, L. Guzzi⁹², M. R. Kim⁹², G. Lavizzari^{92,93}, M. L. Lucchini^{92,93}, M. Malberti⁹², S. Malvezzi⁹², A. Massironi⁹², D. Menasce⁹², L. Moroni⁹², M. Paganoni^{92,93}, S. Palluotto^{92,93}, D. Pedrini⁹², A. Perego^{92,93}, B. S. Pinolini⁹², G. Pizzati^{92,93}, S. Ragazzi^{92,93}, T. Tabarelli di Fatis^{92,93}, S. Buontempo⁹⁴, C. Di Fraia⁹⁴, F. Fabozz^{94,96}, L. Favilla^{94,97}, A. O. M. Iorio^{94,95}, L. Lista^{94,95}, P. Paolucci^{94,96}, B. Rossi⁹⁴, P. Azzi⁹⁸, N. Bacchetta^{98,99}, D. Bisello^{98,100}, P. Bortignon⁹⁸, G. Bortolato⁹⁸, A. C. M. Bulla⁹⁸, P. Checchia⁹⁸, T. Dorigo⁹⁸, F. Gasparini^{98,100}, U. Gasparini^{98,100}, S. Giorgetti⁹⁸, E. Lusian⁹⁸, M. Margoni^{98,100}, A. T. Meneguzzo^{98,100}, F. Montecassiano⁹⁸, J. Pazzini^{98,100}, P. Ronchese^{98,100}, R. Rossin^{98,100}, F. Simonetti^{98,100}, M. Tosi^{98,100}, A. Triossi^{98,100}, S. Ventura⁹⁸, M. Zanetti^{98,100}, P. Zotto^{98,100}, A. Zucchetta^{98,100}, A. Braghieri¹⁰¹, S. Calzaferr¹⁰¹, P. Montagna^{101,102}, M. Pelliccioni¹⁰¹, V. Re¹⁰¹, C. Riccardi^{101,102}, P. Salvini¹⁰¹, I. Vai^{101,102}, P. Vitulo^{101,102}, S. Ajmal^{103,104}, M. E. Ascioti^{103,104}, G. M. Bilei¹⁰³, C. Carrivale^{103,104}, D. Ciangottelli^{103,104}, L. Della Penna^{103,104}, L. Fano^{103,104}, V. Mariani^{103,104}, M. Menichelli¹⁰³, F. Moscattelli¹⁰³, A. Rossi^{103,104}, A. Santocchia^{103,104}, D. Spiga¹⁰³, T. Tedeschi^{103,104}, C. Aime^{105,106}, C. A. Alexe^{105,107}, P. Aasen^{105,106}, P. Azzurri¹⁰⁵, G. Bagliesi¹⁰⁵, R. Bhattacharya¹⁰⁵, L. Bianchini^{105,106}, T. Boccali¹⁰⁵, E. Bossini¹⁰⁵, D. Bruschini^{105,107}, L. Calligaris^{105,106}, R. Castaldi¹⁰⁵, F. Cattafesta^{105,107}, M. A. Ciocci^{105,108}, M. Cipriani^{105,106}, R

F. Cossutti¹¹⁴, K. De Leo¹¹⁴, G. Della Ricca^{114,115}, R. Delli Gatti^{114,115}, S. Dogra¹¹⁶, J. Hong¹¹⁶, J. Kim¹¹⁶, T. Kim¹¹⁶, D. Lee¹¹⁶, H. Lee¹¹⁶, J. Lee¹¹⁶, S. W. Lee¹¹⁶, C. S. Moon¹¹⁶, Y. D. Oh¹¹⁶, S. Sekmen¹¹⁶, B. Tae¹¹⁶, Y. C. Yang¹¹⁶, M. S. Kim¹¹⁷, P. Gwak¹¹⁸, H. Kim¹¹⁸, D. H. Moon¹¹⁸, J. Seo¹¹⁸, E. Asilar¹¹⁹, F. Carnevali¹¹⁹, J. Choi^{119,120}, T. J. Kim¹¹⁹, Y. Ryu¹¹⁹, S. Ha¹²⁰, S. Han¹²⁰, B. Hong¹²⁰, K. Lee¹²⁰, K. S. Lee¹²⁰, J. Yoo¹²⁰, J. Goh¹²¹, J. Shin¹²¹, S. Yang¹²¹, Y. Kang¹²², H. S. Kim¹²², Y. Kim¹²², S. Lee¹²², J. Almond¹²³, J. H. Bhyun¹²³, J. Choi¹²³, J. Choi¹²³, W. Jun¹²³, H. Kim¹²³, J. Kim¹²³, T. Kim¹²³, Y. Kim¹²³, Y. W. Kim¹²³, S. Ko¹²³, H. Lee¹²³, J. Lee¹²³, J. Lee¹²³, B. H. Oh¹²³, S. B. Oh¹²³, J. Shin¹²³, U. K. Yang¹²³, I. Yoon¹²³, W. Wang¹²³, D. Y. Kang¹²⁴, D. Kim¹²⁴, S. Kim¹²⁴, B. Ko¹²⁴, J. S. H. Lee¹²⁴, Y. Lee¹²⁴, I. C. Park¹²⁴, Y. Roh¹²⁴, I. J. Watson¹²⁴, G. Cho¹²⁵, K. Hwang¹²⁵, B. Kim¹²⁵, S. Kim¹²⁵, K. Lee¹²⁵, H. D. Yoo¹²⁵, M. Choi¹²⁶, Y. Lee¹²⁶, I. Yu¹²⁶, T. Beyrouthy¹²⁷, Y. Gharbia¹²⁷, F. Alazemi¹²⁷, K. Dreimanis¹²⁷, O. M. Eberlins¹²⁷, A. Gaile¹²⁹, C. Munoz Diaz¹²⁹, D. Osite¹²⁹, G. Pikurs¹²⁹, R. Plese¹²⁹, A. Potrebko¹²⁹, M. Seidel¹²⁹, D. Sidiropoulos Kontos¹²⁹, N. R. Strautnieks¹³⁰, M. Ambrozias¹³¹, A. Juodagalvis¹³¹, S. Nargelas¹³¹, A. Rinkevicius¹³¹, G. Tamulaitis¹³¹, I. Yusuf¹³², Z. Zolkapi¹³², J. F. Benitez¹³³, A. Castaneda Hernandez¹³³, A. Cota Rodriguez¹³³, L. E. Cuevas Picos¹³³, H. A. Encinas Acosta¹³³, L. G. Gallegos Maríñez¹³³, J. A. Murillo Quijada¹³³, A. Sehrawat¹³³, L. Valencia Palomo¹³³, G. Ayala¹³⁴, H. Castilla-Valdez¹³⁴, H. Crotte Ledesma¹³⁴, R. Lopez-Fernandez¹³⁴, J. Mejia Guisao¹³⁴, R. Reyes-Almanza¹³⁴, A. Sánchez Hernández¹³⁴, C. Oropeza Barrera¹³⁵, D. L. Ramirez Guadarrama¹³⁵, M. Ramirez García¹³⁵, I. Bautista¹³⁶, F. E. Neri Huerta¹³⁶, I. Pedraza¹³⁶, H. A. Salazar Ibarguen¹³⁶, C. Uribe Estrada¹³⁶, I. Bujanja¹³⁷, N. Raicevic¹³⁷, P. H. Buttle¹³⁸, A. Ahmad¹³⁹, M. I. Asghar¹³⁹, A. Awaiz¹³⁹, M. I. M. Awan¹³⁹, W. A. Khan¹³⁹, V. Avati¹⁴⁰, L. Forthomme¹⁴⁰, L. Grzanka¹⁴⁰, M. Malawski¹⁴⁰, K. Piotrkowski¹⁴⁰, M. Bluj¹⁴¹, M. Górski¹⁴¹, M. Kazana¹⁴¹, M. Szeleper¹⁴¹, P. Zalewski¹⁴¹, K. Bunkowski¹⁴², K. Doroba¹⁴², A. Kalinowski¹⁴², M. Konecki¹⁴², J. Krolkowski¹⁴², A. Muhammad¹⁴², P. Fokow¹⁴³, K. Pozniak¹⁴³, W. Zabolotny¹⁴³, M. Araujo¹⁴⁴, D. Bastos¹⁴⁴, C. Beirão Da Cruz E Silva¹⁴⁴, A. Boletti¹⁴⁴, M. Bozzo¹⁴⁴, T. Camporesi¹⁴⁴, G. Da Molin¹⁴⁴, M. Gallinaro¹⁴⁴, J. Hollar¹⁴⁴, N. Leonardo¹⁴⁴, G. B. Marozzo¹⁴⁴, A. Petrill¹⁴⁴, M. Pisano¹⁴⁴, J. Seixas¹⁴⁴, J. Varela¹⁴⁴, J. W. Wulff¹⁴⁴, P. Adzic¹⁴⁵, L. Markovic¹⁴⁵, P. Milenovic¹⁴⁵, V. Milosevic¹⁴⁵, D. Devetak¹⁴⁶, M. Dordevic¹⁴⁶, J. Milosevic¹⁴⁶, L. Nadder¹⁴⁶, V. Rekovic¹⁴⁶, M. Stojanovic¹⁴⁶, M. Alcalde Martinez¹⁴⁷, F. Alcaraz Maestre¹⁴⁷, Cristina F. Bedoya¹⁴⁷, J. A. Brochero Cifuentes¹⁴⁷, Oliver M. Carretero¹⁴⁷, M. Cepeda¹⁴⁷, M. Cerrada¹⁴⁷, N. Colino¹⁴⁷, J. Cuchillo Ortega¹⁴⁷, B. De La Cruz¹⁴⁷, A. Delgado Peris¹⁴⁷, A. Escalante Del Valle¹⁴⁷, D. Fernández Del Val¹⁴⁷, J. P. Fernández Ramos¹⁴⁷, J. Flix¹⁴⁷, M. C. Fouz¹⁴⁷, M. Gonzalez Hernandez¹⁴⁷, O. Gonzalez Lopez¹⁴⁷, S. Goy Lopez¹⁴⁷, J. M. Hernandez¹⁴⁷, M. I. Josa¹⁴⁷, J. Llorente Merino¹⁴⁷, C. Martin Perez¹⁴⁷, E. Martin Viscasillas¹⁴⁷, D. Moran¹⁴⁷, M. M. Morcillo Perez¹⁴⁷, R. Paz Herrera¹⁴⁷, C. Perez Dengra¹⁴⁷, A. Pérez-Callero Yzquierdo¹⁴⁷, J. Puerta Pelayo¹⁴⁷, I. Redondo¹⁴⁷, J. Vazquez Escobar¹⁴⁷, J. F. de Trocóniz¹⁴⁸, B. Alvarez Gonzalez¹⁴⁹, J. Ayllon Torresano¹⁴⁹, A. Cardini¹⁴⁹, J. Cuevas¹⁴⁹, J. Del Riogado Badas¹⁴⁹, D. Estrada Acevedo¹⁴⁹, J. Fernandez Menendez¹⁴⁹, S. Folgueras¹⁴⁹, I. Gonzalez Caballero¹⁴⁹, P. Leguina¹⁴⁹, M. Obeso Menendez¹⁴⁹, E. Palencia Cortezon¹⁴⁹, J. Prado Pico¹⁴⁹, A. Soto Rodriguez¹⁴⁹, C. Vico Villalba¹⁴⁹, P. Vischia¹⁴⁹, S. Blanco Fernández¹⁵⁰, I. J. Cabrillo¹⁵⁰, A. Calderon¹⁵⁰, J. Duarte Campderros¹⁵⁰, M. Fernandez¹⁵⁰, G. Gomez¹⁵⁰, C. Lasasas Garcia¹⁵⁰, R. Lopez Ruiz¹⁵⁰, C. Martinez Rivero¹⁵⁰, P. Martinez Ruiz del Arbol¹⁵⁰, F. Matorras¹⁵⁰, P. Matorras Cuevas¹⁵⁰, E. Navarrete Ramos¹⁵⁰, J. Piedra Gomez¹⁵⁰, C. Quintana San Emeterio¹⁵⁰, L. Scodellaro¹⁵⁰, I. Vila¹⁵⁰, R. Vilar Cortabitarte¹⁵⁰, J. M. Vizan Garcia¹⁵⁰, B. Kailasapathy¹⁵¹, D. D. C. Wickramaratne¹⁵¹, W. G. D. Dharmaratna¹⁵², K. Liyanage¹⁵², N. Perera¹⁵², D. Abbaneo¹⁵⁴, C. Amendola¹⁵⁴, R. Ardino¹⁵⁴, E. Auffray¹⁵⁴, J. Baechler¹⁵⁴, D. Banerjee¹⁵⁴, M. Bianco¹⁵⁴, A. Boccia¹⁵⁴, L. Borghonovi¹⁵⁴, C. Botta¹⁵⁴, A. Bragagnolo¹⁵⁴, C. E. Brown¹⁵⁴, C. Caillio¹⁵⁴, G. Cerminara¹⁵⁴, P. Conno¹⁵⁴, D. d'Enterria¹⁵⁴, A. Dabrowski¹⁵⁴, A. David¹⁵⁴, A. De Roeck¹⁵⁴, M. M. Defranichis¹⁵⁴, M. Deile¹⁵⁴, M. Dobson¹⁵⁴, W. Funk¹⁵⁴, A. Gaddi¹⁵⁴, S. Giani¹⁵⁴, D. Gigli¹⁵⁴, K. Gill¹⁵⁴, F. Glege¹⁵⁴, M. Glowacki¹⁵⁴, A. Gruber¹⁵⁴, J. Hegeman¹⁵⁴, J. K. Heikkilä¹⁵⁴, B. Huber¹⁵⁴, V. Innocente¹⁵⁴, T. James¹⁵⁴, P. Janot¹⁵⁴, O. Kaluzinska¹⁵⁴, O. Karacheban¹⁵⁴, G. Karathanasis¹⁵⁴, S. Laurila¹⁵⁴, P. Leo¹⁵⁴, C. Lourenço¹⁵⁴, A.-M. Lyon¹⁵⁴, M. Magherini¹⁵⁴, L. Malgeri¹⁵⁴, M. Mannelli¹⁵⁴, A. Mehta¹⁵⁴, F. Meijers¹⁵⁴, J. A. Merlin¹⁵⁴, S. Mersi¹⁵⁴, E. Meschi¹⁵⁴, M. Migliorini¹⁵⁴, F. Monti¹⁵⁴, F. Moortgat¹⁵⁴, M. Mulders¹⁵⁴, M. Musich¹⁵⁴, I. Neutelings¹⁵⁴, S. Orfanelli¹⁵⁴, F. Pantaleo¹⁵⁴, M. Parisi¹⁵⁴, G. Petrucciani¹⁵⁴, A. Pfeiffer¹⁵⁴, M. Pierini¹⁵⁴, M. Pitt¹⁵⁴, H. Qu¹⁵⁴, D. Rabady¹⁵⁴, B. Ribeiro Lopes¹⁵⁴, F. Rit¹⁵⁴, P. Rosado¹⁵⁴, M. Rovere¹⁵⁴, H. Sakulin¹⁵⁴, R. Salvatico¹⁵⁴, S. Sanchez Cruz¹⁵⁴, S. Scarfi¹⁵⁴, C. Schwick¹⁵⁴, M. Selvaggi¹⁵⁴, A. Sharma¹⁵⁴, K. Schelch¹⁵⁴, P. Silva¹⁵⁴, P. Sphicas^{154,156}, A. G. Stahl Leiton¹⁵⁴, A. Steen¹⁵⁴, S. Summers¹⁵⁴, D. Treille¹⁵⁴, S. T. Protopop¹⁵⁴, E. Vernazza¹⁵⁴, J. Wanczyk¹⁵⁴, J. Wang¹⁵⁴, S. Wuchter¹⁵⁴, M. Zarecki¹⁵⁴, P. Zehetner¹⁵⁴, P. Zetj¹⁵⁴, G. Zevi Della Porta¹⁵⁴, T. Bevilacqua^{153,154}, L. Caminada^{153,154}, W. Erdmann¹⁵³, R. Horisberger¹⁵³, Q. Ingram¹⁵³, H. C. Kaestli¹⁵³, D. Kotlinski¹⁵³, C. Lange¹⁵³, U. Langenegger¹⁵³, M. Missiroli^{153,154}, L. Noehte^{153,154}, T. Roho¹⁵³, A. Samalan¹⁵³, T. K. Aarrestad¹⁵⁵, M. Backhaus¹⁵⁵, G. Bonomelli¹⁵⁵, C. Cazzaniga¹⁵⁵, K. Datta¹⁵⁵, P. De Bryas Daxnieder¹⁵⁵, D. Archiacchiaco¹⁵⁵, A. De Cosa¹⁵⁵, G. Dissertori¹⁵⁵, M. Dittmar¹⁵⁵, M. Donega¹⁵⁵, F. Eble¹⁵⁵, K. Gedia¹⁵⁵, F. Glessgen¹⁵⁵, C. Grab¹⁵⁵, N. Häringer¹⁵⁵, T. G. Harte¹⁵⁵, W. Luster¹⁵⁵, M. Malucchi¹⁵⁵, R. A. Manzoni¹⁵⁵, M. Marchegiani¹⁵⁵, L. Marchese¹⁵⁵, A. Mascellani¹⁵⁵, F. Nessi-Tedaldi¹⁵⁵, F. Pauss¹⁵⁵, V. Perovic¹⁵⁵, B. Ristic¹⁵⁵, R. Seidita¹⁵⁵, J. Steggen¹⁵⁵, A. Tarabini¹⁵⁵, D. Valsecchi¹⁵⁵, R. Wallny¹⁵⁵, C. AMSler¹⁵⁴, P. Bärttschi¹⁵⁴, F. Bilandzija¹⁵⁴, M. F. Canelli¹⁵⁴, G. Celotto¹⁵⁴, K. Cormier¹⁵⁴, M. Huwiler¹⁵⁴, W. Jin¹⁵⁴, A. Jofrehe¹⁵⁴, B. Kilminster¹⁵⁴, T. H. Kwok¹⁵⁴, S. Leontsinis¹⁵⁴, V. Lukashenko¹⁵⁴, A. Macchiolo¹⁵⁴, F. Meng¹⁵⁴, J. Motta¹⁵⁴, H. A. Reimers¹⁵⁴, P. Robmann¹⁵⁴, M. Senger¹⁵⁴, E. Shokri¹⁵⁴, F. Stäger¹⁵⁴, R. Tramontano¹⁵⁴, D. Bhowmik¹⁵⁶, C. M. Kuo¹⁵⁶, P. K. Rout¹⁵⁶, S. Taj¹⁵⁶, P. C. Tiwari¹⁵⁶, L. Card¹⁵⁷, K. F. Chen¹⁵⁷, Z. G. Chen¹⁵⁷, A. De Iorio¹⁵⁷, W.-S. Hou¹⁵⁷, T. H. Hsu¹⁵⁷, Y. W. Kao¹⁵⁷, S. Karmak¹⁵⁷, G. Kole¹⁵⁷, Y. Y. Li¹⁵⁷, R.-S. Lu¹⁵⁷, E. Paganis¹⁵⁷, X. F. Su¹⁵⁷, J. Thomas-Wilsker¹⁵⁷, L. S. Tsai¹⁵⁷, D. Tsionou¹⁵⁷, H. Y. Wu¹⁵⁷, E. Yazgan¹⁵⁷, C. Asawatangtrakuldee¹⁵⁸, N. Srimanobhan¹⁵⁸, Y. Maghrbi¹⁵⁹, D. Aggelou¹⁶⁰, F. Dolek¹⁶⁰, I. Dumanoglu¹⁶⁰, Y. Guler¹⁶⁰, E. Gurpinar Guler¹⁶⁰, C. Isik¹⁶⁰, O. Kara¹⁶⁰, A. Kayis Topaksu¹⁶⁰, Y. Komurcu¹⁶⁰, G. Onengut¹⁶⁰, K. Ozdemir¹⁶⁰, B. Tali¹⁶⁰, U. G. Tok¹⁶⁰, E. Uslan¹⁶⁰, I. S. Zorbakir¹⁶⁰, M. Yalvac¹⁶¹, B. Akgun¹⁶², I. O. Atakisi¹⁶², E. Gülmez¹⁶², M. Kaya¹⁶², O. Kaya¹⁶², M. A. Sarkisla¹⁶², S. Tekten¹⁶², A. Cakir¹⁶³, K. Cankocak¹⁶³, S. Sen¹⁶³, O. Aydilek¹⁶⁴, B. Hacishingol¹⁶⁴, I. Hos¹⁶⁴, B. Kaynak¹⁶⁴, S. Ozkocucuklu¹⁶⁴, O. Potok¹⁶⁴, H. Sert¹⁶⁴, C. Simsek¹⁶⁴, C. Zorbilmez¹⁶⁴, S. Cerci¹⁶⁵, B. Isildak¹⁶⁵, E. Simsek¹⁶⁵, D. Sunar Cerci¹⁶⁵, T. Yetkin¹⁶⁵, A. Boyaryntsev¹⁶⁶, O. Dadzhanova¹⁶⁶, B. Grynov¹⁶⁶, L. Levchuk¹⁶⁷, J. J. Brooke¹⁶⁸, A. Bundock¹⁶⁸, F. Bury¹⁶⁸, E. Clement¹⁶⁸, D. Cussans¹⁶⁸, D. Dharmender¹⁶⁸, H. Flacher¹⁶⁸, J. Goldstein¹⁶⁸, H. F. Heath¹⁶⁸, M.-L. Holmberg¹⁶⁸, L. Kreczko¹⁶⁸, S. Paramesvaran¹⁶⁸, L. Robertshaw¹⁶⁸, M. S. Sanjani^{168,169}, J. Segal¹⁶⁸, V. J. Smith¹⁶⁸, A. H. Ball¹⁶⁹, K. W. Bell¹⁶⁹, A. Belyaev¹⁶⁹, C. Brew¹⁶⁹, R. M. Brown¹⁶⁹, D. J. A. Cockerill¹⁶⁹, A. Elliot¹⁶⁹, K. V. Ellis¹⁶⁹, J. Gajownik¹⁶⁹, K. Harde¹⁶⁹, S. Harper¹⁶⁹, J. Linacre¹⁶⁹, K. Manolopoulos¹⁶⁹, M. Moallem¹⁶⁹, D. M. Newbold¹⁶⁹, E. Olaiya¹⁶⁹, D. Petyt¹⁶⁹, T. Reis¹⁶⁹, A. R. Sahasransu¹⁶⁹, G. Salvi¹⁶⁹, T. Schuh¹⁶⁹, C. H. Shepherd-Themistocleous¹⁶⁹, I. R. Tomalin¹⁶⁹, K. C. Whalen¹⁶⁹, T. Williams¹⁶⁹, I. Andreu¹⁷⁰, R. Bainbridge¹⁷⁰, P. Bloch¹⁷⁰, O. Buchmuller¹⁷⁰, C. A. Carrillo Montoya¹⁷⁰, D. Colling¹⁷⁰, J. S. Dancu¹⁷⁰, I. Das¹⁷⁰, P. Duane¹⁷⁰, G. Davies¹⁷⁰, M. Della Negra¹⁷⁰, S. Fayer¹⁷⁰, G. Fedji¹⁷⁰, G. Hall¹⁷⁰, H. R. Hoorani¹⁷⁰, A. Howard¹⁷⁰, G. Iles¹⁷⁰, C. R. Knight¹⁷⁰, P. Krueper¹⁷⁰, J. Langford¹⁷⁰, K. H. Law¹⁷⁰, J. León Holgado¹⁷⁰, E. Leutgeb¹⁷⁰, L. Lyons¹⁷⁰, A.-M. Magnan¹⁷⁰, B. Maier¹⁷⁰, S. Mallios¹⁷⁰, A. Mastrorilli¹⁷⁰, M. Mieskolainen¹⁷⁰, J. Nash¹⁷⁰, M. Pesaresi¹⁷⁰, P. B. Predeep¹⁷⁰, B. C. Radburn-Smith¹⁷⁰, A. Richards¹⁷⁰, A. Rose¹⁷⁰, L. Russell¹⁷⁰, K. Savva¹⁷⁰, C. Seez¹⁷⁰, R. Shukla¹⁷⁰, A. Tapper¹⁷⁰, K. Uchida¹⁷⁰, G. P. Uttley¹⁷⁰, T. Virdee¹⁷⁰, M. Vojinovic¹⁷⁰, N. Wardle¹⁷⁰, D. Winterbottom¹⁷⁰, J. E. Cole¹⁷¹, A. Khan¹⁷¹, P. Kyberd¹⁷¹, I. D. Reid¹⁷¹, S. Abdullin¹⁷², A. Brinkerhoff¹⁷², E. Collins¹⁷², M. R. Darwish¹⁷², J. Dittmann¹⁷², K. Hatayama¹⁷², V. Hegde¹⁷², J. Hiltbrand¹⁷², B. McMaster¹⁷², J. Samudio¹⁷², S. Sawant¹⁷², C. Satake¹⁷², J. Wilson¹⁷², J. M. Hogan¹⁷³, R. Bartek¹⁷³, A. Dominguez¹⁷³, S. Raj¹⁷³, A. E. Simsek¹⁷⁴, S. S. Yu¹⁷⁴, B. Bam¹⁷⁵, A. Buchot Perraguin¹⁷⁵, S. Campbell¹⁷⁵, R. Chudasama¹⁷⁵, S. I. Cooper¹⁷⁵, C. Crovella¹⁷⁵, G. Fidalgo¹⁷⁵, S. V. Gleyzer¹⁷⁵, A. Khukhushvili¹⁷⁵, K. Matchev¹⁷⁵, E. Pearson¹⁷⁵, C. U. Perez¹⁷⁵, P. Rumerio¹⁷⁵, E. Usa¹⁷⁵, R. Yi¹⁷⁵, S. Cholak¹⁷⁶, G. De Castro¹⁷⁶, Z. Demiragli¹⁷⁶, C. Erice¹⁷⁶, C. Fangmeier¹⁷⁶, C. Fernandez Madrazo¹⁷⁶, E. Fontanesi¹⁷⁶, J. Fulcher¹⁷⁶, F. Golf¹⁷⁶, S. Jeon¹⁷⁶, J. O'Cañ¹⁷⁶, I. Reed¹⁷⁶, J. Rohlf¹⁷⁶, K. Salyer¹⁷⁶, D. Sperka¹⁷⁶, D. Spitzbart¹⁷⁶, I. Suarez¹⁷⁶, A. Tsatsos¹⁷⁶, E. Wurtz¹⁷⁶, A. G. Zocchini¹⁷⁶, G. Barone¹⁷⁷, G. Benelli¹⁷⁷, D. Cutts¹⁷⁷, S. Ellis¹⁷⁷, L. Gouskos¹⁷⁷, M. Hadley¹⁷⁷, U. Heintz¹⁷⁷, K. W. Ho¹⁷⁷, T. Kwon¹⁷⁷, G. Landsberg¹⁷⁷, K. T. Lau¹⁷⁷, J. Luo¹⁷⁷, S. Mondal¹⁷⁷, J. Roloff¹⁷⁷, T. Russell¹⁷⁷, S. Sagir¹⁷⁷, X. Shen¹⁷⁷, M. Stamenkovic¹⁷⁷, N. Venkatasubramanian¹⁷⁷, S. Abbott¹⁷⁸, B. Barton¹⁷⁸, R. Breedon¹⁷⁸, H. Cai¹⁷⁸, M. Calderon De La Barca Sanchez¹⁷⁸, M. Chertok¹⁷⁸, M. Citron¹⁷⁸, J. Conway¹⁷⁸, P. T. Cox¹⁷⁸, R. Erbacher¹⁷⁸, O. Kukral¹⁷⁸, G. Moellin¹⁷⁸, S. Ostrom¹⁷⁸, I. Salazar Segovia¹⁷⁸, W. Wei¹⁷⁸, S. Yoo¹⁷⁸, K. Adamidis¹⁷⁹, M. Bachtis¹⁷⁹, D. Campos¹⁷⁹, R. Cousins¹⁷⁹, A. Datta¹⁷⁹, G. Flores Avila¹⁷⁹, J. Hause¹⁷⁹, M. Ignatenko¹⁷⁹, M. A. Iqbal¹⁷⁹, T. Lam¹⁷⁹, Y. F. Lo¹⁷⁹, E. Manca¹⁷⁹, A. Nunez Del Prado¹⁷⁹, D. Saltzberg¹⁷⁹, V. Valuev¹⁷⁹, R. Clare¹⁸⁰, J. W. Gary¹⁸⁰, G. Hanson¹⁸⁰, A. Aportela¹⁸⁰, A. Arora¹⁸⁰, J. G. Branson¹⁸⁰, S. Cittolin¹⁸⁰, S. Cooperstein¹⁸⁰, D. Diaz¹⁸⁰, J. Duarte¹⁸⁰, L. Giannini¹⁸⁰, Y. Gu¹⁸⁰, J. Guiang¹⁸⁰, V. Krutelyov¹⁸⁰, R. Lee¹⁸⁰, J. Letts¹⁸⁰, H. Li¹⁸⁰, M. Masciovecchio¹⁸⁰, F. Mokhtar¹⁸⁰, S. Mukherjee¹⁸⁰, M. Pieri¹⁸⁰, D. Primoschi¹⁸⁰, M. Quinlan¹⁸⁰, V. Sharma¹⁸⁰, M. Tadel¹⁸⁰, E. Vourliotis¹⁸⁰, F. Würthwein¹⁸⁰, A. Yagil¹⁸⁰, Z. Zhao¹⁸⁰, A. Barzdukas¹⁸², L. Brennan¹⁸², C. Campagnari¹⁸², S. Carron Montero¹⁸², K. Downham¹⁸², C. Grieco¹⁸², M. M. Hussain¹⁸², J. Incandela¹⁸², J. Kim¹⁸², M. W. K. Lai¹⁸², A. J. Li¹⁸², P. Masterson¹⁸², J. Richman¹⁸², S. N. Santapur¹⁸², U. Sarica¹⁸², R. Schmitz¹⁸², F. Setti¹⁸², J. Shephard¹⁸², D. Stuart¹⁸², T. Á. Vami¹⁸², X. Yan¹⁸², D. Zhang¹⁸², A. Albert¹⁸³, S. Bhattacharya¹⁸³, A. Bornheim¹⁸³, O. Cerri¹⁸³, R. Kansal¹⁸³, J. Mao¹⁸³, H. B. Newman¹⁸³, G. Reales Gutiérrez¹⁸³, T. Sievert¹⁸³, M. Spiropulu¹⁸³, J. R. Vlimant¹⁸³, R. A. Wynne¹⁸³, S. Xie¹⁸³, J. Alison¹⁸⁴, S. An¹⁸⁴, M. Cremonesi¹⁸⁴, V. Dutta¹⁸⁴, E. Y. Ertorer¹⁸⁴, T. Ferguson¹⁸⁴, T. A. Gíomez Espinosa¹⁸⁴, A. Harila¹⁸⁴, A. Kallii Tharaly¹⁸⁴, M. Kanemura¹⁸⁴, C. Liu¹⁸⁴, P. Meiring¹⁸⁴, T. Mudholkar¹⁸⁴, S. Murthy¹⁸⁴, P. Palit¹⁸⁴, K. Park¹⁸⁴, M. Paulini¹⁸⁴, A. Roberts¹⁸⁴, A. Sanchez¹⁸⁴, W. Terribil¹⁸⁴, J. P. Cumalat¹⁸⁵, W. T. Ford¹⁸⁵, A. Hart¹⁸⁵, A. Hassan¹⁸⁵, S. Kwan¹⁸⁵, J. Pearkes¹⁸⁵, C. Savard¹⁸⁵, N. Schonbeck¹⁸⁵, K. Stenson¹⁸⁵, K. A. Ulmer¹⁸⁵, S. R. Wagner¹⁸⁵, N. Zippner¹⁸⁵, D. Zuolo¹⁸⁵, J. Alexander¹⁸⁶, X. Chen¹⁸⁶, D. J. Cranshaw¹⁸⁶, J. Dickinson¹⁸⁶, J. Fan¹⁸⁶, X. Fan¹⁸⁶, J. Grassi¹⁸⁶, S. Hogan¹⁸⁶, P. Kotamnives¹⁸⁶, J. Monroy¹⁸⁶, G. Niendorf¹⁸⁶, M. Oshiro¹⁸⁶, J. R. Patterson¹⁸⁶, M. Reid¹⁸⁶, A. Ryd¹⁸⁶, J. Thom¹⁸⁶, P. Wittich¹⁸⁶, R. Zou¹⁸⁶, L. Zygala¹⁸⁶, M. Albrow¹⁸⁶, M. Alyari¹⁸⁶, O. Amram¹⁸⁶, G. Apollinari¹⁸⁶, A. Apresyan¹⁸⁶, L. A. T. Bauerick¹⁸⁶, D. Berry¹⁸⁶, J. Berryhill¹⁸⁶, P. C. Bhat¹⁸⁶, K. Burkett¹⁸⁶, J. N. Butler¹⁸⁶, A. Canepa¹⁸⁶, G. B. Cerat¹⁸⁶, H. W. K. Cheung¹⁸⁶, F. Chlebana¹⁸⁶, C. Cosby¹⁸⁶, G. Cummings¹⁸⁶, I. Dutta¹⁸⁶, V. D. Elvira¹⁸⁶, J. Freeman¹⁸⁶, A. Gandrakota¹⁸⁶, Z. Gecse¹⁸⁶, L. Gray¹⁸⁶, N. D. Green¹⁸⁶, A. Grummer¹⁸⁶, S. Grünendahl¹⁸⁶, D. Guerrero¹⁸⁶, O. Gutsche¹⁸⁶, R. M. Harris¹⁸⁶, T. C. Herwig¹⁸⁶, J. Hirschauer¹⁸⁶, B. Jayatilaka¹⁸⁶, S. Jindariani¹⁸⁶, M. Johnson¹⁸⁶, U. Joshi¹⁸⁶, T. Kljnsma¹⁸⁶, B. Klima¹⁸⁶, K. H. M. Kwok¹⁸⁶, S. Lammel¹⁸⁶, C. Lee¹⁸⁶, D. Lincoln¹⁸⁶, R. Lipton¹⁸⁶, T. Liu¹⁸⁶, K. Maeshima¹⁸⁶, D. Mason¹⁸⁶, P. M. B. P. Merkle¹⁸⁶, S. Mrenna¹⁸⁶, S. Nahn¹⁸⁶, J. Ngadiuba¹⁸⁶, D. Noonan¹⁸⁶, S. Norberg¹⁸⁶, V. Papadimitriou¹⁸⁶, N. Pastika¹⁸⁶, K. Pedro¹⁸⁶, C. Pena^{186,183}, C. E. Perez Lara¹⁸⁶, F. Raveria¹⁸⁶, A. Reinsvald Hall¹⁸⁶, L. Ristori¹⁸⁶, M. Safdari¹⁸⁶, E. Sexton-Kennedy¹⁸⁶, N. Smith¹⁸⁶, A. S. Saha¹⁸⁶, L. Spiegel¹⁸⁶, S. Stoyne¹⁸⁶, J. Strait¹⁸⁶, L. Taylor¹⁸⁶, S. Tkaczyk¹⁸⁶, N. V. Tran¹⁸⁶, L. Upplegger¹⁸⁶, E. W. Vaandering¹⁸⁶, C. Wang¹⁸⁶, I. Zoia¹⁸⁶, C. Aruta¹⁸⁶, P. Avery¹⁸⁶, I. D. Bourlikov¹⁸⁶, P. Chang¹⁸⁶, V. Cherepanov¹⁸⁶, R. D. Field¹⁸⁶, C. Huh¹⁸⁶, E. Koenig¹⁸⁶, M. Kolosova¹⁸⁶, J. Konigsberg¹⁸⁶, A. Korytov¹⁸⁶, N. Menendez¹⁸⁶, G. Mitselmakher¹⁸⁶, K. Mohrman¹⁸⁶, A. Muthirakalayi Madhu¹⁸⁶, N. Rawal¹⁸⁶, S. Rosenzweig¹⁸⁶, V. Sulimov¹⁸⁶, Y. Takahashi¹⁸⁶, J. Wang¹⁸⁶, T. Adams¹⁸⁷, A. Al Kadhim¹⁸⁸, A. Askew¹⁸⁸, S. Bower¹⁸⁸, R. Hashmi¹⁸⁸, R. S. Kim¹⁸⁸, T. Kolberg¹⁸⁸, G. Martinez¹⁸⁸, M. Mazza¹⁸⁸, H. Prosper¹⁸⁸, P. R. Prova¹⁸⁸, M. Wulansatiti¹⁸⁸, R. Yohay¹⁸⁸, B. Alysufyan¹⁸⁹, S. Butala

Article

W. J. Jackson¹⁹⁵, C. Kapsiak¹⁹⁵, M. Krohn¹⁹⁵, D. Mahon¹⁹⁵, J. Mans¹⁹⁵, B. Marzocchi¹⁹⁵, R. Rusack¹⁹⁵, O. Sancar¹⁹⁵, R. Saradhy¹⁹⁵, N. Strobbe¹⁹⁵, K. Bloom¹⁹⁶, D. R. Claes¹⁹⁶, G. Haza¹⁹⁶, J. Hossain¹⁹⁶, C. Joo¹⁹⁶, I. Kravchenko¹⁹⁶, A. Rohilla¹⁹⁶, J. E. Siado¹⁹⁶, W. Tabb¹⁹⁶, A. Vagnerini¹⁹⁶, A. Wightman¹⁹⁶, F. Yan¹⁹⁶, H. Bandyopadhyay¹⁹⁷, L. Hay¹⁹⁷, H. W. Hsia¹⁹⁷, I. Iashvili¹⁹⁷, A. Kaloogeropoulos¹⁹⁷, A. Kharchilava¹⁹⁷, A. Mandal¹⁹⁷, M. Morris¹⁹⁷, D. Nguyen¹⁹⁷, S. Rappocciolo¹⁹⁷, H. Rejeb Sfar¹⁹⁷, A. Williams¹⁹⁷, P. Young¹⁹⁷, D. Yu¹⁹⁷, G. Alverson¹⁹⁸, E. Barberis¹⁹⁸, J. Bonilla¹⁹⁸, B. Bylisma¹⁹⁸, M. Campana¹⁹⁸, J. Dervan¹⁹⁸, Y. Haddad¹⁹⁸, Y. Han¹⁹⁸, I. Israr¹⁹⁸, A. Krishna¹⁹⁸, M. Lu¹⁹⁸, N. Manganeli¹⁹⁸, R. McCarthy¹⁹⁸, D. M. Morse¹⁹⁸, T. Orimoto¹⁹⁸, A. Parker¹⁹⁸, L. Skinnari¹⁹⁸, C. S. Thoresen¹⁹⁸, E. Tsai¹⁹⁸, D. Wood¹⁹⁸, S. Dittmer¹⁹⁹, K. A. Hahn¹⁹⁹, Y. Liu¹⁹⁹, M. McGinnis¹⁹⁹, Y. Miao¹⁹⁹, D. G. Monk¹⁹⁹, M. H. Schmitt¹⁹⁹, A. Taliere¹⁹⁹, M. Velasco¹⁹⁹, J. Wang¹⁹⁹, G. Agarwal²⁰⁰, R. Band²⁰⁰, R. Bucci²⁰⁰, S. Castells²⁰⁰, A. Das²⁰⁰, A. Ehnis²⁰⁰, R. Goldouzian²⁰⁰, M. Hildreth²⁰⁰, K. Hurtado Anampa²⁰⁰, T. Ivanov²⁰⁰, C. Jessop²⁰⁰, A. Karneyev²⁰⁰, K. Lannon²⁰⁰, J. Lawrence²⁰⁰, N. Loukas²⁰⁰, L. Lutton²⁰⁰, J. Mariano²⁰⁰, N. Marinelli²⁰⁰, I. McAlister²⁰⁰, T. McCauley²⁰⁰, C. Mcgrady²⁰⁰, C. Moore²⁰⁰, Y. Muesienko²⁰⁰, H. Nelson²⁰⁰, M. Oscherson²⁰⁰, A. Piccinelli²⁰⁰, R. Ruchti²⁰⁰, A. Townsend²⁰⁰, Y. Wan²⁰⁰, M. Wayne²⁰⁰, H. Yockey²⁰⁰, A. Basnet²⁰¹, M. Carrigan²⁰¹, R. De Los Santos²⁰¹, L. S. Durkin²⁰¹, C. Hill²⁰¹, M. Joyce²⁰¹, M. Nunez Ornelas²⁰¹, D. A. Wenzl²⁰¹, B. L. Winer²⁰¹, B. R. Yates²⁰¹, H. Bouchamaoui²⁰², K. Coldham²⁰², P. Das²⁰², G. Dezoort²⁰², P. Elmer²⁰², A. Frankenthal²⁰², M. Galli²⁰², B. Greenberg²⁰², N. Haubrich²⁰², K. Kennedy²⁰², G. Kopp²⁰², Y. Lai²⁰², D. Lange²⁰², A. Loeliger²⁰², D. Marlow²⁰², I. Ojalvo²⁰², J. Olsen²⁰², J. F. Simpson²⁰², D. Stickland²⁰², C. Tully²⁰², S. Malik²⁰², R. Sharma²⁰², S. Chandra²⁰², R. Chawla²⁰², A. Gu²⁰², L. Gutay²⁰², M. Jones²⁰², A. W. Jung²⁰², D. Kondratyev²⁰², M. Liu²⁰², G. Negro²⁰², N. Neumeister²⁰², G. Paspalaki²⁰², S. Piperov²⁰², N. R. Saha²⁰², J. F. Schulte²⁰², F. Wang²⁰², A. Wildridge²⁰², W. Xie²⁰², Y. Yao²⁰², Y. Zhong²⁰², N. Parashar²⁰⁴, A. Pathak²⁰⁴, E. Shumka²⁰⁴, D. Acosta²⁰⁴, A. Agrawal²⁰⁵, C. Arbour²⁰⁵, T. Carnahan²⁰⁵, K. M. Ecklund²⁰⁵, P. J. Fernández Manteca²⁰⁵, S. Freed²⁰⁵, P. Gardner²⁰⁵, F. J. M. Geurts²⁰⁵, T. Huang²⁰⁵, I. Krommydas²⁰⁵, N. Lewis²⁰⁵, W. Li²⁰⁵, J. Lin²⁰⁵, O. Miguel Colin²⁰⁵, B. P. Padley²⁰⁵, R. Redjimi²⁰⁵, J. Rotter²⁰⁵, E. Yigitbas²⁰⁵, Y. Zhang²⁰⁵, O. Bessidskaia Bylund²⁰⁶, A. Bodek²⁰⁶, P. de Barbaro²⁰⁶, R. Demina²⁰⁶, A. Garcia-Bellido²⁰⁶, H. S. Hare²⁰⁶, O. Hindrichs²⁰⁶, N. Parmar²⁰⁶, P. Parygin²⁰⁶, H. Seo²⁰⁶, R. Taus²⁰⁶, B. Chiarito²⁰⁷, J. P. Chou²⁰⁷, S. V. Clark²⁰⁷, S. Donnelly²⁰⁷, D. Gadjari²⁰⁷, Y. Gershtein²⁰⁷, E. Halkiadakis²⁰⁷, M. Heindl²⁰⁷, C. Houghton²⁰⁷, D. Jaroslowski²⁰⁷, S. Konstantinou²⁰⁷, I. Laflotte²⁰⁷, A. Lath²⁰⁷, J. Martins²⁰⁷, B. Rand²⁰⁷, J. Reichert²⁰⁷, P. Saha²⁰⁷, S. Salur²⁰⁷, S. Schnetzer²⁰⁷, S. Somalwar²⁰⁷, R. Stone²⁰⁷, S. A. Thayer²⁰⁷, S. Thomas²⁰⁷, J. Vora²⁰⁷, D. Ally²⁰⁸, A. G. Delanno²⁰⁸, S. Fiorendi²⁰⁸, J. Harris²⁰⁸, S. Higginbotham²⁰⁸, T. Holmes²⁰⁸, A. R. Kanuganti²⁰⁸, N. Karunarathna²⁰⁸, J. Lawless²⁰⁸, L. Lee²⁰⁸, E. Nibigira²⁰⁸, B. Skipworth²⁰⁸, S. Spanier²⁰⁸, D. Aebi²⁰⁹, M. Ahmad²⁰⁹, T. Akhter²⁰⁹, K. Androsow²⁰⁹, A. Bolshov²⁰⁹, O. Bouhali²⁰⁹, A. Cagnotta²⁰⁹, V. D'Amante²⁰⁹, R. Eusebi²⁰⁹, P. Flanagan²⁰⁹, J. Gilmore²⁰⁹, Y. Guo²⁰⁹, T. Kamon²⁰⁹, S. Luo²⁰⁹, R. Mueller²⁰⁹, A. Safonov²⁰⁹, N. Akchurin²¹⁰, J. Damgov²¹⁰, Y. Feng²¹⁰, N. Gogate²¹⁰, Y. Kazhykarim²¹⁰, K. Lamichhane²¹⁰, S. W. Lee²¹⁰, C. Madrid²¹⁰, A. Manke²¹⁰, T. Peltola²¹⁰, I. Volobouev²¹⁰, E. Appelt²¹¹, Y. Chen²¹¹, S. Greene²¹¹, A. Gurrola²¹¹, W. Johns²¹¹, R. Kunnawalkam Elayal²¹¹, A. Melo²¹¹, D. Rathjens²¹¹, F. Romeo²¹¹, P. Sheldon²¹¹, S. Tuo²¹¹, J. Velkovska²¹¹, J. Viinikainen²¹¹, J. Zhang²¹¹, B. Cardwell²¹², H. Chung²¹², B. Cox²¹², J. Hakala²¹², R. Hirosky²¹², M. Jose²¹², A. Ledovsky²¹², C. Mantilla²¹², C. Neu²¹², C. Ramón Álvarez²¹², S. Bhattacharya²¹³, P. E. Karchin²¹³, A. Aravind²¹³, S. Banerjee²¹³, K. Black²¹³, T. Bose²¹³, E. Chavez²¹³, S. Dasu²¹³, P. Everaerts²¹³, C. Galloni²¹³, H. He²¹³, M. Herndon²¹³, A. Herve²¹³, C. K. Koraka²¹³, S. Lomte²¹³, R. Loveless²¹³, A. Mallampalli²¹³, A. Mohammadi²¹³, S. Mondal²¹³, T. Nelson²¹³, G. Parida²¹³, L. Pétré²¹³, D. Pinna²¹³, A. Savin²¹³, V. Shang²¹³, V. Sharma²¹³, W. H. Smith²¹³, D. Teague²¹³, H. F. Tsoi²¹³, W. Vetens²¹³, A. Warden²¹³, S. Afanasiev²¹⁵, V. Alexakhin²¹⁵, Yu. Andreev²¹⁵, T. Aushkev²¹⁵, D. Budkousk²¹⁵, R. Chistov²¹⁵, M. Danilov²¹⁵, T. Dimova²¹⁵, A. Ershov²¹⁵, S. Gninenko²¹⁵, I. Gorbunov²¹⁵, A. Gribushin²¹⁵, A. Kamenev²¹⁵, V. Karjavine²¹⁵, M. Kirsanov²¹⁵, V. Klyukhin²¹⁵, O. Kodolova²¹⁵, V. Korenko²¹⁵, A. Kozlyev²¹⁵, N. Krasnikov²¹⁵, A. Lanev²¹⁵, A. Malakhov²¹⁵, V. Matveev²¹⁵, A. Nikitenko^{170,215}, V. Palichik²¹⁵, V. Perelygin²¹⁵, S. Petrushanko²¹⁵, S. Polikarpov²¹⁵, O. Radchenko²¹⁵, M. Savina²¹⁵, V. Shalaev²¹⁵, S. Shmatov²¹⁵, S. Shultha²¹⁵, Y. Skovpen²¹⁵, V. Smirnov²¹⁵, O. Teryaev²¹⁵, I. Tisova²¹⁵, A. Toropin²¹⁵, N. Voytishin²¹⁵, B. S. Yuldashev²¹⁵, A. Zarubin²¹⁵, I. Zhizhin²¹⁵, L. Dudko²¹⁶, K. Ivanov²¹⁶, V. Kim²¹⁶, V. Murzin²¹⁶, V. Oreshkin²¹⁶, D. Sosnov²¹⁶, E. Boos²¹⁶, V. Bunichev²¹⁶, M. Dubinin^{183,216}, V. Savrin²¹⁶ & A. Snigirev²¹⁶

¹Yerevan Physics Institute, Yerevan, Armenia. ²Institut für Hochenergiephysik, Vienna, Austria. ³Universiteit Antwerpen, Antwerpen, Belgium. ⁴Vrije Universiteit Brussel, Brussels, Belgium. ⁵Ghent University, Ghent, Belgium. ⁶Université Libre de Bruxelles, Brussels, Belgium. ⁷Université Catholique de Louvain, Louvain-la-Neuve, Belgium. ⁸Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil. ⁹Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil. ¹⁰Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil. ¹¹Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria. ¹²University of Sofia, Sofia, Bulgaria. ¹³Instituto De Alta Investigación, Universidad de Tarapacá, Arica, Chile. ¹⁴Universidad Técnica Federico Santa María, Valparaíso, Chile. ¹⁵Beihang University, Beijing, China. ¹⁶Department of Physics, Tsinghua University, Beijing, China. ¹⁷Institute of High Energy Physics, Beijing, China. ¹⁸State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China. ¹⁹Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China. ²⁰Sun Yat-Sen University, Guangzhou, China. ²¹University of Science and Technology of China, Hefei, China. ²²Nanjing Normal University, Nanjing, China. ²³The University of Iowa, Iowa City, IA, USA. ²⁴Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China. ²⁵Zhejiang University, Hangzhou, China. ²⁶Universidad de Los Andes, Bogotá, Colombia. ²⁷Universidad de Antioquia, Medellín, Colombia. ²⁸Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture University of Split, Split, Croatia. ²⁹Faculty of Science, University of Split, Split, Croatia. ³⁰Institute Rudjer Boskovic, Zagreb, Croatia. ³¹University of Cyprus, Nicosia, Cyprus. ³²Charles University, Prague, Czech Republic. ³³Escuela Politécnica Nacional, Quito, Ecuador. ³⁴Universidad San Francisco de Quito, Quito, Ecuador. ³⁵Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt. ³⁶Center for High Energy Physics (CHEP-FU), Fayoum University, Fayoum, Egypt. ³⁷National Institute of Chemical Physics and Biophysics, Tallinn, Estonia. ³⁸Department of Physics, University of Helsinki, Helsinki, Finland. ³⁹Helsinki Institute of Physics, Helsinki, Finland. ⁴⁰Lappeenranta-Lahti University of Technology, Lappeenranta, Finland. ⁴¹IRFU, CEA, Université

Paris-Saclay, Gif-sur-Yvette, France. ⁴²Purdue University, West Lafayette IN, USA. ⁴³Laboratoire Leprince-Ringuet, CNRS/IN2P3, École Polytechnique, Institut Polytechnique de Paris, Palaiseau, France. ⁴⁴Université de Strasbourg, CNRS IPHC UMR7178, Strasbourg, France. ⁴⁵Centre de Calcul de l'Institut National de Physique Nucléaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France. ⁴⁶Institut de Physique des des2Infinis de Lyon (IP2I), Villeurbanne, France. ⁴⁷Georgian Technical University, Tbilisi, Georgia. ⁴⁸I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany. ⁴⁹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany. ⁵⁰III. Physikalisches Institut B, RWTH Aachen University, Aachen, Germany. ⁵¹Deutsches Elektronen-Synchrotron, Hamburg, Germany. ⁵²University of Hamburg, Hamburg, Germany. ⁵³Karlsruher Institut fuer Technologie, Karlsruhe, Germany. ⁵⁴CERN, European Organization for Nuclear Research, Geneva, Switzerland. ⁵⁵Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece. ⁵⁶National and Kapodistrian University of Athens, Athens, Greece. ⁵⁷National Technical University of Athens, Athens, Greece. ⁵⁸University of Ioánnina, Ioannina, Greece. ⁵⁹HUN-REN Wigner Research Centre for Physics, Budapest, Hungary. ⁶⁰HUN-REN Institute of Nuclear Research - ATOMKI, Debrecen, Hungary. ⁶¹MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary. ⁶²Faculty of Informatics, University of Debrecen, Debrecen, Hungary. ⁶³Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary. ⁶⁴The University of Kansas, Lawrence, KS, USA. ⁶⁵Panjab University, Chandigarh, India. ⁶⁶University of Delhi, New Delhi, India. ⁶⁷University of Hyderabad, Hyderabad, India. ⁶⁸Indian Institute of Technology Kanpur, Kanpur, India. ⁶⁹Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India. ⁷⁰Indian Institute of Technology Madras, Chennai, India. ⁷¹IISER Mohali, Mohali, India. ⁷²Tata Institute of Fundamental Research-A, Mumbai, India. ⁷³Tata Institute of Fundamental Research-B, Mumbai, India. ⁷⁴National Institute of Science Education and Research, Bhubaneswar, India. ⁷⁵Indian Institute of Science Education and Research (IISER), Pune, India. ⁷⁶Indian Institute of Technology Hyderabad, Telangana, India. ⁷⁷Isfahan University of Technology, Isfahan, Iran. ⁷⁸Institute for Research in Fundamental Sciences (IPM), Tehran, Iran. ⁷⁹University College Dublin, Dublin, Ireland. ⁸⁰INFN Sezione di Bari, Bari, Italy. ⁸¹Università di Bari, Bari, Italy. ⁸²Politecnico di Bari, Bari, Italy. ⁸³INFN Sezione di Bologna, Bologna, Italy. ⁸⁴Università di Bologna, Bologna, Italy. ⁸⁵INFN Sezione di Catania, Catania, Italy. ⁸⁶Università di Catania, Catania, Italy. ⁸⁷INFN Sezione di Firenze, Firenze, Italy. ⁸⁸Università di Firenze, Firenze, Italy. ⁸⁹INFN Laboratori Nazionali di Frascati, Frascati, Italy. ⁹⁰INFN Sezione di Genova, Genova, Italy. ⁹¹Università di Genova, Genova, Italy. ⁹²INFN Sezione di Milano-Bicocca, Milano, Italy. ⁹³Università di Milano-Bicocca, Milano, Italy. ⁹⁴INFN Sezione di Napoli, Napoli, Italy. ⁹⁵Università di Napoli 'Federico II', Napoli, Italy. ⁹⁶Università della Basilicata, Potenza, Italy. ⁹⁷Scuola Superiore Meridionale (SSM), Napoli, Italy. ⁹⁸INFN Sezione di Padova, Padova, Italy. ⁹⁹Fermi National Accelerator Laboratory, Batavia, IL, USA. ¹⁰⁰Università di Padova, Padova, Italy. ¹⁰¹INFN Sezione di Pavia, Pavia, Italy. ¹⁰²Università di Pavia, Pavia, Italy. ¹⁰³INFN Sezione di Perugia, Perugia, Italy. ¹⁰⁴Università di Perugia, Perugia, Italy. ¹⁰⁵INFN Sezione di Pisa, Pisa, Italy. ¹⁰⁶Università di Pisa, Pisa, Italy. ¹⁰⁷Scuola Normale Superiore di Pisa, Pisa, Italy. ¹⁰⁸Università di Siena, Siena, Italy. ¹⁰⁹INFN Sezione di Roma, Roma, Italy. ¹¹⁰Gapienza Università di Roma, Roma, Italy. ¹¹¹INFN Sezione di Torino, Torino, Italy. ¹¹²Università di Torino, Torino, Italy. ¹¹³Università del Piemonte Orientale, Novara, Italy. ¹¹⁴INFN Sezione di Trieste, Trieste, Italy. ¹¹⁵Università di Trieste, Trieste, Italy. ¹¹⁶Kyungpook National University, Daegu, Korea. ¹¹⁷Department of Mathematics and Physics, Gangneung-Wonju National University, Gangneung, Korea. ¹¹⁸Institute for Universe and Elementary Particles, Chonnam National University, Kwangju, Korea. ¹¹⁹Hanyang University, Seoul, Korea. ¹²⁰Korea University, Seoul, Korea. ¹²¹Department of Physics, Kyung Hee University, Seoul, Korea. ¹²²Sejong University, Seoul, Korea. ¹²³Seoul National University, Seoul, Korea. ¹²⁴University of Seoung, Seoul, Korea. ¹²⁵Department of Physics, Yonsei University, Seoul, Korea. ¹²⁶Sungkyunkwan University, Suwon, Korea. ¹²⁷College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait. ¹²⁸College of Science, Department of Physics, Kuwait University, Safat, Kuwait. ¹²⁹Riga Technical University, Riga, Latvia. ¹³⁰University of Latvia (LU), Riga, Latvia. ¹³¹Vilnius University, Vilnius, Lithuania. ¹³²National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia. ¹³³Universidad de Sonora (UNISON), Hermosillo, Mexico. ¹³⁴Centro de Investigación y de Estudios Avanzados del IPN, Mexico City, Mexico. ¹³⁵Universidad Iberoamericana, Mexico City, Mexico. ¹³⁶Benemerita Universidad Autónoma de Puebla, Puebla, Mexico. ¹³⁷University of Montenegro, Podgorica, Montenegro. ¹³⁸University of Canterbury, Christchurch, New Zealand. ¹³⁹National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan. ¹⁴⁰AGH University of Krakow, Krakow, Poland. ¹⁴¹National Centre for Nuclear Research, Swierk, Poland. ¹⁴²Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland. ¹⁴³Warsaw University of Technology, Warsaw, Poland. ¹⁴⁴Laboratório de Instrumentação e Física Experimental de Partículas, Lisbon, Portugal. ¹⁴⁵Faculty of Physics, University of Belgrade, Belgrade, Serbia. ¹⁴⁶VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia. ¹⁴⁷Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain. ¹⁴⁸Universidad Autónoma de Madrid, Madrid, Spain. ¹⁴⁹Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Universidad de Oviedo, Oviedo, Spain. ¹⁵⁰Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain. ¹⁵¹University of Colombo, Colombo, Sri Lanka. ¹⁵²Department of Physics, University of Ruhuna, Matara, Sri Lanka. ¹⁵³PSI Center for Neutron and Muon Sciences, Villigen, Switzerland. ¹⁵⁴Universität Zürich, Zurich, Switzerland. ¹⁵⁵Institute for Particle Physics and Astrophysics (IPA), ETH Zurich, Zurich, Switzerland. ¹⁵⁶National Central University, Chung-Li, Taiwan. ¹⁵⁷National Taiwan University (NTU), Taipei, Taiwan. ¹⁵⁸High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand. ¹⁵⁹Tunis El Manar University, Tunis, Tunisia. ¹⁶⁰Physics Department, Science and Art Faculty, Çukurova University, Adana, Turkey. ¹⁶¹Physics Department, Middle East Technical University, Ankara, Turkey. ¹⁶²Bogazici University, Istanbul, Turkey. ¹⁶³Istanbul Technical University, Istanbul, Turkey. ¹⁶⁴Istanbul University, Istanbul, Turkey. ¹⁶⁵Yildiz Technical University, Istanbul, Turkey. ¹⁶⁶Institute for Scintillation Materials, National Academy of Science of Ukraine, Kharkiv, Ukraine. ¹⁶⁷National Science Centre Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine. ¹⁶⁸University of Bristol, Bristol, UK. ¹⁶⁹Rutherford Appleton Laboratory, Didcot, UK. ¹⁷⁰Imperial College, London, UK. ¹⁷¹Brunel University, Uxbridge, UK. ¹⁷²Baylor University, Waco, TX, USA. ¹⁷³Bethel University, St Paul, MN, USA. ¹⁷⁴Catholic University of America, Washington, DC, USA. ¹⁷⁵The University of Alabama, Tuscaloosa, AL, USA. ¹⁷⁶Boston University, Boston, MA, USA. ¹⁷⁷Brown University, Providence, RI, USA. ¹⁷⁸University of California Davis, Davis, CA, USA. ¹⁷⁹University of California Los Angeles, Los Angeles CA, USA. ¹⁸⁰University of California

Riverside, Riverside, CA, USA. ¹⁸¹University of California San Diego, La Jolla, Ca, USA. ¹⁸²Department of Physics, University of California Santa Barbara, Santa Barbara, CA, USA. ¹⁸³California Institute of Technology, Pasadena, CA, USA. ¹⁸⁴Carnegie Mellon University, Pittsburgh, PA, USA. ¹⁸⁵University of Colorado Boulder, Boulder, CO, USA. ¹⁸⁶Cornell University, Ithaca NY, USA. ¹⁸⁷University of Florida, Gainesville, FL, USA. ¹⁸⁸Florida State University, Tallahassee, FL, USA. ¹⁸⁹Florida Institute of Technology, Melbourne, FL, USA. ¹⁹⁰University of Illinois Chicago, Chicago, IL, USA. ¹⁹¹Johns Hopkins University, Baltimore, MD, USA. ¹⁹²Kansas State University, Manhattan, KS, USA. ¹⁹³University of Maryland, College Park, MD, USA. ¹⁹⁴Massachusetts Institute of Technology, Cambridge, MA, USA. ¹⁹⁵University of Minnesota, Minneapolis, MN, USA. ¹⁹⁶University of Nebraska–Lincoln, Lincoln, NE, USA. ¹⁹⁷State University of New York at Buffalo, Buffalo, NY, USA. ¹⁹⁸Northeastern University, Boston, MA, USA.

¹⁹⁹Northwestern University, Evanston, IL, USA. ²⁰⁰University of Notre Dame, Notre Dame IN, USA. ²⁰¹The Ohio State University, Columbus OH, USA. ²⁰²Princeton University, Princeton, NJ, USA. ²⁰³University of Puerto Rico, Mayaguez, PR, USA. ²⁰⁴Purdue University Northwest, Hammond IN, USA. ²⁰⁵Rice University, Houston, TX, USA. ²⁰⁶University of Rochester, Rochester NY, USA. ²⁰⁷Rutgers, The State University of New Jersey, Piscataway, NJ, USA. ²⁰⁸University of Tennessee, Knoxville, TN, USA. ²⁰⁹Texas A&M University, College Station, TX, USA. ²¹⁰Texas Tech University, Lubbock, TX, USA. ²¹¹Vanderbilt University, Nashville, TN, USA. ²¹²University of Virginia, Charlottesville, VA, USA. ²¹³Wayne State University, Detroit, MI, USA. ²¹⁴University of Wisconsin–Madison, Madison, WI, USA. ²¹⁵An institute or international laboratory covered by a cooperation agreement with CERN, Geneva, Switzerland. ²¹⁶An institute formerly covered by a cooperation agreement with CERN, Geneva, Switzerland.

Modelling of hadron properties

To investigate the spin-parity properties of the resonant structure within the J/Ψ J/Ψ invariant mass spectrum in the range of 6.2–8.0 GeV, we use an approach designed to minimize model dependence. This approach relies on the observed J/Ψ J/Ψ invariant mass spectrum and the momentum of the J/Ψ J/Ψ system in both transverse and longitudinal directions with respect to the beam, while remaining independent of the polarization of the system by relying solely on decay angular information. For a spin-zero state, polarization is not relevant. For states with nonzero spin, we assume the state is produced unpolarized, but vary the polarization to evaluate potential small residual effects on the decay angular distributions due to detector acceptance.

The analysis uses a model that ensures consistency with the observations made by CMS, by using simulation adjustments to accurately capture the observed transverse and longitudinal motion, and parameters of the resonances and backgrounds extracted from ref. 32 and shown in Fig. 1. The background arises from nonresonant contributions, single-parton scattering (SPS) and double-parton scattering, plus an empirical term parameterizing the background near the threshold³². The background is parameterized using MC simulation, with adjustments applied to better match the observed data in both the signal and sideband regions.

We start by considering the spin-0 hypothesis for the X states, which are produced without polarization. The helicity amplitudes $A_{\lambda_1\lambda_2}$ of the two J/Ψ mesons are listed in Table 1. For the pseudoscalar state with $J^P = 0^-$, the amplitudes satisfy $A_{++} = -A_{--}$ and $A_{00} = 0$. By contrast, for the scalar state with $J^P = 0^+$, both A_{++} and A_{--} and A_{00} amplitudes contribute, with no specific prediction for the relative magnitude of A_{00} . We adopt the general amplitude approach^{35,46,51}, in which the spin-0 state amplitude can be written as a sum of three Lorentz-invariant structures,

$$A(X_{J=0} \rightarrow V_1 V_2) = a_1(q^2) m_V^2 \epsilon_1^* \epsilon_2^* + a_2(q^2) f_{\mu\nu}^{*(1)} f^{*(2),\mu\nu} + a_3(q^2) f_{\mu\nu}^{*(1)} \tilde{f}^{*(2),\mu\nu}, \quad (1)$$

and where the field strength tensor of a vector boson V_i with momentum q_i and polarization vector ϵ_i is defined as $f^{(i),\mu\nu} = \epsilon_i^\mu q_i^\nu - \epsilon_i^\nu q_i^\mu$, the conjugate field strength tensor is $\tilde{f}^{(i),\mu\nu} = 1/2 \epsilon^{\mu\nu\alpha\beta} f_{\alpha\beta}^{(i)}$, and $q = q_1 + q_2$.

As the X decay proceeds by the strong interaction, which conserves parity, the first two terms in equation (1) can be interpreted as interactions involving scalar particles, corresponding to models O_m^+ and O_h^+ with couplings a_1 and a_2 , respectively. The third term represents the interaction of a pseudoscalar particle, O^- , associated with the coupling a_3 . At $m_{4ii} = 6.9$ GeV, $|A_{00}|^2 = 52\%$ in model O_m^+ and $|A_{00}|^2 = 19\%$ in model O_h^+ . In the following, we analyse the two models, O_m^+ and O_h^+ , separately. Moreover, we consider their combination, denoted as O_{mix}^+ , in which the relative magnitudes and signs of the couplings a_1 and a_2 are varied. A positive relative sign results in constructive interference, whereas a negative relative sign leads to destructive interference.

The polarization of the spin-1 states varies depending on the production mechanism, in which we consider either unpolarized production or polarization with J_z or $J_{z'} = \pm 1$ (ref. 46) because quark-initiated production dominates. The z and z' axes are defined in Extended Data Figure 1, where the motion of the four-muon system within the laboratory frame leads to appearance of noncollinear proton collisions and defines the z' axis, while the z -axis approximates the proton beam line. In the decay process, four helicity amplitudes contribute, and their relationships are shown in Table 1. This is equivalent to two Lorentz-invariant structures in the decay amplitude

$$A(X_{J=1} \rightarrow V_1 V_2) = b_1(q^2) [(\epsilon_1^* q)(\epsilon_2^* \epsilon_X) + (\epsilon_2^* q)(\epsilon_1^* \epsilon_X)] + b_2(q^2) \epsilon_{\alpha\mu\nu\beta} \epsilon_X^\alpha \epsilon_1^{*\mu} \epsilon_2^{*\nu} \tilde{q}^\beta, \quad (2)$$

where $\tilde{q} = q_1 - q_2$ and ϵ_X is the polarization vector of the spin-1 resonance X. The b_1 and b_2 are the couplings to a vector state with $J^P = 1^-$ and an axial vector with $J^P = 1^+$, respectively.

The polarization of the spin-2 states also depends on the production mechanism, in which we consider either unpolarized production or polarization with J_z or $J_{z'} = 0$ or ± 2 (ref. 46) because gluon-initiated production is expected to dominate. In the decay process, all nine helicity amplitudes $A_{\lambda_1\lambda_2}$ contribute. This corresponds to the Lorentz-invariant structures in the decay amplitude found in refs. 35,46. There are two degrees of freedom in the 2^{++} case, and we use the 2_m^- and 2_h^- models, corresponding to $g_8^{(2)} \neq 0$ and $g_{10}^{(2)} \neq 0$ in refs. 35,46, respectively. The 2_m^- model corresponds to $A_{++} = -A_{--}$, with decay angular distributions identical to those of the 0^- case. For 2_h^- , we have $A_{+0} = A_{0+} = -A_{-0} = -A_{0-}$, with decay angular distributions identical to those of the 1^- case.

We use a single representative model for the $J^{PC} = 2^{++}$ state, corresponding to model $g_1^{(2)} = g_5^{(2)} \neq 0$ denoted as 2_m^+ in refs. 35,46. This 2_m^+ model is chosen because the composition of $A_{\lambda_1\lambda_2}$ amplitudes represents all possible polarizations, in particular those that are unique for spin-2, A_{+-} and A_{-+} and the tensor structure of interactions is minimal, avoiding the inclusion of higher-dimension operators. At $m_{4ii} = 6.9$ GeV, $2 |A_{+-}|^2 = 9\%$, $|A_{00}|^2 = 21\%$, $4 |A_{+0}|^2 = 47\%$ and $2 |A_{+-}|^2 = 23\%$ in model 2_m^+ . Therefore, if the data are consistent with $J^P = 2_m^+$ and not with the other models, it will provide an unambiguous determination of $J \geq 2$. A higher spin scenario ($J > 2$) could also be possible, and it would exhibit angular distributions similar to those of $J = 2$ with the same parity^{35,46}.

As the $J^{PC} = 2^{++}$ state has four degrees of freedom in the amplitude composition, three other possibilities could be chosen that lead to the same observable decay distributions as the O_m^+ , O_h^+ and 1^+ states. Interference between the corresponding 2^{++} amplitude tensor structures is also possible. Therefore, from a purely decay-based analysis, if the data are consistent with either the 0^{++} or 1^{++} model, we cannot rule out a general 2^{++} model. Although analysing polarization information through production-sensitive angular distributions could help resolve this ambiguity, this analysis is more challenging due to uncertainty in the production mechanism. If the resonance is produced unpolarized, such an analysis would not provide additional information. In this study, we check production-sensitive angular distributions for consistency with an unpolarized case, whereas a more detailed analysis is left for future work.

Determining the form factors associated with the tensor structure of interactions, represented by $a_i(q^2)$ in equation (1) for spin-0, $b_i(q^2)$ in equation (2) for spin-1, or equivalent $g_i^{(2)}(q^2)$ in refs. 35,46 for spin-2, with $q^2 = m_{4ii}^2$, is nontrivial and relies on the model of strong interactions. Nevertheless, it is completely separable from the computation of the angular distributions, given a particular tensor structure of interactions. Furthermore, interference between the resonances could result in a complex alteration of the angular distributions. Nonetheless, assuming that all three resonances possess identical quantum numbers and coupling constants, the angular distributions remain unaffected by interference effects observed in the mass spectrum. Thus, in this study, we use an empirical approach to analyse the four-muon mass spectrum observed in the data, separating it from the analysis of angular distributions. This makes our analysis independent of the form factor model. The form factors and interference effects are integrated into the observed mass spectrum, as shown in Fig. 1.

Angular observables

The angular information used in the analysis is shown in Extended Data Fig. 1 (refs. 35,46). Two production axes are defined, corresponding to the two production mechanisms, either the direct short-distance production in parton collisions or the fragmentation of a single parton into the hadron. The z -axis is parallel to the beam line within the frame boosted along the beam line from the laboratory frame, in which the longitudinal momentum of X is zero, potentially reflecting polarization

in the two-parton collisions. The z' -axis aligns with the direction of the X momentum, reflecting potential polarization in the single-parton fragmentation scenario.

The decay angles Φ , θ_1 and θ_2 are defined with reference to Fig. 2. The production angles θ^* and Φ_1 , or alternatively θ'^* and Φ_1' , are defined with respect to the z - or z' -axis, respectively. The angle θ^* is defined between the z -axis and the X decay axis in its rest frame, whereas Φ_1 is the angle between the first J/Ψ decay plane and the production plane, defined by the z -axis and the X decay axis. Extended Data Fig. 1 shows only θ^* and Φ_1 , whereas θ'^* and Φ_1' are defined analogously. These production angles are not used directly in the analysis to avoid dependence on the production model, but they are checked for consistency.

The distributions of the decay and production angles in the range $6.2 < m_{4\mu} < 8.0$ GeV are presented in Extended Data Fig. 2, together with the five signal models shown. It is important to note that the one-dimensional angular distributions in Extended Data Fig. 2 do not capture all the information accessible to the optimal discriminant. Correlations between angles are lost in the projections; for instance, the models $J^{PC} = 1^-+$ and 1^{++} cannot be distinguished in each projection, but they can be separated using the optimal discriminants, which preserve all angular correlations. Furthermore, Extended Data Fig. 2 shows all events in the range $6.2 < m_{4\mu} < 8.0$ GeV together, without accounting for the correlation between the angular distributions and $m_{4\mu}$, or the variation in signal purity with $m_{4\mu}$ due to the resonance structure appearing above the background. As a result, the separation power shown in these illustration plots is considerably diminished compared with that in the full analysis.

To eliminate the dependence on the initial polarization of nonzero-spin states, the production angles are excluded from the data analysis. The assumption is made that the resonances are produced unpolarized, although the variation of this polarization along the production axes is allowed for states with nonzero spin. This allows for the examination of any residual effects stemming from polarization dependence due to detector acceptance effects. Extended Data Fig. 2 shows that the production angular distributions are consistent with unpolarized resonance states along both the z - and z' -axes.

Matrix element approach

The analysis of the multidimensional distributions $\mathcal{P}(\theta_1, \theta_2, \Phi, m_{4\mu})$ is complicated by the complex description and the nontrivial effects of detector reconstruction. Rather than using the three angular observables directly, we construct a single observable that effectively projects the angular information onto one dimension and is optimal for distinguishing between two hypotheses, J_i^P and J_j^P . Using a matrix element likelihood approach^{35,46,51}, a kinematic discriminant is formulated based on the ratio of probabilities for hypotheses J_i^P and J_j^P :

$$\mathcal{D}_{ij}(m_{4\mu}, \Omega) = \frac{\mathcal{P}_j(m_{4\mu}, \Omega)}{\mathcal{P}_i(m_{4\mu}, \Omega) + \mathcal{P}_j(m_{4\mu}, \Omega)}, \quad (3)$$

where \mathcal{P}_i is the normalized probability based on the matrix element squared for a given hypothesis J_i^P . These matrix elements are computed within the same quantum-mechanical formalism as used for the generation of MC events, as detailed in section 'Angular analysis'.

In equation (3), we use only the decay angles $\Omega = \{\cos\theta_1, \cos\theta_2, \Phi\}$. The analysis is conducted using the two-dimensional distributions of $(m_{4\mu}, \mathcal{D}_{ij})$. Production information, including $\{\cos\theta^*, \Phi_1\}$ or $\{\cos\theta'^*, \Phi_1'\}$, can be incorporated into a future analysis if a study of the resonance polarization is desired. The distributions of the discriminants, which were calculated to assess alternative models in comparison to the 2_m^+ model, are presented in Fig. 3 and Extended Data Fig. 3 for all models defined without considering amplitude mixtures. The kinematic distributions in Fig. 3 and Extended Data Figs. 2 and 3 have also been separately examined in three different ranges of $m_{4\mu}$, each corresponding to one of the three resonance structures shown in Fig. 1. The data

remain consistent with the 2_m^+ model across the full mass range as well as within each of the three individual intervals.

The final statistical analysis of the data is carried out using two-dimensional distributions of events, $\mathcal{P}_{ij}(m_{4\mu}, \mathcal{D}_{ij})$, where the predicted $m_{4\mu}$ distribution is modelled in the same way as in Fig. 1 following ref. 32, and the \mathcal{D}_{ij} distribution is obtained in bins of 0.05 GeV in $m_{4\mu}$ from the detailed MC simulation discussed earlier in section 'Angular analysis'. For any given slice of $m_{4\mu}$, only five bins of \mathcal{D}_{ij} are used, meaning that the MC simulation can accurately predict probability distributions using an affordable number of simulated events.

Systematic uncertainties

In the maximum likelihood fit introduced in section 'Quantum number determination' and further detailed in the next section, the likelihood function is maximized with respect to the nuisance parameters, which include those representing systematic uncertainties. The fit results encompass systematic variations in the parameterization of both signal and background models. The yields of the signal and each of the background processes are treated as free parameters, to be fully determined by the fit to the data. Further variations are categorized into two groups: variations in mass shapes and variations in discriminants, applied to both signal and background components.

The study of systematic variations in mass shapes was detailed in ref. 32, which includes resonance parameterization, resolution and efficiency for signal components, as well as different models for background components. All these variations in mass shapes were incorporated into the two-dimensional analysis presented here, through the variation of $\mathcal{P}_k(m_{4\mu})$ in equation (4). Therefore, all uncertainties associated with the parameterization of the $m_{4\mu}$ spectrum as detailed in ref. 32 are included in the analysis.

One of the uncertainties arises from the unknown angular distributions of the background contribution near the kinematic threshold³². To allow maximum flexibility, the discriminant parameterization of this contribution, $T_{ijk}(\mathcal{D}_{ij}|m_{4\mu})$ in equation (4), is expressed as the sum of the SPS background model and the two signal models under investigation, with the relative contributions of all three components determined by freely floating nuisance parameters. This approach is intended to accommodate a broad range of hypotheses for the threshold contribution, including possible variations of the nonresonant background and potential resonance excitations.

Other systematic variations in discriminants account for possible inconsistencies in p_T and p_z distributions between data and MC simulation. The p_T and p_z distributions are validated and tuned through comparisons between data and simulation in both the sideband region (8.0–9.0 GeV) and the signal region (6.2–8.0 GeV). Residual discrepancies in the discriminants due to mismodelling of p_T and p_z are addressed through nuisance parameters, which allow for alternative discriminant parameterizations of $T_{ijk}(\mathcal{D}_{ij}|m_{4\mu})$ in equation (4). Similarly, uncertainties in the kinematic distributions of the nonresonant background components are evaluated through comparisons between data and MC simulations in the sideband region, in which a satisfactory agreement is observed. Remaining discrepancies in the discriminants are handled through nuisance parameters that permit alternative discriminant parameterizations of $T_{ijk}(\mathcal{D}_{ij}|m_{4\mu})$.

The nominal discriminant parameterization of all spin-1 and spin-2 resonances assumes no polarization. However, to evaluate the impact of detector acceptance effects on the results, we introduce an alternative parameterization that assumes polarization of spin-1 and spin-2 resonances along either the z - or z' -axis. This approach models polarized production arising from either two-parton collisions or single-parton fragmentation.

Statistical analysis

We perform a binned extended maximum likelihood fit in which the probability density function is a sum of contributions from all signal

Article

and background processes implemented in the Combine tool (v.10.0.2) (ref. 52). This method mirrors the approach used to determine the spin-parity quantum numbers of the Higgs boson at the LHC⁵³, as detailed in ref. 35. Each process k is characterized by a probability density function \mathcal{P}_{ijk} , used to analyse signal hypotheses i and j . This function depends on two observables, $m_{4\mu}$ and \mathcal{D}_{ij} , and is defined as a template binned in a 36×5 grid:

$$\mathcal{P}_{ijk}(m_{4\mu}, \mathcal{D}_{ij}) = \mathcal{P}_k(m_{4\mu}) T_{ijk}(\mathcal{D}_{ij} | m_{4\mu}), \quad (4)$$

where $\mathcal{P}_k(m_{4\mu})$ represents the probability density function of the invariant mass $m_{4\mu}$, which is independent of the hypotheses being tested. The probability density T_{ijk} is a normalized function of \mathcal{D}_{ij} given a specific value of $m_{4\mu}$, obtained from simulation and including systematic variations through alternative distributions, as described above. We assume the same quantum numbers and couplings for all signal resonances, enabling the use of a shared T_{ijk} for parameterizing the signal.

To distinguish between alternative models, the test statistic $q = -2 \ln(\mathcal{L}_{j^p} / \mathcal{L}_{j^p})$ is defined using the ratio of signal plus background likelihoods for the two signal hypotheses. The likelihood is maximized with respect to the nuisance parameters, which include the yields of signal and background (bkg) processes and constrained parameters describing the systematic uncertainties. To quantify the consistency of the observed test statistic q_{obs} with the model J_i^p , the probability $P = P(q \leq q_{\text{obs}} | J_i^p + \text{bkg})$ is determined under the signal-plus-background hypothesis using pseudo-experiments. This probability is then translated into a Z -score, representing the number of standard deviations using the one-sided Gaussian tail integral.

The consistency of the observed data with the alternative signal hypothesis (J_j^p) is assessed from $P(q \geq q_{\text{obs}} | J_j^p + \text{bkg})$. The sign is positive if the tail extends away or negative if it extends towards the median of the other hypothesis. The CL_s criterion^{54,55}, defined as $\text{CL}_s = P(q \geq q_{\text{obs}} | J_j^p + \text{bkg}) / P(q \geq q_{\text{obs}} | J_i^p + \text{bkg}) < \alpha$, is used for the final inference of whether a particular alternative hypothesis J_j^p is excluded or not with respect to a reference hypothesis J_i^p at a given confidence level $(1 - \alpha)$. Figure 3b shows example q distributions for the 2_m^+ and 0^- models, obtained from repeated pseudo-experiments simulating the expected experimental outcome.

The pairs of spin-parity models are tested among the 0^- , 0_{nr}^+ , 0_{hr}^+ , 1^- , 1^+ , 2_{nr}^- , 2_{hr}^- , and 2_m^+ hypotheses. In all tests involving the 2_m^+ model, it is preferred. Therefore, these tests are presented in Fig. 4 and Extended Data Table 1, which is an extended version of Table 2. The pairwise tests between the other models do not provide any additional useful information, as the data frequently show inconsistencies with both models. In the case of $J^{PC} = 0^{++}$ and 2^{++} scenarios, additional tests are conducted to account for a possible mixture of two tensor structures, as outlined in section ‘Quantum number determination’.

It is important to note that the 2_m^+ model represents only one specific realization of the $J^{PC} = 2^{++}$ hypothesis. In practice, a mixture of amplitudes corresponding to the 2^{++} state, as listed in Table 1, could produce angular distributions that resemble those expected for 0^{++} or 1^{++} , apart from 2_m^+ . As a result, even if the true particle is a 2^{++} state, the P -values reported in Extended Data Table 1 may not remain fully consistent with the 2_m^+ model under such admixtures. These mixed scenarios can be explored in future analyses. Taking this into account, all observed data distributions are found to be compatible with the $J^{PC} = 2^{++}$ hypothesis and show deviations from the predictions of alternative J^{PC} assignments, with confidence levels summarized in Extended Data Table 1.

Data availability

Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the CMS data preservation, re-use and open access policy (<https://opendata.cern.ch/record/415>).

Code availability

The CMS core software is publicly available at GitHub (<https://github.com/cms-sw/cmssw>).

51. Gritsan, A. V. et al. New features in the JHU generator framework: constraining Higgs boson properties from on-shell and off-shell production. *Phys. Rev. D* **102**, 056022 (2020).
52. CMS Collaboration. The CMS statistical analysis and combination tool: COMBINE. *Comput. Softw. Big Sci.* **8**, 19 (2024).
53. CMS Collaboration. Constraints on the spin-parity and anomalous HVV couplings of the Higgs boson in proton collisions at 7 and 8 TeV. *Phys. Rev. D* **92**, 012004 (2015).
54. Junk, T. Confidence level computation for combining searches with small statistics. *Nucl. Instrum. Methods Phys. Res. A* **434**, 435–443 (1999).
55. Read, A. L. Presentation of search results: the CL_s technique. *J. Phys. G* **28**, 2693–2704 (2002).

Acknowledgements We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. Furthermore, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid and other centres for delivering so effectively the computing infrastructure essential to our analyses. We also acknowledge the Advanced Research Computing at Hopkins (ARCH) for providing computing resources essential for this analysis. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia); BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST and NSFC (China); Minciencias (Colombia); MES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHEI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (the United Kingdom); DOE and NSF (the United States). Rachada-pisek Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, 101001205 and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l’Industrie et dans l’Agriculture (FRIA-Belgium); the Beijing Municipal Science and Technology Commission, no. Z191100007219010, the Fundamental Research Funds for the Central Universities, the Ministry of Science and Technology of China, under grant no. 2023YFA1605804, and the Natural Science Foundation of China under grant no. 120611401002 (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany’s Excellence Strategy—EXC 2121 ‘Quantum Universe’—390833306 and under project no. 400140256—GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), project no. 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program—UNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64 and 2021-4.1.2-NEMZ_KI-2024-00036 (Hungary); the Council of Science and Industrial Research, India; ICSC—National Research Centre for High Performance Computing, Big Data and Quantum Computing and FAIR—Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MICIU/AEI/10.13039/501100011033, ERDF/EU, ‘European Union NextGenerationEU/PRTR’ and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund through the Program Management Unit for Human Resources and Institutional Development, Research and Innovation, grant B39G680009 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (the United States).

Author contributions All authors have contributed to the publication, being variously involved in the design and the construction of the detectors, in writing software, calibrating subsystems, operating the detectors and acquiring data, and finally analysing the processed data. The CMS Collaboration members discussed and approved the scientific results. The manuscript was prepared by a subgroup of authors appointed by the Collaboration and subject to an internal collaboration-wide review process. All authors reviewed and approved the final version of the paper.


Competing interests The authors declare no competing interests.

Additional information


Correspondence and requests for materials should be addressed to The CMS Collaboration.

Peer review information Nature thanks Roy Briere, Renu Garg, Tatjana Lenz and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at <http://www.nature.com/reprints>.




Extended Data Fig. 1 | Angular observables. The production and decay of a resonance X in proton collisions $pp \rightarrow X \rightarrow V_1 V_2 \rightarrow 4\mu$ define the angular observables in the centre-of-mass frames of the corresponding particles^{35,46}, where the V_1 and V_2 refer to the J/ψ mesons. The z axis approximates the proton beam line, while the z' axis corresponds to the direction of the four-muon system.



Extended Data Fig. 2 | Angular distributions. Distribution of the decay angles: Φ (upper row), $\cos\theta_1$, $\cos\theta_2$ (second row), production angles defined with respect to axis z : Φ_1 , $\cos\theta^*$ (third row), and defined with respect to axis z' : Φ_1' , $\cos\theta'^*$ (lower row) in the range $6.2 < m_{4\mu} < 8.0$ GeV presented together with

the five J_i^P signal models. The background is subtracted from the data, based on the expected distributions, and systematic uncertainties are not incorporated in these plots. The 0^- and 2_m^- distributions are identical, as are those of 1^- and $2_{\bar{h}}^-$.



Extended Data Fig. 3 | Optimal observables. Distributions of D_{ij} that are optimal for separating the 2_m^+ model against the 0_m^+ (upper left), 0_h^+ (upper right), 1^+ (lower left), and $1^+ [2_h]$ (lower right) models in the range $6.2 < m_{4\mu} < 8.0$ GeV. Distributions for signal only (dashed) and for signal plus background (solid and dash-dot-dotted) models are compared to the experimental data points with

error bars, with uncertainty bands representing post-fit model uncertainties, which are partially correlated with the data. The 1^+ and 2_h^+ distributions are identical, apart from systematic uncertainties arising from polarization effects. The lower panels display the ratios of the data and of the model predictions to the mean expectations from the 2_m^+ model.

Article

Extended Data Table 1 | Results from hypothesis test for pairs of spin-parity models

		Observed		Expected	
		<i>p</i> -value	<i>Z</i> -score	<i>p</i> -value	<i>Z</i> -score
0^- vs 2_m^+	0^-	2.7×10^{-13}	7.2	6.5×10^{-14}	7.4
	2_m^+	0.42	0.2	0.5	0
0_m^+ vs 2_m^+	0_m^+	4.3×10^{-5}	3.9	5.6×10^{-9}	5.7
	2_m^+	0.072	1.5	0.5	0
0_{mix}^+ vs 2_m^+	0_{mix}^+	1.4×10^{-2}	2.2	8.4×10^{-4}	3.1
	2_m^+	0.17	1.0	0.5	0
0_h^+ vs 2_m^+	0_h^+	3.1×10^{-9}	5.8	8.5×10^{-5}	3.8
	2_m^+	0.90	-1.3	0.5	0
1^- vs 2_m^+	1^-	8.0×10^{-8}	5.2	6.4×10^{-9}	5.7
	2_m^+	0.38	0.3	0.5	0
1^+ vs 2_m^+	1^+	4.7×10^{-3}	2.6	2.7×10^{-5}	4.0
	2_m^+	0.052	1.6	0.5	0
2_m^- vs 2_m^+	2_m^-	4.1×10^{-12}	6.8	3.9×10^{-14}	7.5
	2_m^+	0.28	0.6	0.5	0
2_{mix}^- vs 2_m^+	2_{mix}^-	6.5×10^{-4}	3.2	1.5×10^{-4}	3.6
	2_m^+	0.31	0.5	0.5	0
2_h^- vs 2_m^+	2_h^-	2.2×10^{-8}	5.5	6.3×10^{-9}	5.7
	2_m^+	0.43	0.2	0.5	0

This is an extended version of Table 2. The expected P-value is presented based on the assumption of the scenario of 2_m^+ . Results with $Z > 5$ have been derived through Gaussian extrapolation.