



OPEN Sociodemographic and clinical characteristics of Palestinian patients with chronic pain: a cross-sectional study from Al-Shifa Hospital, Gaza

Abdallah AbuJlambo^{1✉}, Maha AbuZarifa², Yara Ashour³, Hanne Lossius⁴ & Guido Veronese⁵

This cross-sectional study examined chronic pain, mental health, and pain catastrophizing among 272 Palestinian patients at Al-Shifa Hospital in Gaza. Participants (median age 43; 57.7% female) completed assessments using the depression, anxiety, and stress (DASS-21) and pain catastrophizing (PCS) scales. Most were married (78.7%), unemployed (50%), and lived on <1,974 ILS/month (94.1%). Lower limb pain (26.8% knee) predominated, with a median pain duration of 4 years. High rates of mental health symptoms were observed: depression (83.1%), anxiety (83.8%), and stress (88.6%), with 56.3% showing extremely severe anxiety. Pain catastrophizing correlated strongly with psychological distress. Findings highlight the need for integrated, trauma-informed pain management addressing both biomedical and psychosocial factors in conflict-affected populations. Future research should explore how sociopolitical stressors and catastrophic thinking interact to exacerbate pain experiences in humanitarian crises.

Pain is defined as an "unpleasant sensory and emotional experience associated with actual or potential tissue damage". Chronic pain affects approximately 30% of the global population^{1,2} with prevalence rates ranging from 10–30% among adults in Europe and impacting over 50 million Americans, often referred to as a "silent epidemic"^{3,4}. In the Middle East, chronic pain affects 19% of the population in Al-Kharj, Saudi Arabia, and 20% in Libya^{5,6}. Across low- to high-income nations, chronic pain remains a persistent public health challenge^{7,8}.

Chronic pain management requires a multimodal approach that integrates pharmaceutical and non-pharmaceutical interventions encompassing psychotherapy, physical rehabilitations and multidisciplinary care models⁹. International guidelines emphasize a holistic approach aiming to reduce pain severity, improve functional outcomes, and address the high burden of comorbid mental health symptoms among those patients¹⁰. However, in a resource-limited environment such as Gaza, chronic pain patients have a fundamental right to access pain medication¹¹, yet Gaza's healthcare system faces significant obstacles in delivering adequate chronic pain services. Issues include limited prioritization of chronic pain within the healthcare system, the absence of specialized pain clinics, fragmented care pathways, and barriers to accessing pharmacological and non-pharmacological treatment¹².

Gaza's population has endured sixteen years of conflict and siege with prolonged socioeconomic hardship, resulting in substantial strain on the healthcare system and limited availability of comprehensive medical services^{13,14}. These harsh conditions lead to a severe shortage of critical medical supplies and reduced access to both pharmacological and non-pharmacological health interventions. In addition, the social stigmas surrounding chronic conditions, including chronic pain, particularly affect vulnerable groups such as women and individuals in financial hardship, which only worsens their suffering^{13,15–17}. Individuals living in these conditions may experience heightened vulnerability to physical and psychological distress, underscoring the importance of examining the clinical and mental health characteristics of patients with chronic pain receiving care in this context.

¹Department of ICU, Al-Shifa Hospital, Gaza, Palestine. ²Al-Quds University, Al-Azhar Branch, Gaza, Palestine. ³Al-Quds University, Gaza, Palestine. ⁴Department of Anesthesia and Intensive Care, Haukeland University Hospital, Bergen, Norway. ⁵Department of Human Science and Education R. Massa, University of Milan-Bicocca, Milan, Italy. ✉email: abdallah.abojlambo@gmail.com

The prolonged conflict and repeated injuries have resulted in a substantial number of wounded people living with long-term physical limitations in Gaza, increasing the demands for ongoing medical and rehabilitative care^{18,19}. Despite this, the available data for describing the clinical characteristics of patients presenting with chronic pain and accompanying psychological symptoms in healthcare settings remains scarce. Some studies highlighted a significant knowledge gap and negative attitudes toward pain management among healthcare staff²⁰. A gap in understanding the demographic, pain-related, and mental health profiles of patients with chronic pain receiving outpatient care. Understanding this gap through descriptive clinical data is an important first step toward informing future research and service planning in this setting.

This research aims to measure sociodemographic variables in a sample of chronic pain syndrome patients, including geographic distribution, education level, monthly income, and comorbidities, as well as pain-related characteristics such as pain anatomy and duration. A key strength of this study is its focus on the mental health aspects of chronic pain syndrome, including depression, anxiety, stress, and pain catastrophizing. Furthermore, the study advocates for the integration of mental health assessments in the care of these patients.

Subject and method

Study participants

This cross-sectional study was conducted among patients with chronic pain syndrome attending the rheumatology, orthopedics, and neurology outpatient clinics at Al-Shifa Hospital, Gaza City, between September 3 and 18, 2022.

The minimum required sample size was calculated using G*Power software based on an F-family test (linear multiple regression) to detect a moderate effect size ($f^2 = 0.15$) in the depression subscale of the DASS-21. The sample size calculation was initially based on planned analytical objectives; however, the current study presents descriptive findings. The power level was set at 0.95 with a significance level of $\alpha < 0.05$, yielding a minimum required sample of 107 participants. Due to the absence of a centralized registry for chronic pain patients in Gaza, a consecutive non-probability sampling method was used. All patients visiting the targeted clinics during the study period who met the eligibility criteria were invited to participate. These clinics—rheumatology, neurology, and diabetic foot—are the primary points of care for individuals suffering from chronic pain in the region.

A total of 318 patients were approached for participation, out of which 299 agreed to take part and completed the questionnaires, resulting in a response rate of 93.7%. Nineteen patients declined, citing personal or time-related constraints. Following the application of predefined exclusion criteria, 27 participants were excluded (15 with a known history of psychiatric disorders, 9 with cancer-related pain, and 3 with gastrointestinal causes of pain), resulting in a final analytic sample of 272 patients.

Inclusion criteria Participants eligible for inclusion in the study were aged 18 years or older, had been diagnosed with chronic pain, defined as pain persisting longer than three months, and were able to understand and communicate in Arabic.

Exclusion criteria Individuals who were excluded from the study if they were under 18 years or had a diagnosis of cancer or cancer-related pain, gastrointestinal pain, or a primary psychiatric disorder such as major depressive disorder, anxiety disorders, bipolar disorder, or psychosis—whether documented in medical records or self-reported by the patient. Additional exclusion criteria included a diagnosis of autoimmune or systemic inflammatory diseases such as rheumatoid arthritis, systemic lupus erythematosus, or polymyositis to isolate pain mechanisms primarily related to non-inflammatory etiologies as well as cases where disc herniation was identified as the primary cause of pain, to minimize neuropathic/radicular pain mechanisms that may differentially affect pain perception and treatment responses compared to generalized musculoskeletal pain. Furthermore, individuals with cognitive impairments or dementia that interfered with their ability to participate in the survey were also excluded from the study.

Data collection

Trained research assistants conducted structured face-to-face interviews using validated Arabic versions of standardized instruments. Data collected included sociodemographic variables, pain-related characteristics (duration, intensity, site), medication use (prescription analgesics, NSAIDs, opioids, or self-medication), and mental health indicators (depression, anxiety, stress, and pain catastrophizing).

Participants were asked how they typically responded to chronic pain. Of the participants, 45.95% reported using painkillers as their primary coping mechanism, with non-steroidal anti-inflammatory drugs (NSAIDs) being the most commonly used category. Mental health history was screened through both self-report and medical records. Any participant with ongoing psychiatric treatment or a known past psychiatric diagnosis was excluded.

Ethical approval and consent

This study was approved by the Palestinian Ministry of Health Research Ethics Committee (Approval No. 688733). Written informed consent was obtained from all participants prior to enrollment. Participants were provided with detailed information about the study's purpose, procedures, risks, and benefits, and were assured of confidentiality and voluntary participation. All methods were performed in accordance with the relevant guidelines and regulations, including the Declaration of Helsinki.

Instruments and study variables

Family income

Participant family income was categorized into five groups: (1) less than 1974 New Israeli Shekels (NIS), (2) 1974–2470 NIS, (3) 2471–2967 NIS, (4) 2968–3464 NIS, and (5) greater than 3465 NIS. This classification was used to capture the economic status of the households.

Working status

Participants self-reported their employment status, classified as: (1) full-time/part-time employment, (2) housewife duties, (3) unemployment due to pain, and (4) retired. This aimed to examine the impact of pain on participants' ability to work or engage in income-generating activities, and how it might influence their daily life and socioeconomic status.

Pain

Pain intensity was measured using the Numerical Rating Scale (NRS) 0–10 scale, with a higher score indicating greater pain²¹. The Arabic NRS (NAPRS) was employed to ensure cultural and linguistic appropriateness in the assessment²². Pain severity was categorized as 1–4 for mild pain, 5–6 for moderate pain, and 7–10 for severe pain²³.

Depression, anxiety, and stress-21 (DASS-21)

The depression, anxiety, and stress scale (DASS-21) was used to assess the psychological distress^{24,25}. The cut-off values were: ≥ 10 for depression, ≥ 8 for anxiety, and ≥ 15 for stress²⁴. The Cronbach's alpha value was 0.877, and individual subscales ranged from 0.743 to 0.762, further confirming the scale's reliability and consistency in this context²⁶.

Pain catastrophizing (PCS)

The pain catastrophizing scale (PCS) 13-item tool that is used to measure catastrophic thoughts and feelings related to pain. A total score ranging from 0 to 52. A score > 30 is considered clinically significant²⁷. The PCS yields three subscales: rumination, magnification, and helplessness²⁸. In this study, the Arabic version of the Pain Catastrophizing Scale (PCS-A) was used, demonstrating excellent internal consistency with a Cronbach's alpha of 0.86. The reliability for the individual subscales was also good: rumination ($\alpha = 0.646$), magnification ($\alpha = 0.709$), and helplessness ($\alpha = 0.783$)²⁶.

Statistical analysis

Descriptive statistical tests were used to summarize the data, with the mean and standard deviation (SD) employed for parametric variables, while the medians and interquartile range (IQR) were used for non-parametric variables. Sociodemographic characteristics, pain intensity, depression, anxiety, stress, and pain catastrophizing scores were described. Statistical analyses were performed using SPSS (VERSION 26), with a significant threshold set at 0.05 for all tests.

Results

Baseline characteristics of study participants

Table 1 summarizes the sociodemographic characteristics of participants ($n=272$). Most patients were female (57.7%) with a median age of 43 (IQR 34–55) for the whole sample. The majority of participants (82.7%) resided in Gaza City, and 78.8% were married. About 50% of the participants were engaged in home duties. Approximately 34.9% had completed high school, and 94.1% had an income below the poverty line. Chronic pain was found to contribute to unemployment in 29.4% of the cases.

Comorbidities

As shown in Fig. 1, among the participants, 27.2% had at least one comorbidity, with a median of 2 comorbidities (IQR 1–3). Over 51.8% of participants had osteoarthritis, and 54% had a spinal cord disorder. Around 20% of participants had diabetes-related chronic pain, with diabetic foot and neuropathy being the main causes. Further details on the nature and prevalence of comorbidities among our sample are illustrated in Fig. 1.

Pain-related characteristics

Table 2 summarizes the pain-related characteristics of participants. The median pain duration among the participants was four years (IQR 1.5–12 years). The majority of participants (82.4%) reported severe pain (7–10) over the last week. The most common site of pain was the lower limbs, particularly the knee joint (27.2%). Generalized pain was reported by 26.1% of the participants. Additionally, 45.95% of the patients used painkillers as their primary coping mechanism for pain.

Mental health measurements

As shown in Fig. 2, a significant proportion of participants exhibited symptoms of depression (83.1%), anxiety (83.8%), and stress (88.6%). Additionally, 67.6% (186 participants) scored 30 or higher on the pain Catastrophizing Scale (PCS), indicating clinically relevant pain catastrophizing. Participants' median scores and interquartile ranges (IQR) were as follows: Numerical Rating Scale (NRS): 10 (8–10), DASS-21: 36 (26–45), and PCS: 39 (29–46). These results, along with their subscales, are illustrated in Fig. 2. In terms of severity, 31.6% of participants demonstrated moderate depression, while 56.3% exhibited severe anxiety, and 45.2% showed severe stress symptom Fig. 3 presents the severity of depression, anxiety, and stress among our sample.

Discussion

This study provides a descriptive profile of patients with chronic pain attending outpatient services at Al-Shifa Hospital in Gaza. Most participants were living below the poverty line, and a substantial proportion reported unemployment, mirroring Gaza's broader economic collapse²⁹. Females represented more than half of the

Characteristics	Number	Percentage
Age (years), median (IQR)	43	(34–55)
Age category		
18–24	29	10.7%
25–34	49	18%
35–44	65	23.9%
45–54	59	21.7%
55–64	41	15.1%
65+	29	10.7%
Gender		
Male	115	42.3%
Female	157	57.7%
Residency place		
Gaza city	225	82.7%
North	28	10.3%
Middle	16	5.9%
South	3	1.1%
Marital status		
Married	214	78.7%
Single	35	12.9%
Widowed	15	5.5%
Divorced	8	2.9%
Educational level		
Not educated	12	4.4%
Preliminary	40	14.7%
Preparatory	76	27.9%
High school	95	34.9%
Bachelor	47	17.3%
Higher	2	0.7%
Work status		
Full/part	42	15.4%
Home duties	136	50%
Unemployed due to chronic pain	80	29.4%
Retired	15	5.1%
Monthly income (ILS)		
< 1974	256	94.1%
1974–2470	7	2.6%
2471–2967	3	1.1%
2968–3464	4	1.5%
>3465	2	0.7%

Table 1. Sociodemographic characteristics of study participants (n = 272). Presents demographic and socioeconomic details, including age, sex, marital status, education level, employment status, and household income.

sample, most participants were married, and fewer than half had completed high school. These findings describe the sociodemographic characteristics of individuals seeking care for chronic pain within this clinical setting.

Our findings expose an alarmingly high prevalence of mental health issues—specifically depression (83.1%), anxiety (83.8%), and stress (88.6%). This is strongly supported by a large body of literature that supports the co-occurrence of psychological distress among individuals with chronic pain^{30–34}. While our findings support the well-established link between chronic pain and mental health globally, the prevalence of depression (83.1%) and anxiety (83.8%) in our sample far exceeds rates reported in a recent global meta-analysis of chronic pain individuals across 50 countries, which reported lower rates for depression and anxiety (39.3% and 40.2%, respectively)³⁵. The high prevalence rates observed in this study may be influenced by contextual, methodological, and sampling factors, and should therefore be interpreted cautiously when compared to international estimates.

Our findings also exceed rates described in several regional studies from the Middle East. For example, depressive symptoms were documented among 71% of patients with chronic pain in Saudi-Arabia, while lower rates (25%) have been observed among the same population from Oman^{36,37}. Similarly, Alamam et al. (2019) found that among patients with chronic lower back pain, there were 33% experienced depression, 47%




Fig. 1.. Prevalence of comorbidities among participants with chronic pain

Pain related characteristics	Number	Percentage
Pain duration (years)		
≤ 10	202	74.3%
11–20	49	18%
21–30	13	4.8%
≥ 31	7	2.6%
Pain anatomy		
Upper shoulders and upper limbs	25	9.2%
Lower limbs	74	26.8%
Lower back, lower spine, sacrum	46	16.9%
Face, mouth, head	11	4%
Lower back and lower limbs	45	16.5%
Generalized pain (≥2 pain sites)	71	26.1%
Pain intensity		
Mild	16	5.9%
Moderate	32	11.8%
Severe	224	82.4%

Table 2.. Pain-related characteristics of study participants. Includes duration, intensity, anatomical distribution of pain, and coping strategies, among Palestinian patients with chronic pain.




Fig. 2.. Median scores of pain intensity, psychological distress (DASS-21), and pain catastrophizing (PCS) among participants.




Fig. 3. Severity levels of depression, anxiety, and stress symptoms based on DASS-21.

anxiety, and 42.6% stress above the normal range. Notably, severe or extreme symptoms were also reported (depression: 13.9%, anxiety: 21.7%, stress: 19.2%)³⁸. Although comparisons should be interpreted cautiously due to differences in populations and assessment methods, these findings suggest a substantial psychological symptom burden among chronic pain patients in this clinical setting.

Such a high prevalence is not unexpected in conflict zones like Gaza, where chronic pain patients face compounded distress due to pain and conflict stressors. The humanitarian crisis could drive high rates of psychological distress across the general population overall and particularly among Palestinians with chronic pain^{39,40}. For instance, a population-based study conducted in Gaza revealed high rates of depression, anxiety, and post-traumatic stress symptoms in the general population^{41–43}. While those studies were not conducted among chronic pain patients, they provide important contextual background for understanding the broader mental health environment in which patients in the current study are living.

A key finding of our study is the high prevalence of pain catastrophizing, with 67.6% scoring ≥ 30 on PCS, a pattern linked to greater pain severity, psychological distress, and disability^{44,45}. Cognitive-behavioral theory conceptualizes catastrophizing as a maladaptive coping mechanism, especially in a chronically threatening environment^{46,47}. In a conflict setting such as Gaza, constant uncertainty, limited access to healthcare intensify helplessness and pain rumination⁴⁸. Here, chronic trauma exposure, poverty, which worsened by severe medical supply shortages, provide important contextual background for understanding the lived experiences of patients in this sample⁴⁹. However, further analytical studies are needed to examine the specific mechanisms linking pain catastrophizing, psychological distress, and contextual stressors in this population.

These findings can be interpreted within the biopsychosocial model of chronic pain, which emphasizes the dynamic interaction between biological, psychological, and social factors in shaping pain experiences. In this context, psychological distress and pain catastrophizing are not merely comorbidities but integral components that influence pain perception and disability. Furthermore, cognitive-behavioral theory suggests that maladaptive thought patterns, such as catastrophizing, may amplify pain perception and emotional distress. In a conflict-affected setting such as Gaza, chronic exposure to trauma, uncertainty, and socioeconomic hardship may intensify these cognitive and emotional responses, thereby exacerbating both pain and psychological burden⁵⁰.

Over time, this adversity transforms pain into existential suffering that reflects systematic trauma^{51–53}. Nearly half of our study participants (45.95%) relied on NSAIDs- highlighting potential gaps in chronic pain management within this clinical setting. This pharmacologic dependence reflects a healthcare system working in a resource-limited environment that is overwhelmed by conflict, prioritizes acute emergencies over chronic care, and lacks pain specialists, rehabilitation service and mental health support⁵⁴. Without multidisciplinary pain management services, patients may become trapped in medication dependence, risking side effects, inadequate pain relief, and diminished mental well-being⁵⁰. Given these findings, there urgent need for trauma-informed, multidisciplinary pain management in Gaza. Interventions must extend beyond pharmacological treatment to address both physical and psychological suffering. Integrating cognitive-behavioral therapy (CBT), stress management techniques, and community-based psychosocial support is crucial in conflict-affected settings^{50,55}. These interventions must be culturally sensitive and specifically tailored to the lived experiences of Palestinians under occupation. The current humanitarian crisis, exacerbated by the ongoing violence, makes such integrated care models more critical than ever.

This study has several limitations. The cross-sectional design limits interpretation to descriptive findings and does not allow causal or directional inferences. Consequently, non-probability sampling from a single hospital may introduce selection bias and restrict generalizability. Data were based largely on self-reported measures, including psychological scales and medication use, which may be subject to recall and response biases. Additionally, the absence of trauma-specific measures such as post-traumatic stress disorder (PTSD) limits a comprehensive understanding of psychological distress in this population, particularly in a conflict-affected setting. Future studies using longitudinal designs and broader clinical and psychosocial assessments are warranted.

Conclusion

This study provides important insights into the clinical and psychological profile of patients living with chronic pain in Gaza, a highly vulnerable and underrepresented population. The findings reveal an exceptionally high

burden of psychological distress and pain catastrophizing, emphasizing the need for integrated, trauma-informed approaches to pain management in conflict-affected settings. Importantly, this study contributes to the literature by highlighting how chronic pain in humanitarian contexts is deeply intertwined with psychological and social stressors, supporting a biopsychosocial understanding of pain beyond purely biomedical models. Future research should extend beyond descriptive designs to explore causal relationships between trauma exposure, psychological distress, and pain outcomes. Longitudinal studies, inclusion of trauma-specific measures such as post-traumatic stress disorder (PTSD), and evaluation of culturally adapted multidisciplinary interventions are essential to improve care for patients in similar settings.

Data availability

The datasets generated and/or analyzed during the current study are not publicly available due to privacy and ethical restrictions but are available from the corresponding author on reasonable request.

Received: 20 July 2025; Accepted: 11 May 2026

Published online: 22 May 2026

References

- Merskey, H. Pain terms: a current list with definitions and notes on usage. *Pain* **3**, 217–221 (1986).
- Elzahaf, R. A., Tashani, O. A., Unsworth, B. A. & Johnson, M. I. The prevalence of chronic pain with an analysis of countries with a Human Development Index less than 0.9: a systematic review without meta-analysis. *Curr. Med. Res. Opin.* **28**, 1221–1229 (2012).
- Pain, D.C.N.-M. Prevalence, impact on daily life, and treatment. *Eur. J. Pain* **10**(4), 287–333 (2006) (ON THE STATE OF PUBLIC HEALTH 1, 68 (1997)).
- Baker, M. *et al.* Pain Proposal: Improving the current and future management of chronic pain. *A European Consensus Report* (2010).
- El-Metwally, A. *et al.* The prevalence of chronic pain and its associated factors among Saudi Al-Kharj population; a cross sectional study. *BMC Musculoskelet. Disord.* **20**, 1–9 (2019).
- Elzahaf, R. A., Johnson, M. I. & Tashani, O. A. The epidemiology of chronic pain in Libya: a cross-sectional telephone survey. *BMC Public Health* **16**, 776 (2016).
- Jackson, T. *et al.* A systematic review and meta-analysis of the global burden of chronic pain without clear etiology in low-and middle-income countries: trends in heterogeneous data and a proposal for new assessment methods. *Anesth. Analg.* **123**, 739–748 (2016).
- Dahlhamer, J. Prevalence of chronic pain and high-impact chronic pain among adults—United States, 2016. *MMWR. Morbidity and Mortality Weekly Report* **67** (2018).
- Treede, R.-D. *et al.* Chronic pain as a symptom or a disease: the IASP Classification of Chronic Pain for the International Classification of Diseases (ICD-11). *Pain* **160**, 19–27 (2019).
- Guideline NG193, N. Chronic pain (primary and secondary) in over 16s: assessment of all chronic pain and management of chronic primary pain. *Methods* **10** (2021).
- Lohman, D., Schleifer, R. & Amon, J. J. Access to pain treatment as a human right. *BMC Med.* **8**, 1–9 (2010).
- World Health Assembly, 63 Health conditions in the occupied Palestinian territory, including east Jerusalem, and in the occupied Syrian Golan. World Health Organization. <https://iris.who.int/handle/10665/3074> (2010).
- Monitor, E. M. H. R. *A Generation Under Blockade: Consequences of Israel's 17-year Blockade of the Gaza Strip* (Geneva or London, 2023).
- Al Bakri, D. *et al.* The war on Gaza and its impact on public health: challenges and pathways to recovery. *Front. Public Health* **13**, 1664850 (2025).
- Bates, K. *et al.* Women's health in the occupied Palestinian territories: contextual influences on subjective and objective health measures. *PLoS ONE* **12**, e0186610 (2017).
- UNCTAD *Economic crisis Worsens in Occupied Palestinian Territory Amid ongoing Gaza Conflict* (2024).
- UNCTAD Press Release No. UNCTAD/PRESS/PR/2023/025. Geneva: UNCTAD; Available from: <https://unctad.org/press-materi-al/prior-current-crisis-decades-long-blockade-hollowed-gazas-economy-leaving-80> (2023).
- Affairs, U. N. Humanitarian Situation Update #245 | Gaza Strip O. H. (2024).
- West, L. J. S. Palestinian Central Bureau of Statistics (PCBS) Press release on the occasion of the International Day of Persons with Disabilities, 03/12/2023 (2016).
- Reyala, M. Nursing students' knowledge and attitude toward pain management in Gaza Strip, Palestine. *J. Clin. Anesth. Pain Manag.* **4**, 101–106 (2020).
- Jt, F. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain* **94**, 149–158 (2001).
- Alghadir, A. H., Anwer, S. & Iqbal, Z. A. The psychometric properties of an Arabic numeric pain rating scale for measuring osteoarthritis knee pain. *Disabil. Rehabil.* **38**, 2392–2397 (2016).
- Boonstra A. M., Stewart, Roy E; Köke, Albère JA; Oosterwijk, René FA; Swaan, Jeannette L; Schreurs, Karlein MG; Schiphorst Preuper, Henrica R. Cut-off points for mild, moderate, and severe pain on the numeric rating scale for pain in patients with chronic Musculoskeletal pain (2016).
- Lovibond, S. H. *Manual for the Depression Anxiety Stress Scales* (Sydney Psychology Foundation, 1995).
- Moussa, M. T., Lovibond, P., Laube, R. & Megahead, H. A. Psychometric properties of an Arabic version of the depression anxiety stress scales (DASS). *Res. Soc. Work Pract.* **27**, 375–386 (2017).
- Taber, K. S. The use of Cronbach's alpha when developing and reporting research instruments in science education. *Res. Sci. Educ.* **48**, 1273–1296 (2018).
- Sullivan, M. J., Bishop, S. R. & Pivik, J. The pain catastrophizing scale: development and validation. *Psychol. Assess.* **7**, 524 (1995).
- Terkawi, A. S. *et al.* Development and validation of Arabic version of the pain catastrophizing scale. *Saudi J. Anaesth.* **11**, S63–S70 (2017).
- (UNCTAD), U. N. C. o. T. a. D. Report on UNCTAD assistance to the Palestinian people: developments in the economy of the Occupied Palestinian Territory. Report No. TD/B/71/3 (as noted in the PDF) (Geneva, UNCTAD Headquarters, 2024).
- Dominick, C. H., Blyth, F. M. & Nicholas, M. K. Unpacking the burden: understanding the relationships between chronic pain and comorbidity in the general population. *Pain* **153**, 293–304 (2012).
- Brown, G. K. A causal analysis of chronic pain and depression. *J. Abnorm. Psychol.* **99**, 127 (1990).
- Fishbain, D. A., Cutler, R., Rosomoff, H. L. & Rosomoff, R. S. Chronic pain-associated depression: Antecedent or consequence of chronic pain? A review. *Clin. J. Pain.* **13**, 116–137 (1997).
- IsHak, W. W. *et al.* Pain and depression: a systematic review. *Harv. Rev. Psychiatry* **26**, 352–363 (2018).

34. Gureje, O. et al. The relation between multiple pains and mental disorders: results from the World Mental Health Surveys. *PAIN* **135**, 82–91 (2008).
35. Aaron, R. V. et al. Prevalence of depression and anxiety among adults with chronic pain: a systematic review and meta-analysis. *JAMA Netw. Open* **8**, e250268–e250268 (2025).
36. Al Shukaili, M. et al. Exploring factors associated with depressive symptoms among patients with chronic pain: a cross-sectional multicenter study. *J. Nervous Mental Dis.* **210**, 45–53 (2022).
37. Al-Maharbi, S. et al. Prevalence of depression and its association with sociodemographic factors in patients with chronic pain: a cross-sectional study in a tertiary care hospital in Saudi Arabia. *Saudi J. Anaesth.* **12**, 419–425 (2018).
38. Alamam, D. M., Moloney, N., Leaver, A., Alsobayel, H. I. & Mackey, M. G. Pain intensity and fear avoidance explain disability related to chronic low back pain in a Saudi Arabian population. *Spine* **44**, E889–E898 (2019).
39. Lim, G. Y. et al. Prevalence of depression in the community from 30 countries between 1994 and 2014. *Sci. Rep.* **8**, 2861 (2018).
40. Lim, I. C. Z. et al. Prevalence of depression, anxiety and post-traumatic stress in war-and conflict-afflicted areas: a meta-analysis. *Front. Psychiatry* **13**, 978703 (2022).
41. Bank, W. Mental health in the West Bank and Gaza (Washington, DC, 2022).
42. Marie, M., SaadAdeen, S. & Battat, M. Anxiety disorders and PTSD in Palestine: a literature review. *BMC Psychiatry* **20**, 509 (2020).
43. Boukari, Y. et al. Gaza, armed conflict and child health. *BMJ Paediatr. Open* **8**, e002407 (2024).
44. Sullivan, M. J. et al. Theoretical perspectives on the relation between catastrophizing and pain. *Clin. J. Pain* **17**, 52–64 (2001).
45. Turk, D. C. & Okifuji, A. Psychological factors in chronic pain: evolution and revolution. *J. Consult. Clin. Psychol.* **70**, 678 (2002).
46. Keefe, F. J., Rumble, M. E., Scipio, C. D., Giordano, L. A. & Perri, L. M. Psychological aspects of persistent pain: current state of the science. *J. Pain* **5**, 195–211 (2004).
47. Turk, D. C. & Gatchel, R. J. *Psychological Approaches to Pain Management: A Practitioner's Handbook* (Guilford Publications, 2018).
48. Zheng, Z. et al. Safety needs mediate stressful events induced mental disorders. *Neural Plast.* **2016**, 8058093 (2016).
49. Hamdan, A. & Mosleh, R. How does the general population approach their pain? A cross-sectional study in Palestine. *SAGE Open Med.* **12**, 20503121231223442 (2024).
50. Gatchel, R. J., Peng, Y. B., Peters, M. L., Fuchs, P. N. & Turk, D. C. The biopsychosocial approach to chronic pain: Scientific advances and future directions. *Psychol. Bull.* **133**, 581 (2007).
51. Zughbur, M. R. et al. Prevalence and correlates of anxiety, depression, and symptoms of trauma among Palestinian adults in Gaza after a year of war: A cross-sectional study. *Confl. Heal.* **19**, 43. <https://doi.org/10.1186/s13031-025-00681-1> (2025).
52. Morina, N., Stam, K., Pollet, T. V. & Priebe, S. Prevalence of depression and posttraumatic stress disorder in adult civilian survivors of war who stay in war-afflicted regions. A systematic review and meta-analysis of epidemiological studies. *J. Affect. Disord.* **239**, 328–338 (2018).
53. Marie, M., Hannigan, B. & Jones, A. Challenges for nurses who work in community mental health centres in the West Bank, Palestine. *Int. J. Ment. Heal. Syst.* **11**, 3 (2017).
54. Abuelaish, I. & Musani, A. Reviving and rebuilding the health system in Gaza. *East. Mediterr. Health J.* **31**, 56–58 (2025).
55. Policy, B. o. H. S., Research, C. o. A. P. & Care. *Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research* (National Academies Press, 2011).

Acknowledgements

Mohammed Imad, Mohammed Elsheikh Khalil, Mohammed alkhaldi, Siham Al-Shawamreh, Islam Ayyash colleagues who helped with the data access and collation.

Author contributions

AA conceptualized the study. AA, YA, and MA prepared and refined the data for analysis, executed a statistical evaluation, and designed graphical representations, and contributed to writing the manuscript. HL and GV evaluated the manuscript, provided key insights on the outcome data, and assisted with data interpretation. All authors had the full access to the study data and take responsibility for its submission.

Declarations

Competing interests

The authors declare no competing interests.

Ethical approval and consent to participate

The Palestinian Ministry of Health Research Ethics Committee granted ethical permission with the Serial Number 688733.

Consent for publication

Not applicable, as this manuscript does not contain any individual person's data in any form (including individual details, images, or videos).

Additional information

Correspondence and requests for materials should be addressed to A.A.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

© The Author(s) 2026