

Search for double- β decay of ^{130}Te to the first 0^+ excited state of ^{130}Xe with the CUORICINO experiment bolometer array

E. Andreotti,^{1,2,*} C. Arnaboldi,³ F. T. Avignone III,⁴ M. Balata,⁵ I. Bandac,⁴ M. Barucci,^{6,7} J. W. Beeman,⁸ F. Bellini,^{9,10} C. Brofferio,^{2,3} A. Bryant,^{11,12} C. Bucci,⁵ L. Canonica,^{13,14} S. Capelli,^{2,3} L. Carbone,² M. Carrettoni,^{2,3} M. Clemenza,^{2,3} O. Cremonesi,² R. J. Creswick,⁴ S. Di Domizio,^{13,14} M. J. Dolinski,^{12,15} L. Ejzak,¹⁶ R. Faccini,^{9,10} H. A. Farach,⁴ E. Ferri,^{2,3} E. Fiorini,^{2,3,†} L. Foggetta,^{1,2,‡} A. Giachero,² L. Gironi,^{2,3} A. Giuliani,^{1,2,§} P. Gorla,^{5,||} E. Guardincerri,^{5,11,14} T. D. Gutierrez,¹⁷ E. E. Haller,^{8,18} K. Kazkaz,¹⁵ L. Kogler,^{11,12} S. Kraft,^{2,3} C. Maiano,^{2,3} C. Martinez,^{4,¶} M. Martinez,^{2,19,**} R. H. Maruyama,¹⁶ S. Newman,^{4,5} S. Nisi,⁵ C. Nones,^{1,2,††} E. B. Norman,^{15,20} A. Nucciotti,^{2,3} F. Orio,^{9,10} M. Pallavicini,^{13,14} V. Palmieri,²¹ L. Pattavina,^{2,3} M. Pavan,^{2,3} M. Pedretti,¹⁵ G. Pessina,² S. Pirro,² E. Previtali,² L. Risegari,^{6,7} C. Rosenfeld,⁴ C. Rusconi,^{1,2} C. Salvioni,^{1,2} S. Sangiorgio,^{16,‡‡} D. Schaeffer,^{2,3} N. D. Scielzo,¹⁵ M. Sisti,^{2,3} A. R. Smith,²² C. Tomei,¹⁰ G. Ventura,^{6,7} and M. Vignati^{9,10}

¹*Dipartimento di Fisica e Matematica, Università dell'Insubria, Como I-22100, Italy*

²*INFN - Sezione di Milano Bicocca, Milano I-20126, Italy*

³*Dipartimento di Fisica, Università di Milano-Bicocca, Milano I-20126, Italy*

⁴*Department of Physics and Astronomy, University of South Carolina, Columbia, South Carolina 29208, USA*

⁵*INFN - Laboratori Nazionali del Gran Sasso, Assergi (L'Aquila) I-67010, Italy*

⁶*Dipartimento di Fisica, Università di Firenze, Firenze I-50125, Italy*

⁷*INFN - Sezione di Firenze, Firenze I-50125, Italy*

⁸*Materials Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

⁹*Dipartimento di Fisica, Sapienza Università di Roma, Roma I-00185, Italy*

¹⁰*INFN - Sezione di Roma, Roma I-00185, Italy*

¹¹*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

¹²*Department of Physics, University of California, Berkeley, California 94720, USA*

¹³*Dipartimento di Fisica, Università di Genova, Genova I-16146, Italy*

¹⁴*INFN - Sezione di Genova, Genova I-16146, Italy*

¹⁵*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*

¹⁶*Department of Physics, University of Wisconsin, Madison, Wisconsin 53706, USA*

¹⁷*Physics Department, California Polytechnic State University, San Luis Obispo, California 93407, USA*

¹⁸*Department of Materials Science and Engineering, University of California, Berkeley, California 94720, USA*

¹⁹*Laboratorio de Fisica Nuclear y Astroparticulas, Universidad de Zaragoza, Zaragoza 50009, Spain*

²⁰*Department of Nuclear Engineering, University of California, Berkeley, California 94720, USA*

²¹*INFN - Laboratori Nazionali di Legnaro, Legnaro (Padova) I-35020, Italy*

²²*EH&S Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

(Received 23 August 2011; revised manuscript received 30 January 2012; published 16 April 2012)

The CUORICINO experiment was an array of 62 TeO₂ single-crystal bolometers with a total ^{130}Te mass of 11.3 kg. The experiment finished in 2008 after more than 3 yr of active operating time. Searches for both 0ν and 2ν double- β decay to the first excited 0^+ state in ^{130}Xe were performed by studying different coincidence scenarios. The analysis was based on data representing a total exposure of $N(^{130}\text{Te}) \cdot t = 9.5 \times 10^{25}$ yr. No evidence for a signal was found. The resulting lower limits on the half-lives are $T_{\frac{1}{2}}^{2\nu}(^{130}\text{Te} \rightarrow ^{130}\text{Xe}^*) > 1.3 \times 10^{23}$ yr (90% C.L.), and $T_{\frac{1}{2}}^{0\nu}(^{130}\text{Te} \rightarrow ^{130}\text{Xe}^*) > 9.4 \times 10^{23}$ yr (90% CL).

DOI: [10.1103/PhysRevC.85.045503](https://doi.org/10.1103/PhysRevC.85.045503)

PACS number(s): 23.40.Hc, 23.40.Bw, 21.10.Tg, 27.60.+j

I. INTRODUCTION

Two-neutrino double-beta ($2\nu\beta\beta$) decay and neutrinoless double-beta ($0\nu\beta\beta$) decay have been known for over 70 years now [1,2] (a recent review can be found in Ref. [3]). While experimental evidence for $2\nu\beta\beta$ -decay has been found there is still no observation for the $0\nu\beta\beta$ decay, however several

*Present address: Joint Research Center, Institute for Reference Materials and Measurement, 2440 Geel, Belgium.

†ettore.fiorini@mib.infn.it

‡Present address: Laboratoire de l'Accélérateur Linéaire, Centre Scientifique d'Orsay, 91898 Orsay, France.

§Present address: Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, 91405 Orsay Campus, France.

||Present address: INFN - Sezione di Roma Tor Vergata, Roma I-00133, Italy.

¶Present address: Queen's University, Kingston, ON K7L 3N6, Canada.

**Present address: Institut d'Astrophysique Spatial, 91045 Orsay, France.

††Present address: CEA / Saclay, 91191 Gif-sur-Yvette, France.

‡‡Present address: Lawrence Livermore National Laboratory, Livermore, CA 94550, USA.

TABLE I. Theoretical evaluations (for $m_{\beta\beta} = 1$ eV) and experimental best limits (90% CL) for the half-life of ^{130}Te $0\nu\beta\beta$ and $2\nu\beta\beta$ decay.

Decay	Transition	Theoretical (yr)	Experimental (yr)
0ν	$0^+ \rightarrow 0_1^+$	7.5×10^{25} [17,18]	$> 3.1 \times 10^{22}$ [19]
	$0^+ \rightarrow 0^+$	$(1.6-15) \times 10^{23}$ [20]	$> 2.8 \times 10^{24}$ [21]
2ν	$0^+ \rightarrow 0_1^+$	$(5.1-14) \times 10^{22}$ [15,16] ^a	$> 2.3 \times 10^{21}$ [16]
	$0^+ \rightarrow 0^+$	$(1.7-70) \times 10^{19}$ [20]	7.0×10^{20} [22]

^aCorrected values for [15] (discussion in text).

while the high-energy γ (1257.41 keV) escapes. Scenario 3 is the opposite of scenario 2: the high-energy γ (1257.41 keV) is trapped in the original crystal with the β s, while the low-energy one (536.09 keV) escapes. The signatures and the corresponding efficiencies are reported in Table II. A further explanation of the calculation of the efficiencies can be found in Sec. III.

The first-level analysis of the CUORICINO data is common to all physics processes to be studied and is described in detail in Ref. [21]. It starts from raw events and ends with a set of energy-calibrated hits associated with a time, a crystal, and other ancillary information, such as pulse shape parameters. In this phase of the analysis, a channel- and time-dependent energy threshold is applied to the data, based on the performance of each bolometer.

For the processes studied in this paper, the analysis consists of defining signatures according to the three scenarios reported in Table II, using them to select events from the CUORICINO data and evaluating the corresponding efficiencies from GEANT4-based Monte Carlo simulations [23].

Event selection criteria can be grouped into three categories: global, event based, and coincidence based. Global and event-based cuts are not specific to this analysis, and here we only outline them briefly (refer to [21] for details). Defined *a priori*, global cuts are used to discard time windows in which one or more detectors performed poorly. This could happen because of external noise or cryogenic instabilities, which in turn result in a bad energy resolution. Event-based cuts allow the exclusion of nonphysical pulses (electronic spikes or cryogenic-induced pulses) and physical pulses for which the




FIG. 3. (Color online) Possible capture scenarios. The blue (dashed) lines represent the 1257.41 keV γ , while the red lines represent the 536.09 keV one. For each scenario, the available energy for the emitted β s is 734.0 keV.

energy is not estimated correctly (pile-up or excessive noise superimposed on the pulse).

Coincidence-based cuts rely on the properties of a group of events that occurred within a fixed time window. Events can be selected based on the number of involved crystals, the spatial distance among them, the sum energy or the energy of the single hits. In this paper, a 100 ms time window was used to define coincident events. Physical coincidences induced by ^{130}Xe de-excitation occur on much shorter time scales, but such a large time window must be chosen to account for the slow response of the bolometers.

The coincidence-based event selection criteria were decided based on the scenarios described at the beginning of this section. Because the two electrons emitted in the 2ν decay have a continuous spectrum, a wide energy window must be chosen

TABLE II. Signatures and efficiencies for the three scenarios for 0ν and 2ν decay. We denote with the + sign the coincidence of energies released in different crystals. Efficiencies labeled as MC were computed based on Monte Carlo simulations. Instrumental efficiencies were computed based on CUORICINO data. Total efficiencies are given by the product of MC and instrumental efficiencies, times a factor of 0.86 to account for the branching ratio of the considered decay scheme (see Fig. 2).

Decay mode	Scenario	Signature (energies in keV)	Efficiency		
			MC	Instrumental	Total
0ν	1	$734 (\beta) + 536 (\gamma) + 1257 (\gamma)$	$(0.60 \pm 0.02)\%$	$(86 \pm 2)\%$	$(0.44 \pm 0.02)\%$
	2	$1257 (\gamma) + 1270 (\beta + \gamma)$	$(2.29 \pm 0.04)\%$	$(90 \pm 1)\%$	$(1.77 \pm 0.04)\%$
	3	$536 (\gamma) + 1991 (\beta + \gamma)$	$(1.41 \pm 0.03)\%$	$(90 \pm 1)\%$	$(1.09 \pm 0.03)\%$
2ν	1	$(0-734) (\beta) + 536 (\gamma) + 1257 (\gamma)$	$(0.53 \pm 0.02)\%$	$(86 \pm 2)\%$	$(0.39 \pm 0.02)\%$
	2	$(536-1270) (\beta + \gamma) + 1257 (\gamma)$	$(3.04 \pm 0.04)\%$	$(90 \pm 1)\%$	$(2.35 \pm 0.04)\%$
	3	$(1257-1991) (\beta + \gamma) + 536 (\gamma)$	$(1.28 \pm 0.03)\%$	$(90 \pm 1)\%$	$(0.99 \pm 0.03)\%$




FIG. 4. (Color online) CUORICINO energy spectra after the event selection cuts applied for the 0ν (left) and 2ν (right) analyses. For the 0ν decay, the signal was expected at 1257.41 keV (a), 1270 keV (b), and 1991 keV (c) for scenarios 1, 2, and 3, respectively. For the 2ν decay, the signal was expected at 1257.41 keV for scenarios 1 (d) and 2 (e) and at 536.09 keV for scenario 3 (f).

for one of the crystals. This has the effect of introducing a much bigger background than is present in the analysis of the 0ν decay mode. As a consequence, besides the criteria reported in Table II, additional restrictions were applied to the events to be included in the 2ν analysis. To reduce random coincidences, a cut was imposed on the distance between the crystals involved in the events, as it was seen from the simulation that there is a low chance for the investigated processes to involve crystals that are far apart from each other. The most relevant background from physical processes is due to γ rays that undergo a Compton interaction in one crystal and are then absorbed in another crystal. While the sum energy of these events is

fixed, the energy released in each crystal has a continuous distribution. To reduce this background, events whose sum energy fell into a window of ± 8 keV around the most intense γ lines (1729.60 keV, 1764.49 keV, 1847.42 keV, 2118.5 keV, 2204.21 keV, and 2447.86 keV from ^{214}Bi , 2505 keV from ^{60}Co , and 2615 keV from ^{208}Tl) were removed.

III. ANALYSIS

As stated in Sec. II, Monte Carlo simulations were used to calculate the efficiencies for the processes studied in this

paper. This was achieved by comparing the number of events passing the coincidence cuts to the total number of simulated events. The relatively low efficiencies reported in Table II arise from the fact that most of the γ s escape the crystals undetected and are absorbed by inert materials surrounding them. Moreover, the signatures sought only consider the case of photons that are completely absorbed in one crystal, thus rejecting events in which at least one photon is absorbed in one crystal after undergoing a Compton interaction in a different one. The computed values reported in the last column of Table II also include inefficiencies due to event-based cuts, channel- and time-dependent energy thresholds, and discarded time windows in which one or more detectors were not performing properly (global cuts). Inefficiencies induced by channel based cuts were evaluated on the CUORICINO data in the same way discussed in Ref. [21]. The effect of global cuts was taken into account by removing the simulated events lying in the time windows that were discarded from the real CUORICINO data, after said time windows were rescaled by the ratio between the total duration of the simulation and the real CUORICINO live time. The same procedure was used to associate energy thresholds to the simulation. Because the effect of global cuts was taken into consideration when determining the efficiencies, the exposure used in this work corresponds to the complete CUORICINO statistics without any subtractions: $N(^{130}\text{Te}) \cdot t = 9.5 \times 10^{25}$ yr.

Figure 4 shows the energy spectra obtained from the CUORICINO data after applying the event selection cuts described in Sec. II. For each scenario, the spectrum was built as follows. Coincidence cuts were applied based on Table II, requiring that the accepted events be in coincidence with events satisfying each component hit of the signature *except for* the hit corresponding to the highest-energy γ . The signal search could then consist of a search in the resultant spectrum for evidence of the highest-energy γ of the signature, which is the component with the lowest background. Moreover, the acceptance width for each cut was enlarged by ± 10 keV with respect to the energies and energy ranges listed in Table II, to account for the finite energy resolution of the detectors ($\sigma \simeq 2$ keV; see discussion below). The energy windows used for the spectra were chosen to be much larger than the detector resolution, but small enough that at most one radioactive background peak was included, and the continuum could be assumed to be flat or linear.

No evidence for a signal was found in any of the energy spectra. For the zero-neutrino decay mode, the background is negligible, and no fit was performed. In this case, a condition of zero signal and zero background was assumed. In contrast, the background is not negligible for the 2ν decay mode, and therefore a Bayesian maximum likelihood fit was performed for the 2ν analyses. The best-fit curves are represented by the blue lines in Fig. 4. Depending on the scenario, different background models were adopted for the 2ν spectra. The continuum was fitted with a constant (scenarios 1 and 2) or linear shape (scenario 3), while the possible additional peaks (1238 keV from ^{214}Bi for scenario 2, 511 keV for scenario 3) were fitted with a Gaussian shape. The free parameters in the fit were as follows: the number of signal counts, the number of events from the flat background and the number of counts

TABLE III. 2ν analysis best-fit values for the number of signal (N_S) and background (N_B) counts. For N_S , both statistical and systematic uncertainties are reported.

Scenario	N_S [counts]	N_B [counts/keV]
1	$1.1 \pm 1.4 \pm 0.29$	0.12 ± 0.03
2	$-0.4 \pm 6.6 \pm 2.6$	6.31 ± 0.41
3	$-3.0 \pm 6.8 \pm 2.8$	6.73 ± 0.33

under the additional background peaks (scenarios 2 and 3). The energy resolution was fixed to $\sigma = 1.8$ keV. It was evaluated on the 511 keV peak and on the two ^{60}Co peaks at 1173 keV and 1332 keV that are visible in the CUORICINO energy spectrum (see Fig. 1), and it was found to be comparable for all three peaks. A summary of the best-fit values for the $2\nu\beta\beta$ searches is reported in Table III. Systematic uncertainties were evaluated by repeating the fitting procedure with different background models, fitting ranges and energy resolutions, and, compared to statistical uncertainties, they were found to be negligible.

IV. RESULTS

For each decay mode and for each of the three scenarios, the posterior probability density function (p.d.f.) for the number of signal counts, $P(N_S)$, was extracted using a Bayesian approach and assuming flat priors in the physical region ($N_S > 0$). For the 0ν decay mode, because there was no evidence of a signal and the background was negligible, a Poisson p.d.f. for zero observed events was assumed for all three scenarios. For the 2ν decay mode, the p.d.f.'s were obtained as a result of the maximum likelihood fits on the spectra shown in Fig. 4. For each decay mode, a global p.d.f. for the decay rate was obtained as the product of the three individual p.d.f.'s, $P_{\text{tot}}(\Gamma) = \prod_i P_i(\Gamma)$. In this formula $P_i(\Gamma) = P_i(N_S) \cdot \varepsilon_i \cdot N(^{130}\text{Te}) \cdot t$, where the index i runs over the three scenarios and ε_i is the corresponding detection efficiency from Table II. Systematic uncertainties were included in the $P_i(\Gamma)$ according to the procedure described in Ref. [21]. This resulted in the following half-life lower limits:

$$\begin{aligned} T_{1/2}(2\nu\beta\beta^*) &> 1.3 \times 10^{23} \text{ yr, } 90\% \text{ CL,} \\ T_{1/2}(0\nu\beta\beta^*) &> 9.4 \times 10^{23} \text{ yr, } 90\% \text{ CL.} \end{aligned}$$

These new limits represent an improvement of almost 2 orders of magnitude, for both the 0ν and 2ν processes, with respect to the results of past experiments. It is worth noting that the new lower limit on the half-life of the 2ν decay mode is close to the upper bound of the theoretical calculation presented in Table I. A more clear picture will be available once CUORE, CUORICINO's successor [24], comes online. This is due to the increase in target mass and improved background reduction that will be achieved in CUORE.

ACKNOWLEDGMENTS

The CUORICINO Collaboration owes many thanks to the directors and staff of the Laboratori Nazionali del Gran Sasso

over the years of the development, construction, and operation of CUORICINO, and to the technical staffs of our laboratories. In particular we would like to thank R. Gaigher, R. Mazza, P. Nuvolone, M. Perego, B. Romualdi, L. Tatananni, and A. Rotilio for continuous and constructive help in various stages of this experiment. We are grateful to our colleagues Y. G. Kolomenski and L. Zanotti for help and fruitful dis-

cussions. The CUORICINO experiment was supported by the Istituto Nazionale di Fisica Nucleare (INFN), the Commission of the European Community under Contract No. HPRN-CT-2002-00322, by the US Department of Energy under Contract No. DE-AC03-76-SF00098, and DOE W-7405-Eng-48, and by the National Science Foundation Grant Nos. PHY-0139294 and PHY-0500337.

-
- [1] M. Goepfert-Mayer, *Phys. Rev.* **48**, 512 (1935).
[2] G. Racah, *Nuovo Cimento* **14**, 322 (1937).
[3] F. T. Avignone, S. R. Elliott, and J. Engel, *Rev. Mod. Phys.* **80**, 481 (2008).
[4] V. Rodin, A. Faessler, F. Simkovic, and P. Vogel, *Nucl. Phys. A* **766**, 107 (2006).
[5] V. Rodin, A. Faessler, F. Simkovic, and P. Vogel, *Nucl. Phys. A* **793**, 213 (2007).
[6] MEDEX Conference (2011).
[7] C. Enss and D. McCammon, *J. Low Temp. Phys.* **151**, 5 (2008).
[8] N. E. Booth, B. Cabrera, and E. Fiorini, *Annu. Rev. Nucl. Part. Sci.* **46**, 471 (1996).
[9] E. E. Haller *et al.*, *Neutron Transmutation Doping in Semiconductor Materials* (Plenum Press, New York, 1984), p. 21.
[10] C. Arnaboldi *et al.*, *Phys. Rev. C* **78**, 035502 (2008).
[11] M. Redshaw, B. J. Mount, E. G. Myers, and F. T. Avignone, *Phys. Rev. Lett.* **102**, 212502 (2009).
[12] N. D. Scielzo *et al.*, *Phys. Rev. C* **80**, 025501 (2009).
[13] S. Rahaman, V.-V. Elomaa, T. Eronen, J. Hakala, A. Jokinen, A. Kankainen, J. Rissanen, J. Suhonen, C. Weber, and J. Äystö, *Phys. Lett. B* **703**, 412 (2011).
[14] B. Singh, *Nucl. Data Sheets* **93**, 33 (2001).
[15] J. Toivanen and J. Suhonen, *Phys. Rev. C* **55**, 2314 (1997).
[16] A. S. Barabash *et al.*, *Eur. Phys. J. A* **11**, 143 (2001).
[17] J. Suhonen, *Phys. Lett. B* **477**, 99 (2000).
[18] J. Suhonen and M. Aunola, *Nucl. Phys. A* **723**, 271 (2003).
[19] C. Arnaboldi *et al.*, *Phys. Lett. B* **557**, 167 (2003).
[20] V. Tretyak and Y. G. Zdesenko, *At. Data Nucl. Data Tables* **80**, 83 (2002).
[21] E. Andreotti *et al.*, *Astropart. Phys.* **34**, 822 (2011).
[22] R. Arnold *et al.* (NEMO-3 Collaboration), *Phys. Rev. Lett.* **107**, 062504 (2011).
[23] C. Bucci *et al.*, *Eur. Phys. J. A* **41**, 155 (2009).
[24] C. Arnaboldi *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **518**, 775 (2004).