

COHERENT SYSTEMS AND MODULAR SUBVARIETIES OF $SU_C(r)$

MICHELE BOLOGNESI

*IRMAR, Université de Rennes 1
263 Av. du Général Leclerc, 35042 Rennes, France
michele.bolognesi@univ-rennes1.fr*

SONIA BRIVIO

*Dipartimento di Matematica, Università di Pavia
Via Ferrata 1, 27100 Pavia, Italy
sonia.brivio@unipv.it*

Received 2 February 2010

Accepted 21 September 2011

Published 24 February 2012

Let C be an algebraic smooth complex curve of genus $g > 1$. The object of this paper is the study of the birational structure of certain moduli spaces of vector bundles and of coherent systems on C and the comparison of different type of notions of stability arising in moduli theory. Notably we show that in certain cases these moduli spaces are birationally equivalent to fibrations over simple projective varieties, whose fibers are GIT quotients $(\mathbb{P}^{r-1})^{r,g} // \text{PGL}(r)$, where r is the rank of the considered vector bundles. This allows us to compare different definitions of (semi-)stability (slope stability, α -stability, GIT stability) for vector bundles, coherent systems and point sets, and derive relations between them. In certain cases of vector bundles of low rank when C has small genus, our construction produces families of classical modular varieties contained in the Coble hypersurfaces.

Keywords: Moduli spaces of coherent systems; vector bundles; algebraic curves; Coble hypersurfaces.

Mathematics Subject Classification 2010: 14H60, 14H10

1. Introduction

Let C be a genus $g > 1$ smooth complex algebraic curve, if $g \neq 2$ we will also assume that C is non-hyperelliptic.

Let $\mathcal{U}_C(r, 0)$ be the moduli space of rank r semi-stable vector bundles on C with degree zero determinant and let us denote as usual by $SU_C(r)$ the moduli subspace given by vector bundles with trivial determinant. These moduli spaces appeared first in the second half of the last century thanks to the foundational works of Narasimhan and Ramanan [27] and Mumford and Newstead [26] and very often

their study has gone along the study of the famous theta-map

$$\theta : \mathcal{SU}_C(r) \dashrightarrow |r\Theta|;$$

$$E \mapsto \Theta_E := \{L \in \text{Pic}^{g-1} \mid h^0(C, E \otimes L) \neq 0\}.$$

While we know quite a good deal about θ for low genera and ranks, as the rank or the genus grows our knowledge decreases dramatically, see Sec. 2 for a complete picture of known results. The question of the rational type of these varieties is even more daunting. When rank and degree are coprime the situation is completely settled [23] but when the degree is zero (or degree and rank are not coprime) the open problems are still quite numerous. It is known that all the spaces $\mathcal{SU}_C(r)$ are unirational but the rationality is clear only for $r = 2$, $g = 2$, when in fact the moduli space is isomorphic to \mathbb{P}^3 , [27]. Some first good ideas about the birational structure for $g = 2$ were developed in [2]. Then the $r = 2$ case was analyzed in any genus by Alzati and Bolognesi in [1] with the help of polynomial maps classifying extensions in the spirit of [6]. In this paper we give a description for the higher rank cases via a new construction.

Our approach consists in studying the birational structure of $\mathcal{U}_C(r, 0)$ and $\mathcal{SU}_C(r)$ via a study of similar moduli spaces of augmented vector bundles, notably the moduli space of coherent systems. By a coherent system on a curve we mean a vector bundle together with a linear subspace of given dimension of its space of global sections. Coherent systems come with a notion of stability that depends on a real parameter α , that leads to a finite family of moduli spaces depending on the value of α . Hence typically one will write $G_\alpha(r, d, k)$ for a moduli space of coherent systems, where α is the real parameter, r is the rank of vector bundles, d their degree and k the prescribed dimension of the space of sections (see Sec. 4 for details). It turns out that for $\alpha > g(r - 1)$ the moduli space $G_\alpha(r, rg, r)$ has a natural structure of a fibration and, moreover it is birational to $\mathcal{U}_C(r, 0)$. The first main theorem of this paper is then the following.

Theorem 1.1. *Let C be a smooth complex curve of genus $g > 1$, non-hyperelliptic if $g > 2$, and let $\alpha > g(r - 1)$. Then $G_\alpha(r, rg, r)$ is birational to a fibration over $C^{(rg)}$ whose fibers are GIT quotients $(\mathbb{P}^{r-1})^{rg} // \text{PGL}(r)$.*

Of course, since if $\alpha > g(r - 1)$, $G_\alpha(r, rg, r)$ is birationally equivalent to $\mathcal{U}_C(r, 0)$, a corresponding result holds also for $\mathcal{U}_C(r, 0)$. Notably, if we consider the moduli subspace $\mathcal{SU}_C(r) \subset \mathcal{U}_C(r, 0)$ of vector bundles with fixed trivial determinant we get the following.

Theorem 1.2. *The moduli space $\mathcal{SU}_C(r)$ is birational to a fibration over $\mathbb{P}^{(r-1)g}$ whose fibers are GIT quotients $(\mathbb{P}^{r-1})^{rg} // \text{PGL}(r)$.*

Theorem 1.2 allows us to give a more precise explicit description of the projective geometry of the fibration of $\mathcal{SU}_C(r)$ in the case $r = 3, g = 2$. In fact $\mathcal{SU}_C(3)$ is

a double covering of \mathbb{P}^8 branched along a hypersurface of degree six \mathcal{C}_6 called the *Coble–Dolgachev sextic* [25]. Our result is the following.

Theorem 1.3. *The Coble–Dolgachev sextic \mathcal{C}_6 is birational to a fibration over \mathbb{P}^4 whose fibers are Igusa quartics. More precisely, \mathcal{C}_6 contains a four-dimensional family of Igusa quartics parametrized by an open subset of \mathbb{P}^4 .*

We recall that an Igusa quartic is a modular quartic hypersurface in \mathbb{P}^4 that is related to some classical GIT quotients (see e.g. [15]) and moduli spaces. Its dual variety is a cubic three-fold called the *Segre cubic*, that is isomorphic to the GIT quotient $(\mathbb{P}^1)^6//\mathrm{PGL}(2)$.

If $r = 2$ and $g = 3$, then $SU_C(2)$ is embedded by θ in \mathbb{P}^7 as a remarkable quartic hypersurface \mathcal{C}_4 called the *Coble quartic* [27]. Our methods also allow us to give a quick proof of the following fact.

Proposition 1.1. *The Coble quartic \mathcal{C}_4 is birational to a fibration over \mathbb{P}^3 whose fibers are Segre cubics. There exists a three-dimensional family of Segre cubics contained in \mathcal{C}_4 .*

We underline that the cases of \mathcal{C}_4 and \mathcal{C}_6 are particularly interesting because one can interpret explicitly the beautiful projective geometry of the Igusa quartic and the Segre cubic in terms of vector bundles on C (see Sec. 6). We hope that these results could help to shed some new light on the question of rationality of $SU_C(r)$ and on the properties of the theta map.

On the other hand, a side result of Theorem 1.1 and of its proof is that we get a bijection between the general vector bundle (or coherent system) and a set of rg points in \mathbb{P}^{r-1} . It then makes perfect sense to compare the GIT stability of a set of points in $(\mathbb{P}^{r-1})^{rg}/\mathrm{PGL}(r)$ with the slope stability of vector bundles and the α -stability of coherent systems. This is discussed in Sec. 6.

A little caution: in some points of the paper it is important to distinguish a vector bundle E from its S-equivalence class $[E]$. We have tried to keep the two distinguished notations when it is necessary, using just the vector bundle one when it has no importance, even if this sometimes may offend the good taste of the reader.

Description of contents. In Sec. 2 we collect a few results on the theta map. In Sec. 3 we outline the relation between theta maps and theta-linear systems by introducing the theta divisor of a vector bundle with integral slope. In Sec. 4 we introduce generically generated coherent systems and their moduli spaces plus some properties of the evaluation and the determinant map for a coherent system of any rank. In Sec. 5 we introduce the *fundamental divisor* of a coherent system. This definition allows us to define the *fundamental map* in Sec. 6, the fibers of this map give us the fibration we look for. Then we prove Theorem 1.1 and discuss briefly the relation between α -stability of coherent systems, slope stability of vector bundles and GIT stability of point sets in the projective space. Finally, in Sec. 7 we restrict our analysis to moduli of vector bundles. Notably we apply our results to

the cases $g = 2, r = 3$ and $g = 3, r = 2$ and construct explicit families of invariant hypersurfaces contained in the moduli spaces.

2. The Theta Map

Let C be a smooth complex algebraic curve of genus $g \geq 2$, we assume that it is non-hyperelliptic if $g > 2$. Let $\text{Pic}^d(C)$ be the Picard variety parametrizing line bundles of degree d on C , $\text{Pic}^0(C)$ will often be denoted by $J(C)$. Let $\Theta \subset \text{Pic}^{g-1}(C)$ be the canonical theta divisor

$$\Theta := \{L \in \text{Pic}^{g-1}(C) \mid h^0(C, L) \neq 0\}.$$

For $r \geq 2$, let $\mathcal{S}U_C(r)$ denote the coarse moduli space of semi-stable vector bundles of rank r and trivial determinant on C . It is a normal, projective variety of dimension $(r^2 - 1)(g - 1)$. It is well known that $\mathcal{S}U_C(r)$ is locally factorial and that $\text{Pic}(\mathcal{S}U_C(r)) = \mathbb{Z} [16]$, generated by a line bundle \mathcal{L} called the *determinant bundle*. On the other hand, for $E \in \mathcal{S}U_C(r)$ we define

$$\Theta_E := \{L \in \text{Pic}^{g-1}(C) \mid h^0(C, E \otimes L) \neq 0\}.$$

This is either a divisor in the linear system $|r\Theta|$ or the whole $\text{Pic}^{g-1}(C)$. For E a general bundle Θ_E is a divisor, *the theta divisor* of E . This means that one can define the rational *theta map* of $\mathcal{S}U_C(r)$:

$$\theta : \mathcal{S}U_C(r) \dashrightarrow |r\Theta| \tag{2.1}$$

sending E to its theta divisor Θ_E . The relation between the theta map and the determinant bundle is given by the following fundamental result.

Theorem 2.1 ([5]). *There is a canonical isomorphism $|r\Theta| \xrightarrow{\sim} |\mathcal{L}|^*$ which identifies θ with the rational map $\varphi_{\mathcal{L}} : \mathcal{S}U_C(r) \dashrightarrow |\mathcal{L}|^*$ associated to the determinant line bundle.*

The cases when θ is a morphism or finite are of course very appealing. Notably, θ is an embedding for $r = 2$, [10, 21, 27] and it is a morphism when $r = 3$ for $g \leq 3$ and for a general curve of genus $g > 3$, [4, 36]. Finally, θ is generically finite for $g = 2$, [4, 11] and we know its degree for $r \leq 4$, [25, 35]. There are also good descriptions of the image of θ for $r = 2, g = 2, 3$ [28, 34], $r = 3, g = 2$ [29, 31], $r = 2, g = 4$ [33]. Moreover, it has recently been shown in [12] that if C is general and $g \gg r$ then θ is generically injective.

3. Vector Bundles and Theta Linear Systems

The notion of theta divisor can be extended to vector bundles with integral slope, in this paper we will consider bundles with slope $g = g(C)$. Let $\mathcal{U}_C(r, rg)$ be the moduli space of semi-stable vector bundles as above. The tensor product defines a

natural map:

$$\begin{aligned} t: SU_C(r) \times \text{Pic}^g(C) &\rightarrow \mathcal{U}_C(r, rg); \\ (E, \mathcal{O}_C(D)) &\mapsto E \otimes \mathcal{O}_C(D), \end{aligned}$$

which is étale, Galois, with Galois group $J(C)[r]$, the group of r -torsion points of the Jacobian of C .

Moreover, if one restricts t to $SU_C(r) \times \mathcal{O}_C(D)$ this yields an isomorphism $t_D: SU_C(r) \rightarrow SU_C(r, \mathcal{O}_C(rD))$, where the latter is the moduli space of rank r semi-stable vector bundles with determinant $\mathcal{O}_C(rD)$.

Definition 3.1. Let $F \in \mathcal{U}_C(r, rg)$, then we define the *theta divisor* of F as follows:

$$\Theta_F := \{L \in \text{Pic}^1(C) \mid h^0(C, F \otimes L^{-1}) \neq 0\}.$$

Let $E \in SU_C(r)$ and $\mathcal{O}_C(D) \in \text{Pic}^g(C)$. If $F = E \otimes \mathcal{O}_C(D)$, then we have that $\Theta_F = \mathcal{O}_C(D) - \Theta_E$, thus Θ_F is a divisor if and only if Θ_E is a divisor. We define

$$\Theta_D := \{L \in \text{Pic}^1(C) \mid h^0(\mathcal{O}_C(D) \otimes L^{-1}) \geq 1\}. \quad (3.1)$$

Then, for any $r \geq 1$, we have a natural isomorphism $\sigma_D: |r\Theta| \rightarrow |r\Theta_D|$ given by the translation $M \mapsto \mathcal{O}_C(D) - M$; moreover, if $\mathcal{O}_C(rD_1) \simeq \mathcal{O}_C(rD_2)$, then $|r\Theta_{D_1}| = |r\Theta_{D_2}|$. So we conclude that if $F \in \mathcal{U}_C(r, rg)$ admits a theta divisor, then $\Theta_F \in |r\Theta_D|$, for any line bundle $\mathcal{O}_C(D) \in \text{Pic}^g(C)$ which is an r th-root of $\det F$. In this way we obtain a family of theta linear systems over the Picard variety $\text{Pic}^{rg}(C)$, as the following shows.

Lemma 3.1. *There exists a projective bundle \mathcal{T} over $\text{Pic}^{rg}(C)$:*

$$p: \mathcal{T} \rightarrow \text{Pic}^{rg}(C),$$

whose fiber over $\mathcal{O}_C(M) \in \text{Pic}^{rg}(C)$ is the linear system $|r\Theta_D|$, where $\mathcal{O}_C(D) \in \text{Pic}^g(C)$ is any r th-root of $\mathcal{O}_C(M)$.

Proof. Remark first that the linear system $|r\Theta_D|$ is well-defined since it does not depend on which r th-root $\mathcal{O}_C(D)$ of $\mathcal{O}_C(M)$ we choose. Let us now consider the following map:

$$\begin{aligned} \delta: \text{Pic}^g(C) \times \text{Pic}^1(C) &\rightarrow \text{Pic}^{g-1}(C); \\ (\mathcal{O}_C(D), L) &\mapsto \mathcal{O}_C(D) \otimes L^{-1}. \end{aligned}$$

For any $\mathcal{O}_C(D) \in \text{Pic}^g(C)$, we have: $\delta^* r\Theta|_{\mathcal{O}_C(D) \times \text{Pic}^1(C)} \simeq \mathcal{O}_{\text{Pic}^1(C)}(r\Theta_D)$. Let $p_1: \text{Pic}^g(C) \times \text{Pic}^1(C) \rightarrow \text{Pic}^g(C)$ be the projection onto the first factor. Consider the sheaf $\mathcal{F} := p_{1*} \mathcal{O}_{\text{Pic}^g(C) \times \text{Pic}^1(C)}(\delta^*(r\Theta))$. It is locally free and its fiber at $\mathcal{O}_C(D) \in \text{Pic}^g(C)$ is canonically identified with the following vector space

$$H^0(\text{Pic}^1(C), \mathcal{O}_{\text{Pic}^1(C)}(r\Theta_D)).$$

Let $\tilde{\mathcal{T}}$ be the projective bundle $\mathbb{P}(\mathcal{F})$ on $\text{Pic}^g(C)$.

Moreover the vector bundle \mathcal{F} is $J[r](C)$ -equivariant, hence by easy descent theory (see [37, Theorem 4.46]) it passes to the quotient by $J[r](C)$, i.e. the image of the cover $\rho: \text{Pic}^g(C) \rightarrow \text{Pic}^{rg}(C)$ given by taking the r th power of each $L \in \text{Pic}^g(C)$. The projectivization of the obtained bundle is the projective bundle \mathcal{T} we are looking for. We denote by $p: \mathcal{T} \rightarrow \text{Pic}^{rg}(C)$ the natural projection on the base of the projective bundle. \square

The previous arguments allow us to define the rational *theta map* of $\mathcal{U}_C(r, rg)$:

$$\theta_{r,g}: \mathcal{U}_C(r, rg) \dashrightarrow \mathcal{T}; \tag{3.2}$$

$$F \mapsto \Theta_F. \tag{3.3}$$

Finally, let us denote by θ_D the restriction of $\theta_{r,g}$ to $S\mathcal{U}_C(r, \mathcal{O}_C(rD))$. Then we have the following commutative diagram:

$$\begin{array}{ccc} S\mathcal{U}_C(r) & \xrightarrow{t_D} & S\mathcal{U}_C(r, \mathcal{O}_C(rD)) \\ \downarrow \theta & & \downarrow \theta_D \\ |r\Theta| & \xrightarrow{\sigma_D} & |r\Theta_D| \end{array}$$

since t_D and σ_D are isomorphisms, we can identify the two theta maps. Finally remark that the composed map $p \circ \theta_{r,g}$ is precisely the natural map which associates to each vector bundle F its determinant line bundle $\det(F)$.

4. Generically Generated Coherent Systems

A pair (F, V) is a *coherent system* of type (r, d, k) on the curve C if F is a vector bundle of rank r and degree d on C and $V \subseteq H^0(F)$ is a linear subspace of dimension k . A coherent system (F, V) is *generically generated* if the evaluation map

$$\text{ev}_{F,V}: V \otimes \mathcal{O}_C \rightarrow F, \quad (s, x) \rightarrow s(x)$$

has torsion cokernel. A proper coherent subsystem of (F, V) is a pair (G, W) where G is a nonzero sub-bundle of F and $W \subseteq V \cap H^0(G)$, with $(G, W) \neq (F, V)$.

For any real number α , we define the α -slope of a coherent system (F, V) of type (r, d, k) as follows:

$$\mu_\alpha(F, V) = \frac{d}{r} + \alpha \frac{k}{r}.$$

Definition 4.1. A coherent system (F, V) is α -stable (respectively, α -semi-stable) if for any proper coherent subsystem (G, W) of (F, V) we have:

$$\mu_\alpha(G, W) < \mu_\alpha(F, V) \quad (\text{respectively } \leq).$$

Every α -semi-stable coherent system (F, V) admits a Jordan–Hölder fibration:

$$0 = (F_0, V_0) \subset (F_1, V_1) \subset \cdots \subset (F_n, V_n) = (F, V),$$

with $\mu_\alpha(F_j, V_j) = \mu_\alpha(F, V)$, $\forall j = 1, \dots, n$ and such that each coherent quotient system

$$(G_j, W_j) = \frac{(F_j, V_j)}{(F_{j-1}, V_{j-1})}$$

is α -stable. This defines the graded coherent system:

$$\text{gr}(F, V) = \bigoplus_{j=1}^n (G_j, W_j).$$

Finally, we say that two α -semi-stable coherent systems are *S-equivalent* if and only if their graded coherent systems are isomorphic.

Let $G_\alpha(r, d, k)$ be the moduli space parametrizing α -stable coherent systems of type (r, d, k) , its compactification $\bar{G}_\alpha(r, d, k)$ is a projective scheme parametrizing S-equivalence classes of α -semi-stable coherent systems of type (r, d, k) , see [22] for details. For $k \geq 1$, it follows easily from the definitions that, if $\bar{G}_\alpha(r, d, k) \neq \emptyset$, then $\alpha \geq 0$ and $d \geq 0$; if $G_\alpha(r, d, k) \neq \emptyset$, then $\alpha > 0$.

Definition 4.2. A positive real number α is said to be a *virtual critical value* for coherent systems of type (r, d, k) if it is numerically possible for a system (F, V) to have a proper subsystem (G, W) of type (r', d', k') such that $\mu_\alpha(F, V) = \mu_\alpha(G, W)$ with $\frac{k}{r} \neq \frac{k'}{r'}$. If there is a coherent system (F, V) and a subsystem (F', V') such that this actually holds, we say that α is an *actual critical value*.

It is well known ([9, Secs. 2.1 and 4]) that, for coherent systems of type (r, d, k) , the actual critical values form a finite set:

$$0 = \alpha_0 < \alpha_1 < \cdots < \alpha_L,$$

and that within the interval (α_i, α_{i+1}) the property of α -stability of a pair is independent of α . This means that $G_\alpha(r, d, k)$ is isomorphic to $G_{\alpha'}(r, d, k)$ whenever α and α' are contained in the same open interval (α_i, α_{i+1}) . The same isomorphism holds for the respective compactifications. It is customary to call $G_L(r, d, k)$ the *terminal* moduli space, i.e. the one that comes within the range $(\alpha_L, +\infty)$.

In this paper we will consider coherent systems of type (r, rg, r) . The following properties hold, thanks to [9, Theorems 4.4, 4.6 and 5.6].

Proposition 4.1. *Let $r \geq 2$. For $\alpha > g(r - 1)$ the moduli spaces $G_\alpha(r, rg, r)$ stabilize, i.e. we have:*

$$G_\alpha(r, rg, r) = G_L(r, rg, r), \quad \text{if } \alpha > g(r - 1).$$

The moduli space $G_L(r, rg, r)$ is a smooth quasi-projective variety of dimension $r^2(g - 1) + 1$ and its compactification $\bar{G}_L(r, rg, r)$ is irreducible. Moreover, after $g(r - 1)$, each α -semi-stable (F, V) is generically generated.

One relation between the stability of a coherent system and that of the underlying vector bundle is given by the following lemma.

Lemma 4.1. *Let (F, V) be a coherent system of type (r, rg, r) , which is generically generated. Then we have the following properties:*

- (1) *If F is semi-stable, then (F, V) is α -semi-stable for any $\alpha \geq 0$;*
- (2) *If F is stable, then (F, V) is α -stable for any $\alpha > 0$;*
- (3) *If $\alpha > g(r - 1)$, then either (F, V) is α -semi-stable or there exists a subsystem (G, W) of type (s, d, k) with $s = k$ and $\frac{d}{s} > g$.*

Proof. Let (G, W) be a proper coherent subsystem of (F, V) of type (s, d, k) :

$$1 \leq s \leq r, \quad 0 \leq k \leq r, \quad (G, W) \neq (F, V).$$

We show that (F, V) generically generated implies $\frac{k}{s} \leq 1$. We can consider the map $\text{ev}_{F,W} : W \otimes \mathcal{O}_C \rightarrow F$, its image is a subsheaf $\text{Im}(\text{ev}_{F,W})$ of F . The inclusion $W \subset V$ implies $\text{Im}(\text{ev}_{F,W})$ has generically rank k . Finally, $W \subset H^0(G)$ so $\text{Im}(\text{ev}_{F,W})$ is a subsheaf of G , which implies $k \leq s$.

(1)–(2) Suppose that F is semi-stable. Since for any proper coherent subsystem (G, W) of type (s, d, k) we have $\frac{k}{s} \leq 1$ then for any α we have: $\mu_\alpha(G, W) \leq \mu_\alpha(F, V)$. In particular, if F is stable, then it is easy to see that (F, V) is α -stable if $\alpha > 0$.

(3) Let (G, W) be a coherent subsystem of type (s, d, k) which contradicts the α -semi-stability of (F, V) . As we have seen, $\frac{k}{s} \leq 1$. If $\frac{k}{s} = 1$, then $\frac{d}{s} > g$. If F is not semi-stable and (F, V) is not α -semi-stable. If $\frac{k}{s} < 1$, then $\frac{F}{G}$ is generically generated too, hence $\text{deg}(\frac{F}{G}) \geq 0$. This implies $d \leq rg$. So we have:

$$g + \alpha < \frac{d}{s} + \alpha \frac{k}{s} \leq \frac{rg}{s} + \alpha \frac{k}{s},$$

which implies:

$$\alpha < \frac{g(r - s)}{s - k} \leq g(r - 1). \quad \square$$

Lemma 4.2. *There is a natural birational map $b : \mathcal{U}_C(r, rg) \dashrightarrow \bar{G}_L(r, rg, r)$.*

Proof. Let $U_{r,rg} \subset \mathcal{U}_C(r, rg)$ be the subset of stable points $[F]$ satisfying the following properties:

- (1) $h^0(F) = r$;
- (2) the determinant map $d_F : \wedge^r H^0(F) \rightarrow H^0(\det F)$ is not the zero map.

The conditions that define the set $U_{r,rg} \subset \mathcal{U}_C(r, rg)$ are clearly open; we prove that $U_{r,rg}$ is not empty. Let $F_0 = L_1 \oplus L_2 \oplus \dots \oplus L_r$, with L_i a line bundle of degree g and $h^0(L_i) = 1$ for any $i = 1, \dots, r$. Then F_0 satisfies (1) and (2). Let F_t be a stable deformation of F_0 along a one parameter family T . For a generic $\bar{t} \in T$,

$F_{\bar{t}}$ satisfies (2). Moreover, by semicontinuity $h^0(F_{\bar{t}}) \leq r$ and by Riemann–Roch $h^0(F_{\bar{t}}) \geq r$, so $F_{\bar{t}}$ satisfies (1) too, hence $[F_{\bar{t}}] \in U_{r,rg}$.

Let $[F] \in U_{r,rg}$. The pair $(F, H^0(F))$ is a coherent system of type (r, rg, r) , by Property (2) it is generically generated and by Lemma 4.1(2) it is α -stable. So we have a morphism

$$b: U_{r,rg} \rightarrow \bar{G}_L(r, rg, r); \quad (4.1)$$

$$[F] \rightarrow [(F, H^0(F))], \quad (4.2)$$

it is easy to see that it is an isomorphism onto the image. Since $\dim \mathcal{U}_C(r, rg) = \dim \bar{G}_L(r, rg, r)$ and $\bar{G}_L(r, rg, r)$ is irreducible, we conclude that b induces a birational map between the moduli spaces $\mathcal{U}_C(r, rg)$ and $\bar{G}_L(r, rg, r)$. \square

5. The Fundamental Divisor of a Coherent System

Let (F, V) be an α -stable coherent system of type (r, rg, r) on the curve C . Assume that it is generically generated, then the map $\text{ev}_{F,V}: V \otimes \mathcal{O}_C \rightarrow F$ is a generically surjective map between two vector bundles of the same rank, so its degeneracy locus is an effective divisor on the curve C :

$$\mathbb{D}_{(F,V)} \in |\det F|.$$

Moreover, let us consider the restriction to $\wedge^r V$ of the determinant map of F

$$d_{F,V}: \wedge^r V \rightarrow H^0(C, \det F);$$

$$s_1 \wedge s_2 \cdots \wedge s_r \mapsto (x \mapsto s_1(x) \wedge s_2(x) \wedge \cdots \wedge s_r(x)).$$

This is not zero so its image is a line generated by a nonzero global section σ of $H^0(\det F)$:

$$d_{F,V}(\wedge^r V) = \text{Span}(\sigma), \quad \sigma \in H^0(C, \det F).$$

It is easy to see that $\mathbb{D}_{(F,V)} = \text{Zeros}(\sigma)$.

Definition 5.1. We call $\mathbb{D}_{(F,V)}$ the *fundamental divisor* of (F, V) .

Let $(F, V) = (F_1, V_1) \oplus (F_2, V_2)$, assume that for $i = 1, 2$, the pair (F_i, V_i) is an α -stable coherent system, which is generically generated too. Then we have $\mathbb{D}_{(F,V)} = \mathbb{D}_{(F_1,V_1)} + \mathbb{D}_{(F_2,V_2)}$.

Definition 5.2. Let $[F, V]$ be the S-equivalence class of an α -semi-stable coherent system (F, V) . We define the *fundamental divisor* of $[F, V]$ as

$$\mathbb{D}_{[F,V]} := \mathbb{D}_{\text{gr}(F,V)}.$$

Note that if (F, V) is α -stable, then $\text{gr}(F, V) = (F, V)$, so that $\mathbb{D}_{[F,V]} = \mathbb{D}_{(F,V)}$.

Every coherent system $(F, L) \in \bar{G}_L(r, rg, r)$ is generically generated (see Proposition 4.1), hence we can define the following map:

$$\Phi: \bar{G}_L(r, rg, r) \rightarrow C^{(rg)}, \tag{5.1}$$

$$[F, V] \mapsto \mathbb{D}_{\text{gr}(F, V)}. \tag{5.2}$$

Definition 5.3. We call Φ the *fundamental map* of generically generated coherent systems of type (r, rg, r) .

6. The Fundamental Map and Its Fibers

The aim of this section is the description of the fibers of Φ . We start by showing some basic properties of the map Φ itself.

Theorem 6.1. *For any $r \geq 2$, $\Phi: \bar{G}_L(r, rg, r) \rightarrow C^{(rg)}$ is a surjective morphism.*

Proof. Let $(\mathcal{F}, \mathcal{V})$ be a flat family of α -semi-stable generically generated coherent systems of type (r, rg, r) over a scheme S . Then \mathcal{F} is a rank r vector bundle on $C \times S$, $\mathcal{F}|_{C \times s} = F_s$ is a vector bundle of rank r and degree rg on C . Let p_1 and p_2 be the natural projections of $C \times S$ onto its factors, $\mathcal{V} \subset (p_2)_*\mathcal{F}$ is a vector bundle of rank r on S , with fiber V_s at the point s . Finally, (F_s, V_s) is a generically generated coherent system of type (r, rg, r) on the curve C , which is α -semi-stable, for any $s \in S$. Let $\mu: S \rightarrow \bar{G}_L(r, rg, r)$ be the morphism defined by the family, sending $s \rightarrow (F_s, V_s)$.

Let $E_v: (p_1)^*\mathcal{V} \rightarrow \mathcal{F}$ be the natural evaluation map, note that it is a map between two vector bundles of the same rank, moreover for any $s \in S$ we have: $E_v|_{C \times s} = \text{ev}_{F_s, V_s}$. This implies that the degeneracy locus of E_v is a relative divisor \mathcal{D} on C over S of relative degree rg . For any $s \in S$ we have $\mathcal{D}|_{C \times s} = \mathbb{D}_{F_s, V_s}$. This induces a morphism $\Phi_S: S \rightarrow C^{(rg)}$, sending $s \rightarrow \mathbb{D}_{F_s, V_s}$, such that $\Phi_S = \Phi \circ \mu$. This proves that Φ is a morphism.

Let us come to surjectivity. Let G be a point set in $C^{(rg)}$, note that G can be written as the sum of r effective divisors of degree g :

$$G = G_1 + G_2 + \dots + G_r.$$

For any $i = 1, \dots, r$, let $\sigma_i \in H^0(\mathcal{O}_C(G_i))$ be a nonzero global section such that $G_i = \text{Zeros}(\sigma_i)$ and let $V_i = \text{Span}(\sigma_i) \subset H^0(\mathcal{O}_C(G_i))$. Now let us define the following pair:

$$F: = \bigoplus_{i=1}^r \mathcal{O}_C(G_i), \quad V: = \bigoplus_{i=1}^r V_i.$$

Then (F, V) is a generically generated coherent system of type (r, rg, r) on C and F is semi-stable. By Lemma 4.1(1), (F, V) is α -semi-stable and $\mathbb{D}_{F, V} = G$. This implies that $\Phi([F, V]) = G$ and proves the surjectivity. □

Let $(\mathbb{P}^{r-1})^{rg} // \text{PGL}(r)$ denote the GIT quotient of $(\mathbb{P}^{r-1})^{rg}$ with respect to the diagonal action of $\text{PGL}(r)$. We recall that the notion of GIT stability of a point set $v \in (\mathbb{P}^{r-1})^{rg}$ has a nice geometric formulation in terms of the dimension of the linear span in \mathbb{P}^{r-1} of subsets of v (see [15, Theorem 1, p. 23]).

Proposition 6.1. *The point set $v = (v_1, \dots, v_{rg}) \in (\mathbb{P}^{r-1})^{rg}$ is GIT semi-stable (respectively stable) if and only if for any subset $\{v_1, \dots, v_k\}$ of v we have*

$$\dim(\text{Span}(v_1, \dots, v_k)) \geq \frac{k}{g} \quad (\text{respectively } >).$$

The first important feature of the map Φ is the following.

Theorem 6.2. *The general fiber of Φ is isomorphic to $(\mathbb{P}^{r-1})^{rg} // \text{PGL}(r)$.*

Proof. Let $B \in C^{(rg)}$, we assume that B is not contained in the big diagonal Δ , that is

$$B = \sum_{i=1}^{rg} x_i, \quad x_i \neq x_j, \quad \forall i \neq j, \quad x_i \in C, \quad \forall i.$$

Then the fiber of Φ at B is the following:

$$\Phi^{-1}(B) = \{[(F, V)] \in \tilde{G}_L(r, rg, r) \mid \mathbb{D}_{\text{gr}(F, V)} = B\}.$$

Let $[F, V] \in \Phi^{-1}(B)$, from now on we will write for simplicity $\text{gr}(F, V) := (F_g, V_g)$. Since B is the degeneracy locus of the evaluation map ev_{F_g, V_g} , by dualizing we find the following exact sequence:

$$0 \rightarrow F_g^* \rightarrow V_g^* \otimes \mathcal{O}_C \rightarrow \mathcal{O}_B \rightarrow 0. \quad (6.1)$$

Hence, up to the choice of a basis of V_g , F_g^* is the kernel of a surjective morphism $v_{F_g, V_g} \in \text{Hom}(V_g^* \otimes \mathcal{O}_C, \mathcal{O}_B)$:

$$v_{F_g, V_g} = (v_1, \dots, v_{rg}), \quad v_i \neq 0, \quad v_i \in \text{Hom}(V_g^* \otimes \mathcal{O}_C, \mathcal{O}_{x_i}) \simeq V_g.$$

This means that (F_g, V_g) defines a point (that we will still denote by v_{F_g, V_g}) in the product space $(\mathbb{P}(V_g))^{rg} \cong (\mathbb{P}^{r-1})^{rg}$, and $\text{PGL}(r)$ acts diagonally on this space via the choice of the basis of V_g .

Claim 1. *$v_{F_g, V_g} \in (\mathbb{P}^{r-1})^{rg}$ is GIT semi-stable with respect to the diagonal action of $\text{PGL}(r)$. If (F_g, V_g) is α -stable then v_{F_g, V_g} is GIT stable.*

Proof. Let $w = \{v_1, \dots, v_d\}$ be a subset of v_{F_g, V_g} , set $W := \text{Span}(v_1, \dots, v_d)$. Let x_i be the point of B that corresponds to v_i , for $i = 1, \dots, d$. Then $W \subset V_g$, so we get a commutative diagram as follows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & F_g^* & \longrightarrow & V_g^* \otimes \mathcal{O}_C & \xrightarrow{v_{F_g, V_g}} & \mathcal{O}_B \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & G^* & \longrightarrow & W^* \otimes \mathcal{O}_C & \xrightarrow{w} & \bigoplus_{i=1}^d \mathcal{O}_{x_i} \longrightarrow 0 \end{array}$$

for some vector bundle G^* with $\text{rk}(G^*) = \dim(W) = s$. Note that the pair (G, W) is a coherent system of type (s, d, s) , which is a proper subsystem of (F_g, V_g) . Since this is α -semi-stable, we have:

$$\mu_\alpha(G, W) = \frac{d}{s} + \alpha \leq \mu_\alpha(F_g, V_g) = g + \alpha,$$

which implies $s \geq \frac{d}{g}$, which gives the GIT semi-stability of v_{F_g, V_g} . The stable case is described in the same way but with strict inequalities. \square

Let V be a vector space of dimension r and $\mathbb{P}(V)$ the associated projective space. By mimicking sequence (6.1), we can construct a flat family of coherent systems of type (r, rg, r) on C over $(\mathbb{P}(V))^{rg}$. Let $v = (v_1, \dots, v_{rg}) \in \mathbb{P}(V)^{rg}$, v defines a surjective morphism of sheaves $V^* \otimes \mathcal{O}_C \rightarrow \mathcal{O}_B$ as follows: it is the zero map out of the support of B and it is obtained by taking one lift of v_i to V^* and applying it on the fiber of $V^* \otimes \mathcal{O}_C$ over $x_i \in B$. The morphism depends on the choice of the lift but the kernel of the sequence

$$0 \rightarrow \ker(v) \rightarrow V^* \otimes \mathcal{O}_C \xrightarrow{v} \mathcal{O}_B \rightarrow 0, \tag{6.2}$$

is well defined over $\mathbb{P}(V)^{rg}$. This implies that $F_v := \ker(v)^*$, for $v \in \mathbb{P}(V)^{rg}$, is a family of rank r vector bundles on C with determinant $\mathcal{O}_C(B)$. Note that F_v is generically generated by a linear subspace of $H^0(C, F_v)$ of dimension r : we will denote it by V_v . The pair (F_v, V_v) is a coherent system of type (r, rg, r) , generically generated and $\mathbb{D}_{F_v, V_v} = B$. Moreover the family (F_v, V_v) is invariant under the diagonal action of $\text{PGL}(r)$ on $\mathbb{P}(V)^{rg}$.

Claim 2. *Let $v \in \mathbb{P}(V)^{rg}$ be GIT semi-stable (respectively stable), then the pair (F_v, V_v) is α -semi-stable (respectively stable) for $\alpha > g(r - 1)$, hence $[F_v, V_v] \in \bar{G}_L(r, rg, r)$.*

Proof. Let $\alpha > g(r - 1)$. Since (F_v, V_v) is generically generated, we can apply Lemma 4.1(3): either (F_v, V_v) is α -semi-stable, or there exists a proper subsystem (G, W) of type (s, d, k) with $s = k$ and $\frac{d}{s} > g$. Then (G, W) is generically generated too, and we have a commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & W \otimes \mathcal{O}_C & \longrightarrow & G & \xrightarrow{v_{G,W}} & \bigoplus_{i=1}^d \mathcal{O}_{x_i} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & V \otimes \mathcal{O}_C & \longrightarrow & F_v & \xrightarrow{v=v_{F_v,V}} & \mathcal{Q}_B \longrightarrow 0 \end{array}$$

Let $v_{G,W} = (v_1, \dots, v_d)$ and $\text{Span}(v_1, \dots, v_d) \subset W \subset V$ with $\dim W = s$. Since $v \in \mathbb{P}(V)^{rg}$ is GIT semi-stable then we have that $\dim W \geq \frac{d}{g}$, which contradicts the hypothesis. This proves that (F_v, V_v) is α -semi-stable.

Suppose now that we have a stable point set v , and that there exists a proper coherent subsystem $(G, W) \subset (F_v, V_v)$ of type (s, d, k) such that $\mu_\alpha(G, W) = \mu_\alpha(F_v, V_v)$. Since α is not a critical value, $\frac{k}{s} = 1$, and thus $s = \frac{d}{g}$

(see Definition 4.2). But, keeping the same notation of the first part of the proof, if v is stable, then $\dim(W) = s > \frac{d}{g}$. Hence (F_v, V_v) is α -stable. \square

If $\alpha > g(r - 1)$, Claim 2 and the fact that the family of coherent systems over $\mathbb{P}(V)^{rg}$ is $\mathrm{PGL}(r)$ -invariant allow us to define a morphism

$$\mathbb{P}(V)^{rg} // \mathrm{PGL}(r) \rightarrow \Phi^{-1}(B).$$

Indeed, the family over $\mathbb{P}(V)^{rg}$ induces a morphism from the semi-stable locus to $\bar{G}_L(r, rg, r)$. This morphism is $\mathrm{PGL}(r)$ -invariant hence it factors through $\mathbb{P}(V)^{rg} // \mathrm{PGL}(r)$. Furthermore, by construction its inverse is the map sending each system $[F, V] \mapsto [v_{F_g, V_g}]$ (see Claim 1). Hence we conclude that the general fiber of Φ is isomorphic to $\mathbb{P}(V)^{rg} // \mathrm{PGL}(r)$. \square

Let us denote $\alpha_1 > 0$ the smaller positive critical value for coherent systems of a given type (n, d, k) . The stability condition induced by values of α such that $0 < \alpha < \alpha_1$ is called 0^+ -stability (see [8, pp. 4–5]). It is well known that if a coherent system (F, V) is 0^+ -stable then the underlying vector bundle F is slope-semistable. On the other hand, if the underlying vector bundle F is stable, then (F, V) is 0^+ -stable. In the following example, we suppose $r = 2$ and $g(C) > 3$. We produce a coherent system on C of type $(2, 2g, 2)$ which comes from a GIT stable point and is α -semi-stable for any $\alpha \geq 1$, but the underlying vector bundle is unstable, i.e. the coherent system is not 0^+ -semi-stable.

Example 6.1. Let C be a smooth curve of genus $g > 3$. Let L_1 be a special line bundle of degree $g - 1$ with $h^0(C, L_1) = 2$. Let $\{t_1, t_2\}$ be a basis of $H^0(C, L_1)$. We can assume that the zeros of t_1 are all simple and distinct. By definition, given a scalar λ , one may find at most a finite set (possibly empty) of points $x \in C$ such that $t_1(x) = \lambda t_2(x)$. Hence we are allowed to choose a set of $g + 1$ distinct points z_1, \dots, z_{g+1} such that: $t_1(z_i) \neq 0$, $t_1(z_i) = \lambda_i t_2(z_i)$, with $\lambda_i \neq \lambda_j$ for $1 \leq i \neq j \leq g + 1$ and $\lambda_i \neq 0$. Let $M := z_1 + \dots + z_{g+1}$, $L_2 := \mathcal{O}_C(M)$ and let s be a nonzero global section of $H^0(C, L_2)$ such that $M = \mathrm{Zeros}(s)$. Now let us consider the coherent system (F, V) of type $(2, 2g, 2)$ defined as follows:

$$F := L_1 \oplus L_2 \quad V := \langle t_1, t_2 + s \rangle.$$

Note that the evaluation map

$$\mathrm{ev}_{F, V} : V \otimes \mathcal{O}_C \rightarrow F,$$

is generically surjective, degenerating on the divisor $B := \mathrm{Zeros}(t_1) + \mathrm{Zeros}(s)$.

We show that $(F, V) \in \bar{G}_L(r, rg, r)$, hence $(F, V) \in \Phi^{-1}(B)$, but the underlying vector bundle is not semi-stable: in fact $L_2 \subset F$ is a destabilizing sub-bundle. Notably, we prove that (F, V) is α -semi-stable for any $\alpha \geq 1$. Let (G, W) be a coherent subsystem of type (s, d, k) . It is easy to see that $\mu_\alpha(G, W) > \mu_\alpha(F, V)$ implies that $s = 1$, $d = g + 1$, $k = 0$ and $\alpha < 1$. Note that $\alpha = 1$ is a critical value for coherent systems of type $(2, 2g, 2)$.

Finally, let $v = (v_1, v_2, \dots, v_{2g}) \in \mathbb{P}(V)^{2g}$ be the point set defined by the pair (F, V) , as in the proof of Theorem 6.2. By the assumptions we made choosing M and the sections t_i and s , the point set v has the following shape: for $i = 1, \dots, g - 1$, $x_i \in \text{Zeros}(t_1)$, so $[v_i] = [1 : 0]$; for $i = g, \dots, 2g$, $x_i \in \text{Zeros}(s)$, so we have: $[v_i] = [\lambda_i : 1]$, with $\lambda_i \neq 0$. Such a point set $v \in \mathbb{P}(V)^{2g}$ is clearly stable.

As the reader may expect, given the birationality result between the moduli space $\mathcal{U}_C(r, rg)$ and $G_L(r, rg, r)$, the fundamental map gives information also on the geometry of $\mathcal{U}_C(r, rg)$ and of the theta map.

Theorem 6.3. *Let b be the birational map defined in (4.1):*

$$b: \mathcal{U}_C(r, rg) \dashrightarrow \bar{G}_L(r, rg, r), \quad [F] \rightarrow [(F, H^0(F))].$$

Let b_D be its restriction to $S\mathcal{U}_C(r, \mathcal{O}_C(rD))$ and let θ_{rg} be the theta map of $\mathcal{U}_C(r, rg)$ defined in (3.2) and θ_D its restriction to $S\mathcal{U}_C(r, \mathcal{O}_C(rD))$, then we have the following commutative diagrams of rational maps

$$\begin{array}{ccc} \mathcal{U}_C(r, rg) & \xrightarrow{b} & \bar{G}_L(r, rg, r) & & S\mathcal{U}_C(r, \mathcal{O}_C(rD)) & \xrightarrow{b_D} & b_D(S\mathcal{U}_C(r, \mathcal{O}_C(rD))) \\ \theta_{rg} \downarrow & & \downarrow \Phi & & \theta_D \downarrow & & \downarrow \Phi \\ \mathcal{T} & \xrightarrow{a^*} & C^{(rg)} & & |r\Theta_D| & \xrightarrow{a^*} & |\mathcal{O}_C(rD)| \end{array}$$

where a^* is the pull-back of theta divisors via the Abel map $a: C \rightarrow \text{Pic}^{(1)}(C)$. Notably, the restricted map

$$\pi_e = a^*|_{|r\Theta_D|}: |r\Theta_D| \rightarrow |\mathcal{O}_C(rD)|$$

is a linear projection with center $L_e := \{M \in |r\Theta_D| : a(C) \subset M\}$.

Proof. Let $[F] \in \mathcal{U}_C(r, rg)$. Then $[F]$ is contained in the regular locus of θ_{rg} if and only if $[F]$ admits a theta divisor

$$\Theta_F = \{L \in \text{Pic}^{(1)}(C) \mid h^0(C, F \otimes L^{-1}) \geq 1\}.$$

Furthermore, the divisor Θ_F lies in the regular locus of a^* if it does not contain the image of C via a . Remark in fact that in this case, the pull back of Θ_F via a^* is an effective divisor of degree rg . We have:

$$a^*(\Theta_F) = \{x \in C \mid h^0(C, F \otimes \mathcal{O}_C(-x)) \geq 1\}.$$

This implies that $h^0(F) = r$ and the evaluation map $\text{ev}_{F, H^0(F)}: H^0(F) \otimes \mathcal{O}_C \rightarrow F$ is generically surjective, hence $b([F])$ is defined. In order to show that the two diagrams commute it is enough to remark that if $x \in \mathbb{D}_{(F, H^0(F))}$ then there exist at least one nonzero section in $h^0(C, F \otimes \mathcal{O}_C(-x))$, hence $\mathbb{D}_{(F, H^0(F))}$ and $a^*(\Theta_F)$ coincide. □

As a corollary of the results of this section we have the following theorem.

Theorem 6.4. *For any $r \geq 2$ and $g \geq 2$, the moduli space $\mathcal{U}_C(r, rg)$ is birational to a fibration over $C^{(rg)}$ whose fibers are $(\mathbb{P}^{r-1})^{rg} // \mathrm{PGL}(r)$ and the moduli space $SU_C(r, \mathcal{O}_C(rD))$ is birational to a fibration over $\mathbb{P}^{(r-1)g} = |\mathcal{O}_C(rD)|$ whose fibers are $(\mathbb{P}^{r-1})^{rg} // \mathrm{PGL}(r)$.*

7. Application to the Coble Hypersurfaces

As Example 6.1 showed, in the case of vector bundles we cannot expect any isomorphism between the fibers of Φ and GIT quotients. Nevertheless, for low rank and genus, the geometry of moduli spaces of vector bundles is simple enough that, by taking the closure of the fibers, we still find, at least as projective varieties, families of GIT quotients contained in moduli spaces of vector bundles. This section is devoted to the construction and study of such families of projective varieties.

When the genus of C is 2 or 3 and the rank is small enough, the moduli spaces $SU_C(r)$ have very nice explicit descriptions related to certain hypersurfaces, called the *Coble hypersurfaces*. The main theorems of this section show how these hypersurfaces contain large families of projective classical modular varieties related to invariant theory, namely the *Segre cubic* and the *Igusa quartic*.

We know a purely vector bundle-theoretic construction (i.e. without the use of the theta map) of such families, but it is rather involved and, in our opinion, less instructive. For this reason and for the beauty of the objects of study, we have preferred to give a construction that relies on the theta map and the projective geometry of the Coble hypersurfaces.

7.1. The Coble sextic

In this subsection we assume that C is a curve of genus 2 and we consider the moduli space $SU_C(3)$ of semi-stable vector bundles on C with rank 3 and trivial determinant. The theta map

$$\theta : SU_C(3) \rightarrow |3\Theta| \simeq \mathbb{P}^8$$

is a finite morphism of degree 2 and the branch locus is a sextic hypersurface \mathcal{C}_6 , called the Coble–Dolgachev sextic [25]. The Jacobian variety $J(C)$ is embedded in $|3\Theta|^*$ as a degree 18 surface, and there exists a unique cubic hypersurface $\mathcal{C}_3 \subset |3\Theta|^*$ whose singular locus coincides with $J(C)$: the *Coble cubic*, [13]. It was conjectured by Dolgachev, and subsequently proved in [31] and independently in [29], that \mathcal{C}_6 is the dual variety of \mathcal{C}_3 .

Let us consider the moduli space $SU(3, 3K_C)$ and let

$$\Phi_{K_C} : SU(3, 3K_C) \dashrightarrow |3K_C|,$$

be the rational map sending $[F]$ to $\mathbb{D}_{\mathrm{gr}(F), H^0(\mathrm{gr}(F))}$. It is the composition of b_{K_C} with the fundamental map Φ . By Theorem 6.3, we have the following result.

Proposition 7.1. *The map Φ_{K_C} is the composition of the theta map*

$$\theta_{K_C} : SU(3, 3K_C) \rightarrow |3\Theta_{K_C}|,$$

with the linear projection $\pi_e : |3\Theta_{K_C}| \dashrightarrow |3K_C|$, whose center $L_e \simeq \mathbb{P}^3$, is the linear subsystem of theta divisors corresponding to decomposable bundles of type $E \oplus K_C$, with $E \in SU_C(2, 2K_C)$.

Proof. Since $\dim|3K_C| = 4$, the center of the projection π_e is a three-dimensional linear subspace L_e , which is the image by the theta map of the indeterminacy locus of Φ_{K_C} . Let us consider the subset $\mathcal{I} \subset SU(3, 3K_C)$ of semi-stable bundles $[F] = [E \oplus K_C]$, $[E] \in SU_C(2, 2K_C)$. Note that $\mathcal{I} \simeq SU_C(2, 2K_C) \simeq \mathbb{P}^3$ [28]. Moreover, since $h^0(\text{gr}(E \oplus K_C)) \geq 4$, then \mathcal{I} is contained in the indeterminacy locus of Φ_{K_C} . Finally, let h be the hyperelliptic involution on C , it defines the involution

$$\begin{aligned} \sigma : SU(3, 3K_C) &\rightarrow SU(3, 3K_C); \\ F &\mapsto h^*F^*; \end{aligned}$$

which is associated to the 2:1 covering given by θ_{K_C} . Then \mathcal{I} is contained in the fixed locus of σ , [29, Secs. 3 and 4]. The image via θ_{K_C} of this locus is $L_e \simeq \mathbb{P}^3$ and it is actually the center of the projection. \square

By Theorem 6.3, a general fiber of Φ_{K_C} is birational to the quotient $(\mathbb{P}^2)^6 // \text{PGL}(3)$. This is a degree two covering of \mathbb{P}^4 branched along a Σ_6 -invariant quartic hypersurface $\mathcal{I}_4 \subset \mathbb{P}^4$. \mathcal{I}_4 is known as the *Igusa quartic*, it is the Satake compactification of the moduli space $\mathcal{A}_2(2)$ of principally polarized abelian surfaces with a level two structure [20], embedded in \mathbb{P}^4 by fourth powers of theta-constants. The involution that defines the covering is the Gale transform (also called association, for details see [15, 17]), which is defined as follows.

Definition 7.1 ([17, Definition 1.1]). Let $r, s \in \mathbb{Z}$. Set $\gamma = r + s + 2$, and let $\Gamma \subset \mathbb{P}^r$, $\Gamma' \subset \mathbb{P}^s$ be ordered nondegenerate sets of γ points represented by $\gamma \times (r+1)$ and $\gamma \times (s+1)$ matrices G and G' , respectively. We say that Γ' is the Gale transform of Γ if there exists a nonsingular diagonal $\gamma \times \gamma$ matrix D such that $G^T \cdot D \cdot G' = 0$, where G^T is the transposed matrix of G .

The Gale transform acts trivially on the sets of 6 points in \mathbb{P}^2 that lie on a smooth conic. The branch locus of the double covering is then, roughly speaking, the moduli space of 6 points on a conic and henceforth a birational model of the moduli space of 6 points on a line. One can say even more, in fact the GIT compactification of the moduli space of 6 points on a line is a cubic three-fold in \mathbb{P}^4 , called the Segre cubic, and its projectively dual variety is the Igusa quartic (see [18, 24] for details). From the projective geometry point of view the singular locus of \mathcal{I}_4 is an abstract configuration of lines and points that make up a 15_3 configuration. This means the following: there are 15 distinguished lines and 15 distinguished points. Each line

contains 3 of the points and through each point pass 3 lines (see [14, Sec. 9] for more). Moreover \mathcal{L}_4 is the only hypersurface with such a singular locus in the pencil of Σ_6 -invariant quartics in \mathbb{P}^4 (see [19, Example 7]).

Now let us recall some results from the literature about $SU(3, \mathcal{O}_C)$. Of course the twist by K_C is an isomorphism and it is easy to understand the analogous result for $SU_C(3, 3K_C)$. Since \mathbb{P}^8 is smooth, the image of the singular locus of $SU(3, \mathcal{O}_C)$ and the singular locus of the branch locus coincide, i.e. $\text{Sing}(\mathcal{C}_6) = \theta(\text{Sing}(SU(3)))$. On $\text{Pic}^1(C)$ we have the involution $\lambda : L \mapsto K_C \otimes L^{-1}$ that leaves Θ invariant. Hence λ induces an action on all the powers of Θ and in particular, on $|\mathcal{C}_6|$. The linear system $|\mathcal{C}_6|$ decomposes in two eigenspaces, respectively four- and three-dimensional. We call \mathbb{P}_e^4 the four-dimensional eigenspace. It turns out that it cuts out on \mathcal{C}_6 a reducible variety given by a double \mathbb{P}^3 (which is indeed contained in $\text{Sing}(\mathcal{C}_6)$) and a quartic hypersurface $I \subset \mathbb{P}^4$. After the twist by K_C , the first component is precisely L_e , whereas the quartic three-fold is an Igusa quartic.

Lemma 7.1. *The intersection of the closure of the general fiber of π_e with $\text{Sing}(\mathcal{C}_6)$ is a 15_3 configuration of lines and points.*

Proof. Recall that $\text{Sing}(\mathcal{C}_6)$ is the locus of theta divisors corresponding to decomposable bundles, we will prove the claim by constructing explicitly these bundles. Let us denote by Δ_{3K_C} the closed subset of $|3K_C|$ given by the intersection with the big diagonal of $C^{(6)}$. Let us take $G = q_1 + \dots + q_6 \in |3K_C| - \Delta_{3K_C}$, and let us consider the fiber of Φ_{K_C} over G . In order to guarantee the semi-stability of the vector bundles, the only totally decomposable bundles in the fiber of G are all the 15 obtained by permuting the q_i 's in $\mathcal{O}_C(q_1 + q_2) \oplus \mathcal{O}_C(q_3 + q_4) \oplus \mathcal{O}_C(q_5 + q_6)$. Let us now consider the bundles that decompose as the direct sum of a line bundle L and a rank two indecomposable bundle. By the previous argument of semi-stability then L must be of the type $\mathcal{O}_C(q_i + q_j)$ for some $i, j \in \{1, \dots, 6\}$ and E should have fundamental divisor $\mathbb{D}_E = \sum_{k \neq i, j} q_k$. Call F the line bundle $\mathcal{O}_C(\sum_{k \neq i, j} q_k) \cong 3K_C - q_i - q_j$. It is easy to see that $SU_C(2, F) \cong SU_C(2, \mathcal{O}_C) \cong \mathbb{P}^3$, the isomorphism being given by the tensor product by a square root F' of F . Now we recall from [7] the following description of the fundamental map $\Phi_{F'} : SU_C(2, F) \dashrightarrow |F|$. The linear system $|F|$ is a \mathbb{P}^2 and the fibers of $\Phi_{F'}$ are just lines passing through $D \in |F|$ and the origin $[\mathcal{O}_C \oplus \mathcal{O}_C]$. Now the composition of the following embedding

$$\zeta : SU_C(2, F) \hookrightarrow SU_C(3, 3K), \tag{7.1}$$

$$[E] \mapsto [\mathcal{O}_C(q_i + q_j) \oplus E], \tag{7.2}$$

with the theta map is linear. In fact $\zeta(SU_C(2, F))$ is contained in the branch locus and the associated 3Θ divisors form a three-dimensional linear subsystem isomorphic to $|\mathcal{C}_6|$. Then the image of ζ intersects the closure $\overline{\Phi_{K_C}^{-1}(G)}$ of the fiber over G exactly along the fiber of $\Phi_{F'}$ over the divisor $\sum_{k \neq i, j} q_k \in |F|$. By [7] we know that this is a line and it is not difficult to see that it contains 3 of the 15 totally decomposable bundles. On the other hand each totally decomposable bundle with fundamental divisor G is contained in three lines of this kind. \square

Remark 7.1. When the divisor G is taken in Δ_{3K_C} then the 15_3 configuration degenerates because some of the points and some of the lines coincide.

Theorem 7.1. *The closure of the general fiber of Φ_{K_C} is the GIT quotient $(\mathbb{P}^2)^6 // \text{PGL}(3)$.*

Proof. We recall that L_e is contained in $\text{Sing}(\mathcal{C}_6)$ and in particular scheme-theoretically it is contained twice in \mathcal{C}_6 . Since Φ_{K_C} factors through the projection with center L_e , then $\overline{\Phi_{K_C}^{-1}(G)}$, for $G \in |3K_C| - \Delta_{3K_C}$, is a degree two cover of $\mathbb{P}_G^4 := \overline{\pi_e^{-1}(G)}$ ramified along the intersection of \mathcal{C}_6 with \mathbb{P}_G^4 which is residual to $2L_e$. This intersection is then a quartic hypersurface in \mathbb{P}_G^4 . Notably, since $L_e \subset \mathbb{P}_e^4$, there exist a point $T \in |3K_C|$ such that $\overline{\pi_e^{-1}(T)} \cap \mathcal{C}_6$ is an Igusa quartic (see [32, Proposition 5.2] or [30, Sec. 4]).

Let us now blow up $|3\Theta|$ along L_e and call the resulting variety $\tilde{\mathbb{P}}^8$. This contains canonically the blown up Coble sextic, which we denote by $\tilde{\mathcal{C}}_6$. Then the rational map π_e resolves in a proper, flat map $\tilde{\pi}_e$ as in the following diagram.

$$\begin{array}{ccc} \tilde{\mathbb{P}}^8 & & \\ \downarrow & \searrow \tilde{\pi}_e & \\ |3\Theta| - \frac{\pi_e}{\gg} & & |3K_C| \end{array}$$

This says that the restriction of $\tilde{\pi}_e$ to $\tilde{\mathcal{C}}_6$ is a flat family of quartic three-folds over $|3K_C|$ and for any $B \in |3K_C|$ we have an isomorphism $\overline{\pi_e^{-1}(B)} \cap \mathcal{C}_6 \cong \tilde{\pi}_e^{-1}(B) \cap \tilde{\mathcal{C}}_6$. Hence also one fiber of $\tilde{\pi}_e|_{\tilde{\mathcal{C}}_6}$ is an Igusa quartic. Since the ideal of the singular locus of \mathcal{I}_4 is generated by the four polar cubics ([18, Lemma 3.3.13]) then the Igusa quartic has no infinitesimal deformations, i.e. it is rigid. This implies that the generic member of the flat family of quartics over $|3K_C|$ is an Igusa quartic. This in turn implies that the closure of the generic fiber of Φ_{K_C} is $(\mathbb{P}^2)^6 // \text{PGL}(3)$. \square

Corollary 7.1. *The Coble sextic \mathcal{C}_6 is birational to a fibration over \mathbb{P}^4 whose fibers are Igusa quartics.*

Corollary 7.2. *Along the generic fiber of Φ_{K_C} , the involution of the degree two covering $\mathcal{SU}(3, \mathcal{O}_C) \rightarrow |3\Theta|$ coincides with the involution given by the association isomorphism on $(\mathbb{P}^2)^6 // \text{PGL}(3)$.*

The fact that the intersection of $\text{Sing}(\mathcal{C}_6)$ with the fibers over the open set $|3K_C| - \Delta_{3K_C}$ is precisely a 15_3 configuration makes us argue that $|3K_C| - \Delta_{3K_C}$ should be the open locus where by rigidity the family of quartic three-folds is isotrivially isomorphic to the Igusa quartic. As already seen in Remark 7.1, if B is an effective divisor out of this locus, then $\overline{\pi_e^{-1}(B)} \cap \text{Sing}(\mathcal{C}_6)$ is a *degenerate* 15_3 configuration, in the sense that some of the 15 points and lines coincide. We are not able to prove the following, but it is tempting to say that this is all the singular locus of the special quartic three-folds over Δ_{3K} . These would give very

interesting examples of *degenerate Igusa quartics*. It would be interesting to study projective properties of these fibers such as the relation with the Segre cubics or with the Mumford–Knudsen compactification $\overline{\mathcal{M}}_{0,6}$ of the moduli space of 6 points on a line. For instance, do they come from linear systems on $\overline{\mathcal{M}}_{0,6}$? If it is so, what linear systems on $\overline{\mathcal{M}}_{0,6}$ do they come from?

The rational dual map of the Coble sextic has been thoroughly studied and described in [29, 31]. Let us denote by X_0, \dots, X_8 the coordinates on $\mathbb{P}^8 \cong |3\Theta|$ and by $F(X_0 : \dots : X_8)$ the degree six polynomial defining \mathcal{C}_6 . Then the dual map is defined as follows:

$$\mathcal{D}_6 : \mathcal{C}_6 \dashrightarrow \mathcal{C}_3;$$

$$x \mapsto \left[\frac{\partial F}{\partial X_0} : \dots : \frac{\partial F}{\partial X_8} \right].$$

The polar linear system is given by quintics that vanish along $\text{Sing}(\mathcal{C}_6)$. Now fix a general divisor $B \in |3K|$ and let \mathcal{I}_B be the Igusa quartic defined by $(\mathcal{C}_6 \cap \overline{\pi_e^{-1}(B)}) - 2L_e$. Let us consider the restriction of \mathcal{D}_6 to $\mathcal{I}_B \subset \overline{\pi_e^{-1}(B)} =: \mathbb{P}_B^4$ and denote by A the 15_3 configuration of points and lines in \mathbb{P}_B^4 . Now let H be the class of L_e in $\text{Pic}(\mathbb{P}_B^4)$ and consider the four-dimensional linear system $|\mathcal{I}_A(3) + 2H|$ on \mathbb{P}_B . We can show the following proposition.

Proposition 7.2. *The restricted dual map $\mathcal{D}_{6|\mathcal{I}_B}$ is given by a linear system $|\mathcal{D}_{\mathcal{I}_B}|$ that contains $|\mathcal{I}_A(3) + 2H|$ as a linear subsystem.*

Remark that this means that for the general fiber \mathcal{I}_B , there exists a canonical way to project the image $\mathcal{D}(\mathcal{I}_B) \subset \mathcal{C}_3$ to a \mathbb{P}_B^4 where the image of \mathcal{I}_B is a Segre cubic. This is summarized in the following corollary.

Corollary 7.3. *The Coble cubic is birational to a fibration in Segre cubics over \mathbb{P}^4 .*

Remark 7.2. The birationality in itself is trivial, since \mathcal{C}_3 is birational to \mathcal{C}_6 which is birational to a fibration in Igusa quartics (which are in turn all birational to the Segre cubic) over \mathbb{P}^4 . The projections on the linear systems $|\mathcal{I}_A(3) + 2H|$ give a constructive canonical way to realize it.

7.2. The Coble quartic

In this subsection we assume that C is a curve of genus 3 and we consider the moduli space $SU(2, \mathcal{O}_C)$. We recall that the Kummer variety $\text{Kum}(C) := J(C)/\pm \text{Id}$ of C is contained naturally in the 2Θ -linear series, whereas the moduli space $SU(2, \mathcal{O}_C)$ is embedded by θ in $\mathbb{P}^7 \cong |2\Theta|$ as the unique quartic hypersurface \mathcal{C}_4 singular along $\text{Kum}(C)$. This hypersurface is called the *Coble quartic*. It is also known [34] that the Coble quartic is projectively self dual.

Now we recall some properties of the *Segre cubic*. This is a nodal (and hence rational) cubic three-fold S_3 in \mathbb{P}^4 whose singular locus is given by ten double points.

There is a natural action of Σ_6 on this projective space and S_3 is invariant with respect to this action. S_3 is in fact the GIT quotient $(\mathbb{P}^1)^6 // \text{PGL}(2)$ [15]. Moreover, S_3 realizes the so-called *Varchenko bound*, i.e. it has the maximum number of double points (ten) that a cubic three-fold with isolated singularities may have and this property identifies the Segre cubic in a unique way, up to projective equivalence. As already stated it is the projective dual variety of the Igusa quartic.

Our construction allows us to give a simple proof of the following result from [1].

Proposition 7.3. *The moduli space $SU(2, \mathcal{O}_C)$ is birational to a fibration over \mathbb{P}^3 whose fibers are Segre cubics.*

Proof. First of all we twist all vector bundles by a degree 3 divisor D , thus obtaining the isomorphic moduli space $SU(2, \mathcal{O}_C(2D))$. Let $\Phi_D : SU(2, \mathcal{O}_C(2D)) \rightarrow |\mathcal{O}_C(2D)| \cong \mathbb{P}^3$ be the composition of b_D and the fundamental map Φ . By Theorem 6.3, since θ_D is an embedding, we can identify $SU(2, \mathcal{O}_C(2D))$ with its image $\mathcal{C}_4 \subset \mathbb{P}^7$ and Φ_D with the linear projection onto $|2D|$. The center of the projection is the linear span of the locus of vector bundles E such that $h^0(C, E \otimes \mathcal{O}_C(D)) > 2$. Suppose E is stable. Since it has rank 2 and trivial determinant, then $E \cong E^*$ and by an easy Riemann–Roch computation we find that $h^0(C, E \otimes \mathcal{O}_C(D)) > 2$ if and only if $h^0(C, E \otimes \mathcal{O}_C(K - D)) > 0$. This is equivalent to the fact the E lies in the $\mathbb{P}^3 \cong |3K - 2D|^* \subset \mathcal{C}_4$ that parametrizes vector bundles E that can be written as an extension of the following type

$$0 \rightarrow \mathcal{O}_C(D - K) \rightarrow E \rightarrow \mathcal{O}_C(K - D) \rightarrow 0.$$

Let us denote by \mathbb{P}_c^3 this projective space. \mathcal{C}_4 contains \mathbb{P}_c^3 with multiplicity one. This implies that the closure of any fiber of the projection $\Phi_D : \mathcal{C}_4 \dashrightarrow |2D|$ is a cubic three-fold contained in the \mathbb{P}^4 spanned by \mathbb{P}_c^3 and a point of $|2D|$. Let us denote as usual by Δ_D the intersection of the large diagonal with the linear system $|2D| \subset C^{(6)}$. Then suppose we fix a $B \in |2D| - \Delta_D$. Let us consider the intersection of the fiber of Φ_D over B with the strictly semi-stable locus. By semi-stability it is easy to see that these points correspond to the partitions of the 6 points of B in complementary subsets of 3 elements each. We have ten of them. As stated above, a cubic three-fold cannot have more than ten ordinary double points and the Segre cubic is uniquely defined by this singular locus up to projective equivalence. \square

Also in this case, if $B \in \Delta_D$ then the intersection $\overline{\Phi_D^{-1}(B)} \cap \text{Kum}(C)$ is set-theoretically a finite set of points of cardinality strictly smaller than 10: the singular locus seems to degenerate. It is tempting, as in the case of Igusa quartics, to say that some of these points have multiplicity bigger than one and we obtain *degenerate Segre cubics* over Δ_D .

As we have already remarked, in the case of \mathcal{C}_4 the polar map is also well known and described. Let Y_i be the coordinates on $\mathbb{P}^7 \cong |2\Theta|$ and $G(Y_0 : \dots : Y_7)$ the quartic equation defining \mathcal{C}_4 , then the (self) polar rational map of \mathcal{C}_4 is defined in

the following way.

$$\mathcal{D}_4 : \mathcal{C}_4 \dashrightarrow \mathcal{C}_4;$$

$$x \mapsto \left[\frac{\partial G}{\partial Y_0} : \cdots : \frac{\partial G}{\partial Y_7} \right].$$

Let $B \in |2D| - \Delta_D$ and let \mathbb{P}_B^4 be the linear span of the point corresponding to B and of \mathbb{P}_c^3 . It turns out that the restriction of \mathcal{D}_4 to \mathbb{P}_B^4 behaves in a way very similar to the case of the sextic (see Proposition 7.2). Let $S_{3B} \subset \mathbb{P}_B^4$ be the Segre cubic such that $\mathcal{C}_4 \cap \mathbb{P}_B^4 = S_{3B} \cup \mathbb{P}_c^3$. We denote by J the set of 10 nodes of S_{3B} . Then the linear series $|\mathcal{I}_J(2)|$ on S_{3B} is the polar system of the Segre cubic.

Proposition 7.4. *The restricted dual map $\mathcal{D}_4|_{S_{3B}}$ is given by a linear system $|\mathcal{D}_{S_3}|$ that contains $|\mathcal{I}_J(2) + H|$ as a linear subsystem.*

As in the case of \mathcal{C}_6 this implies that we have a canonical way to construct the birational map of the following corollary via the polar map \mathcal{D}_4 .

Corollary 7.4. *The Coble quartic is birational to a fibration in Igusa quartics over \mathbb{P}^3 .*

Acknowledgments

The inspiration of this work came from a conversation that the first author had with Norbert Hoffmann. This work also benefitted from remarks and discussions with Christian Pauly, Alessandro Verra and Angela Ortega. Arnaud Beauville brought to our attention some existing literature. Quang Minh and Cristian Anghel kindly let us have copies of their papers. Duco Van Straten and Edoardo Sernesi gave us good advice about the deformation theory of singular projective hypersurfaces. Finally, we acknowledge the referee for his careful work and suggestions.

References

- [1] A. Alzati and M. Bolognesi, A structure theorem for $SU_C(2)$ and the moduli of pointed genus zero curves, preprint (2009), [hyyp://arxiv.org/0903.5515](http://arxiv.org/0903.5515), 1–20.
- [2] C. Anghel, Fibrés vectoriels semi-stables sur une courbe de genre deux et association des points dans l'espace projectif, *Serdica Math. J.* **30**(2–3) (2004) 103–110.
- [3] E. Arbarello, M. Cornalba, P. A. Griffiths and J. Harris, *Geometry of Algebraic Curves*, Vol. I, Grundlehren der Mathematischen Wissenschaften, Vol. 267 (Springer-Verlag, New York, 1985).
- [4] A. Beauville, Vector bundles and theta functions on curves of genus 2 and 3, *Amer. J. Math.* **128**(3) (2006) 607–618.
- [5] A. Beauville, M. S. Narasimhan and S. Ramanan, Spectral curves and the generalized theta divisor, *J. Reine Angew. Math.* **398** (1989) 169–179.
- [6] A. Bertram, Moduli of rank-2 vector bundles, theta divisors, and the geometry of curves in the projective space, *J. Differential Geom.* **35**(2) (1992) 429–469.
- [7] M. Bolognesi, A conic bundle degeneration on the Kummer surface, *Math. Z.* **261**(1) (2009) 149–168.

- [8] S. B. Bradlow, O. Garcia-Prada, V. Mercat, V. Muñoz and P. E. Newstead, Moduli spaces of coherent systems of small slope on algebraic curves, *Comm. Algebra* **37**(8) (2009) 2649–2678.
- [9] S. B. Bradlow, O. Garcia-Prada, V. Muñoz and P. E. Newstead, Coherent systems and Brill–Noether theory, *Internat. J. Math.* **14**(7) (2003) 683–733.
- [10] S. Brivio and A. Verra, The theta divisor of $SU_C(2, 2d)^s$ is very ample if C is not hyperelliptic, *Duke Math. J.* **82** (1996) 503–552.
- [11] S. Brivio and A. Verra, The Brill–Noether curve of a stable vector bundle on a genus two curve, in *Algebraic Cycles and Motives*, Vol. 2, London Mathematical Society Lecture Note Series, Vol. 344 (Cambridge University Press, Cambridge, 2007), pp. 73–93.
- [12] S. Brivio and A. Verra, Plucker forms and the theta map, to appear in *Amer. J. Math.*
- [13] A. B. Coble, *Algebraic Geometry and Theta Functions*, Colloquium Publications, Vol. 10 (American Mathematical Society, Providence, RI, 1982). Reprint of the 1929 edition.
- [14] I. Dolgachev, Abstract configurations in algebraic geometry, in *The Fano Conference* (Università di Torino, Turin, 2004), pp. 423–462.
- [15] I. Dolgachev and D. Ortland, *Point Sets in the Projective Spaces and Theta Functions*, Asterisque, Vol. 165 (Société Mathématique de France, 1988).
- [16] J.-M. Drezet and M. S. Narasimhan, Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques, *Invent. Math.* **97**(1) (1989) 53–94.
- [17] D. Eisenbud and S. Popescu, The projective geometry of the Gale transform, *J. Algebra* **230**(1) (2000) 127–173.
- [18] B. Hunt, *The Geometry of Some Special Arithmetic Quotients*, Lecture Notes in Mathematics, Vol. 1637 (Springer, Berlin, 1996).
- [19] B. Hunt, Nice modular varieties, *Experiment. Math.* **9**(4) (2000) 613–622.
- [20] J.-I. Igusa, On the graded ring of theta-constants, *Amer. J. Math.* **86** (1964) 219–246.
- [21] E. Izadi and L. Van Geemen, The tangent space to the moduli space of vector bundles on a curve and the singular locus of the theta divisor of the Jacobian, *J. Algebr. Geom.* **10**(1) (2001) 133–177.
- [22] A. D. King and P. E. Newstead, Moduli of Brill–Noether pairs on algebraic curves, *Internat. J. Math.* **6**(5) (1995) 733–748.
- [23] A. King and A. Schofield, Rationality of moduli of vector bundles on curves, *Indag. Math. (N.S.)* **10**(4) (1999) 519–535.
- [24] K. Koike, Remarks on the Segre cubic, *Arch. Math. (Basel)* **81**(2) (2003) 155–160.
- [25] Y. Laszlo, Local structure of the moduli space of vector bundles over curves, *Comment. Math. Helv.* **71**(3) (1996) 373–401.
- [26] D. Mumford and P. Newstead, Periods of a moduli space of vector bundles over curves, *Amer. J. Math.* **90** (1968) 1200–1208.
- [27] M. S. Narasimhan and S. Ramanan, Moduli of vector bundles on a compact Riemann surface, *Ann. of Math. (2)* **89** (1969) 14–51.
- [28] M. S. Narasimhan and S. Ramanan, Vector bundles on curves, in *Algebraic Geometry* (Oxford University Press, London, 1969), pp. 335–346.
- [29] Q. M. Nguyen, Vector bundles, dualities and classical geometry on a curve of genus two, *Internat. J. Math.* **18**(5) (2007) 535–558.
- [30] Q. M. Nguyen and S. Rams, On the geometry of the Coble–Dolgachev sextic, *Le Matematiche (Catania)* **58**(2) (2003), (2005) 257–275.
- [31] A. Ortega, On the moduli space of rank 3 vector bundles on a genus 2 curve and the Coble cubic, *J. Algebr. Geom.* **14**(2) (2005) 327–356.

- [32] W. Oxbury and C. Pauly, $SU_C(2)$ -Verlinde spaces as theta spaces on Pryms, *Internat. J. Math.* **7**(3) (1996) 393–410.
- [33] W. Oxbury and C. Pauly, Heisenberg invariant quartics and $SU_C(2)$ for a curve of genus four, *Math. Proc. Cambridge Philos. Soc.* **125**(2) (1999) 295–319.
- [34] C. Pauly, Self-duality of Coble’s quartic hypersurface and applications, *Michigan Math. J.* **50**(3) (2002) 551–574.
- [35] C. Pauly, Rank four vector bundles without theta divisor over a curve of genus two, to appear in *Adv. Geom.*
- [36] M. Raynaud, Sections des fibrés vectoriels sur une courbe, *Bull. Soc. Math. France* **110**(1) (1982) 103–125.
- [37] A. Vistoli, Grothendieck topologies, fibered categories and descent theory, in *Fundamental Algebraic Geometry*, Mathematical Surveys Monographs, Vol. 123 (American Mathematical Society, Providence, RI, 2005), pp. 1–104.