



US 20130041013A1

(19) **United States**

(12) **Patent Application Publication**
Lavitrano et al.

(10) **Pub. No.: US 2013/0041013 A1**

(43) **Pub. Date: Feb. 14, 2013**

(54) **ISOFORM OF BRUTON'S TYROSINE KINASE (BTK) PROTEIN**

Publication Classification

(75) Inventors: **Marialuisa Lavitrano**, Milano (IT);
Emanuela Grassilli, Monza (IT);
Kristian Helin, Charlottenlund (DK)

(51) **Int. Cl.**
A61K 31/713 (2006.01)
G01N 33/574 (2006.01)
C12Q 1/68 (2006.01)
A61P 35/00 (2006.01)

(73) Assignee: **BIONSIL S.R.L.**, Milano (IT)

(52) **U.S. Cl.** **514/44 A**; 435/7.4; 435/6.12

(21) Appl. No.: **13/532,677**

(57) **ABSTRACT**

(22) Filed: **Jun. 25, 2012**

The use of compounds is described which are capable of functionally blocking at least one of the genes chosen from the group composed of EphA1, EphA2, EphA8, EphB2, CSF1R, VEGFR2, RAMP2, RAMP3, CLRN1, MAPK4, PIK3C2A, PIK3CG, GSK3alpha, GSK3beta, IRAK3, DAPK1, JAK1, PIM1, TRB3, BTG1, LATS1, LIMK2, MYLK, PAK1, PAK2, CDC2, BTK, PNRC2, NCOA4, NR2C1, TPR, RBBP8, TRPC7, FXYD1, ERNI, PRSS16, RPS3, CCL23 and SERPINE1, for the manufacture of a medicament destined to diminish the resistance to chemotherapeutic drugs in the therapeutic treatment of epithelial tumour pathologies. Also described is a method for the determination of the drug resistance in tumour cells, as well as a method for the identification of tumour stem cells.

Related U.S. Application Data

(62) Division of application No. 12/531,061, filed on Sep. 25, 2009, now Pat. No. 8,232,085, filed as application No. PCT/EP2008/053099 on Mar. 14, 2008.

Foreign Application Priority Data

Mar. 14, 2007 (IT) PD2007A000088
Apr. 13, 2007 (EP) EP07106119.6




Fig. 1




Fig. 2a




Fig. 2b




Fig. 3a

Fig. 3b

Fig. 3c




Fig. 4a

Fig. 4b




Fig. 13




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10a





Fig. 10b

Fig. 11






Fig. 12a




Fig. 12b






Fig. 14



CLUSTAL 2.0.3 multiple sequence alignment

```

NM_000061 alternative -----
TTTGGTGGACTCTGCTACGTAGTGGCGTTCAGTGAAGGGAGCAGTGTTTTCCAGATC 60

NM_000061 alternative -----
CTCTGGCCTCCCCGTCCCCGAGGGAAGCCAGGACTAGGGTCGAATGAAGGGTCTCCAC 120

NM_000061 alternative -----CTCAGACTG 9
CTCCACGTTCCATTCTCTGTTCCACCTCAAGGTCAGTGGGAACACCTTTCGCAGCAAAC 180
** ****

NM_000061 alternative TCCTTCCTCTCTGGACTGTAAGAATATGTCTCCAGGGCCAGT--GTCTGCTGCG-ATCGA 66
--CTAATTCATGAAGACCTGGAGGAGCCAATTGTCCAGTTCATCACATGGCCA 238
** * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *


NM_000061 alternative GTCCACCTTCCAAGTCTGGCATCTCAATGCATCTGGGAAGCTACCTGCATTAAGTCAG 126
GTTGGTCCATTCAACAAATGGTTATTGGATGC--CCATTATGTGGCAGGCACTGTCCGG 296
** * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *

NM_000061 alternative GACTGAGCACACAGGTGAACCTCCAGAAAGAAGAAGCTATGGCCGAGTGATTCTGGAGAG 186
GGGAGAGCACACAGGTGAACCTCCAGAAAGAAGAAGCTATGGCCGAGTGATTCTGGAGAG 356
* * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *
    
```

Fig. 17

FIG. 18

BLAST EXONS 1-5 NM_000061 vs GENOMIC



BLAST EXONS 1-5 ALTERNATIVE BTK vs GENOMIC





FIG. 19



ISOFORM OF BRUTON'S TYROSINE KINASE (BTK) PROTEIN

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application is a division of U.S. Ser. No. 12/531,061, filed Sep. 25, 2009, which is a national stage filing under 35 U.S.C. 371 of International Application PCT/EP2008/053099, filed on Mar. 14, 2008, which claims priority from Italian Application No. PD2007A000088, filed on Mar. 14, 2007, and EP Application No. 07106119.6, filed on Apr. 13, 2007. The entire content of each of the referenced Application is incorporated herein by reference. International Application PCT/EP2008/053099 was published under PCT Article 21(2) in English.

TECHNICAL FIELD

[0002] The present invention regards the use of compounds for the production of a medicament capable of modulating, and in particular diminishing the drug resistance in human epithelial tumour cells, according to the characteristics stated in the preamble of the main claim. It is also intended for a determination method of the drug resistance in tumour cells as well as a method for the identification of tumour stem cells.

TECHNOLOGICAL BACKGROUND

[0003] One of the most commonly followed strategies in the therapy of neoplastic pathologies foresees the use of drugs (chemotherapy) capable of damaging the DNA of the tumour cells, so as to induce the natural process of apoptosis in the same.

[0004] Nevertheless, it is known that the tumour cells may respond in an unexpected manner to the drug therapy, showing, on the contrary, a strong resistance to the same. It is also known that one of the main reasons for the drug resistance shown by the tumour cells is the incapacity of the cells to begin the process of apoptosis, even in the presence of considerable DNA damage.

[0005] This phenomenon was traced back to a functional alteration (mutation or deletion) of the gene p53, which is no longer capable of initiating the process of cellular apoptosis, thus leading the cells to resist the drug action. The drug resistance levels of the tumour cells can be very high. For example, in the case of tumour cells of the colon-rectum in advanced phase, a drug therapy based on 5-fluorouracil (5FU) shows an effective response only for 10-15% of the cells, and a combination of 5FU with new drugs such as irinotecan and oxaliplatin leads to an increase of the cell mortality up to 40-50%, a value which is still entirely unsatisfactory for an effective therapeutic action towards the neoplastic pathologies.

[0006] In recent years, a cell phenomenon was discovered called "RNA interference" (RNAi) by means of which the gene expression is silenced in a specific manner. By taking advantage of such process, it is possible to obtain the selective silencing of genes with unknown function, thus permitting the definition of its specific function through the study of the obtained phenotype. By applying RNAi techniques and studying the phenotypic results, it is moreover possible to assign new functions to already known genes.

[0007] The genes involved in the phenomenon of drug resistance of the tumour cells are today largely unknown.

[0008] There is very much the need, therefore, to identify new genes involved in the drug-resistance and to design methods and compounds capable of substantially diminishing the tumour cells' resistance to drugs.

DESCRIPTION OF THE INVENTION

[0009] The problem underlying the present invention is that of making available compounds capable of reducing the drug resistance of the tumour cells of epithelial type, in order to permit the manufacture of medicaments destined for the therapy of related neoplastic pathologies.

[0010] This problem is resolved by the present finding by means of the use of compounds capable of functionally blocking one or more genes chosen from the group identified in the attached claims.

[0011] In a second aspect thereof, this invention moreover provides a method for the determination of the drug resistance of epithelial tumour cell lines.

[0012] In a further aspect thereof, this invention moreover provides a method for the identification of tumour stem cells.

BRIEF DESCRIPTION OF THE DRAWINGS

[0013] The characteristics and advantages of the invention will be clearer from the detailed description which follows of the tests and results which have led to its definition, reported with reference to the drawing set wherein:

[0014] FIG. 1 is a graph illustrating the results of inhibition of the capacity to form drug-resistant tumour cell colonies of the colon treated with the compounds according to the invention and a chemotherapeutic drug,

[0015] FIGS. 2a and 2b are graphs showing the results of an analysis of reversion of the resistance of a tumour cell line to two different chemotherapeutic drugs by means of silencing of the alpha and beta isoforms of the gene GSK3,

[0016] FIGS. 3a-3c are graphs illustrating the results of the tests of reversion of the resistance to a chemotherapeutic drug of three different tumour cell lines of the colon by means of silencing of the gene GSK3beta,

[0017] FIG. 4a is a graph illustrating the results of the tests of reversion of the resistance to a different chemotherapeutic drug of two different tumour cell lines of the colon by means of silencing of the gene GSK3beta,

[0018] FIG. 4b is a graph illustrating the results of an analysis of reversion of the resistance of a tumour cell line to the combination of two different chemotherapeutic drugs by means of the silencing of the beta isoform of the gene GSK3,

[0019] FIG. 5 is a collection of images illustrating the location of cytochrome C for 5FU-induced cell death in the absence of GSK3.

[0020] FIG. 6 is a graph illustrating the results of caspase 3 and caspase 7 activation assays in response to 5FU treatment of HCT116 and GSK3alpha, GSK3beta gene silenced HCT116p53KO colon carcinoma cells,

[0021] FIG. 7 is a collection of images illustrating the translocation of AIF to the nucleus during 5FU-induced cell death in the absence of GSK3,

[0022] FIG. 8 is a graph illustrating the reversion of the resistance to a chemotherapeutic drug of a tumour cell line carried out by means of functional blocking of the gene BTK obtained with different methods,

[0023] FIG. 9 is a graph illustrating the percentage of cell death induced by 5FU upon BTK overexpression in HCT116 colon carcinoma cells,

[0024] FIGS. 10*a* and 10*b* are graphs illustrating the reversion of the resistance to chemotherapeutic drugs by means of the functional blocking of the gene BTK on different colon tumour cell lines,

[0025] FIG. 11 is a western blot analysis showing BTK expression in several tumoral cell lines derived from different epithelial cancers,

[0026] FIGS. 12*a* and 12*b* are graphs illustrating the reversion of the resistance to chemotherapeutic drugs by means of the functional blocking of the gene BTK on epithelial tumour cell lines other than colon,

[0027] FIG. 13 is collection of images illustrating the cytoplasmic accumulation of cytochrome C in 5FU-treated cells upon BTK inhibition,

[0028] FIG. 14 is a graph illustrating the results of fluorimetric caspase activation assays after 72 hrs and 96 hrs in HCT116 and HCT116p53KO colon carcinoma cells treated with 5FU, BTK inhibition, or the combination of the two,

[0029] FIG. 15*a* is western blot image showing that antibody sc-1696 specifically recognizes BTK protein and that in epithelial carcinoma cells (HCT116p53KO) BTK has an apparent molecular weight of ~67 kDa and that the same form of the protein is present, together with the “classic” 77 kDa form, also in leukemic cells (Nalm6),

[0030] FIG. 15*b* is an immunoprecipitation-western blot image confirming that also a different antibody (BL7) identify BTK protein as being ~67 kDa in HCT116p53KO cells and that this isoform is present in leukemic cells (Nalm6) together with the “classic” 77 kDa form,

[0031] FIG. 16 is an image illustrating the results of a PCR experiment showing that the 5' end upstream of nucleotide 202 in the BTK-encoding mRNA from HCT116p53KO cells is absent or different from the 5' end of BTK mRNA from peripheral blood mononuclear cells (PBMC),

[0032] FIG. 17 is an image illustrating the results of ClustalW alignment of BTK transcript, identified by GenBank Accession #NM_000061, and the novel transcript identified in HCT116p53KO cells by 5'RACE PCR followed by cloning and sequencing (indicated as alternative), the box outlining the part of BTK sequence (starting from nucleotide 131 of the known transcript) common between BTK mRNA deriving from #NM_000061 and the novel BTK transcript found in HCT116p53KO,

[0033] FIG. 18 is an image illustrating the results of a Blast alignment vs a genomic database of exons 1-5 of BTK transcript identified by GenBank Accession # NM_000061 and the same exons of the novel BTK transcript, the dots indicating the position on chromosome X of the different exons of BTK,

[0034] FIG. 19 is an image illustrating the results of a nested PCR experiment showing that the novel BTK transcript is expressed in Formalin-Fixed Paraffin-Embedded (FFPE) tumoral tissues from coloncarcinoma patients.

IDENTIFICATION METHODOLOGY OF THE GENES MODULATING THE DRUG RESISTANCE IN EPITHELIAL TUMOUR CELLS AND RELATED VALIDATION TESTS

[0035] The abovementioned technical problem was tackled by subjecting several epithelial tumour cell lines to a series of tests and analyses geared to identify and characterise the genes capable of giving rise to the phenotypic expression of interest, i.e. the reversion of the resistance to the apoptosis induced by chemotherapeutic drugs.

[0036] Such identification was conducted by means of phenotypic screening of a representative cell line of the epithelial tumour cells following the selective silencing of an extended group of genes by means of RNAi.

[0037] This complex screening work was made possible due to a library of retroviral pRetroSuper (pRS) vectors recently made by the Bernards laboratories. It is known that such vectors are capable of expressing a specific oligonucleotide molecule in a stable manner, known as small interference RNA, in brief siRNA, capable of blocking the translation process of a specific messenger RNA (mRNA) in the corresponding protein. Such mechanism, in brief, foresees that the siRNA molecule (or rather a filament thereof) is associated with the RISC enzymatic complex, activating it, such that the latter can recognise and bind to the mRNA complementary to the siRNA associated thereto and then degrade it. It follows that the mRNA, identified in a specific manner by the siRNA, cannot be translated in the corresponding amino acid chain, thus obtaining the silencing of the gene from which the mRNA was transcribed. The interference process is specific, so that a siRNA molecule is normally capable of degrading only one mRNA and therefore silencing one single gene. On the other hand, it is instead possible that the same mRNA can be degraded, via RISC, by different siRNA.

[0038] This mechanism is one of the different possible modes for functionally blocking a gene (functional knock-out). However, the consequent degradation of siRNA via RISC produces only a transient effect on protein levels, thus allowing only short term experiments whose results may be not relevant in the long term. To overcome this problem, short hairpin RNAs (shRNAs), a sequence of RNA folded in the shape of a hairpin, were used as a mechanism for functional knockout. In brief, a vector is used to introduce and incorporate shRNA into the cell chromosome. Transcription of the DNA produces shRNA which is subsequently cleaved by the cellular machinery, DICER, into siRNA. The siRNA then functions as previously mentioned. The incorporation of the shRNA vector into the cell chromosome allows for the gene silencing to be inherited by daughter cells. Thus the usage of shRNAs allows for functional knock out cells that can be used for long term experiments.

[0039] It is also known in the literature that the efficiency of transfection for siRNAs is never 100%. Therefore, the use of transfected siRNAs does not assure that all treated cells are successfully deprived of a target protein. The absence of a selectable marker further creates a problem in working with homogeneous populations of cells. Using a selectable marker, such as a puromycin resistance gene, along with a retroviral library, allows for recovery of only the cells bearing shRNAs. Moreover, following shRNA expression selection, only genes whose silencing is compatible with normal cell survival and proliferation are selected for. The absence of these genes does not influence normal cell physiology but only the response to anticancer drugs, a very important consideration while developing anticancer therapy.

[0040] It is nevertheless important to specify that the same effect can, in general, be reached by acting in any other step of a gene's protein coding process, such as for example the step of transcription of the gene in mRNA, or the step of mRNA transduction, or by means of inhibition of the protein resulting from the coding process.

[0041] It is evident that the essential and critical step in the resolution of the problem is represented by the identification of the responsible gene or genes of the desired phenotype.

[0042] The retroviral library arranged by the Bernards laboratories consists of about 25,000 different elements, capable of silencing about 8,300 genes of the human genome, with a ratio of about 3 different vectors for each gene.

[0043] The tests were initially conducted *in vitro* and subsequently validated *ex vivo* on different epithelial tumour cell lines, characterised by the lack of or by the mutation of the gene p53 and hence provided with a marked resistance to chemotherapeutic drugs.

[0044] In detail, a human RNAi library (NKi library) was established, consisting of 8,300 targeted genes for silencing. The targeted genes included kinases, phosphatases, oncogenes, tumor suppressors, transcription factors, genes involved in transformation, metastasis, cell cycle, differentiation, apoptosis, metabolic and anabolic processes. A protocol was followed similar to that mentioned in a prior publication (Berns et al., NATURE vol. 428, 25 March 2004). The contents of this publication are incorporated by reference into this application. The mRNA sequence for each targeted gene was selected from UniGene. The sequences were masked using RepeatMasker to remove repetitive sequences and searched with NCBI BLAST against UniVec to mask for vector contamination. Three different 19 nucleotide (19-mer) sequences for silencing each targeted gene were designed, for a total of approximately 25,000 59-mer oligonucleotides that specify short hairpin RNAs (shRNAs). The 19-mer sequences were selected using a selection criteria as mentioned in the Berns publication wherein, a) there were no stretches of four or more consecutive T or A residues (to avoid premature polymerase III transcription termination signals); b) to have 30-70% overall GC content; c) to lie within the coding sequence of the target gene; d) to begin with a G or C residue (consistent with recently established rules for strand bias); e) to begin after an AA dimer in the 5' flanking sequence; f) to end just prior to a TT, TG or GT doublet in the 3' flanking sequence; g) to not contain XhoI or EcoRI restriction enzyme sites to facilitate subsequent shuttling of the knockdown cassette into vector backbones; h) to share minimal sequence identity with other genes; i) to target all transcript variants represented by RefSeq mRNAs; and j) to not overlap with other 19-mers selected from the same target sequence. The 59-mer oligonucleotides were designed as to contain a 19-mer sequence, its complimentary 19-mer sequence, pol_III transcription initiation site, pol_III termination site, and HindIII/BglIII cloning sites. Utilizing the HindIII/BglIII cloning sites, the oligonucleotides were ligated into pRetroSuper (pRS) retroviral vectors, which included a selection cassette for puromycin resistance. The DNA from the three different vectors that targeted the same gene was pooled and virus was produced to infect target cells.

EXAMPLE 1

[0045] The tumour cell line initially used was HCT116p53KO (which differs from the wild type wt HCT116 due to the lack of gene p53), related to the colon tumour, while further detailed studies of specific genes were carried out on other drug-resistant tumour cell lines of the colon, such as DLD-1 and SW480, as well as on other tumour cell lines of the lung and ovary. In particular, all cell lines, object of the analysis, were related to tumours of epithelial type.

[0046] Preliminarily, cell lines HCT116p53KO, DLD-1 and SW480 were treated with common chemotherapeutic drugs in order to confirm their resistance to the drug-induced apoptosis.

[0047] The chemotherapeutic drugs usable in accordance with the present invention can be of any type suitable for inducing the apoptosis process in the affected tumour cells, such as for example an antimetabolite or any DNA-damaging agent comprising the inhibitors of the topoisomerase I, inhibitors of the topoisomerase II, the platinum coordination compounds and alkylating agents.

[0048] The aforesaid preliminary tests have shown how, after treatment for 72 hours in 200 μ M 5FU, the cell mortality was less than 10%, against a wt HCT116 cell mortality of greater than 95%.

[0049] Supplementary tests of colony forming assays (CFA), have moreover demonstrated how such drug resistance was of non-transitory type. Once the resistance to apoptosis induced by chemotherapeutic drugs was confirmed, 200 \times 10⁶ HCT116p53KO cells were infected with the above-identified pRetroSuper library, provided by the Bernards laboratories. Each vector of this library was advantageously equipped with a selection cassette for a puromycin resistance gene, so that it was possible to select the HCT116p53KO cells actually infected by the vectors of the library through treatment with puromycin (2 mg/l in the culture medium for two days).

[0050] At the end of the antibiotic treatment, the still-living cells were then collected, which therefore comprised all of the cells infected by the retroviruses of the library whose silenced genes were not incompatible with the cellular survival.

[0051] The cells thus collected were then treated with 200 μ M 5FU for 72 hours, while at the same time wt HCT116 cells and uninfected HCT116p53KO cells were also subjected to the same treatment, as controls.

[0052] At the end of the treatment, it was found that about half of the cells were floating in the culture medium, therefore dead. Such cells represented the sought-after phenotype, so that they were collected and subjected to the necessary treatment for the identification of the genes silenced by the retroviral library.

[0053] In brief, such treatments comprised of the extraction of the DNA and the amplification by means of PCR (Polymerase Chain Reaction) of a region of 643 base pairs containing the region H1 of a specific promoter and the adjacent region coding the nucleotide sequence of interest. PCR amplification was performed using a pRS-fw primer: 5'-CCCTGAACTCCTCGTTCGACC-3' and pRS-rev primer: 5'-GAGACGTGCTACTTCCATTGTC-3'. The products obtained by the amplification were then bound in pRS retroviral vectors and the infection process of HCT116p53KO cells was completely repeated with the new vectors, so as to refine the screening. The products were digested with EcoRI/XhoI and recloned into pRS.

[0054] The products obtained from the second amplification treatment with PCR on the DNA harvested from the dead cells after new treatment with 5FU were newly isolated and bound in retroviral vectors pRS and then used for transforming DH5 α bacteria whose respective plasmids were sequenced for the identification of the genes which gave rise to the phenotypic expression of interest.

[0055] Once the single genes were obtained and identified whose silencing by means of interference with RNA gave rise

to the reversion of the resistance of the tumour cells tested to the 5FU, the separated and independent validation of the single genes took place.

[0056] Firstly, the validation was carried out in vitro on HCT116p53KO tumour cell samples, each of which separately infected with one of the previously-made plasmids, so as to silence in a specific and stable manner one of the genes indicated by the preceding screening and verify its capacity to modulate the resistance to drugs. The samples were then selected with puromycin, placed in Petri dishes at 50% confluence and treated with 200 μ M of 5FU for 12 hours.

[0057] The evaluation of the reversion of the drug resistance was carried out by means of observation of the formation of colonies according to the protocol defined by the CFA and their comparison with a wt HCT116 sample and an uninfected HCT116p53KO sample.

[0058] In FIG. 1, the results are reported in graph form which were obtained from this first in vitro validation. As clearly shown by the graph, a high percentage of identified genes are capable, when functionally blocked, of consistently reverting the colony forming ability of 5FU-treated HCT116p53KO cells. HCT116 and KO represent positive and negative controls.

[0059] This first validation has in particular permitted identifying a group of genes whose specific silencing has given rise to a 5FU-induced inhibition of the growth of the tumour colonies greater than 50% with respect to the HCT116p53KO sample.

[0060] These genes, themselves known and characterised, are listed below with their official symbol together with their identification number (between parentheses) reported on the NCBI Entrez Gene data bank: EphA1 (2041), EphA2 (1969), EphA8 (2046), EphB2 (2048), CSF1R (1436), VEGFR2 (3791), RAMP2 (10266), RAMP3 (10268), CLRN1 (7401), MAPK4 (5596), PIK3C2A (5286), PIK3CG (5294), GSK3beta (2932), IRAK3 (11213), DAPK1 (1612), JAK1 (3716), CHEK1 (1111), PIM1 (5292), TRB3 (57761), BTG1 (694), LATS1 (9113), LIMK2 (3985), MYLK (4638), PAK1 (5058), PAK2 (5062), CDC2 (983), BTK (695), PNRC2 (55629), NCOA4 (8031), NR2C1 (7181), TPR (7185), RBBP8 (5932), TRPC7 (57113), FXYD1 (5348), ERN1 (2081), PRSS16 (10279), RPS3 (6188), CCL23 (6368) and SERPINE1 (5054).

[0061] Among the above-listed genes, a first subgroup is also identifiable of genes whose silencing advantageously leads to an over 75% inhibition of the tumour cells growth.

[0062] Such first subgroup is formed by the following genes:

[0063] EphA1 (2041), EphA2 (1969), EphA8 (2046), EphB2 (2048), CSF1R (1436), VEGFR2 (3791), RAMP2 (10266), RAMP3 (10268), MAPK4 (5596), PIK3C2A (5286), PIK3CG (5294), GSK3beta (2932), IRAK3 (11213), DAPK1 (1612), JAK1 (3716), CHEK1 (1111), PIM1 (5292), TRB3 (57761), BTG1 (694), LATS1 (9113), LIMK2 (3985), BTK (695), PNRC2 (55629), NCOA4 (8031), NR2C1 (7181), TPR (7185), TRPC7 (57113), FXYD1 (5348), ERN1 (2081), RPS3 (6188) and SERPINE1 (5054).

[0064] In a more advantageous manner, a second subgroup was further identified of genes whose silencing advantageously leads to an over 95% inhibition of the growth of the tumour cells.

[0065] Such subgroup is formed by the following genes:

[0066] EphA1 (2041), EphA2 (1969), EphA8 (2046), RAMP3 (10268), PIK3C2A (5286), GSK3beta (2932),

IRAK3 (11213), DAPK1 (1612), CHEK1 (1111), PIM1 (5292), BTK (695), NCOA4 (8031), TPR (7185).

[0067] The gene GSK3alpha (2931), isoform of the gene GSK3beta, must be added to the above-listed genes; in separate tests whose results are shown by the graphs of FIGS. 2a and 2b it has shown an optimal efficiency in the reversion of the resistance both to 5FU and to the oxaliplatin in HCT116p53KO cells, entirely comparable to its beta isoform. In detail, FIG. 2a compares the percentages of cell deaths in the absence (symbol “-”) and presence (symbol “+”) of 200 μ M 5FU (72 hr treatment) upon wild type (wt) HCT116 cells, HCT116p53KO drug resistant cells, GSK3alpha and GSK3beta silenced gene cells. In comparison to the HCT116p53KO drug resistant cells, HCT116p53KO cells with GSK3alpha and GSK3beta genes silenced resulted in a high percentage of tumour cell death in the presence of 5FU. FIG. 2b compares the percentages of cell deaths in the absence (symbol “-”) and presence (symbol “+”) of 50 μ M oxaliplatin (72 hr treatment) upon wt HCT116 cells, HCT116p53KO drug resistant cells, GSK3alpha and GSK3beta silenced gene cells. In comparison to the HCT116p53KO drug resistant cells, HCT116p53KO cells with GSK3alpha and GSK3beta genes silenced also resulted in a high percentage of tumour cell death in the presence of oxaliplatin.

[0068] The formal confirmation of the silencing of the specific genes by means of interference was carried out through Western Blot analysis of the levels of the protein coded by them, if the antibody was commercially available (EphA1, EphA2, CSF1R, VEGFR, GSK3, JAK1, CHEK1, LIMK2, CDC2, BTK). Western Blot analysis was performed by lysing the puromycin-selected cells in E1A buffer (50 mM Hepes pH 8; 500 mM NaCl; 0.1% NP 40; DTT 1M; EDTA 1 mM). 8-12% SDS-polyacrylamide gel electrophoresis was used to separate 30 μ g of protein, which was later transferred to polyvinylidene difluoride membranes. Antibodies were then used to probe the Western Blots.

[0069] The effectiveness of the plasmids capable of silencing the genes belonging to the above-identified group was further tested on DLD-1 and SW480, other tumour cell lines of the colon known for possessing mutated p53 and for their resistance to drugs.

[0070] The capacity to diminish the resistance to chemotherapeutic drugs was generally confirmed, even with different performances. In particular, the inhibition percentage of the growth of colonies after drug treatment over all three cell lines was optimal when the following genes were silenced:

[0071] EphA1 (2041), EphA2 (1969), EphA8 (2046), EphB2 (2048), CSF1R (1436), VEGFR2 (3791), PIK3C2A (5286), PIK3CG (5294), GSK3alpha (2931), GSK3beta (2932), IRAK3 (11213), CDC2 (983), CHEK1 (1111), LATS1 (9113), TRB3 (57761), JAK1 (3716), BTK (695), PIM1 (5292), LIMK2 (3985), PAK2 (5062).

EXAMPLE 2

[0072] As an example, in FIGS. 3a-3c graphs are reported following silencing tests of the gene GSK3beta in the three tested tumour cell lines, in which the fraction of dead cells after the treatment with 5FU in samples infected with vectors capable of silencing the aforesaid gene is reported and compared with samples infected with empty vectors. In detail, FIG. 3a compares the percentages of cell deaths in the absence (symbol “-”) and presence (symbol “+”) of 200 μ M 5FU (72 hr treatment) upon wt HCT116 cells,

HCT116p53KO drug resistant cells, and GSK3beta silenced gene cells. In comparison to the HCT116p53KO drug resistant cells, HCT116p53KO cells with GSK3beta gene silenced resulted in a high percentage of tumour cell death in the presence of 5FU. FIG. 3b compares the percentages of cell deaths in the absence (symbol “-”) and presence (symbol “+”) of 200 μ M 5FU (72 hr treatment) upon wt DLD-1 cells and GSK3beta silenced gene DLD-1 cells. In comparison to the wt DLD-1 cells, GSK3beta silenced gene DLD-1 cells resulted in a high percentage of tumour cell death in the presence of 5FU. FIG. 3c compares the percentages of cell deaths in the absence (symbol “-”) and presence (symbol “+”) of 200 μ M 5FU (72 hr treatment) upon wt SW480 cells and GSK3beta silenced gene SW480 cells. In comparison to the wt SW480 cells, GSK3beta silenced gene SW480 cells resulted in a high percentage of tumour cell death in the presence of 5FU.

[0073] In addition to 5FU, representative example of the family of chemotherapeutic drugs of the antimetabolic type, the reversion of the resistance to drugs was also tested on chemotherapeutic drugs of different types, such as oxaliplatin.

[0074] In FIG. 4a, the fractions of dead cells induced by a treatment with oxaliplatin (50 μ M) are reported in samples of colon tumour cell lines DLD-1 and SW480, respectively infected with empty vectors and with vectors silencing the gene GSK3beta.

[0075] The substantial diminution of the resistance to the apoptosis induced by oxaliplatin in the sample in which GSK3beta was silenced is evident. In FIG. 4b, a graph is reported in which the results of an analogous test are indicated on the cell line SW480, in which the drug used was a combination of 5FU and oxaliplatin.

[0076] As mentioned previously and shown in FIGS. 2a and 2b, GSK3alpha silencing has the same effect as GSK3beta in modulating the apoptotic response to 5FU and oxaliplatin.

[0077] Further study was conducted to determine if 5FU-induced cell death in the absence of GSK3 was cytochrome C-dependent or independent. Utilizing anti-cytochrome C and DAPI staining, as shown in FIG. 5, it was found that 5FU-induced cell death in the absence of GSK3 is cytochrome C-independent. More specifically, utilizing anti-AIF and DAPI staining, as shown in FIG. 7, it was discovered that in the absence of GSK3, AIF translocates to the nucleus resulting in cell death. This was further supported by the finding that caspases 3 and 7 were not activated during 5FU-induced cell death in GSK3 silenced cells, as shown in FIG. 6.

EXAMPLE 3


[0078] Another particularly representative example of the above-identified gene group is constituted by the gene BTK, on which several investigations were undertaken.

[0079] BTK kinase is a cytoplasmic protein tyrosine kinase crucial for B-cell development and differentiation. BTK mutation is in fact responsible for X-linked agammaglobulinemia (XLA), a primary immunodeficiency mainly characterized by lack of mature B cells as well as low levels of immunoglobulins. In B cells BTK has been reported as having either pro-apoptotic or anti-apoptotic functions. Moreover, BTK has been so far assumed as being expressed only in some bone marrow-derived lineages such as B and mast cells, erythroid progenitors, platelets. Our finding that 18 BTK is a gene whose silencing reverts resistance to the cytotoxic

action of 5FU demonstrated for the first time that BTK is expressed also in cell types others than cells of the hematopoietic lineage. Firstly, the effectiveness was tested of the reversion of the resistance to drugs in HCT116p53KO tumour cells treated with an inhibitor compound of the protein BTK, such to demonstrate how the functional blocking of the gene of interest can be carried out in alternative ways to the silencing by means of RNAi.

[0080] The compound employed in these tests was (2Z)-2-cyan-N-(2,5-dibromophenyl)-3-hydroxy-2-butenamide, known as LFM-A13, whose structural formula is reported below in Formula 1.

Formula 1



[0081] In FIG. 8, the results of the comparison test are reported in a graph of the functional blocking of the BTK gene by means of plasmids, siRNA or LFM-A13 in samples of HCT116p53KO cells treated with 5FU. It can be easily noted how the obtained level of reversion to the drug resistance, expressed by means of the percentage fraction of dead cells, is entirely comparable. In detail, FIG. 8 compares the percentages of cell deaths in the presence of 200 μ M 5FU (72 hr treatment) upon wt HCT116 cells and HCT116p53KO drug resistant cells with or without (symbol “empty” and “-”) specific depletions of BTK following transient siRNA transfection (symbol “BTKi”), stable retroviral-mediated RNA interference (symbol “shBTK”), and using LFM-A13, an inhibitor compound of BTK.

[0082] According to the protective effect of BTK revealed by the above described inhibition experiments, BTK overexpression protects sensitive HCT116 wt from 5FU-induced cell death. In detail, FIG. 9 compares the percentages of cell deaths in the presence of 200 μ M 5FU (72 hr treatment) upon wt HCT116 cells infected with empty pBabe vector, wt HCT116 cells infected with pBabe BTK vector and HCT116p53KO drug resistant cells.

[0083] It was also determined that BTK inhibition reverts resistance also to oxaliplatin (FIG. 10a) and in DLD-1 (FIG. 10b).

[0084] The diminution of the resistance to chemotherapeutic drugs following the functional blocking of the gene BTK by means of LFM-A13 was further confirmed by tests conducted in vitro on epithelial tumour cell lines different from those of the colon. In FIG. 11 levels of BTK have been investigated, by means of Western blot, in several different epithelial carcinoma cell lines showing that the kinase is expressed in most of them. In particular, in FIGS. 12a and 12b, graphs were reported of the reversion tests obtained on SKOV cell lines (related to an ovarian tumour) and A549 cell lines (related to a lung tumour).

[0085] In detail, FIG. 12a compares the percentages of cell deaths in the absence (symbol “NT” and “-”) and presence of 200 μ M 5FU, 50 μ M OxPt, and the combination of the two upon resistant ovarian (SKOV) cells with or without BTK inhibition through the usage of LFM-A13. In comparison to the SKOV cells without LFM-A13, the SKOV cells with

LMFA13 resulted in a high percentage of tumour cell death. FIG. 12b compares the percentages of cell deaths in the absence (symbol “NT” and “-”) and presence of 200 μ M 5FU, 50 μ M OxPt, and the combination of the two upon resistant lung (A549) cells with or without BTK inhibition through the usage of LMFA13. In comparison to the A549 cells without LMFA13, the A549 cells with LMFA13 resulted in a high percentage of tumour cell death.

[0086] In both cases, it is noted how the treatment of the cell lines with LFM-A13, functional inhibitor of the BTK gene, leads to a considerable increase of the cellular mortality after exposure to 5FU or oxalipatin or both drugs in combination.

[0087] Moreover, the high effectiveness of the action of diminution of the drug-resistance also in these cell lines, confirms that the validity of the results obtained in the preceding tests can be extended at least to all types of epithelial tumour, such as lung tumour, ovarian tumour and breast tumour.

[0088] The optimal results pointed out above have suggested validating the gene BTK also through ex-vivo analysis on epithelial human tumour samples.

[0089] Firstly, it was verified through Western Blot that the BTK protein levels were high in 30% of the samples of ovarian tumour cells drawn from patients in advanced stage of disease and/or resistant to chemotherapeutic drugs.

[0090] Secondly, examinations were conducted on colon tumour stem cells isolated from patients in order to verify if the investigated genes were expressed (and in what measure) also in this cell type. In fact, according to recent studies (Dean et al., 2005), these stem cells would be the responsible principals of the drug resistance.

[0091] In all four tumour stem cell lines analysed, isolated from different patients, the expression of the protein BTK is very high, at least 4-5 times greater than the expression detected on the colon carcinoma cell lines used in the functional experiments, suggesting that the determination of the BTK levels can advantageously be used as a method for defining the stem cell properties of the tumour cells examined, and consequently also their resistance to chemotherapeutic drugs.

[0092] The results have shown that BTK levels determine the sensitivity of tumor cells to 5FU and that BTK inhibition reverses drug-resistance. Anti-cytochrome C immunostaining in 5FU-treated cells (FIG. 13) showed cytoplasmic accumulation after 5FU treatment in resistant cells only when BTK was inhibited, supporting the finding that reversal of drug-resistance upon BTK inhibition is due to activation of apoptosis. The same finding is also supported by the graph reported in FIG. 14, evaluating the level of caspase 3/7 activation upon 5FU treatment in HCT116p53KO resistant cells in the presence or in the absence of BTK inhibitor LFM-A13. High levels of caspase 3/7 activation, measured by means of a luminometric assay as RLU/number of cells, are observed in 5FU-treated HCT116p53KO cells only when BTK is inhibited.

[0093] The predicted and reported molecular weight of BTK protein is 77 kDa. The protein identified in western blot as BTK by a specific antibody (sc-1696, from Santa Cruz Biotechnology) in HCT116p53KO and in all other epithelial carcinoma cell lines tested (6 breast cell lines, 3 ovary cell lines, 7 lung cell lines, 5 colon cell lines), in contrast, has an apparent molecular weight around 65-68 kDa (FIG. 11, FIG. 15a) suggesting that in epithelial cell 22 lines a shorter BTK isoform is expressed.

[0094] To confirm these results an immunoprecipitation analysis was carried out using two different and specific BTK antibodies (the above cited sc-1696 and BL7, kind gift of Dr. Mike Tomlinson, University of Birmingham, UK). The results in FIG. 15b show that only this novel isoform is expressed in epithelial cell lines whereas in a leukemic cell line (Nalm6), already known from the literature to express the classical 77 kDa form, both isoforms are present.

[0095] Therefore, a bioinformatic analysis of BTK coding sequence (cds) has been carried out and, accordingly, a second nucleotide triplet ATG (nt 428-430), in frame with the one known to be used to translate traslatic BTK (nt 164-166) and susceptible to start the translation of a protein has been identified. The expected molecular weight of the putative protein translated starting from this second ATG is 67 kDa, consistent with the apparent molecular weight of the band identified by different BTK antibodies in epithelial carcinoma cells.

[0096] In order to identify which portion is missing in the mRNA coding for the shorter and novel isoform of the BTK protein, PCR experiments using different primers pairs (annealing at different parts of the cds as indicated in the upper diagram of FIG. 16) have been performed. As shown in FIG. 16, these experiments indicate that the 5' end, upstream of nucleotide 202, is absent or different in HCT116p53KO cells. 5'RACE/sequencing experiments have been performed on mRNA from HCT116p53KO cells in order to determine the identity of the unknown 5' end. Subsequently, alignment analysis between the cDNA derived from mRNA from HCT116p53KO cells and the cDNA deriving from the standard BTK mRNA, identified by GenBank accession #NM_000061, using ClustalW computer program (results shown in FIG. 17) demonstrated that the sequence upstream of the second exon (starting at nt 134 of the cds) is different from what reported in literature, i.e., epithelial colon carcinoma cells express a different first exon.

[0097] BLAST analysis using a genomic database (FIG. 18) localize the first exon of NM_000061 BTK transcript at 101160K on the chromosome X contig, in correspondence with the beginning of BTK locus. At variance, the first exon of the novel BTK transcript alignes 15192 by 5' of the first known BTK exon, immediately downstream of the RPL36A locus, suggesting that it corresponds to a hitherto unrecognized BTK exon. Moreover, this novel exon is present in HCT116p53KO instead of the first “classical” exon suggesting that this is an “alternative” first exon, whose usage gives raise to a different BTK mRNA, transcribed in cells expressing this shorter BTK isoform.

[0098] PCR/sequencing experiments demonstrated the expression of this “alternative” BTK mRNA not only in all coloncarcinoma cell lines tested (HCT116p53KO, DLD-1 and SW480) but also in 9/9 FFPE-samples from colon carcinoma patients (FIG. 19).

[0099] It should be noted that the first exon of BTK, as identified by GenBank accession # NM_000061, corresponds to the 5'UTR of the mRNA, being the ATG triplet encoding the first Met amino acid of BTK protein located in the second exon. 5'UTRs usually perform regulatory functions such as directing cap-dependent or IRES-mediated cap-independent translation. A different first exon, as identified in the novel transcript, can therefore dictate whether a different ATG (in this case an ATG located in the 4th exon) has to be used to start the translation of BTK protein, and therefore regulate the expression of different isoforms.

[0100] The nucleotide sequence of the first exon corresponding to the 5'UTR of the novel mRNA expressed by the BTK gene is reported in SEQ ID NO:1, attached to the present description.

[0101] The amino acid sequence of the novel isoform of BTK protein coded by the novel transcript is reported in SEQ ID NO:2, attached to the present description.

[0102] The above results suggest a method for determining the resistance of tumour cells to chemotherapeutic drugs as well as a method for the identification of the presence of tumour stem cells, wherein the expression of gene BTK comprises the steps of verifying the presence of the novel isoform of the BTK protein.

[0103] The presence of the novel isoform of the BTK protein may be controlled verifying the presence of the protein, for instance using western blot or immunoprecipitation or immunochemistry or immunofluorescence analysis, or, preferably, verifying the presence of the mRNA having the alternative first exon, whose cDNA shows the nucleotide sequence defined in SEQ ID NO:1. The latter may be advantageously carried out by means of PCR analysis, preferably using primers having a sequence included in SEQ ID NO:1.

[0104] This new method is expected to show relevant advantages, with respect to the known prior art, particularly when used for analysing tumoral tissues taken from human patients.

[0105] Actually, it is well known that tumoral tissues taken from human patients may contain an effective amount of lymphocytes which may also express BTK protein, thus disturbing the search of BTK protein expressed by the tumoral cells.

[0106] However, BTK protein expressed by lymphocytes is the "classical" isoform of BTK protein having a molecular weight of 77 KDa, so that the search of the novel isoform of BTK protein may be carried out without any interference, with a simple PCR analysis looking for the presence of BTK mRNA having the alternative first exon.

[0107] The present invention therefore resolves the above-lamented problem with reference to the mentioned prior art, offering at the same time numerous other advantages, including making possible the development of diagnostic methods capable of predicting the therapeutic response so to refine not only the diagnostics but above all direct the best therapeutic choice.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 3

<210> SEQ ID NO 1

<211> LENGTH: 300

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

ttttggtgga ctctgctacg tagtggcggt cagtgaaggg agcagtggtt ttcccagatc    60
ctctggcctc cccgtccccg agggaagcca ggactagggt cgaatgaagg ggtcctccac    120
ctccacgttc cattcctggt ccacctcaag gtcactggga acacctttcg cagcaaactg    180
ctaattcaat gaagacctgg agggagccaa ttgttccagt tcatctatca catggccagt    240
tggtcattc aacaaatggt tattggatgc ccattatgtg gcaggcactg ttccggggga    300

```

<210> SEQ ID NO 2

<211> LENGTH: 571

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Met Glu Gln Ile Ser Ile Ile Glu Arg Phe Pro Tyr Pro Phe Gln Val
 1           5           10          15
Val Tyr Asp Glu Gly Pro Leu Tyr Val Phe Ser Pro Thr Glu Glu Leu
          20          25          30
Arg Lys Arg Trp Ile His Gln Leu Lys Asn Val Ile Arg Tyr Asn Ser
          35          40          45
Asp Leu Val Gln Lys Tyr His Pro Cys Phe Trp Ile Asp Gly Gln Tyr
          50          55          60
Leu Cys Cys Ser Gln Thr Ala Lys Asn Ala Met Gly Cys Gln Ile Leu
          65          70          75          80
Glu Asn Arg Asn Gly Ser Leu Lys Pro Gly Ser Ser His Arg Lys Thr
          85          90          95
Lys Lys Pro Leu Pro Pro Thr Pro Glu Glu Asp Gln Ile Leu Lys Lys

```

-continued

100					105					110					
Pro	Leu	Pro	Pro	Glu	Pro	Ala	Ala	Ala	Pro	Val	Ser	Thr	Ser	Glu	Leu
	115						120					125			
Lys	Lys	Val	Val	Ala	Leu	Tyr	Asp	Tyr	Met	Pro	Met	Asn	Ala	Asn	Asp
	130					135					140				
Leu	Gln	Leu	Arg	Lys	Gly	Asp	Glu	Tyr	Phe	Ile	Leu	Glu	Glu	Ser	Asn
	145					150					155				160
Leu	Pro	Trp	Trp	Arg	Ala	Arg	Asp	Lys	Asn	Gly	Gln	Glu	Gly	Tyr	Ile
				165					170					175	
Pro	Ser	Asn	Tyr	Val	Thr	Glu	Ala	Glu	Asp	Ser	Ile	Glu	Met	Tyr	Glu
			180					185					190		
Trp	Tyr	Ser	Lys	His	Met	Thr	Arg	Ser	Gln	Ala	Glu	Gln	Leu	Leu	Lys
		195					200					205			
Gln	Glu	Gly	Lys	Glu	Gly	Gly	Phe	Ile	Val	Arg	Asp	Ser	Ser	Lys	Ala
	210					215					220				
Gly	Lys	Tyr	Thr	Val	Ser	Val	Phe	Ala	Lys	Ser	Thr	Gly	Asp	Pro	Gln
	225					230					235				240
Gly	Val	Ile	Arg	His	Tyr	Val	Val	Cys	Ser	Thr	Pro	Gln	Ser	Gln	Tyr
				245					250					255	
Tyr	Leu	Ala	Glu	Lys	His	Leu	Phe	Ser	Thr	Ile	Pro	Glu	Leu	Ile	Asn
			260					265					270		
Tyr	His	Gln	His	Asn	Ser	Ala	Gly	Leu	Ile	Ser	Arg	Leu	Lys	Tyr	Pro
		275					280					285			
Val	Ser	Gln	Gln	Asn	Lys	Asn	Ala	Pro	Ser	Thr	Ala	Gly	Leu	Gly	Tyr
		290					295					300			
Gly	Ser	Trp	Glu	Ile	Asp	Pro	Lys	Asp	Leu	Thr	Phe	Leu	Lys	Glu	Leu
	305					310					315				320
Gly	Thr	Gly	Gln	Phe	Gly	Val	Val	Lys	Tyr	Gly	Lys	Trp	Arg	Gly	Gln
				325					330					335	
Tyr	Asp	Val	Ala	Ile	Lys	Met	Ile	Lys	Glu	Gly	Ser	Met	Ser	Glu	Asp
			340					345					350		
Glu	Phe	Ile	Glu	Glu	Ala	Lys	Val	Met	Met	Asn	Leu	Ser	His	Glu	Lys
		355					360					365			
Leu	Val	Gln	Leu	Tyr	Gly	Val	Cys	Thr	Lys	Gln	Arg	Pro	Ile	Phe	Ile
		370					375					380			
Ile	Thr	Glu	Tyr	Met	Ala	Asn	Gly	Cys	Leu	Leu	Asn	Tyr	Leu	Arg	Glu
		385					390					395			400
Met	Arg	His	Arg	Phe	Gln	Thr	Gln	Gln	Leu	Leu	Glu	Met	Cys	Lys	Asp
				405					410					415	
Val	Cys	Glu	Ala	Met	Glu	Tyr	Leu	Glu	Ser	Lys	Gln	Phe	Leu	His	Arg
			420					425					430		
Asp	Leu	Ala	Ala	Arg	Asn	Cys	Leu	Val	Asn	Asp	Gln	Gly	Val	Val	Lys
		435					440					445			
Val	Ser	Asp	Phe	Gly	Leu	Ser	Arg	Tyr	Val	Leu	Asp	Asp	Glu	Tyr	Thr
		450					455					460			
Ser	Ser	Val	Gly	Ser	Lys	Phe	Pro	Val	Arg	Trp	Ser	Pro	Pro	Glu	Val
		465				470					475				480
Leu	Met	Tyr	Ser	Lys	Phe	Ser	Ser	Lys	Ser	Asp	Ile	Trp	Ala	Phe	Gly
				485					490					495	
Val	Leu	Met	Trp	Glu	Ile	Tyr	Ser	Leu	Gly	Lys	Met	Pro	Tyr	Glu	Arg
			500					505						510	

-continued

Phe Thr Asn Ser Glu Thr Ala Glu His Ile Ala Gln Gly Leu Arg Leu
515 520 525

Tyr Arg Pro His Leu Ala Ser Glu Lys Val Tyr Thr Ile Met Tyr Ser
530 535 540

Cys Trp His Glu Lys Ala Asp Glu Arg Pro Thr Phe Lys Ile Leu Leu
545 550 555 560

Ser Asn Ile Leu Asp Val Met Asp Glu Glu Ser
565 570

<210> SEQ ID NO 3
<211> LENGTH: 659
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Met Ala Ala Val Ile Leu Glu Ser Ile Phe Leu Lys Arg Ser Gln Gln
1 5 10 15

Lys Lys Lys Thr Ser Pro Leu Asn Phe Lys Lys Arg Leu Phe Leu Leu
20 25 30

Thr Val His Lys Leu Ser Tyr Tyr Glu Tyr Asp Phe Glu Arg Gly Arg
35 40 45

Arg Gly Ser Lys Lys Gly Ser Ile Asp Val Glu Lys Ile Thr Cys Val
50 55 60

Glu Thr Val Val Pro Glu Lys Asn Pro Pro Pro Glu Arg Gln Ile Pro
65 70 75 80

Arg Arg Gly Glu Glu Ser Ser Glu Met Glu Gln Ile Ser Ile Ile Glu
85 90 95

Arg Phe Pro Tyr Pro Phe Gln Val Val Tyr Asp Glu Gly Pro Leu Tyr
100 105 110

Val Phe Ser Pro Thr Glu Glu Leu Arg Lys Arg Trp Ile His Gln Leu
115 120 125

Lys Asn Val Ile Arg Tyr Asn Ser Asp Leu Val Gln Lys Tyr His Pro
130 135 140

Cys Phe Trp Ile Asp Gly Gln Tyr Leu Cys Cys Ser Gln Thr Ala Lys
145 150 155 160

Asn Ala Met Gly Cys Gln Ile Leu Glu Asn Arg Asn Gly Ser Leu Lys
165 170 175

Pro Gly Ser Ser His Arg Lys Thr Lys Lys Pro Leu Pro Pro Thr Pro
180 185 190

Glu Glu Asp Gln Ile Leu Lys Lys Pro Leu Pro Pro Glu Pro Ala Ala
195 200 205

Ala Pro Val Ser Thr Ser Glu Leu Lys Lys Val Val Ala Leu Tyr Asp
210 215 220

Tyr Met Pro Met Asn Ala Asn Asp Leu Gln Leu Arg Lys Gly Asp Glu
225 230 235 240

Tyr Phe Ile Leu Glu Glu Ser Asn Leu Pro Trp Trp Arg Ala Arg Asp
245 250 255

Lys Asn Gly Gln Glu Gly Tyr Ile Pro Ser Asn Tyr Val Thr Glu Ala
260 265 270

Glu Asp Ser Ile Glu Met Tyr Glu Trp Tyr Ser Lys His Met Thr Arg
275 280 285

Ser Gln Ala Glu Gln Leu Leu Lys Gln Glu Gly Lys Glu Gly Gly Phe
290 295 300

-continued

Ile Val Arg Asp Ser Ser Lys Ala Gly Lys Tyr Thr Val Ser Val Phe
305 310 315 320

Ala Lys Ser Thr Gly Asp Pro Gln Gly Val Ile Arg His Tyr Val Val
325 330 335

Cys Ser Thr Pro Gln Ser Gln Tyr Tyr Leu Ala Glu Lys His Leu Phe
340 345 350

Ser Thr Ile Pro Glu Leu Ile Asn Tyr His Gln His Asn Ser Ala Gly
355 360 365

Leu Ile Ser Arg Leu Lys Tyr Pro Val Ser Gln Gln Asn Lys Asn Ala
370 375 380

Pro Ser Thr Ala Gly Leu Gly Tyr Gly Ser Trp Glu Ile Asp Pro Lys
385 390 395 400

Asp Leu Thr Phe Leu Lys Glu Leu Gly Thr Gly Gln Phe Gly Val Val
405 410 415

Lys Tyr Gly Lys Trp Arg Gly Gln Tyr Asp Val Ala Ile Lys Met Ile
420 425 430

Lys Glu Gly Ser Met Ser Glu Asp Glu Phe Ile Glu Glu Ala Lys Val
435 440 445

Met Met Asn Leu Ser His Glu Lys Leu Val Gln Leu Tyr Gly Val Cys
450 455 460

Thr Lys Gln Arg Pro Ile Phe Ile Ile Thr Glu Tyr Met Ala Asn Gly
465 470 475 480

Cys Leu Leu Asn Tyr Leu Arg Glu Met Arg His Arg Phe Gln Thr Gln
485 490 495

Gln Leu Leu Glu Met Cys Lys Asp Val Cys Glu Ala Met Glu Tyr Leu
500 505 510

Glu Ser Lys Gln Phe Leu His Arg Asp Leu Ala Ala Arg Asn Cys Leu
515 520 525

Val Asn Asp Gln Gly Val Val Lys Val Ser Asp Phe Gly Leu Ser Arg
530 535 540

Tyr Val Leu Asp Asp Glu Tyr Thr Ser Ser Val Gly Ser Lys Phe Pro
545 550 555 560

Val Arg Trp Ser Pro Pro Glu Val Leu Met Tyr Ser Lys Phe Ser Ser
565 570 575

Lys Ser Asp Ile Trp Ala Phe Gly Val Leu Met Trp Glu Ile Tyr Ser
580 585 590

Leu Gly Lys Met Pro Tyr Glu Arg Phe Thr Asn Ser Glu Thr Ala Glu
595 600 605

His Ile Ala Gln Gly Leu Arg Leu Tyr Arg Pro His Leu Ala Ser Glu
610 615 620

Lys Val Tyr Thr Ile Met Tyr Ser Cys Trp His Glu Lys Ala Asp Glu
625 630 635 640

Arg Pro Thr Phe Lys Ile Leu Leu Ser Asn Ile Leu Asp Val Met Asp
645 650 655

Glu Glu Ser

1-49. (canceled)

50. A method for identifying the presence of tumour cells comprising the step of measuring the level of expression of mRNA and/or the isolated active protein of a GSK3 gene, wherein said gene is the GSK3alpha or the GSK3beta isoform.

51. A method for predicting the resistance of tumour cells to chemotherapeutic drugs comprising the step of measuring the level of expression of mRNA and/or of the isolated active protein of a GSK3 gene, wherein said gene is the GSK3alpha or the GSK3beta isoform.

52. A method for predicting the reversal of resistance of tumour cells to chemotherapeutic drugs comprising the steps of a) measuring the level of expression of mRNA and/or of the isolated active protein of a GSK3 gene, b) functionally and selectively blocking the gene corresponding to said protein, c) testing the percentage of tumour cell death in the presence of chemotherapeutic drug.

53. The method according to claim **52**, wherein said GSK3 gene is the GSK3alpha or the GSK3beta isoform.

54. The method according to claim **50**, wherein said tumour cells are of epithelial type.

55. The method according to claim **50**, wherein said tumour cells are stem cells.

56. A method for reversing resistance of a tumour cell or a tumour stem cell to a chemotherapeutic drug in a tumour of a subject, the method comprising

administering to the tumour of the subject a therapeutic amount of a composition comprising a small interfering RNA (siRNA) or a small hairpin RNA (shRNA),

wherein the small interfering RNA (siRNA) or the small hairpin RNA (shRNA) inhibits expression of messenger RNA (mRNA) encoding GSK3; and

wherein inhibition of the expression of the messenger RNA (mRNA) encoding GSK3 reverses resistance of the tumour cell or the tumour stem cell to the chemotherapeutic drug.

57. A pharmaceutical composition comprising a compound capable of functionally and selectively blocking the GSK3alpha and/or the GSK3beta and a pharmaceutically acceptable vehicle.

58. A pharmaceutical composition comprising a compound capable of functionally and selectively blocking an

expression of GSK3 and a chemotherapeutic drug for the treatment of an epithelial tumour.

59. The pharmaceutical composition according to claim **57**, wherein said GSK3 is a GSK3alpha or a GSK3beta.

60. The pharmaceutical composition according to claim **57**, wherein said compound capable of functionally blocking said GSK3 gene is an oligonucleotide molecule of small interference RNA (siRNA).

61. The pharmaceutical composition according to claim **57**, wherein said compound capable of functionally blocking said GSK3 is a compound capable of expressing an oligonucleotide molecule of small interference RNA (siRNA).

62. The pharmaceutical composition according to claim **57**, wherein said compound capable of functionally blocking said GSK3 gene is LiCl, tideglusib.

63. The method according to claim **58**, wherein the chemotherapeutic drug is an anti-metabolite agent.

64. The method according to claim **58**, wherein the chemotherapeutic drug is a DNA-damaging agent.

65. The method according to claim **64**, wherein the DNA-damaging agent is selected from the group consisting of a topoisomerase I inhibitor, a topoisomerase II inhibitor, a platinum coordination compound, an alkylating agent, and a combination thereof.

66. The method according to claim **58**, wherein the chemotherapeutic drug is 5-fluorouracil.

67. The method according to claim **58**, wherein the chemotherapeutic drug is oxaliplatin alone or in admixture with 5-fluorouracil (5FU).

68. A diagnostic kit for identifying the presence of tumour cells in human tissues comprising means for measuring the level of expression of a mRNA and/or the protein and/or different functional products (products phosphorylated in specific Serine or Tyrosine residues) of a GSK3 gene.

69. A diagnostic kit for predicting the resistance of tumour cells to chemotherapeutic drugs comprising means for measuring the level of expression of mRNA and/or the protein and/or different functional products (products phosphorylated in specific Serine or Tyrosine residues) of a GSK3 gene, in the presence of substances able to functionally and selectively block said gene and of a chemotherapeutic drug.

70. The diagnostic kit according to claim **69**, wherein said gene is the GSK3alpha or the GSK3beta isoform.

* * * * *