

COVERING OF ELLIPTIC CURVES AND THE KERNEL OF THE PRYM MAP

FILIPPO F. FAVALE AND SARA TORELLI

ABSTRACT. Motivated by a conjecture of Xiao, we study families of coverings of elliptic curves and their corresponding Prym map Φ . More precisely, we describe the codifferential of the period map P associated to Φ in terms of the residue of meromorphic 1-forms and then we use it to give a characterization for the coverings for which the dimension of $\text{Ker}(dP)$ is the least possible. This is useful in order to exclude the existence of non isotrivial fibrations with maximal relative irregularity and thus also in order to give counterexamples to the Xiao's conjecture mentioned above. The first counterexample to the original conjecture, due to Pirola, is then analysed in our framework.

INTRODUCTION

Hurwitz spaces were classically introduced and studied by Clebsch and Hurwitz (see [Cle72] and [Hur91]) as spaces parametrizing branched coverings of \mathbb{P}^1 . Nowadays, the term Hurwitz space refers to a variety which parametrizes, up to equivalence, coverings $\pi : F \rightarrow E$ of curves with some geometric restrictions. In this article we will use a local version of Hurwitz spaces, namely a *local family of coverings*, whose seminal idea can be found in [Kan04]. Roughly, given a fixed covering $\pi : F \rightarrow E$ where E is an elliptic curve, one is able to construct a map $p : \mathcal{F} \rightarrow \mathcal{E}$ of curves over \mathcal{H} , where \mathcal{H} is a contractible open set. Then \mathcal{H} is a parameter space for smooth coverings which share the same degree and the same ramification indices with π .

Attached to a local family of coverings $p : \mathcal{F} \rightarrow \mathcal{E}$ with parameter space \mathcal{H} there is the Prym map Φ , which associates to a $b \in \mathcal{H}$ the generalized Prym variety of $\pi_b = \pi|_{\mathcal{F}_b} : \mathcal{F}_b \rightarrow \mathcal{E}_b$, i.e. the connected component containing 0 of the kernel of the norm map $\text{Nm}(\pi_b)$. The Prym map is, in some sense, the analogous of the Torelli map T from \mathcal{M}_g , the moduli space of curves of genus g , to \mathcal{A}_g , the moduli space of principally polarized abelian varieties of dimension g .

A celebrated theorem, the infinitesimal Torelli theorem, states that the differential of the Torelli map is injective outside the hyperelliptic locus of \mathcal{M}_g and it should be interesting to have a similar theorem also for Prym maps or, at least, to their lifting P to a period domain. We will show that, in our case, i.e. when the base E is an elliptic curve, the dimension of the kernel of $d_o P^\vee$ is at least 1 as a consequence of how the local families that we will use are constructed. Roughly, by composing a covering with a traslation of the base we always have coverings with the same Prym, so there is a tangent direction in the parameter space along which the Prym map is constant. Hence a question analogous to the one answered by the infinitesimal Torelli is

Date: March 20, 2017.

2010 Mathematics Subject Classification. 14H40, 14H30, 14B10, 14J99.

Given a family of coverings with central fiber π , which conditions can we put on π in order to have that $\text{Ker}(d_\circ P)$ has dimension 1?

It is already known that an infinitesimal Torelli-like theorem for the Prym map cannot hold without restrictions as there are examples of coverings $\pi : F \rightarrow E$ (moreover with F non hyperelliptic) for which there are two independent directions along which $d_\circ P$ is 0. One of these examples, due to Pirola, will be analyzed in Section 4. This paper is devoted to the study of the Prym map $\Phi : \mathcal{H} \rightarrow \mathcal{A}$ in the cases for which \mathcal{H} parametrizes coverings over an elliptic curve.

A further motivation to study this kind of problems comes from a conjecture about fibered surfaces. Recall that, given a fibration $f : S \rightarrow B$ of a smooth compact surface S over a smooth compact curve B , the *relative irregularity* q_f is defined to be the difference $q(S) - g(B)$. A modified version of a conjecture of Xiao states that, if f is not isotrivial, then

$$(1) \quad q_f \leq \left\lceil \frac{g+1}{2} \right\rceil.$$

The original conjecture was without the round up and has been modified after a counterexample of Pirola, the one that we will present in Section 4. To have an insight of what is known about the relative irregularity and about recent results about an upper bound a good reference is [BGN15]. The link between the world of non isotrivial fibrations and the one of the families of coverings we will define is broadly given as follows. The fibration f induces a surjective map $\text{alb}(f) : \text{Alb}(S) \rightarrow \text{Alb}(B) = J(B)$ with $\dim(\text{Ker}(\text{alb}(f))) = q_f$, which has a connected component containing 0. We shall denote it with K_f . If B^0 is the open subset of B over which the fibration has smooth fibers, we denote by F_b the fiber over $b \in B^0$. Via the map $F_b \hookrightarrow S$ we have a map $JF_b \rightarrow \text{Alb}(S)$ whose image is, up to translation, exactly K_f . Dualizing we have a map

$$K_f^\vee \hookrightarrow JF_b^\vee = JF_b$$

Note that K_f^\vee doesn't depend on b whereas F_b strongly depends on it. In particular we have proved that the Jacobian of every smooth fiber of a non isotrivial fibration contains a fixed abelian variety of dimension q_f . Assume now that we are in an extreme case, i.e., assume that $q_f = g - 1$. Since in this article we are only interested in non isotrivial fibrations, we will call fibration with *maximal relative irregularity* those with $q_f = g - 1$. In fact, every fibration satisfies $0 \leq q_f \leq g$ and the equality $q_f = g$ holds if and only if the fibration is trivial (this follows from a result of Beauville: see the appendix of [Deb82] for details). In this case $\dim(K_f^\vee) = q_f = g - 1$ and we can consider the quotient JF_b/K_f^\vee which will be an abelian variety of dimension $g - q_f = 1$: an elliptic curve E_b .

$$\Phi(\pi_b) = K_f^\vee \hookrightarrow JF_b \xrightarrow{p_b} E_b$$

$$\begin{array}{ccc} & \uparrow & \nearrow \pi_b \\ & F_b & \xrightarrow{\pi_b} E_b \end{array} \quad \parallel$$

Moreover, in this case K_f^\vee is the connected component through the origin of the kernel of the norm map associated to the ramified covering $\pi_b : F_b \rightarrow E_b$, i.e. the Prym variety $\Phi(\pi_b)$. Hence, an eventual counterexample to the modified version of the conjecture of

Xiao, under the additional assumption $q_f = g - 1$, would give a family of coverings of elliptic curves with constant Prym variety. At the moment the question

Is there a non isotrivial fibration (with maximal relative irregularity q_f or not) giving a counterexample to the modified Xiao's conjecture?

is still completely open but, by answering precisely to our first question one should be able to construct counterexamples or to prove that, at least for the case of maximal relative irregularity, such examples cannot exist. It is worth to mention that, by the original work of Xiao (see [Xia87]), a non isotrivial fibration with maximal relative irregularity can exist only if $g \leq 7$.

The paper is organized as follows. In Section 1 we recall some definitions and facts about Prym varieties associated to ramified coverings and Prym maps that we are going to use extensively in what follows. In Section 2 we will extend the techniques developed in [Kan04] for coverings with simple ramification to the case of arbitrary one. The main result is this theorem

Theorem (2.3). *With the notations of section 2, for any $\varphi \in \text{Sym}^2(H^0(\omega_F)^-)$ we have*

$$(2) \quad d_o P^\vee(\varphi) = \sum_{j=1}^n \text{Res}_{a_j} \left(\frac{m(\varphi)}{\pi^* \alpha} \right) dt_j + \left(\sum_{k=0}^n \frac{m(\varphi)}{\pi^* \alpha^2}(x_k) \right) ds.$$

that describe the (dual of the) differential of the Period map in terms of residues of some meromorphic forms. In Section 3, given a covering $\pi : F \rightarrow E$ and assuming that F is not hyperelliptic, we prove Theorem 3.4, a geometric criterion on the canonical model F that is a sufficient condition in order to have $\dim(\text{Ker}(d_o P)) = 1$. Finally, in Section 4, we analyze in our framework the family that was constructed in [Pir92]. We will prove, using our framework, that the existence of the family is consistent with our Theorem as well as other interesting geometric aspects that may suggest a different way to approach, in the future, the problem of finding an answer to the second question by starting from the geometry of canonical models.

Acknowledgements. The authors are both supported by INdAM - GNSAGA. The first author is supported by FIRB2012 "Moduli spaces and applications" and by IMUB during his stay at the University of Barcelona. The seminal ideas at the base of this article were born in the framework of the PRAGMATIC project of year 2016. We would like to thank Miguel Ángel Barja, Víctor González-Alonso, Juan Carlos Naranjo and Gian Pietro Pirola for introducing us to the subject and for all the helpful remarks on the problem and the manuscript.

1. SOME PRELIMINARIES

In this section we recall some definitions that we are going to use in the following sections.

Let F, E be two smooth curves of genus $g \geq 2$ and 1 respectively and consider the covering $\pi : F \rightarrow E$. One can consider the Albanese variety associated to F , which coincides with its Jacobian, because F is a curve. Namely

$$(3) \quad J(F) = \frac{H^0(\omega_F)^\vee}{H_1(F, \mathbb{Z})} = \text{Alb}(F) = \frac{H^1(\mathcal{O}_F)}{H^1(F, \mathbb{Z})}.$$

This is a principally polarized abelian variety of dimension g . As E has genus 1 we have $E = J(E) = \text{Alb}(E)$. By the universal property of $\text{Alb}(F)$ there is a map $\text{alb}(\pi)$ such that the diagram

$$\begin{array}{ccc} F & \xrightarrow{\pi} & E \\ \downarrow & \circlearrowleft & \downarrow = \\ JF & \xrightarrow{\text{alb}(\pi)} & JE \end{array}$$

commutes, where the map $F \rightarrow JF$ is the Albanese map of F , also called the Abel-Jacobi map. The map $\text{alb}(\pi)$ is also called the *norm map of π* , Nm_π , and it is surjective. The *generalized Prym variety* associated to $\pi : F \rightarrow E$ (or simply *Prym variety*) is the connected component of $\text{Ker}(\text{alb}(\pi))$ that contains the 0, i.e.

$$(4) \quad P(\pi) = \text{Ker}(\text{alb}(\pi))_0.$$

$P(\pi)$ is an abelian variety of dimension $g - 1$ with a natural polarization Θ_P given by $\Theta_{JF}|_P$ via the embedding

$$P(\pi) \hookrightarrow JF.$$

The map $\pi : F \rightarrow E$ induces a map $tr_\pi : H^0(\omega_F) \rightarrow H^0(\omega_E)$ called the *trace* of π (see Appendix A of [Kan04] for the definition). The trace satisfies

$$tr_\pi \circ \pi^* = \text{Deg}(\pi) \text{Id}_{H^0(\omega_E)}.$$

If we define

$$(5) \quad H^0(\omega_F)^- = \text{Ker}(tr_\pi)$$

we have a canonical splitting

$$(6) \quad H^0(\omega_F) = \pi^* H^0(\omega_E) \oplus H^0(\omega_F)^-$$

and we can identify the quotient $H^0(\omega_F)/\pi^* H^0(\omega_E)$ with $H^0(\omega_F)^-$. In particular, the tangent bundle of $P(\pi)$ can be described as

$$(7) \quad TP(\pi) = \left(\frac{H^0(\omega_F)}{\pi^* H^0(\omega_E)} \right)^\vee \otimes \mathcal{O}_{P(\pi)} = (H^0(\omega_F)^-)^\vee \otimes \mathcal{O}_{P(\pi)}.$$

Now we will introduce the families of coverings of elliptic curves we are interested in. Fix a smooth curve F of genus $g \geq 2$ and consider a degree d covering $\pi : F \rightarrow E$, where E is an elliptic curve. Denote with

$$R = \sum_{j=1}^n (n_j - 1) a_j$$

the ramification divisor and call b_j the branch point corresponding to the ramification point a_j , i.e. $\pi(a_j) = b_j$. Thus n_j is the degree of π when restricted to a suitable neighborhood of a_j .

Fix a generator α of $H^0(\omega_E)$. Choose a suitable set $\{\Delta_j\}$ of coordinate neighborhoods centered in the points b_j and call w_j the corresponding coordinate on E . This is not needed at the moment but observe that we can assume that $\alpha|_{\Delta_j} = dw_j$. We can chose a collection of pairwise disjoint coordinate neighborhoods (U_j, z_j) centered in a_j in such a way that $w_j = \pi|_{U_j}(z_j) = z_j^{n_j}$.

Denote by \mathcal{H}_E the polydisc $\prod_{j=1}^n \Delta_j$ and consider the coordinates $t = (t_j)_{j=1}^n$ defined by the relation

$$t_j(P_1, \dots, P_n) = w_j(P_j).$$

We can consider, as in Section 4.1 of [Kan04], a family

$$(\Psi, f) : \mathcal{F} \rightarrow E \times \mathcal{H}_E$$

of d -sheeted branched coverings deforming π parametrized by \mathcal{H}_E such that

$$(8) \quad w_j = \Psi|_{U_j}(z_j, t) = z_j^{n_j} + t_j.$$

In this way, to each $b' \in \mathcal{H}_E$, it is associated a covering $\pi_{b'} : F_{b'} \rightarrow E$ which is a deformation of π , the central fiber. Note that (8) forces the ramification orders to remain constant and allows different branch points to move independently. This is what we will call in the following *local family of coverings over E with central fiber π parametrized by \mathcal{H}_E* .

The tangent space to \mathcal{H}_E in $b = (b_1, \dots, b_n) \in \mathcal{H}_E$ is

$$T_b \mathcal{H}_E \simeq \bigoplus_{j=1}^n T_{b_j} E \simeq \bigoplus_{j=1}^n \mathbb{C} \frac{\partial}{\partial t_j},$$

where the tangent vectors on the right are evaluated in 0.

We can also take into account the deformation of the elliptic curve. Indeed, following [ACG11], if one chooses $c \in E$ not among the b_j and considers a small coordinate neighborhood (N, v) of c (eventually shrinking Δ_j in such a way that for all j they are disjoint from N), one can consider the associated Schiffer variation $\mathcal{E} \rightarrow N$ of E with coordinate s . Observe that we can assume $\alpha|_N = dv$. Taking into account also the movement of the branch points one has a family $f : \mathcal{F} \rightarrow \mathcal{H}_E \times N$ of curves of genus g that fits into the diagram

$$(9) \quad \begin{array}{ccc} \mathcal{F} & \xrightarrow{p} & \mathcal{E} \\ f \downarrow & & \downarrow \\ \mathcal{H}_E \times N & \longrightarrow & N \end{array}$$

For a choice $(b', s') \in \mathcal{H} = \mathcal{H}_E \times N$ we have an elliptic curve $E_{s'}$, the fiber of the map $\mathcal{E} \rightarrow N$ over s' , a curve $F_{(b', s')}$ of genus g and a covering

$$\pi_{(b', s')} = p|_{F_{(b', s')}} : F_{(b', s')} \rightarrow E_{s'}.$$

For this reason, the map p is what we will call *local family of coverings with central fiber π parametrized by \mathcal{H}* or, simply, *local family of coverings*. The tangent space to \mathcal{H} in (b, s) is

$$T_{(b, s)} \mathcal{H} \simeq \left(\bigoplus_{j=1}^n \mathbb{C} \frac{\partial}{\partial t_j} \right) \oplus \mathbb{C} \frac{\partial}{\partial s}.$$

and, clearly, contains $T_b \mathcal{H}_E$ in a natural way. We stress that, through the whole article, unless otherwise stated, we will always refer to the families of coverings constructed in this sections.

If we have a family of coverings parametrized by \mathcal{H} , for each (b, s) we can construct the Prym variety associated to the covering. Moreover, the type of polarization remains

constant. Hence we can consider the *Prym map*

$$(10) \quad \begin{aligned} \mathcal{H} &\xrightarrow{\Phi} \mathcal{A}_{g-1} \\ (b, s) &\longmapsto [P(\pi_{(b,s)})] \end{aligned}$$

where \mathcal{A}_{g-1} is the moduli space of abelian varieties with polarization (which will be omitted) equal to the one of the central fiber. In the same way one has the Prym map Φ_E associated to a local family of coverings over E .

To avoid technical subtleties around singular points of \mathcal{A}_{g-1} , we will consider the period map $P : \mathcal{H} \rightarrow \mathbb{D}$ (or $P_E : \mathcal{H}_E \rightarrow \mathbb{D}$) instead of the Prym map Φ (respectively Φ_E), where \mathbb{D} is a suitable period domain for \mathcal{A}_{g-1} . The interested reader is referred to [Kan04, Section 3] for technical details.

Through the whole article, giving two sections $s_1, s_2 \in H^0(\mathcal{O}_X(D))$ we will write $s_1 \hat{\otimes} s_2$ to mean their symmetric product, i.e.

$$\frac{1}{2}(s_1 \otimes s_2 + s_2 \otimes s_1) \in \text{Sym}^2(H^0(\mathcal{O}_X(D))).$$

If $s_i \in H^0(\mathcal{O}_X(D_i))$, $s_1 \cdot s_2$ will mean the evaluation of $s_1 \otimes s_2$ in $H^0(\mathcal{O}_X(D_1 + D_2))$ under the multiplication map.

2. A DIRECT FORMULA FOR THE CODIFFERENTIAL OF THE PRYM MAP

In this section we will prove an explicit formula for the codifferential of the period map in terms of the residue at the ramification points of some forms. The framework is similar to the one in [Kan04] with the main difference being that we don't restrict ourselves to the case of simple ramification. First of all we introduce some notations.

Fix an elliptic curve E and let $\pi : F \rightarrow E$ be a covering of E with F of genus g . Consider

$$(\Psi, f) : \mathcal{F} \rightarrow E \times \mathcal{H}_E,$$

the local family of coverings with fixed base E , central fiber π and parameter space \mathcal{H}_E constructed in Section 1. By construction, it induces a family $f : \mathcal{F} \rightarrow \mathcal{H}_E$ with central fiber $\mathcal{F}_o = F$. If we consider a minimal versal deformation $f' : \mathcal{F}' \rightarrow M$ of F then the previous family is induced by f' by means of a pullback. More precisely there exists a holomorphic map $h_E : \mathcal{H}_E \rightarrow M$ such that

$$(11) \quad \begin{array}{ccc} \mathcal{F} & \longrightarrow & \mathcal{F}' \\ f \downarrow & & \downarrow f' \\ \mathcal{H}_E & \xrightarrow{h_E} & M \end{array}$$

is commutative. Being f' a minimal versal deformation we have

$$T_o M \simeq H^1(T_F) \simeq H^0(\omega_F^{\otimes 2})^\vee.$$

Moreover, under this identification, if we take a tangent vector v in $T_o \mathcal{H}_E$ and evaluate dh_E in v we get the Kodaira-Spencer map KS_E associated to $\mathcal{F} \rightarrow \mathcal{H}_E$ evaluated in v .

We are able to prove the first important part of Theorem 2.3.

Proposition 2.1. *Using the identifications introduced above, we have that*

$$dh_E^\vee : T_o^\vee M \rightarrow T_o^\vee \mathcal{H}_E$$

can be written as

$$(12) \quad dh_E^\vee(\varphi) = \sum_{j=1}^n \gamma_j dt_j \quad \text{where} \quad \gamma_j = 2\pi i \operatorname{Res}_{a_j} \left(\frac{\varphi}{\pi^* \alpha} \right)$$

and $\varphi \in T_o^\vee M = H^0(\omega_F^{\otimes 2})$.

Proof. For every $\varphi \in H^0(\omega_F^{\otimes 2})$ we have that $dh_E^\vee(\varphi)$ is identified, as cotangent vector on M in o , by the complex numbers γ_j such that

$$dh_E^\vee(\varphi) = \sum_{j=1}^n \gamma_j dt_j.$$

By construction, we can obtain these numbers simply by pairing $dh_E^\vee(\varphi)$ against $\frac{\partial}{\partial t_j}$:

$$\gamma_j = dh_E^\vee(\varphi) \left(\frac{\partial}{\partial t_j} \right) = \varphi \left(dh_E \left(\frac{\partial}{\partial t_j} \right) \right) = \varphi \left(KS_E \left(\frac{\partial}{\partial t_j} \right) \right).$$

In order to develop the computation we may proceed using a description of KS_E in terms of the Čech cohomology (details of this can be found in [Hor73]). To do it consider the exact sequence

$$(13) \quad 0 \longrightarrow T_F \xrightarrow{d\pi} \pi^* T_E \xrightarrow{\psi} \mathcal{R} \longrightarrow 0$$

and let δ be the coboundary map $H^0(\mathcal{R}) \rightarrow H^1(T_F)$. Then KS_E factors as $\delta \circ \tau = KS_E$ where $\tau : T_b H \rightarrow H^0(\mathcal{R})$ is the *characteristic map* of the family (see [Hor73] for the definition and the proof of this fact). Hence we can unfold the calculation using these exact sequences.

If one restricts the exact sequence (13) on U_j (or some sufficiently small subset of this coordinate neighborhood), it can be identified with

$$(14) \quad 0 \longrightarrow \mathcal{O}_{U_j} \frac{\partial}{\partial z_j} \xrightarrow{d\pi} \mathcal{O}_{U_j} \frac{\partial}{\partial w_j} \xrightarrow{\psi} \mathcal{R}|_{U_j} \longrightarrow 0.$$

The first map sends $\frac{\partial}{\partial z_j}$ to $n_j z_j^{n_j-1} \frac{\partial}{\partial w_j}$ while the second one is simply the restriction to the ramification locus. Let $\mathcal{U} = \{U_0, U_1, \dots, U_n\}$ where U_j for $j = 1, \dots, n$ are the neighborhoods defined above and $U_0 = F \setminus \{a_j\}$. Let, as usual, $U_{\alpha, \beta}$, be a shorthand for $U_\alpha \cap U_\beta$ with $\alpha < \beta$. If $\eta = [\eta_j] \in H^0(\mathcal{U}, \mathcal{R})$ with $\eta_0 = 0$ and $\eta_j = p_j(z_j) \frac{\partial}{\partial w_j}$ we have

$$\delta(\eta) = [\lambda_{\alpha, \beta}] \quad \text{with} \quad \lambda_{0, j} = \frac{p_j(z_j)}{n_j z_j^{n_j-1}} \frac{\partial}{\partial z_j}$$

for $j > 0$ and $\lambda_{\alpha, \beta} = 0$ if $\alpha, \beta > 0$. Following [Hor73] and using Equation (8) we have

$$(15) \quad \tau \left(\frac{\partial}{\partial t_j} \right) = [\tau_k^{(j)}] \quad \text{with} \quad \tau_k^{(j)} = \begin{cases} 0 & k \neq j \\ \frac{\partial}{\partial w_j} & k = j. \end{cases}$$

Hence we have

$$KS_E \left(\frac{\partial}{\partial t_j} \right) = \delta \left(\tau \left(\frac{\partial}{\partial t_j} \right) \right) = [\chi_{\alpha, \beta}^{(j)}] \quad \text{with} \quad \chi_{\alpha, \beta}^{(j)} = \begin{cases} \frac{1}{n_j z_j^{n_j-1}} \frac{\partial}{\partial z_j} & (\alpha, \beta) = (0, j) \\ 0 & \text{otherwise.} \end{cases}$$

If $\varphi \in H^0(\omega_F^{\otimes 2})$ we can represent it as Čech-cocycle as $[\phi_j]$ where

$$\phi_0 = \phi|_{U_0} \quad \text{and} \quad \phi_j = q_j(z_j)dz_j^2$$

are the local expressions of φ in coordinates around a_j . The numbers we are interested in are simply the ones obtained by considering the perfect pairing

$$(16) \quad H^0(\omega_F^{\otimes 2}) \otimes H^1(T_F) \longrightarrow H^1(\omega_F) \xrightarrow{\cong} \mathbb{C}$$

applied to $KS_E\left(\frac{\partial}{\partial t_j}\right)$ and φ . Using Čech cohomology, the image in $H^1(\omega_F)$ of our product is given by the Čech class $[\epsilon_{\alpha,\beta}^{(j)}]$ with

$$\epsilon_{\alpha,\beta}^{(j)} = \begin{cases} \frac{q_j(z_j)}{n_j z_j^{n_j-1}} dz_j & (\alpha, \beta) = (0, j) \\ 0 & \text{otherwise.} \end{cases}$$

What remains to be proven is the analogous to the calculation of [Kan04] for the case of simple ramification: roughly, one can adapt the techniques of [ACG11, pag. 14-15] to develop the last isomorphism of (16) in order to finally get

$$\gamma_j = 2\pi i \operatorname{Res}_0 \frac{q_j(z_j)dz_j^2}{n_j z_j^{n_j-1} dz_j} = 2\pi i \operatorname{Res}_{a_j} \frac{\varphi}{\pi^* \alpha}.$$

□

Consider now the family $p : \mathcal{F} \rightarrow \mathcal{E}$ with central fiber $\pi : F \rightarrow E$ and parameter space $\mathcal{H} = \mathcal{H}_E \times N$ as defined in Section 1. As before, we have an induced deformation $f : \mathcal{F} \rightarrow \mathcal{H}$ of F , its associated Kodaira-Spencer map KS and, when a minimal versal deformation $f' : \mathcal{F}' \rightarrow M$ of F is chosen, an holomorphic map $h : \mathcal{H} \rightarrow M$ such that

$$(17) \quad \begin{array}{ccc} \mathcal{F} & \longrightarrow & \mathcal{F}' \\ f \downarrow & & \downarrow f' \\ \mathcal{H} & \xrightarrow{h} & M \end{array}$$

is commutative. Again, as $T_o M \simeq H^1(T_F)$, we can identify dh with KS . We will denote by x_1, \dots, x_d the points of the fiber of π over the point c which, by construction, are all different.

Proposition 2.2. *Using the identifications introduced above, we have that*

$$dh^\vee : T_o^\vee M \rightarrow T_o^\vee \mathcal{H}$$

can be written for any $\varphi \in H^0(\omega_F^{\otimes 2}) = T_o^\vee M$ as $dh^\vee(\varphi) = \sum_{j=1}^n \gamma_j dt_j + \gamma ds$ where

$$(18) \quad \gamma_j = 2\pi i \operatorname{Res}_{a_j} \left(\frac{\varphi}{\pi^* \alpha} \right) \quad \text{and} \quad \gamma = 2\pi i \sum_{k=1}^d \frac{\varphi}{\pi^* \alpha}(x_k).$$

Proof. As before, by duality,

$$dh^\vee(\varphi) = \varphi \circ dh = \varphi \circ KS.$$

It is then clear that the formula for γ_j follows directly from Proposition 2.1. The one that gives γ , as it involves calculations done far from the ramification points, doesn't depend on the type of the ramifications. Hence, the one given in [Kan04] when π as only simple ramification is still valid. □

Recall that we have a decomposition of $H^0(\omega_F)$ given by $H^0(\omega_F)^- \oplus \pi^*H^0(\omega_E)$ where the first space is the vector space of 1-forms on F with trivial trace. This induces a decomposition on $\text{Sym}^2(H^0(\omega_F))$. Unless otherwise specified, consider $\text{Sym}^2(H^0(\omega_F)^-)$ as a subspace of $\text{Sym}^2(H^0(\omega_F))$ in the natural way. Let $m : \text{Sym}^2(H^0(\omega_F)^2) \rightarrow H^0(\omega_F^{\otimes 2})$ be the multiplication map. Denote by $P : \mathcal{H} \rightarrow \mathbb{D}$ the period map associated to the Prym map $\Phi : \mathcal{H} \rightarrow \mathcal{A}_{g-1}$ where \mathbb{D} is a suitable period domain. We are ready to prove Theorem 2.3.

Theorem 2.3. *With the notation introduced in this section, for any $\varphi \in \text{Sym}^2(H^0(\omega_F)^-)$ we have*

$$(19) \quad d_o P^\vee(\varphi) = \sum_{j=1}^n \text{Res}_{a_j} \left(\frac{m(\varphi)}{\pi^* \alpha} \right) dt_j + \left(\sum_{k=1}^d \frac{m(\varphi)}{\pi^* \alpha^2}(x_k) \right) ds.$$

Proof. Theorem 3.21 of [Kan04] expresses the codifferential of the period map calculated in $\varphi \in \text{Sym}^2(H^0(\omega_F)^-)$ and paired with $\frac{\partial}{\partial t_j}$ as

$$\varphi \left(KS \left(\frac{\partial}{\partial t_j} \right) \right)$$

without any restriction on the ramification type. In particular, this formula, together with Proposition 2.2 ends the proof of the Theorem. \square

Remark 2.4. *As a consequence of the last Theorem we can conclude that, if we fix E , the codifferential $d_o P_E^\vee : \text{Sym}^2(H^0(\omega_F)^-) \rightarrow T_o^\vee \mathcal{H}_E$ factors as*

$$(20) \quad \begin{array}{ccc} H^0(\omega_F^{\otimes 2}) & \xleftarrow{dT^\vee} & \text{Sym}^2(H^0(\omega_F)) \\ dh_E^\vee \downarrow & & \uparrow \sigma \\ T_o^\vee \mathcal{H}_E & \xleftarrow{d_o P_E^\vee} & \text{Sym}^2(H^0(\omega_F)^-) \end{array}$$

where T is the Torelli map (so that $m = dT^\vee$) and σ is the lifting of the projection of $\text{Sym}^2(H^0(\omega_F)) \rightarrow \text{Sym}^2(H^0(\omega_F)^-)$ induced by the decomposition $H^0(\omega_F) = H^0(\omega_F)^- \oplus \pi^*H^0(\omega_E)$. The commutativity of the diagram is a consequence of Proposition 2.1 as, for any $\varphi \in H^0(\omega_F) \hat{\otimes} \pi^*H^0(\omega_E)$, we have that $\varphi/\pi^*\alpha$ is holomorphic and hence has residue zero everywhere.

3. A GEOMETRIC APPROACH VIA THE CANONICAL EMBEDDING

In this section we will use the technical result of the previous section in order to prove that $\dim(\text{Ker}(d_o P_E)) = 1$ for arbitrary ramification types and a geometric criterion to determine whether $\dim(\text{Ker}(d_o P)) = 1$ or not. First we fix some notation and facts about the canonical curves that we are going to use extensively in the following.

As F has genus $g \geq 3$ and is not hyperelliptic, we may identify it with its canonical model in $\mathbb{P} = \mathbb{P}H^0(\omega_F)^\vee$. This is a non-degenerate curve of degree $2g - 2$, which is also projectively normal by a classical result of Max Noether (see, for example, [ACGH85]). One of the consequences of this fact is that the multiplication map $m_k : \text{Sym}^k H^0(\omega_F) \rightarrow H^0(\omega_F^{\otimes k})$ is surjective. As before we will denote m_2 simply by m . We will use frequently the natural identifications $H^0(\mathcal{O}_{\mathbb{P}}(d)) = \text{Sym}^d H^0(\omega_F)$ which enable us to identify $\mathbb{P}(\text{Ker}(m_d))$

with the space of hypersurfaces of degree d in $\mathbb{P}H^0(\omega_F)^\vee$ that contain F . By abuse of notation we will simply say that an element in $\text{Sym}^d H^0(\omega_F)$ is an hypersurface of degree d if no confusion arises. In particular, if I_F is the ideal sheaf of F in $\mathbb{P}H^0(\omega_F)^\vee$, then $\text{Ker}(m) = H^0(I_F(2))$ gives the set of all quadrics in $\mathbb{P}H^0(\omega_F)^\vee$ containing the curve F , and has dimension $\frac{(g-2)(g-3)}{2}$.

Recall that the decomposition

$$H^0(\omega_F) = H^0(\omega_F)^- \oplus \pi^* H^0(\omega_E)$$

where the first space is the space of forms with zero trace.

Since elements in $H^0(\omega_F)$ are linear equations on $\mathbb{P}H^0(\omega_F)^\vee$, all the hyperplanes defined by elements in $H^0(\omega_F)^-$ intersect in a single point q^- of \mathbb{P} which is a point really important in what will follows. We have also a particular hyperplane, the one defined by the subspace $\pi^* H^0(\omega_E)$ which will be denoted by H^- . More precisely,

$$q^- = \mathbb{P}((H^0(\omega_F)^-)^{\perp}) \quad \text{and} \quad H^- = \mathbb{P}((\pi^* H^0(\omega_E))^{\perp})$$

As before, we will fix a generator α of $H^0(\omega_E)$ so that

$$(21) \quad \text{Sym}^2(H^0(\omega_F)) = \text{Sym}^2(H^0(\omega_F)^-) \oplus (\pi^* \alpha \hat{\otimes} H^0(\omega_F)).$$

Given a quadric Q in \mathbb{P} we will denote by $G_Q \in \text{Sym}^2(H^0(\omega_F))$ one of its equations and by $G_Q^- \in \text{Sym}^2(H^0(\omega_F)^-)$ and $\omega_Q \in H^0(\omega_F)$ the only elements such that

$$G_Q = G_Q^- + \pi^* \alpha \hat{\otimes} \omega_Q$$

under the decomposition (21). Finally, given a quadric Q , we will denote by Q^- the cone given by the equation G_Q^- , i.e. the quadric such that $G_{Q^-} = G_{Q^-}^- = G_Q^-$.

In order to prove Theorem 3.2 we will need the following result:

Lemma 3.1. *We have a natural inclusion of $H^0(I_F(2))$ in $\text{Ker}(d_o P_E^\vee)$.*

Proof. Recall that, fixed a family of coverings with base E and central fiber $\pi : F \rightarrow E$, by fixing a minimal versal deformation $\mathcal{F}' \rightarrow M$ of F , we can construct $h_E : \mathcal{H} \rightarrow M$ like in diagram (11). As observed in Remark 2.4 we have a commutative diagram

$$(22) \quad \begin{array}{ccccccc} & & & & 0 & & \\ & & & & \uparrow & & \\ 0 & \longrightarrow & \text{Ker } d_o P_E^\vee & \xrightarrow{j} & \text{Sym}^2(H^0(\omega_F)^-) & \xrightarrow{d_o P_E^\vee} & T_o^\vee \mathcal{H}_E \\ & & \uparrow \gamma & & \uparrow pr & & \uparrow dh_E^\vee \\ 0 & \longrightarrow & H^0(I_F(2)) & \xrightarrow{\iota} & \text{Sym}^2(H^0(\omega_F)) & \xrightarrow{m} & H^0(\omega_F^{\otimes 2}) \longrightarrow 0 \\ & & \downarrow \downarrow & & \downarrow \downarrow & & \\ & & & & H^0(\omega_F) \hat{\otimes} \pi^* H^0(\omega_E) & & \\ & & & & \uparrow & & \\ & & & & 0 & & \\ & & & & 10 & & \end{array}$$

It is easy to see that the image of $pr \circ \iota$ lives in $\text{Ker}(d_o P_E^\vee)$ so we have a well defined map $\gamma : H^0(I_F(2)) \rightarrow \text{Ker}(d_o P_E^\vee)$. We want to prove that this map is indeed injective. This follows from the geometry of the problem. Indeed, if a quadric Q contains F , i.e. if the quadric has equation

$$G_Q = G_Q^- + \pi^* \alpha \hat{\otimes} \omega_Q \in H^0(I_F(2)),$$

and if $\gamma(G_Q) = 0$ then we have that the quadric has equation $\pi^* \alpha \hat{\otimes} \omega_Q$. But this is impossible because such a quadric the union of two planes (one of which is H^-) and the canonical curve is non-degenerate. Hence γ is injective. \square

Theorem 3.2. *Let $\pi : F \rightarrow E$ be a covering with F non-hyperelliptic, consider a local family of coverings with base E and parameter space \mathcal{H}_E constructed in Section 1. Let P_E be the period mapping associated to the Prym map Φ_E . Then $\dim(\text{Ker}(d_o P_E)) = 1$.*

Proof. First of all, observe that for dimensional reasons, one has $\dim(\text{Ker}(d_o P_E)) = 1$ if and only if

$$\dim(\text{Ker}(d_o P_E^\vee)) = \frac{g(g-1)}{2} - n + 1.$$

From the splitting $H^0(\omega_F) = H^0(\omega_F)^- \oplus \pi^* H^0(\omega_E)$ we have the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^0(\omega_F) & \xrightarrow{\hat{\otimes} \pi^* \alpha} & \text{Sym}^2(H^0(\omega_F)) & \xrightarrow{pr} & \text{Sym}^2(H^0(\omega_F)^-) \longrightarrow 0 \\ & & & & & \searrow \Psi & \downarrow d_o P_E^\vee \\ & & & & & & T_o \mathcal{H}_E \end{array}$$

with Ψ defined by extending the formula in Theorem 2.3 to $\text{Sym}^2(H^0(\omega_F))$. This can be done because, as previously observed (see Remark 2.4), $d_o P_E^\vee(H^0(\omega_F) \hat{\otimes} \pi^* \alpha) = \{0\}$. In particular, we have the relation

$$(23) \quad \dim(\text{Ker}(d_o P_E^\vee)) = \dim(\text{Ker}(\Psi)) - \dim(\text{Ker}(pr)) = \dim(\text{Ker}(\Psi)) - g.$$

By definition, Ψ factors through the multiplication map m as $\Psi = \bar{\Psi} \circ m$. The map $\bar{\Psi}$ is well defined as, by Lemma 3.1, $\text{Ker}(m) \subset \text{Ker}(\Psi)$.

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^0(\omega_F) & \xrightarrow{\hat{\otimes} \pi^* \alpha} & \text{Sym}^2(H^0(\omega_F)) & \xrightarrow{pr} & \text{Sym}^2(H^0(\omega_F)^-) \longrightarrow 0 \\ & & \downarrow m & & \searrow \Psi & & \downarrow d_o P_E^\vee \\ & & H^0(\omega_F^{\otimes 2}) & \xrightarrow{\bar{\Psi}} & T_o \mathcal{H}_E & & \end{array}$$

Being m surjective (as F is non-hyperelliptic) we obtain the further relation

$$(24) \quad \dim(\text{Ker}(\Psi)) = \dim(\text{Ker}(\bar{\Psi})) + \dim(\text{Ker}(m)) = \dim(\text{Ker}(\bar{\Psi})) + \frac{(g-2)(g-3)}{2}.$$

As the divisor associated to $\pi^* \alpha$ is exactly R , the ramification divisor, we have that $\omega_F = \mathcal{O}_F(R)$ and there is an exact sequence

$$(25) \quad 0 \longrightarrow \omega_F \xrightarrow{\pi^* \alpha} \omega_F^{\otimes 2} \longrightarrow \omega_F^{\otimes 2}|_R \longrightarrow 0$$

which yields, denoting with V the quotient $H^0(\omega_F^{\otimes 2})/(H^0(\omega_F) \cdot \pi^*\alpha)$, the exact sequences

$$(26) \quad \begin{aligned} 0 &\longrightarrow H^0(\omega_F) \xrightarrow{\cdot\pi^*\alpha} H^0(\omega_F^{\otimes 2}) \xrightarrow{\epsilon} V \longrightarrow 0 \\ 0 &\longrightarrow V \xrightarrow{\zeta} H^0(\omega_F^{\otimes 2}|_R) \longrightarrow H^1(\omega_F) \longrightarrow 0 \end{aligned}$$

Let $\eta \in \text{Ker}(\epsilon)$. We want to prove that $\bar{\Psi}(\eta) = 0$. This is easily proven: write η as $\omega \cdot \pi^*\alpha$ and observe that

$$\bar{\Psi}(\eta) = (\bar{\Psi} \circ m)(\omega \hat{\otimes} \pi^*\alpha) = (d_o P_E^\vee \circ pr)(\omega \hat{\otimes} \pi^*\alpha) = 0$$

because $\omega \hat{\otimes} \pi^*\alpha \in \text{Ker}(pr)$. In particular, $\text{Ker}(\epsilon) \subset \text{Ker}(m)$ and we can define a map $\lambda : V \rightarrow T_o^\vee \mathcal{H}_E$ such that $\bar{\Psi} = \lambda \circ \epsilon$. Moreover

$$(27) \quad \dim(\text{Ker}(\bar{\Psi})) = \dim(\text{Ker}(\lambda)) + g.$$

Using the second exact sequence in 26 we can also define a map $\mu : H^0(\omega_F^{\otimes 2}|_R) \rightarrow T_o \mathcal{H}_E$ such that $\mu \circ \zeta = \lambda$.

$$(28) \quad \begin{array}{ccccc} & & H^0(I_F(2)) & & \\ & & \downarrow & & \\ H^0(\omega_F) & \xrightarrow{\hat{\otimes} \pi^*\alpha} & \text{Sym}^2(H^0(\omega_F)) & \xrightarrow{pr} & \text{Sym}^2(H^0(\omega_F)^-) \\ & \searrow \cdot\pi^*\alpha & \downarrow m & \searrow \Psi & \downarrow d_o P_E^\vee \\ & & H^0(\omega_F^{\otimes 2}) & \xrightarrow{\bar{\Psi}} & T_o^\vee \mathcal{H}_E \\ & & \downarrow \epsilon & \nearrow \lambda & \uparrow \mu \\ & & V & \xrightarrow{\zeta} & H^0(\omega_F^{\otimes 2}|_R) \longrightarrow H^1(\omega_F) \end{array}$$

Note that we have several ways to define μ . Since $\omega_F^{\otimes 2}|_R = \omega_F(R)|_R$ the global sections of $\omega_F^{\otimes 2}|_R$ are just collections of meromorphic tails on the points of ramification, i.e. elements

$$\left\{ \sum_{j=1}^{n_k-1} \beta_{jk} \frac{dz_k}{z_k^j} \right\}_{a_k \in R}$$

where n_k is the ramification index of the point a_k . In particular, we can define μ as the map which gives the residue in the corresponding point of the meromorphic tail. This ensures that the diagram is commutative. In addition, μ is surjective (this because the image of a collection of meromorphic tails $\{s_k\}$, one for each point of ramification, with $\beta_{1m} = \delta_{km}$, generates the image), and as a consequence, $\zeta|_{\text{Ker}(\lambda)}$ is an isomorphism between $\text{Ker}(\lambda)$ and $\text{Ker}(\mu)$. Hence,

$$(29) \quad \begin{aligned} \dim \text{Ker}(d_o P_E^\vee) &= \dim(\Psi) - g = \dim(\bar{\Psi}) + \dim(\text{Ker}(m)) - g = \\ &= \dim(\lambda) + \dim(\text{Ker}(m)) = \dim(\mu) + \dim(\text{Ker}(m)) = \\ &= h^0(\omega_F(R)|_R) - \dim T_o^\vee \mathcal{H}_E + \dim(\text{Ker}(m)) = \frac{g(g-1)}{2} - n + 1 \end{aligned}$$

as wanted. \square

Now we will prove the first main theorem:

Theorem 3.3. *Let $\pi : F \rightarrow E$ be a covering with F non-hyperelliptic, consider the local family of coverings with parameter space \mathcal{H} constructed in Section 1. Let P be the period mapping associated to the Prym map $\Phi : \mathcal{H} \rightarrow \mathcal{A}_{g-1}$. Using the same notations of Theorem 2.3 we have*

$$\dim(\text{Ker}(d_o P)) = 1 \quad \iff \quad \exists \beta \in \text{Ker}(d_o P_E^\vee) \quad | \quad \sum_{k=1}^d \frac{m(\beta)}{\pi^* \alpha^2}(x_k) \neq 0.$$

Proof. First of all consider the diagrams

$$\begin{array}{ccc} T_o \mathcal{H} & \xrightarrow{d_o P} & T_{P(o)} \mathbb{D} \\ \uparrow & \nearrow d_o P_E & \\ T_o \mathcal{H}_E & & \end{array} \quad \begin{array}{ccc} T_o^\vee \mathcal{H} & \xleftarrow{d_o P^\vee} & T_{P(o)}^\vee \mathbb{D} \\ \downarrow & \nwarrow d_o P_E^\vee & \\ T_o^\vee \mathcal{H}_E & & \end{array}$$

and observe that one always has

$$\text{Ker}(d_o P_E) \subseteq \text{Ker}(d_o P) \quad \text{Ker}(d_o P^\vee) \subseteq \text{Ker}(d_o P_E^\vee).$$

Moreover, the codimensions are at most 1. If one considers the exact sequences

$$0 \longrightarrow \text{Ker}(d_o P_E) \longrightarrow T_o \mathcal{H}_E \longrightarrow T_{P(o)} \mathbb{D} \longrightarrow \text{Ker}(d_o P_E^\vee)^\vee \longrightarrow 0$$

$$0 \longrightarrow \text{Ker}(d_o P) \longrightarrow T_o \mathcal{H} \longrightarrow T_{P(o)} \mathbb{D} \longrightarrow \text{Ker}(d_o P^\vee)^\vee \longrightarrow 0$$

it is clear that $\text{Ker}(d_o P_E) = \text{Ker}(d_o P)$ if and only if $\text{Ker}(d_o P^\vee) \subsetneq \text{Ker}(d_o P_E^\vee)$. Hence we have

$$\dim(\text{Ker}(d_o P)) = 1 \iff \text{Ker}(d_o P^\vee) \subsetneq \text{Ker}(d_o P_E^\vee).$$

This is true if and only if there exists an element $\beta \in \text{Ker}(d_o P_E^\vee)$ on which $d_o P^\vee$ doesn't vanish. This can only be possible if $d_o P^\vee(\beta)$ is not zero on $\frac{\partial}{\partial s}$, where s is the parameter taking into account the moduli of the elliptic curve. By using Theorem 2.3 we have

$$d_o P^\vee(\beta) = \sum_{k=1}^d \frac{m(\beta)}{\pi^* \alpha^2}(x_k)$$

and this concludes the proof. \square

This result improves the one in [Kan04] where it is proved only for simple ramification. In the same work is proved that, for simple ramification, having the sum in Theorem 3.3 different from zero for some $\beta \in \text{Ker}(d_o P_E^\vee)$ is equivalent to ask that the intersection of the quadrics that contain the canonical model of F doesn't contain the point q^- defined before. Unfortunately, in the case of arbitrary ramification, we are not able to prove this equivalence but only one implication.

Theorem 3.4. *With the same hypothesis of Theorem 3.3, if we identify F with its canonical model in $\mathbb{P}H^0(\omega_F)^\vee$, then we have*

$$(30) \quad q^- \notin \bigcap_{F \subset Q} Q \implies \dim(\text{Ker}(d_o P)) = 1,$$

where Q ranges in the set of quadrics of $\mathbb{P}H^0(\omega_F)^\vee$ containing F .

The proof of the theorem uses some arguments developed in [Kan04] that we have summarized in the following Lemma.

Lemma 3.5. *Let Q be a quadric of $\mathbb{P}H^0(\omega_F)^\vee$ containing F and denote by $G_Q = G_Q^- + \pi^*\alpha \hat{\otimes} \omega_Q$ one of its equations. Then*

$$\sum_{k=1}^d \frac{m(G_Q^-)}{\pi^*\alpha^2}(x_k) = 0 \iff G_Q(q^-) = 0 \iff q^- \in Q.$$

Proof. The last statement is clear by definition so we really need to prove only the first one. First of all observe that we can choose the coordinate s in such a way that α is locally given by ds . Then, as $G_Q^- = G_Q - \pi^*\alpha \hat{\otimes} \omega_Q$ and $Q \in H^0(I_F(2)) = \text{Ker}(m)$ by hypothesis, one has

$$\sum_{k=1}^d \frac{m(G_Q^-)}{\pi^*\alpha^2}(x_k) = - \sum_{k=1}^d \frac{m(\pi^*\alpha \hat{\otimes} \omega_Q)}{\pi^*\alpha^2}(x_k) = - \frac{\text{Tr}_\pi(\omega_Q)}{\alpha}(c).$$

But $\text{Tr}_\pi(\omega_Q)$ is an element of $H^0(\omega_E)$ so it is equal to $r \cdot \alpha$ for some r . Thus we have

$$\sum_{k=1}^d \frac{m(G_Q^-)}{\pi^*\alpha^2}(x_k) = -r$$

which is zero if and only if ω_Q has trace 0, i.e. if and only if $\omega_Q \in H^0(\omega_F)^-$. This happens if and only if $(\pi^*\alpha)^{\otimes 2}$ doesn't appear in the equation of Q , i.e. if and only if $q^- \in Q$. \square

Using Lemma 3.1 and Lemma 3.5 the proof of Theorem 3.4 is straightforward.

Proof of Theorem 3.4. Assume that

$$q^- \notin \bigcap_{F \subset Q} Q.$$

Then, there exists a quadric which contains F but doesn't contain q^- . Denote by G_Q its equation. By Lemma 3.1 we know that $\beta = \gamma(G_Q) = G_Q^- \in \text{Ker}(d_o P_E^\vee)$ and by Lemma 3.5 we have that

$$\sum_{k=1}^d \frac{m(\beta)}{\pi^*\alpha^2}(x_k) \neq 0.$$

Hence, using Theorem 3.3 we have the thesis. \square

Remark 3.6. *In [Kan04], with different methods, it is proved that $H^0(I_F(2)) = \text{Ker}(d_o P_E^\vee)$ if the ramification is simple. This fact is exactly what allows to prove the converse implication of Theorem 3.4.*

Remark 3.7. *Notice that $H^0(I_F(2)) = \text{Ker}(d_o P_E^\vee)$ if and only if all the ramification indices are equal to 2. Indeed, denote by R_{red} the reduced divisor whose support equals the support of the ramification divisor. Let \bar{R} be $R - R_{red}$. From Riemann-Hurwitz we have*

$$2g - 2 = \deg(R) = \deg(R_{red}) + \deg(\bar{R}) = n + \deg(\bar{R}).$$

Hence, from Equation (29) one has

$$\dim \text{Ker } d_o P_E^\vee = h^0(I_F(2)) + \deg(\bar{R}).$$

As $\bar{R} \geq 0$ and is trivial if and only if all the ramification indices are equal to 2 the claim follows. In particular, the converse implication of (30) in Theorem 3.4 holds for coverings whose ramification indices are all equal to 2.

We conclude this section by proving the existence of an exact sequence which should help to measure, in a more intrinsic way, how much $H^0(I_F(2))$ and $\text{Ker}(d_o P_E^\vee)$ differ.

Proposition 3.8. *Under the hypothesis of Theorem 3.4 there is an exact sequence*

$$(31) \quad 0 \longrightarrow H^0(I_F(2)) \xrightarrow{\gamma} \text{Ker}(d_o P_E^\vee) \longrightarrow \frac{\text{Ker}(dh^\vee)}{H^0(\omega_F) \hat{\otimes} \pi^* H^0(\omega_E)} \longrightarrow 0.$$

Proof. Starting from diagram (22) it is easy to see that the composition of the inclusion of $H^0(\omega_F) \hat{\otimes} \pi^* H^0(\omega_E)$ with m has image in $H^0(\omega_F^{\otimes 2})$ but also in the kernel of dh^\vee . Hence there is a map

$$\epsilon : H^0(\omega_F) \hat{\otimes} \pi^* H^0(\omega_E) \rightarrow \text{Ker}(dh^\vee),$$

which is easily proven to be injective as we have done with γ . We can also complete the diagram on the right by adding two (trivial) vertical arrows. The complete diagram looks like this

$$(32) \quad \begin{array}{ccccccccc} & & & & 0 & & & & \\ & & & & \uparrow & & & & \\ 0 & \longrightarrow & \text{Ker } d_o P_E^\vee & \xrightarrow{j} & \text{Sym}^2(H^0(\omega_F)^-) & \xrightarrow{d_o P_E^\vee} & T_o^\vee \mathcal{H}_E & \longrightarrow & (\text{Ker } d_o P_E)^\vee \longrightarrow 0 \\ & & \uparrow \gamma & & \uparrow pr & & \uparrow dh^\vee & & \uparrow \\ 0 & \longrightarrow & H^0(I_F(2)) & \xrightarrow{\iota} & \text{Sym}^2(H^0(\omega_F)) & \xrightarrow{m} & H^0(\omega_F^{\otimes 2}) & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ & & 0 & \longrightarrow & H^0(\omega_F) \hat{\otimes} \pi^* H^0(\omega_E) & \xrightarrow{\epsilon} & \text{Ker } dh^\vee & \longrightarrow & \text{Coker } \epsilon \longrightarrow 0 \\ & & & & \uparrow & & \uparrow & & \\ & & & & 0 & & 0 & & \end{array}$$

By using the snake lemma on the central columns one obtain the wanted sequence. \square

4. AN INTERESTING FAMILY OF CURVES

In this section we review the first example, due to Pirola, of a non-trivial family of coverings of elliptic curves with 2 independent directions along which the Prym map Φ is constant. Hence the kernel of the differential of the Period map associated to Φ has dimension greater than 1. The existence of the family is proved in [Pir92] but the proof is not constructive and uses a framework different from ours. After some notations and a brief idea of how to prove the existence of this family (for details, see [Pir92]), we will prove that q^- belongs to the only quadric that contains F and that for all the elements of $\text{Ker}(d_o P_E^\vee)$ the sum in Theorem 3.3 is 0.

In order to prove the existence of such a family, let $G \simeq \mathbb{Z}_3$ and consider the space \mathcal{H}^G of Galois coverings $\pi : F \rightarrow E$ of degree 3 with ramification given by 3 points (so the number of branch points is exactly 3 and the genus of F is 4) modulo the identifications given by

a commutative diagram like

$$\begin{array}{ccc} F_1 & \xrightarrow{\cong} & F_2 \\ \pi_1 \downarrow & & \downarrow \pi_2 \\ E_1 & \xrightarrow{\cong} & E_2 \end{array}$$

With this type of identification of two coverings the dimension of \mathcal{H}^G is 3. Note that, with this definition, a covering $\pi : F \rightarrow E$ and the covering obtained by composing π with a translation of E are equivalent: they represent the same point in \mathcal{H}^G .

Fix a generator g of G and ρ , a primitive root of 1 of order 3. If V is a vector space on which G acts, we will denote by V_{ρ^k} the subspace where g acts as the multiplication by ρ^k . As π is the quotient by the group G , the G -action on F induces several other G -actions. We will do now a small list of the one that we are going to use in this section.

a) The canonical action on $H^0(\omega_F)$ via pullback: by changing, if necessary, g with g^2 , we have

$$(33) \quad H^0(\omega_F) = H^0(\omega_F)_1 \oplus H^0(\omega_F)_\rho \oplus H^0(\omega_F)_{\rho^2} = \pi^* H^0(\omega_E) \oplus \mathbb{C}_\rho^2 \oplus \mathbb{C}_{\rho^2}^1.$$

b) A canonical G -action on $H^0(\omega_F)^-$ which is simply the restriction of the canonical representation on $H^0(\omega_F)$.

c) An action on $\text{Sym}^2(H^0(\omega_F))$, whose decomposition in irreducible subrepresentations is given by

$$\text{Sym}^2(H^0(\omega_F)) = \mathbb{C}_1^3 \oplus \mathbb{C}_\rho^3 \oplus \mathbb{C}_{\rho^2}^4.$$

d) an action on $H^0(\omega_F^{\otimes 2})$ using the surjectivity of m by imposing that m becomes a morphism of G -vector spaces and hence on its dual $H^1(T_F)$.

e) An action on $H^0(I_F(2))$ as the kernel of m .

f) An action on the Prym $\Phi(\pi)$: this is induced at level of tangent spaces (as the tangent space $T_0\Phi(\pi)$ is $H^0(\omega_F)^-$) and it is compatible with the quotient by the periods' lattice.

g) An action of G on $\mathbb{P}H^0(\omega_F)^\vee = \mathbb{P}$ as every automorphism of F , seen as a canonical curve in \mathbb{P} lifts to an automorphism of the whole space.

All these actions, by construction, are compatible via the usual identification. For example, if we interpret $H^0(\omega_F)$ as the space of equations of hyperplanes of \mathbb{P} an invariant hyperplane in \mathbb{P} has an equation which is an eigenvector of g in $H^0(\omega_F)$.

One has a Prym map $\tilde{\Phi} : \mathcal{H}^G \rightarrow \mathcal{A}_{g-1}$ and a period map $\tilde{P} : \mathcal{H}^G \rightarrow \mathbb{D}$. We stress that, by construction, if we prove that $\dim(\text{Ker}(d_o\tilde{P})) > k$ then, the period map P associated to the Prym map of a local family of coverings with π as central fiber will have kernel of dimension at least $k + 1$.

The rough idea to prove that there exists a family of coverings in \mathcal{H}^G which gets contracted by $\tilde{\Phi}$ is to observe, as we have done in f), that the Prym map $\tilde{\Phi}$ factors through the inclusion of \mathcal{A}_{g-1}^G , the space of abelian varieties of dimension $g - 1$ with an action of G , in \mathcal{A}_{g-1} . If we denote by \mathbb{D}^G a period domain for \mathcal{A}_{g-1}^G we have an analogous period map $\tilde{P}^G : \mathcal{H}^G \rightarrow \mathbb{D}^G$. We want to get a bound on the dimension of the image $d\tilde{P}^G$.

Clearly, the image of $d\tilde{P}^G$ has dimension at most the dimension of

$$T\mathbb{D}^G = \text{Sym}^2(H^0(\omega_F)^-)^G$$

and the same bound holds, by construction, for the dimension of the image of $d\tilde{P}$. As $\text{Sym}^2(H^0(\omega_F)^-)^G$ is isomorphic, by b), to

$$H^0(\omega_F)_\rho \hat{\otimes} H^0(\omega_F)_{\rho^2},$$

we have that its dimension is 2. As $T_{[\pi]}\mathcal{H}^G$ has dimension 3 this implies that the kernel of $d\tilde{P}$ has dimension at least 1 and the existence of the family is proved.

Proposition 4.1. *Let $\pi : F \rightarrow E$ with $[\pi] \in \mathcal{H}^G$, identify F with its canonical model and let Q be the only quadric containing F . Then $q^- \in Q$ and either*

$$H^0(I_F(2)) \subset \text{Sym}^2(H^0(\omega_F))_\rho \quad \text{or} \quad H^0(I_F(2)) \subset \text{Sym}^2(H^0(\omega_F))_{\rho^2}.$$

Proof. There exists only a quadric containing F because $g(F) = 4$. More precisely F is the complete intersection of a quadric Q and a cubic surface C . Let $G_Q \in H^0(I_F(2))$ be an equation for Q . Being F invariant under the G -action introduced in g), we have that the orbit of G_Q under the action given in e), is simply given by itself plus, possibly, some of its multiple by elements in \mathbb{C}^* . The key point now is to see that $H^0(I_F(2))_1 = 0$. In order to prove this observe that, by construction, we have an exact sequence of G -vector spaces given by

$$0 \longrightarrow H^0(I_F(2)) \longrightarrow \text{Sym}^2(H^0(\omega_F)) \longrightarrow H^0(\omega_F^{\otimes 2}) \longrightarrow 0$$

Hence, by taking invariant parts and dimensions we have

$$\dim(H^0(I_F(2))^G) = \dim(\text{Sym}^2(H^0(\omega_F))^G) - \dim(H^0(\omega_F^{\otimes 2})^G).$$

As claimed in [Pir92], we can identify $T_{[\pi]}\mathcal{H}^G$ with $H^1(T_F)^G = (H^0(\omega_F^{\otimes 2})^\vee)^G$. Hence, we have $\dim(H^0(\omega_F^{\otimes 2})^G) = 3$. Using c) we have that also $\dim(\text{Sym}^2(H^0(\omega_F))^G) = 3$ so, as claimed, $H^0(I_F(2))_1 = 0$.

As consequence we have either $G_Q \in \text{Sym}^2(H^0(\omega_F))_\rho$ or $G_Q \in \text{Sym}^2(H^0(\omega_F))_{\rho^2}$. Note that in both cases, as $\pi^*H^0(\omega_E)^2 \subset \text{Sym}^2(H^0(\omega_F))_1$, we have $G_Q(q^-) = 0$ so $q^- \in Q$. \square

Lemma 4.2. *Let $\pi : F \rightarrow E$ with $[\pi] \in \mathcal{H}^G$ and assume that F is not hyperelliptic. Denote by a_1, a_2 and a_3 the 3 ramification points of π . Let L be a g_3^1 . Then:*

- L is G -invariant, i.e. $g^*L = L$;
- $h^0(\mathcal{O}_F(3a_i)) = 1$;
- If L' is a g_3^1 then $L \simeq L'$, i.e. there is only one g_3^1 on F .

Proof. Recall that every curve of genus 4 is trigonal and moreover, the number of g_3^1 is at most 2. If there is only one g_3^1 clearly it is G -invariant. If there are 2, as G has order 3 and acts on a set of two elements, it has to fix both of them.

Now let's prove that $h^0(\mathcal{O}_F(3a_i)) = 1$. The Riemann-Roch formula for $\mathcal{O}_F(3a_i)$ is

$$h^0(\mathcal{O}_F(3a_i)) - h^1(\mathcal{O}_F(3a_i)) = \deg(h^0(\mathcal{O}_F(3a_i))) - 4 + 1 = 0$$

so, by Serre duality, we have

$$h^0(\mathcal{O}_F(3a_i)) = h^0(\omega_F(-3a_i)).$$

From

$$0 \rightarrow \omega_F(-3a_i) \rightarrow \omega_F(-2a_i) \rightarrow \omega_F(-2a_i)|_{a_i} \rightarrow 0$$

one has $H^0(\omega_F(-3a_i)) \leq H^0(\omega_F(-2a_i))$. In particular, as F is not hyperelliptic we obtain that the dimension of $H^0(\omega_F(-3a_i))$ is either 1 or 2. Moreover, $h^0(\omega_F(-3a_i)) = 2$ if and only if $H^0(\omega_F(-3a_i)) = H^0(\omega_F(-2a_i))$. But this cannot happen as the pullback η of a non-zero holomorphic form on E has a zero of multiplicity 2 exactly in the ramification points so there is at least one element in $H^0(\omega_F(-2a_i)) \setminus H^0(\omega_F(-3a_i))$. Hence $h^0(\mathcal{O}_F(3a_i)) = 1$ as claimed.

Recall that on F there are at most two g_3^1 and they are related by

$$L \otimes L' = \omega_F = \mathcal{O}_F(2a_1 + 2a_2 + 2a_3).$$

Hence we will conclude by proving that $L = \mathcal{O}_F(a_1 + a_2 + a_3)$. Let A, B in F such that $L \simeq \mathcal{O}_F(a_1 + A + B)$. As a_1 is invariant and the same holds for L , we have that $\mathcal{O}_F(g(A) + g(B)) = \mathcal{O}_F(A + B)$. Therefore, as F is not hyperelliptic, also the equality of divisors $g(A) + g(B) = A + B$ has to hold. Moreover, as g has order 3, it cannot exchange A and B : we have proved that A and B are ramification points. If we assume that $L \neq \mathcal{O}_F(a_1 + a_2 + a_3)$ there are several possibilities:

$A = a_1 = B$: This is impossible as we would have

$$2 = h^0(L) = h^0(\mathcal{O}_F(3a_1)) = 1.$$

$A = a_1 \neq B$: Assume that $A = a_1$ and $B = a_2$ so that $L \simeq \mathcal{O}_F(2a_1 + a_2)$. Let $C, D \in F$ such that $\mathcal{O}_F(2a_1 + a_2) \simeq \mathcal{O}_F(a_3 + C + D)$. As before, we have that C and D are ramification points and as F is not hyperelliptic the only possible option is to have $C = D = a_3$. But then, again, we have a contradiction

$$2 = h^0(\mathcal{O}_F(L)) = h^0(\mathcal{O}_F(2a_1 + a_2)) = h^0(\mathcal{O}_F(3a_3)) = 1.$$

$A \neq a_1 = B$: This case is analogous to the previous one.

$A = B \neq a_1$: This case is analogous to the second one.

Hence, we have proved that $L = \mathcal{O}_F(a_1 + a_2 + a_3)$ and thus that $L \simeq L'$ and there is only a g_3^1 on F . \square

Proposition 4.3. *Let $\pi : F \rightarrow E$ with $[\pi] \in \mathcal{H}^G$ and assume that F is not hyperelliptic. Denote by Q be the only quadric containing the canonical curve F . Then Q is a quadric cone with vertex V and $V \notin F$. Moreover, the hyperplane H^- is tangent to the cone and the 3 ramification points of π lie on a line on the cone.*

Proof. Recall that if the quadric Q containing F is smooth, then F can be seen as a curve of bidegree $(3, 3)$ in $\mathbb{P}^1 \times \mathbb{P}^1$ and the projections on each factor give two different g_3^1 . If, instead, Q is a cone (these are the only possible cases as F is non degenerate) there exists only one g_3^1 . Hence, by Lemma 4.2, we can conclude that Q is a cone. If V is the vertex, it is clear that $V \notin F$ as, otherwise F would be singular.

Now we will prove that the ramification points are on a line in the canonical model of F . By what we have seen in this section we have a decomposition of $H^0(\omega_F)$ into subrepresentations with $H^0(\omega_F)_1 = \pi^*H^0(\omega_E)$. We can assume, as before, that $H^0(\omega_F)_\rho$ has dimension 2. Denote respectively with $\{u_0\}$, $\{u_1, u_2\}$ and $\{u_3\}$ a basis for $\pi^*H^0(\omega_E)$, $H^0(\omega_F)_\rho$ and $H^0(\omega_F)_{\rho^2}$. By abuse of notation we will write $u_i u_j$ to mean $u_i \hat{\otimes} u_j$. With these coordinates, the hyperplane H^- has equation $u_0 = 0$ and $q^- = (1 : 0 : 0 : 0)$. As

$$(34) \quad \text{Sym}^2(H^0(\omega_F)) = \langle u_0^2, u_1 u_3, u_2 u_3 \rangle_1 \oplus \langle u_0 u_1, u_0 u_2, u_3^2 \rangle_\rho \oplus \langle u_0 u_3, u_1^2, u_2^2, u_1 u_2 \rangle_{\rho^2}$$

We know by Proposition 4.1 that an equation G_Q of Q is either an element of $\text{Sym}^2(H^0(\omega_F))_\rho$ or of $\text{Sym}^2(H^0(\omega_F))_{\rho^2}$. In the first case the generic element of $\text{Sym}^2(H^0(\omega_F))_\rho$ is a quadric cone and has equation

$$u_3^2 + u_0(au_1 + bu_2) = 0.$$

Moreover, it is easy to see that H^- is tangent to the cone along the line $L_1 = \{u_3 = u_0 = 0\}$. In the second case the generic element of $\text{Sym}^2(H^0(\omega_F))_{\rho^2}$ is a smooth quadric but it is easy to see that the generic singular element is a cone with equation

$$u_3u_0 + (au_1 + bu_2)^2 = 0.$$

As before, H^- is a plane tangent to Q along the line $L_2 = \{u_0 = au_1 + bu_2 = 0\}$. So, in both cases, as the ramification points of the canonical curve F are given exactly as $H^- \cap F$, they are on a line as claimed. \square

Now we are going to calculate the sum in Theorem 3.3 and to see that it is zero for each element in $\text{Ker}(d_\circ P_E^\vee)$.

Proposition 4.4. *Let $\pi : F \rightarrow E$ be a Galois covering of degree 3 as before and consider a local family of coverings with central fiber π . Then, if*

$$\nu(\beta) = \sum_{k=1}^3 \frac{m(\beta)}{\pi^*\alpha^2}(x_k),$$

one has $\nu(\beta) = 0$ for all $\beta \in \text{Ker}(d_\circ P_E^\vee)$.

Proof. $\text{Ker}(d_\circ P_E^\vee)$ is a subspace of $\text{Sym}^2(H^0(\omega_F)^-)$. If $\beta \in \text{Ker}(d_\circ P_E^\vee)$ we can decompose it as

$$\beta = \beta_1 + \beta_\rho + \beta_{\rho^2}$$

with $\beta_\mu \in \text{Sym}^2(H^0(\omega_F)^-)_\mu$. First of all we will prove that $\nu(\beta_\rho) = \nu(\beta_{\rho^2}) = 0$.

As c is not a branch point, we have that the fiber $\pi^{-1}(c) = \{x_1, x_2, x_3\}$ over c is equal to the orbit of each of its points: $\pi^{-1}(c) = \{x_1, g(x_1), g^2(x_1)\}$. Hence

$$\nu(\beta) = \sum_{k=1}^3 \frac{m(\beta)}{\pi^*\alpha^2}(x_k) = \sum_{k=0}^2 \frac{m(\beta)}{\pi^*\alpha^2}(g^k(x_1)) = \sum_{k=0}^2 \frac{m(\beta \circ g^k)}{\pi^*\alpha^2}(x_1).$$

If we assume that β is in the eigenspace $\text{Sym}^2(H^0(\omega_F))_\mu$ of g^* then

$$\nu(\beta) = \frac{m\left(\sum_{k=1}^d (g^*)^k(\beta)\right)}{\pi^*\alpha^2}(x_1) = \frac{m\left(\sum_{k=0}^2 \mu^k \beta\right)}{\pi^*\alpha^2}(x_1) = \left(\sum_{k=0}^2 \mu^k\right) \frac{m(\beta)}{\pi^*\alpha^2}(x_1).$$

Hence, if $\mu \neq 1$, we have $\lambda(\beta) = 0$ as claimed.

Hence we have that $\nu(\beta) = \nu(\beta_1)$ so it is enough to prove that

$$\text{ker}(d_\circ P_E^\vee) \subseteq \text{Sym}^2(H^0(\omega_F)^-)_\rho \oplus \text{Sym}^2(H^0(\omega_F)^-)_{\rho^2}$$

i.e., that $\beta_1 = 0$.

Let a be a ramification point and consider holomorphic coordinates (U, z) centered in a and (V, w) centered in $\pi(a) = b$. Assume, moreover, that $\alpha|_V = dw$, the relation $w = z^3$ holds and the action of $g \in G$ near a is given by $z \mapsto \rho z$ for $\rho \neq 1$ such that $\rho^3 = 1$. By changing ρ with ρ^2 we can assume, moreover, that the decomposition of $H^0(\omega_F)$ in

invariant subspaces with respect to the action of G is the one given in Equation (33). Consider $\eta \in H^0(\omega_F)$. Near a we can write

$$\eta|_U = \left(\sum_{j \geq 0} \eta_j z^j \right) dz \quad \text{and} \quad g^* \eta|_U = \rho \left(\sum_{j \geq 0} \eta_j \rho^j z^j \right) dz.$$

In particular, $\eta \in H^0(\omega_F)^G$ if and only if, near a we have

$$\eta|_U = \left(\sum_{j \geq 0} \eta_{2+3j} z^{2+3j} \right) dz$$

and an analogous decomposition holds near the other ramification points. Similarly, we have

$$\eta|_U = \left(\sum_{j \geq 0} \eta_{3j} z^{3j} \right) dz \quad \text{and} \quad \eta|_U = \left(\sum_{j \geq 0} \eta_{1+3j} z^{1+3j} \right) dz$$

if $\eta \in H^0(\omega_F)_\rho$ and $\eta \in H^0(\omega_F)_{\rho^2}$ respectively.

As

$$\text{Sym}^2(H^0(\omega_F)^-)_1 = H^0(\omega_F)_\rho \otimes H^0(\omega_F)_{\rho^2}, \quad \text{Sym}^2(H^0(\omega_F)^-)_\rho = H^0(\omega_F)_{\rho^2}^{\otimes 2},$$

and

$$\text{Sym}^2(H^0(\omega_F)^-)_{\rho^2} = \text{Sym}^2(H^0(\omega_F)_\rho),$$

if $\varphi \in \text{Sym}^2(H^0(\omega_F)^-)_\mu$ we can write it in coordinate near a as

$$\varphi|_U = z (\varphi_0 + \varphi_1 z^3 + o(z^5)) dz^2$$

for $\mu = 1$ and as

$$\varphi|_U = (\varphi_0 + \varphi_1 z^3 + o(z^5)) dz^2 \quad \text{and} \quad \varphi|_U = z^2 (\varphi_0 + \varphi_1 z^3 + o(z^5)) dz^2$$

if $\mu = \rho$ and $\mu = \rho^2$, respectively. In the latter cases, we have that the residue of $\varphi/\pi^* \alpha$ in a is 0 as $\varphi/\pi^* \alpha$ is either holomorphic or has a pole of order 2 with coefficient of degree -1 equal to 0. Hence

$$\text{Sym}^2(H^0(\omega_F)^-)_\rho \oplus \text{Sym}^2(H^0(\omega_F)^-)_{\rho^2} \subseteq \text{Ker}(d_o P_E^\vee).$$

By Theorem 3.2 and using Diagram (32) we obtain $\dim(\text{Ker}(d_o P_E^\vee)) = 4$. This is equal to the dimension of $\text{Sym}^2(H^0(\omega_F)^-)_\rho \oplus \text{Sym}^2(H^0(\omega_F)^-)_{\rho^2}$ so

$$\text{Sym}^2(H^0(\omega_F)^-)_\rho \oplus \text{Sym}^2(H^0(\omega_F)^-)_{\rho^2} = \text{Ker}(d_o P_E^\vee).$$

In particular, $\beta_1 = 0$ and $\nu(\beta) = 0$ as claimed. □

REFERENCES

- [ACG11] Arbarello E., Cornalba M., Griffiths P. A. *Geometry of algebraic curves. Vol. II.* With a contribution by Joseph Daniel Harris. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 268. Springer, Heidelberg, 2011.
- [ACGH85] Arbarello E., Cornalba M., Griffiths P. A., Harris J. *Geometry of algebraic curves. Vol. I.* Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 267. Springer-Verlag, New York, 1985
- [AP16] Albano A., Pirola G. P. *Dihedral monodromy and Xiao fibrations* Ann. Mat. Pura Appl. (4) 195 (2016), no. 4, 1255–1268

- [BGN15] Barja M. A., Gonzalez-Alonso V., Naranjo, J. C. *Xiao's conjecture for general fibred surfaces* To appear in Journal für die reine und angewandte Mathematik.
- [BHPV04] Barth, W.P., Hulek, K., Peters, C.A.M., Van de Ven, A. *Compact complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 4, second edn. Springer-Verlag, Berlin (2004).
- [Cle72] A. Clebsch *Zur Theorie der Riemann'schen Flächen*, Math. Ann. 6 (1872), 216–230
- [Deb82] O. Debarre, *Inégalités numériques pour les surfaces de type général* (French. English summary) [Numerical inequalities for surfaces of general type] With an appendix by A. Beauville. Bull. Soc. Math. France 110 (1982), no. 3, 319–346.
- [Hor73] Horikawa E. *On deformations of holomorphic maps I*. J. Math. Soc. Japan 25 (1973), 372–396.
- [Hor74] Horikawa E. *On deformations of holomorphic maps II*. J. Math. Soc. Japan 26 (1974), 647–667.
- [Hur91] A. Hurwitz, *Über Riemann'sche Flächen mit gegebenen Verzweigungspunkten*, Math. Ann. 39 (1891), 1–61.
- [Isk77] Iskovskih V. A. *Fano threefolds I*. Math. USSR-Izv. 11 (1977), no. 3, 485–527 (1978)
- [Kan04] Kanev V., *Hurwitz spaces of triple coverings of elliptic curves and moduli spaces of abelian threefolds*. Ann. Mat. Pura Appl. (4) 183 (2004), no. 3, 333–374.
- [Mir95] Miranda R. *Algebraic curves and Riemann surfaces* Graduate Studies in Mathematics, Vol 5, American Mathematical Society, Providence, RI, 1995
- [Pir92] Pirola G. P. *On a conjecture of Xiao* J. Reine Angew. Math. 431 (1992), 75–89.
- [Xia87] Xiao G. *Fibered algebraic surfaces with low slope* Math. Ann. 276 (1987), 449–466.

(Filippo F. Favale) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TRENTO, VIA SOMMARIVE 14,
I-38123 TRENTO, ITALY

E-mail address: `filippo.favale@unitn.it`

(Sara Torelli) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PAVIA, VIA FERRATA, 5, I-27100
PAVIA, ITALY

E-mail address: `sara.torelli02@ateneopv.it`