



Lefschetz properties for jacobian rings of cubic fourfolds and other Artinian algebras

Davide Bricalli¹ · Filippo Francesco Favale²

Received: 2 May 2022 / Accepted: 25 October 2022
© The Author(s) 2022

Abstract

In this paper, we exploit some geometric-differential techniques to prove the strong Lefschetz property in degree 1 for a complete intersection standard Artinian Gorenstein algebra of codimension 6 presented by quadrics. We prove also some strong Lefschetz properties for the same kind of Artinian algebras in higher codimensions. Moreover, we analyze some loci that come naturally into the picture of “special” Artinian algebras: for them we give some geometric descriptions and show a connection between the non emptiness of the so-called non-Lefschetz locus in degree 1 and the “lifting” of a weak Lefschetz property to an algebra from one of its quotients.

Keywords Artinian Gorenstein algebras · Cubic fourfolds · Jacobian rings · Lefschetz properties · Complete intersections

Mathematics Subject Classification Primary 14J35; Secondary 13E10 · 14J70 · 14M10 · 13F20 · 14N05

Introduction

In this article we exploit and generalize some techniques that the authors have developed in previous works on these topics, in collaboration with Gian Pietro Pirola (see [9] and [4]) in order to prove the validity of some Lefschetz properties for particular Artinian algebras. More precisely, we deal with standard Artinian Gorenstein graded algebras over a field \mathbb{K} of characteristic 0. For brevity, we refer to such an algebra using the acronym SAGA. SAGAs are generated in degree 1 and satisfy a sort of “inner duality” property induced by the products of elements in complementary degrees (see Sect. 1.1 for details). We recall that the codimension

✉ Filippo Francesco Favale
filippo.favale@unipv.it

Davide Bricalli
d.bricalli1@campus.unimib.it

¹ Dipartimento di Matematica e Applicazioni, Università degli Studi di Milano-Bicocca, Via Roberto Cozzi, 55, I-20125 Milano, Italy

² Dipartimento di Matematica, Università degli Studi di Pavia, Via Ferrata, 5, I-27100 Pavia, Italy

of a standard Artinian graded algebra R is the dimension of R^1 , i.e. its degree-1 part, as vector space over \mathbb{K} .

Classical examples of these algebras are jacobian rings of smooth hypersurfaces in projective spaces. These are constructed by considering the smooth hypersurface $X = V(F) \subseteq \mathbb{P}^n$ with F homogeneous in $S = \mathbb{K}[x_0, \dots, x_n]$ and by taking the quotient $R = S/I$ where I is the ideal generated by the partial derivatives of F (the so-called jacobian ideal of F). The importance of these objects lies, for example, in the strong geometric relations between X and its jacobian ring. Just to mention some, in the seminal works [5, 6, 12, 13], Carlson, Griffiths, Green and Harris proved that a portion of the primitive part of the Dolbeault cohomology is codified in R and that R plays a crucial role in the study of the infinitesimal variation of Hodge structure of X .

This construction is a special case of a more general one. Indeed, one can also consider the quotient S/I , where I is generated by a regular sequence of length $n + 1$ of homogeneous polynomials, i.e. $I = (f_0, \dots, f_n) \subset \mathbb{K}[x_0, \dots, x_n]$, where $V(f_0, \dots, f_n) \subset \mathbb{P}^n$ is the empty set. This kind of algebras are SAGAs and are also complete intersection rings so it makes sense to refer to them as *complete intersection SAGAs*. For brevity one often says (see, for example, [19]) that R is a *complete intersection SAGA presented by forms of degree e* , if the generators of the ideal I have all the same degree, equal to e . Our main result (Theorem 3.1) will deal with the case $e = 2$, i.e. with complete intersection SAGAs presented by forms of degree 2.

Lefschetz properties for an Artinian algebra were defined in the 80's by taking inspiration from the Hard Lefschetz Theorem for a Kähler variety X of complex dimension n . We recall that this theorem states that for all $k \leq n$ the cup product map $\omega^k \cdot : H^{n-k}(X, \mathbb{C}) \rightarrow H^{n+k}(X, \mathbb{C})$ is an isomorphism for $\omega \in H^2(X, \mathbb{C})$ generic (more precisely, for any Kähler form in $H_{\bar{\partial}}^{1,1}(X)$). Then, the even cohomology ring $R = \bigoplus_{i=0}^n R^i = \bigoplus_{i=0}^n H^{2i}(X, \mathbb{C})$ of X is a Gorenstein Artinian algebra. In particular, R is such that for $x \in R^1$ general, the multiplication map $x^{n-2i} \cdot : R^i \rightarrow R^{n-i}$ is an isomorphism for $i \leq n/2$. This is, roughly, the definition of the *strong Lefschetz property (SLP)* for an Artinian algebra (for details see Definition 1.2). Starting from this idea a weaker version has been introduced by following similar ideas: an algebra R has the *weak Lefschetz property (WLP)* if the multiplication map $x \cdot : R^k \rightarrow R^{k+1}$ is of maximal rank for all $k \geq 0$ and $x \in R^1$ general. For a comprehensive treatment of Lefschetz properties the interested reader can refer to [14].

In the recent years, there has been a growing interest in Artinian algebras and their Lefschetz properties. We summarize here some results and conjectures relevant with respect to the topics of our article. In [15] it is shown that any Artinian standard algebra R of codimension 2 satisfies *SLP*. In the same article, it is also proved that if R has codimension 3 and is a complete intersection then it satisfies *WLP*. Again in codimension 3, if one restricts its attention to jacobian rings of curves of degree d in \mathbb{P}^2 , the validity of *SLP* is known only up to degree 4 with the case for $d = 4$ treated in the very recent work [7]. In codimension 4, again in [7], the authors prove that jacobian rings of cubic surfaces have the *SLP*. In particular, nothing is known for Lefschetz properties for jacobian rings when the degree increases: quintic curves in \mathbb{P}^2 (regarding *SLP*) and quartic surfaces in \mathbb{P}^3 (regarding both *WLP* and *SLP*) are the first open cases. For what concerns the case of complete intersection SAGAs presented by quadrics, in [19] it is shown that *WLP* holds in degree 1 (i.e. $x \cdot : R^1 \rightarrow R^2$ is injective for $x \in R^1$ general) in any codimension. If one wonders if all SAGAs have to satisfy *WLP*, this is known to be false as there are counterexamples (which are not complete intersection) in higher codimension [10] but it is conjectured to hold for complete intersection SAGAs. For codimension bigger than or equal to 5 very little is known. For codimension

5, *WLP* has been proved in [2] for complete intersection SAGAs presented by quadrics whereas, in [4] the authors proved *SLP* for the same SAGAs (in particular, for jacobian rings of smooth cubic threefolds). The cases of higher codimensions or higher degrees are, at the moment, completely open.

Just to mention some other papers, which treat problems related to Lefschetz properties for Artinian Algebras, we cite, for example, [1, 16–18] and [8]. These articles are really interesting on their own since the approaches used to study, roughly, the “same” problems are really different and involve a lot of different techniques.

In this work, we follow the “geometric-differential” approach used in [2] and [4] by exploiting and generalising the techniques of the latter, in order to prove, partially, the *SLP* for the first open case with low degree in higher codimension. More precisely, we deal with a complete intersection SAGA R presented by quadrics of codimension 6 and we prove our first main result:

Theorem (Theorem 3.1) *Let R be a SAGA as above. Then R satisfies the *SLP* in degree 1, i.e. for x general in R^1 the multiplication map $x^4 \cdot : R^1 \rightarrow R^5$ is a bijection.*

This result is a step towards the evidence of a well known conjecture, according to which, in characteristic 0 all complete intersection SAGAs should satisfy the Lefschetz properties (see [14, Conjecture 3.46], for example). Beyond being the first open case, the interest for these SAGAs lies in the fact that they involve Lefschetz properties for the jacobian rings of cubic fourfolds, which are very interesting on their own for what concerns several other unrelated topics (such as, for example, the well known Kuznetsov’s conjecture about the rationality of the cubic fourfolds).

We stress that Lefschetz properties, at least for a *general* complete intersection SAGA S/I with fixed degrees of the generators of I , are known to hold. For this reason, it is interesting to analyze “special” SAGAs. With this idea, we proceed by obtaining two interesting results concerning special SAGAs. The first one is a collection of geometric properties satisfied by the subschemes

$$\mathcal{N}_i := \{[x] \in \mathbb{P}(R^1) \mid x^i = 0\}$$

that we call *Nilpotent loci*. Roughly, when R is “general”, \mathcal{N}_i is empty for low values of i , then, in case of non-emptiness, we do expect that these loci reflect some properties of “special” SAGAs. The most relevant results are Proposition 2.1, which has as a consequence, a characterization of Fermat hypersurfaces (see 2.2) and Theorem 2.6 which analyzes the geometric behaviour of these loci when they contain a linear space.

For the second result, we consider what are called the *non-Lefschetz loci*, i.e. the subschemes parametrising linear forms for which the injectivity of the multiplication map fails (see Definition 1.4). These are really interesting for the geometry and the algebraic structure of a SAGA. Just to mention some articles which study these loci, one can consider [3] and [2]. Our contribution for this topic is the following lifting criterion:

Theorem (Theorem 4.2) *Let $R = S/I$ be a complete intersection SAGA of codimension $n + 1 \geq 6$ presented by quadrics and assume that z is a non-Lefschetz element for R . If $\bar{R} = R/(z)$ satisfies the *WLP* in degree 2, then the same holds for R .*

This is a result for “special” SAGAs since, as for the Nilpotent loci, when R is general, all $[x] \in \mathbb{P}(R^1)$ are Lefschetz elements. This unexpected theorem gives a sort of converse for results which prove that Lefschetz properties are inherited by suitable quotients. This

happens, for example, for *SLP* when one takes the quotient by the conductor of a Lefschetz element ([14, Proposition 3.11]) or, for the *WLP* as shown in the recent article [11].

The plan of the article is the following. In Section 1, we set our notation and we introduce the main characters of the paper. Moreover, we establish a general framework that will be used in several specific situations through the paper, and we prove the first general technical results. In Section 2, we analyze the nilpotent loci \mathcal{N}_i for a SAGA R regardless of the validity of some Lefschetz properties. We study them from a geometric point of view and we recollect some examples which show how the nature of these loci can be very different from one case to another. In Section 3 we prove the first main result, i.e. Theorem 3.1. In Section 4, we analyze the weak Lefschetz property for complete intersection SAGAs presented by quadrics and prove the “lifting” of the *WLP* to R from suitable quotients, i.e. Theorem 4.2. Finally, one might wonder how much the different techniques developed in this paper could be exploited to study the next open cases. In Section 5 we try to answer to this question, by generalizing the validity of some strong Lefschetz properties to complete intersection SAGAs presented by quadrics of higher codimension.

1 Notations, preliminaries and technical results

Through all the article we will assume that \mathbb{K} is an algebraically closed field of characteristic 0. First of all, let us recall two basic definitions that will play an important role in what follows (the reader can refer to [14] for a deeper treatment).

Definition 1.1 Consider an Artinian graded \mathbb{K} -algebra $R = \bigoplus_{i=0}^N R^i$. Then

- the dimension of R^1 as \mathbb{K} -vector space is called **codimension** of R ;
- R is **standard** if it is generated by R^1 as \mathbb{K} -algebra;
- R is said to have the **Gorenstein duality** if $R^N \simeq \mathbb{K}$ and the multiplication map $R^j \times R^{N-j} \rightarrow R^N$ is a perfect pairing whenever $0 \leq j \leq N$.

If R is a graded Artinian algebra, having the Gorenstein duality is equivalent to ask that R is Gorenstein. In this case, R^N is called **socle** of R and $R^0 \simeq R^N$. A standard Artinian Gorenstein algebra will be called, for brevity, **SAGA**.

Examples of SAGAs are algebras that can be written as $\mathbb{K}[x_0, \dots, x_n]/I$ with I an homogeneous ideal generated by a regular sequence of polynomials of length $n + 1$. We will be interested in the case where the generators of I have all the same degree e : for brevity, in this situation we say that R is a **complete intersection SAGA presented by forms of degree e** . Relevant examples of this kind of SAGAs are jacobian rings of smooth hypersurfaces of \mathbb{P}^n .

We now introduce Lefschetz properties for a SAGA R . These involve the multiplication maps by suitable elements of R . In the following we will often deal with kernels of such maps so it is convenient to set

$$K_q^a := \ker(q \cdot : R^a \rightarrow R^{a+s}) \quad \text{for } q \in R^s.$$

Definition 1.2 Consider a SAGA $R = \bigoplus_{i=0}^N R^i$. It is said to satisfy the

- weak Lefschetz property in degree k , (*WLP* $_k$ in short) if there exists $x \in R^1$ such that $x \cdot : R^k \rightarrow R^{k+1}$ has maximal rank;
- strong Lefschetz property in degree k at range s , (*SLP* $_k(s)$ in short) if there exists $x \in R^1$ such that $x^s \cdot : R^k \rightarrow R^{k+s}$ has maximal rank.

The algebra R has the strong Lefschetz property in degree k (SLP_k) if $SLP_k(s)$ holds for all s . We also say that R satisfies **weak (strong) Lefschetz property - WLP** (respectively SLP) in short - if it satisfies WLP_k (respectively SLP_k) for all k .

Remark 1.3 For SAGAs, the above definition of SLP is equivalent to ask $SLP_k(N - 2k)$ for all $k \leq N/2$ (see Definition 3.18 and subsequent discussion in [14]). Moreover, notice that if $k \leq N/2$ and $1 \leq s \leq N - 2k$, $SLP_k(s)$ implies $SLP_k(s - 1)$. Note that, by definition, WLP_k is equivalent to $SLP_k(1)$ and, by Gorenstein duality, WLP_k holds if and only if WLP_{N-k-1} holds. Finally, for $k \leq N/2$, WLP_k implies WLP_{k-1} by [18, Proposition 2.1].

Elements of $\mathbb{P}(R^1)$ for which the multiplication map is not of maximal rank will be important in Section 4 so it is convenient to introduce the following subschemes of $\mathbb{P}(R^1)$.

Definition 1.4 Let R be any SAGA with socle in degree N . For $1 \leq a \leq N - 1$ we define the **Lefschetz locus in degree a** to be

$$\mathcal{L}_a := \{[x] \in \mathbb{P}(R^1) \mid x \cdot : R^a \rightarrow R^{a+1} \text{ has maximal rank}\} \subset \mathbb{P}(R^1).$$

An element $[x] \in \mathbb{P}(R^1)$ (or, equivalently, $x \in R^1 \setminus \{0\}$) is called **Lefschetz element in degree a** if $[x] \in \mathcal{L}_a$. On the contrary, elements not in \mathcal{L}_a are called **non-Lefschetz elements (in degree a)**.

Geometric results on these loci can be found, for example, in [2] and [3].

In the next sections, we will make a large use of the framework that we present here in a general setting. This kind of construction has been used in some articles (see, for example [4] and [2]) in order to study from a geometric perspective SAGAs which do not satisfy some Lefschetz properties.

Let R be a SAGA of codimension $n + 1$ and socle in degree N . For $1 \leq a, b \leq N$ we set $\Gamma_{i,j}^{(a,b)} = \{([x], [y]) \in \mathbb{P}(R^a) \times \mathbb{P}(R^b) \mid x^i y^j = 0\}$ and $\mathcal{N}_k^{(a)} = \{[x] \in \mathbb{P}(R^a) \mid x^k = 0\}$.

We will refer to $\mathcal{N}_k^{(a)}$ as **nilpotent loci** of $\mathbb{P}(R^a)$. For brevity, we will set $\mathcal{N}_k^{(1)} = \mathcal{N}_k$. We will denote by p_1 and p_2 the standard projections from $\Gamma_{i,j}^{(a,b)}$ to $\mathbb{P}(R^a)$ and $\mathbb{P}(R^b)$, respectively. Notice that when we consider $\Gamma_{s,1}^{(a,b)}$, we have

$$p_1^{-1}([x]) = \{([x], [y]) \mid x^s y = 0\} = [x] \times \mathbb{P}(K_{x^s}^b)$$

so all the fibers of p_1 are projective spaces (not necessarily of the same dimension).

We will often assume the failure of some Lefschetz property and this assumption will reflect on a condition on the projection p_1 . More precisely, we make the following observations.

- Assume that $b \leq N/2$ and that $SLP_b(s)$ does not hold. Then, for all $[x] \in \mathbb{P}(R^1)$ we have that the multiplication map $x^s \cdot : R^b \rightarrow R^{b+s}$ is not injective. In particular, there exists $[y] \in \mathbb{P}(R^b)$ such that $x^s y = 0$ in R^{b+s} . This shows that the failure of $SLP_b(s)$ is equivalent to ask that $p_1 : \Gamma_{s,1}^{(1,b)} \rightarrow \mathbb{P}(R^1)$ is surjective.
- Assume that $p_1 : \Gamma_{s,1}^{(a,b)} \rightarrow \mathbb{P}(R^a)$ is surjective. Then, as observed above, we have that all the fibers of p_1 are projective spaces and this implies that there exists a unique irreducible component Θ of $\Gamma_{s,1}^{(a,b)}$ which dominates $\mathbb{P}(R^a)$ via p_1 . In this case, we set

$$\pi_i = p_i|_{\Theta}, \quad Y = p_2(\Theta) = \pi_2(\Theta) \quad \text{and} \quad F_y = \pi_1(\pi_2^{-1}([y])) \text{ for all } [y] \in Y.$$

Construction 1.5 To summarize, if R is a SAGA of codimension $n + 1$ and socle in degree N and we assume that $SLP_b(s)$ does not hold for R , we can construct the loci $\Gamma_{s,1}^{(1,b)}$, Θ , Y and F_y as above and we have the following diagram

$$\begin{array}{ccc}
 F_y \times [y] & \xrightarrow{\pi_2} & [y] \\
 \downarrow \cong & \searrow & \downarrow \\
 & \Theta & Y \\
 & \searrow \pi_2 & \downarrow \\
 & & \mathbb{P}(R^b) \\
 & \searrow \pi_1 & \downarrow p_2 \\
 & & \mathbb{P}(R^1) \\
 F_y \subset & \xrightarrow{\quad} & \mathbb{P}(R^1)
 \end{array} \tag{1}$$

We stress that, in this case, as Θ is the unique irreducible component which dominates $\mathbb{P}(R^1)$, we have

$$\pi_1^{-1}([x]) = p_1^{-1}([x]) = [x] \times \mathbb{P}(K_{x^s}^b) \text{ for general } [x] \in \mathbb{P}(R^1).$$

On the contrary, for specific $[x] \in \mathbb{P}(R^1)$, it can happen that $\pi_1^{-1}([x]) \subsetneq p_1^{-1}([x])$ and that $\pi_1^{-1}([x])$ is not a projective subspace of $[x] \times \mathbb{P}(R^b)$.

In [4] and in [9] the authors have developed a technical lemma which gives strong informations on the locus Θ introduced above. We present now a generalised version of it.

Proposition 1.6 Let T be an irreducible variety in $\mathbb{P}(R^a) \times \mathbb{P}(R^b)$ such that $p_{1|T} : T \rightarrow \mathbb{P}(R^a)$ is surjective. Assume that $T \subseteq \{x^i y^j = 0\} = \Gamma_{i,j}^{(a,b)}$ with $i, j \geq 1$. Then

- (1) For all $v \in R^a$ one has $vx^{i-1}y^{j+1} = 0$;
- (2) If $a(i) + b(j + 1) \leq N$, then all points of T satisfy also $x^{i-1}y^{j+1} = 0$.

In particular, if $a = b = 1$ and $k \leq N - 2$, if $T \subseteq \Gamma_{k,1}^{(1,1)}$ then $T \subseteq \Gamma_{i,j}^{(1,1)}$ for all i, j such that $i + j = k + 1$ and $j \geq 1$.

Proof It is enough to prove the claim for a general smooth point $p = ([x], [y]) \in T$. For any $v \in R^a, t \in \mathbb{K}$ consider $x' = x + tv \in R^a$. Since $p_1 : T \rightarrow \mathbb{P}(R^a)$ is surjective by hypothesis, we have that there exists y' in $R^b \setminus \{0\}$ such that $(x')^i (y')^j = 0$. Then we can define $\beta(t)$ such that $\beta(0) = y$ and $(x + tv)^i (\beta(t))^j = 0$ for all $t \in \mathbb{K}$. We can consider the expansion of β and write this relation as

$$0 \equiv (x + tv)^i (y + tw + t^2(\dots))^j = x^i y^j + t(ivx^{i-1}y^j + jwx^i y^{j-1}) + t^2(\dots).$$

In particular we have $ivx^{i-1}y^j + jwx^i y^{j-1} = 0$ for all $v \in R^a$. If we multiply by y we have $ivx^{i-1}y^{j+1} = 0$ for all $v \in R^a$ which yields the first claim since $i \geq 1$ by hypothesis.

For the second claim, consider the multiplication map $R^a \times R^{(i-1)a+(j+1)b} \rightarrow R^{ia+jb+b}$ and notice that it is non degenerate by the assumption $ia+bj+b \leq N$. Hence, if $ivx^{i-1}y^{j+1} = 0$ for all $v \in R^a$, then one has also $x^{i-1}y^{j+1} = 0$ as claimed. \square

Proposition 1.6 gives some easy but relevant relations involving the dimension of Y, Θ and F_y for $[y] \in Y$ general. We summarize these results in the following

Lemma 1.7 Assume that R is any SAGA of codimension $n + 1$ with socle in degree N and assume that $SLP_b(k)$ does not hold for some integer k such that $1 \leq k \leq N - 2b$. Then, defining $\Gamma_{k,1}^{(1,b)}, \Theta, Y$ and F_y as in 1.5, we have the following properties:

- (a) $\dim(Y) + \dim(F_y) = \dim(\Theta) \geq n = \dim(\mathbb{P}(R^1))$;
- (b) if $b(k + 1) \leq N - 1$ and $b \geq 2$ then $0 \leq \dim(F_y) \leq n - 1$ and $1 \leq \dim(Y) \leq \dim(\mathbb{P}(R^b)) - 1$;
- (c) if $b = 1$ and $k = N - 2$ then $1 \leq \dim(Y) \leq n - 2$ and $2 \leq \dim(F_y) \leq n - 1$.

Proof One can prove (a) and (b) by following the proof of [4, Proposition 2.6] and using Proposition 1.6. The last point has been proved in [4, Corollary 2.11]. □

Through the whole article, if $p \in X$ is a smooth point we will denote by $T_{X,p}$ the differential tangent space. The notation $T_p(X)$ will be used when X is inside a projective space \mathbb{P}^n to mean its embedded Zariski tangent space. In particular, if $[x] \in X \subseteq \mathbb{P}^n$ is a smooth point, and \tilde{X} is the affine cone of X , we have $\mathbb{P}(T_{\tilde{X},x}) = T_{[x]}(X)$. Let us now introduce a simple description of the tangent space for the nilpotent loci which generalize [4, Corollary 4.2]. The proof is analogous with minimal changes.

Lemma 1.8 *Let R be any SAGA. Then for $[\eta] \in \mathcal{N}_k^{(a)}$ general we have $T_{[\eta]}(\mathcal{N}_k^{(a)}) \subseteq \mathbb{P}(K_{\eta^{k-1}}^a)$. If, moreover, $\eta^{k-1} \neq 0$, we have an equality.*

Let us now show a technical result that holds for any SAGA when we deny the $SLP_1(k)$ for some $1 \leq k \leq N/2$. Then, as in 1.5 we can construct $\Theta \subseteq \Gamma_{k,1}^{(1,1)}$, Y and F_y .

Lemma 1.9 *With notations as above, if $p = ([x], [y]) \in \Theta$ is a general point, we have:*

- (a) $y \in T_{F_y,[x]}$ and $x \notin T_{Y,[y]}$;
- (b) $T_{\tilde{F}_y,x}^1 \subseteq K_{x^\alpha y^\beta}^1$ whenever $\alpha + \beta = k$ and $\beta \geq 1$, where \tilde{F}_y denotes the affine cone over F_y .

Proof For (a), let $p = ([x], [y]) \in \Theta$ be a general point. By [4, Proposition 2.6] we have that F_y is a cone and $[y]$ is a vertex for it, so that the line $\langle [x], [y] \rangle$ is contained in F_y . This means that $[y]$ is a tangent vector in $[x]$, i.e. $[y] \in T_{F_y,[x]}$.

For the second claim of (a), let us suppose by contradiction that $[x] \in T_{Y,[y]}$. Let $\tilde{\Theta}$ be the lifting to $R^1 \times R^1$ of $\Theta \subset \mathbb{P}(R^1) \times \mathbb{P}(R^1)$ and let $\tilde{\pi}_2$ be the projection on the second factor from $\tilde{\Theta}$. By construction, we have that $\tilde{\Theta} \subseteq \{(x, y) \mid x^k y = 0\}$. Let \tilde{Y} be the affine cone of Y . Since $\pi_2 : \Theta \rightarrow Y$ is surjective, we have $\tilde{\pi}_2(\tilde{\Theta}) = \tilde{Y}$ and that, for $p = (x, y) \in \tilde{\Theta}$ general,

$$d_p \tilde{\pi}_2 : T_{\tilde{\Theta},p} \rightarrow T_{\tilde{Y},y}$$

is surjective.

By assumption we have also that $x \in T_{\tilde{Y},y}$ so there exists a tangent vector to $\tilde{\Theta}$ at p of the form (v, x) . Since points of $\tilde{\Theta}$ satisfy $x^k y = 0$ we have

$$0 \equiv (x + tv + t^2(\dots))^k (y + tx + t^2(\dots)) \pmod{t^2}$$

which yields $x^{k+1} = 0$. This is impossible by the generality of x : $[x] \notin T_{Y,[y]}$.

For (b), first of all, notice that by Proposition 1.6 we have $\tilde{F}_y \subseteq \{x \in R^1 \mid x^i y^j = 0\}$ for all $i + j = k + 1$ and $i, j \geq 1$. Hence, for p general, if an element $v \in R^1$ belongs to $T_{\tilde{F}_y,x}$ then the following relation must be satisfied

$$0 \equiv (x + tv + t^2(\dots))^i y^j = itvx^{i-1}y^j + t^2(\dots) \pmod{t^2}.$$

Hence, we have that $v \in K_{x^\alpha y^\beta}^1$, with $\alpha + \beta = k$ and $\beta \geq 1$. □

Let us now consider a complete intersection SAGA R : we recall another technical result, dealing with the kernels of the multiplication maps presented above, that will be used extensively in the whole paper.

Proposition 1.10 (Proposition 4.1 of [4]) *Let R be any complete intersection SAGA of codimension $n + 1$ presented in degree $d - 1$. Assume that $1 \leq s \leq N - 1$. The following properties hold:*

- (a) *If $[\eta] \in \mathbb{P}(R^s)$, then $s \geq (d - 2) \dim(K_\eta^1)$;*
- (b) *Let $[\eta], [\zeta] \in \mathbb{P}(R^s)$ and assume $s = (d - 2) \dim(K_\eta^1) = (d - 2) \dim(K_\zeta^1)$. Then $K_\eta^1 = K_\zeta^1$ if and only if $[\eta] = [\zeta]$.*

Finally, let us now focus on the specific case that will be treated in the next sections, that is the case where R is a complete intersection SAGA of codimension $n + 1$ presented in degree $d - 1 = 2$.

Remark 1.11 If $d = 3$, by Proposition 1.10 we get that for $[\eta] \in \mathbb{P}(R^s)$ we have $\dim(K_\eta^1) \leq s$. From this, together with Lemma 1.8, we also get that $\dim(\mathcal{N}_k) \leq k - 2$.

2 Nihilpotent loci and geometric properties

In this section we will study geometric properties of the nihilpotent loci $\mathcal{N}_k \subseteq \mathbb{P}(R^1)$ where R is a complete intersection SAGA of codimension $n + 1$ presented by quadrics. Any R as above has socle in degree $N = n + 1$. We stress that we don't make any assumptions about the validity of any weak or strong Lefschetz property for R in this section.

We recall that the nihilpotent loci (in $\mathbb{P}(R^1)$) are defined as

$$\mathcal{N}_k = \{[x] \in \mathbb{P}(R^1) \mid x^k = 0\}.$$

If $X \subseteq \mathbb{P}^r$ is non-empty, we denote by $\text{Sec}^k(X) \subseteq \mathbb{P}^r$ the k -secant variety associated to X , i.e.

$$\text{Sec}^k(X) := \overline{\bigcup_{p_1, \dots, p_k \in X} \langle p_1, \dots, p_k \rangle}$$

where $\langle p_1, \dots, p_k \rangle$ is the linear span of the points p_1, \dots, p_k . For brevity, we set $\text{Sec}^2(X) := \text{Sec}(X)$. The interested reader can refer to [20, Chapter 1] for various properties of these classical loci (although the definition considered is slightly different from the one adopted by us).

We stress that, like the non-Lefschetz loci $\mathbb{P}(R^1) \setminus \mathcal{L}_k$, the nihilpotent loci \mathcal{N}_k are expected to be empty when k is small for R "general". Hence it is interesting to study these loci when R is "special". For example, these loci give a lot of information for SAGAs for which some Lefschetz properties do not hold.

Let us start by analyzing the locus $\mathcal{N}_2 \subseteq \mathbb{P}(R^1) \simeq \mathbb{P}^n$. We recall that by Lemma 1.8 we have that $\dim(\mathcal{N}_2) \leq 0$ so it is either empty or it is the union of a finite number of points. These points have to satisfy the following:

Proposition 2.1 *Assume that $[t_1], \dots, [t_k] \in \mathcal{N}_2$ are distinct points. Then $\prod_{i=1}^k t_i \neq 0$ in R and $[t_1], \dots, [t_k]$ are in general position in $\mathbb{P}(R^1)$. In particular, $\#\mathcal{N}_2 \leq n + 1 = N$.*

Proof The statement is trivially true for $k = 1$. If $k = 2$ the only statement one has to check is that $t_1 t_2 \neq 0$. This is true since $K_{t_1}^1 = \langle t_1 \rangle$ by Proposition 1.10 and $[t_1] \neq [t_2]$. We will then proceed by induction assuming that the claim is true till $k - 1$.

Let $T = \{[t_1], \dots, [t_k]\}$ be a set of k distinct points of \mathcal{N}_2 . By contradiction, let us assume that either (A_1) or (A_2) holds, where

- (A_1) $\{t_1, \dots, t_k\}$ are linearly dependent
- (A_2) $\prod_{i=1}^k t_i = 0$.

First of all, we claim that (A_2) is equivalent to (A_1) . By induction hypothesis, for $\{z_1, \dots, z_{k-1}\} \subset \{t_1, \dots, t_k\}$ with $[z_i] \neq [z_j]$ for all $i \neq j$, we have $\prod_{i=1}^{k-1} z_i \neq 0$, so $K_{z_1 \dots z_{k-1}}^1$ has dimension at most $k - 1$ by Proposition 1.10. Since $z_i^2 = 0$ by assumption, we have $K_{z_1 \dots z_{k-1}}^1 = \langle z_1, \dots, z_{k-1} \rangle$. Then, (A_1) holds if and only if we have, up to a permutation of the elements, $t_k \in \langle t_1, \dots, t_{k-1} \rangle = K_{t_1 \dots t_{k-1}}^1$ and this is equivalent to $\prod_{i=1}^k t_i = 0$, i.e. (A_2) .

Hence, let us suppose that $t_k \in \langle t_1, \dots, t_{k-1} \rangle$, so we can write $t_k = \sum_{i=1}^{k-1} a_i t_i$. Then we have

$$0 = t_k^2 = 2 \sum_{1 \leq i < j \leq k-1} a_i a_j t_i t_j.$$

If $k = 3$ we have $0 = t_3^2 = 2a_1 a_2 t_1 t_2$ so, since $t_1 t_2 \neq 0$ by induction hypothesis, we have either $a_1 = 0$ or $a_2 = 0$. This implies either $\{t_1, t_3\}$ or $\{t_2, t_3\}$ linearly dependent, and we get a contradiction since this is against the induction hypothesis. If $k \geq 4$, by multiplying by $\prod_{i=1}^{k-3} t_i$, we get

$$0 = 2a_{k-2} a_{k-1} \prod_{i=1}^{k-1} t_i.$$

Since $\prod_{i=1}^{k-1} t_i \neq 0$ by induction hypothesis, we have either $a_{k-2} = 0$ or $a_{k-1} = 0$ and we have a contradiction as in the case $k = 3$. □

By considering the Fermat hypersurface $X = V(F)$ in \mathbb{P}^n , one can easily see that, for the Jacobian ring $R = S/J(F)$, the set \mathcal{N}_2 consists of exactly $n + 1$ independent points. However, also the converse is true, as shown by the following:

Corollary 2.2 *Assume that $\#\mathcal{N}_2 = n + 1$. Then R is the Jacobian ring of a cubic hypersurface X projectively equivalent to the Fermat cubic hypersurface in \mathbb{P}^n .*

Proof By assumption we have that $\mathcal{N}_2 = \{[t_0], \dots, [t_n]\}$. By Proposition 2.1, $\{t_0, \dots, t_n\}$ are $n + 1$ linearly independent forms so $R = S/I$ with $S = \mathbb{K}[t_0, \dots, t_n]$. On the other hand, in S we have $t_i^2 \in I$ and $\{t_0^2, \dots, t_n^2\}$ is a regular sequence which generates I as ideal of S . Then, if we set $F = \sum_{i=0}^n t_i^3$, we have that I is the Jacobian ideal of the Fermat cubic hypersurface $X = V(F)$ as claimed. □

Remark 2.3 We have $\text{Sec}^k(\mathcal{N}_2) \subseteq \mathcal{N}_{k+1}$. Indeed, if $[t_1], \dots, [t_k] \in \mathcal{N}_2$ we have $t_i^2 = 0$. In particular, every monomial of degree $k + 1$ in the variables t_i is identically 0. Then $(\sum_{i=1}^k a_i t_i)^{k+1} \equiv 0$ for all $a_1, \dots, a_k \in \mathbb{K}$ so $\text{Sec}^k(\mathcal{N}_2) \subseteq \mathcal{N}_{k+1}$. More generally,

$$\text{whenever } r > k(a - 1) \text{ one has } \text{Sec}^k(\mathcal{N}_a) \subseteq \mathcal{N}_r.$$

Indeed, consider $[t_1], \dots, [t_k] \in \mathcal{N}_a$ and let $m = \prod_{i=1}^k t_i^{\alpha_i}$ with $\sum_{i=1}^k \alpha_i = r$. We have $m = 0$ if there exists i such that $\alpha_i \geq a$. On the other hand, this always happens if $r > k(a - 1)$: if $\alpha_i \leq (a - 1)$ for all i , we would have

$$r = \sum_{i=1}^k \alpha_i \leq \sum_{i=1}^k (a - 1) = k(a - 1) < r$$

which gives a contradiction.

Lemma 2.4 *If L is a line contained in \mathcal{N}_3 , then $L \subseteq \text{Sec}(\mathcal{N}_2)$, i.e. a line in \mathcal{N}_3 is a line joining two different points of \mathcal{N}_2 .*

Proof Assume that L is a line in \mathcal{N}_3 . Since the dimension of \mathcal{N}_3 is at most 1 by Remark 1.11, we have that L is a component of \mathcal{N}_3 . As $\dim(\mathcal{N}_2) \leq 0$ and $\dim(K_x^1) \leq 1$ for any $x \in R^1$ by Proposition 1.10(a), we can find $[v], [w] \in L$ such that $[v] \neq [w], [v], [w] \notin \mathcal{N}_2$ and $vw \neq 0$. By hypothesis, we have that $(v + tw)^3 = 0$ for all $t \in \mathbb{K}$ so $v^3 = v^2w = vw^2 = w^3 = 0$. Then, $K_{v^2}^1, K_{w^2}^1$ and K_{vw}^1 contain $\langle v, w \rangle$. On the other hand, these subspaces have dimension at most 2 by Proposition 1.10(a) so they coincide with $\langle v, w \rangle$. By Proposition 1.10(b), there exist $\lambda, \mu \in \mathbb{K}$ such that

$$v^2 = \lambda w^2 \quad \text{and} \quad vw = \mu w^2 \tag{2}$$

so we have $(v + tw)^2 = v^2 + 2tvw + t^2w^2 = w^2(t^2 + 2\mu t + \lambda)$.

We claim that $t^2 + 2\mu t + \lambda$ has two distinct roots so L is indeed a line contained in $\text{Sec}(\mathcal{N}_2)$. Assume, on the contrary, that $t^2 + 2\mu t + \lambda$ is a square. This implies that $\mu^2 = \lambda$. Then, from the Equations (2), we obtain

$$v(v - \mu w) = 0 \quad w(v - \mu w) = 0$$

so $v - \mu w \in K_v^1 \cap K_w^1$. By Proposition 1.10 we can conclude that $[v] = [w]$ which is against our assumptions. \square

We will generalize this result in Theorem 2.6 by considering suitable linear subspaces contained in \mathcal{N}_k . We need first the following technical lemma.

Lemma 2.5 *Let $k \geq 2$ and let T be an hypersurface in $\mathbb{P}^k \subset \mathbb{P}(R^1)$. Assume either that*

- (1) $0 \leq s \leq k - 1$ or
- (2) $s = k$ and the support of T is not contained in the union of 2 different hyperplanes.

Then there exist $[x_0], \dots, [x_s] \in T$ which are linearly independent and such that $\prod_{i=0}^s x_i \neq 0$.

Proof The statement of the lemma is clearly true for $s = 0$. We will proceed by induction on $s \leq k$. Then assume that there are $[x_0], \dots, [x_{s-1}] \in T$ which are linearly independent and with $y = x_0 \cdots x_{s-1} \neq 0$. Consider the linear spaces $\tau_1 = \langle x_0, \dots, x_{s-1} \rangle$ and $\tau_2 = \mathbb{P}(K_y^1)$. We are done if we prove that $U = T \setminus (\tau_1 \cup \tau_2)$ is not empty. By construction we have $\dim(\tau_1) = s - 1$ and $\dim(\tau_2) \leq s - 1$ by Proposition 1.10. Hence, if $s < k$, U is an open dense subset of T . If $s = k$ and the support of T is not contained in the union of 2 different hyperplanes, there exists an irreducible component C of T which is different from τ_1 and τ_2 . Then $C \setminus (\tau_1 \cup \tau_2)$ is not empty so U is again not empty as claimed. \square

Theorem 2.6 *Assume that π is a $(k - 1)$ -plane contained in \mathcal{N}_{k+1} . Then*

- (A) $T_k = \pi \cap \mathcal{N}_k$ is an hypersurface of degree k (with possible multiple components) in π ;
- (B) there exist $[x_0], \dots, [x_{k-1}]$ in T_k which are linearly independent, $\prod_{i=0}^{k-1} x_i \neq 0$.

In particular, T_k is non degenerate in π and $\pi \subseteq \text{Sec}^k(\mathcal{N}_k)$.

Proof Notice, first of all, that $\dim(\mathcal{N}_k) \leq k - 2$ by Remark 1.11, so $\pi \setminus \mathcal{N}_k \neq \emptyset$. Then, we can find $\{x_0, \dots, x_{k-1}\}$ linearly independent which span π and such that $x_{k-1} \notin \mathcal{N}_k$. Since $\pi \subseteq \mathcal{N}_{k+1}$, we have that

$$(\alpha_0 x_0 + \dots + \alpha_{k-1} x_{k-1})^{k+1} \equiv 0 \quad \forall \alpha_0, \dots, \alpha_{k-1} \in \mathbb{K}$$

and this is equivalent to say that all monomials of degree $k + 1$ in the variables x_0, \dots, x_{k-1} are 0. Then, if m is a monomial of degree k in these variables, either $m = 0$ or $K_m^1 = \langle x_0, \dots, x_{k-1} \rangle$, by Proposition 1.10. In particular, we have that for each monomial of degree k there exists $\lambda_m \in \mathbb{K}$ with $m = \lambda_m x_{k-1}^k$ (recall that we assumed $x_{k-1} \notin \mathcal{N}_k$). Then

$$(\alpha_0 x_0 + \dots + \alpha_{k-1} x_{k-1})^k = p_k(\underline{\alpha}) x_{k-1}^k \tag{3}$$

where $p_k(\underline{\alpha})$ is a homogeneous polynomial of degree k in the variables $\alpha_0, \dots, \alpha_{k-1}$. It is not 0 since the coefficient of α_{k-1}^k is 1 by construction. By Equation (3), $T_k = \pi \cap \mathcal{N}_k$ is described by the vanishing of $p_k(\underline{\alpha})$. In particular, \mathcal{N}_k is not empty and we have also proved (A).

For (B), if T_k has support which is not contained in 2 different hyperplanes, the thesis follows directly from Lemma 2.5 so we have to discuss only the cases

$$(B_1) : \text{Supp}(T_k) = H_1 \cup H_2 \quad \text{and} \quad (B_2) : \text{Supp}(T_k) = H_1$$

where H_1 and H_2 are distinct hyperplanes.

In both cases (B_1) and (B_2) , there is an hyperplane H_1 of π contained in T_k . We recall that T_k is contained in \mathcal{N}_k by construction. By Lemma 2.5 applied to $H_1 \subset \pi$ we can find $[x_0], \dots, [x_{k-2}] \in H_1$ which are linearly independent and such that $y = \prod_{i=0}^{k-2} x_i \neq 0$. Since $H_1 = \langle [x_0], \dots, [x_{k-2}] \rangle$ and $H_1 \subset \mathcal{N}_k$ we have that all monomials of degree k in the variables x_0, \dots, x_{k-2} are 0. Then, by Proposition 1.10, $K_y^1 = \langle x_0, \dots, x_{k-2} \rangle$ so $H_1 = \mathbb{P}(K_y^1)$.

If we are in case (B_1) we can then choose x_{k-1} in $H_2 \setminus H_1$ and $\{x_0, \dots, x_{k-2}, x_{k-1}\}$ is a set of points with the desired properties. We claim now that case (B_2) can not occur. Assume, by contradiction, that $\text{Supp}(T_k)$ is the hyperplane $H_1 = \mathbb{P}(K_y^1)$. Then for any x_{k-1} in $\pi \setminus H_1$ we have that $\pi = \langle x_0, \dots, x_{k-2}, x_{k-1} \rangle$, $x_{k-1}^k \neq 0$ and $y x_{k-1} \neq 0$. With this choice of the x_i 's, the polynomial $p_k(\underline{\alpha})$ of Equation (3) is proportional to α_{k-1}^k since $T_k = \pi \cap \mathcal{N}_k$ has support on H_1 . On the other hand the coefficient of $\prod_{i=0}^{k-1} \alpha_i$ can not be zero since $\prod_{i=0}^{k-1} x_i = y x_{k-1} \neq 0$. □

We conclude this section by presenting some examples in order to make the phenomenology of the nilpotent loci clearer (some computations have been made by using the computer algebra software Magma). We set $S = \mathbb{K}[x_0, \dots, x_n] = \bigoplus_{k \geq 0} S^k$ and we define $\{w_0, \dots, w_n\}$ to be the projective coordinates on $\mathbb{P}(R^1)$ induced by the basis $\{x_0, \dots, x_n\}$ of $R^1 = S^1$.

Example 2.7 Let X be the Fermat cubic in \mathbb{P}^n and consider the Jacobian ring R of X . For any $2 \leq k \leq n$ we have $(\sum_i w_i x_i)^k \in J^k$ if and only if all monomials in the $\{w_i\}$ of degree k without multiple factors vanish. This is true whenever any set of $n - k + 1$ variables is zero. With these arguments one can prove that \mathcal{N}_k is the union of the coordinated planes of dimension $k - 2$. In particular, $\mathcal{N}_k = \text{Sec}^k(\mathcal{N}_2)$ and $\text{Sing}(\mathcal{N}_k) = \mathcal{N}_{k-1}$ for $k \geq 2$.

Example 2.8 Consider the smooth cubic surface $X = V(f)$ with $f = x_0^3 + x_1^3 + x_2^3 + x_3^3 + 6x_0 x_1 x_2$ and consider the Jacobian ring R of X . One has that there are 4 points in \mathcal{N}_2 so, by

Corollary 2.2, X is the Fermat cubic up to a projective transformation. Indeed, if λ is a non trivial third root of 1, we have

$$(x_0 + x_1 + x_2)^3 + (x_0 - (\lambda + 1)x_1 + \lambda x_2)^3 + (x_0 + \lambda x_1 - (\lambda + 1)x_2)^3 + 3x_3^3 = 3f.$$

Example 2.9 Consider the smooth cubic surface $X = V(f)$ with $f = x_0^3 + x_1^3 + x_2^3 + x_3^3 + 3x_0x_1x_2$ and consider the Jacobian ring R of X . If $P = [0 : 0 : 0 : 1]$ and $C = V(w_3, g)$ is the smooth plane cubic with $g := w_0^3 + w_1^3 + w_2^3 - 6w_0w_1w_2$, we have (considering the reduced structure)

$$\mathcal{N}_2 = \{P\} \quad \mathcal{N}_3 = \{P\} \cup C \quad \mathcal{N}_4 = V(w_3) \cup V(g).$$

In particular, \mathcal{N}_4 is the union of the plane containing the cubic curve C and the cone with vertex P generated by C . Notice that \mathcal{N}_3 does not have pure dimension.

Example 2.10 Consider the smooth cubic surface $X = V(f)$ with $f = x_0^3 + x_1^3 + x_2^3 + x_3^3 + x_0(x_1^2 + x_2^2 + x_3^2)$ and let R be its Jacobian ring. One can show that, in this case, \mathcal{N}_2 and \mathcal{N}_3 are both empty whereas \mathcal{N}_4 is a smooth quartic hypersurface.

Example 2.11 Consider the regular sequence $\{x_0^2, x_1^2, x_2^2, x_3^2 + 2x_0x_1\}$ in $S = \mathbb{K}[x_0, \dots, x_3]$, the ideal J spanned by it and set $R = S/J$. Notice that J is not the Jacobian ideal of a cubic surface. Let P_i be the coordinated points and consider the conic $C = V(w_2, g)$ with $g := w_3^2 - 3w_0w_1$. Then we have

$$\mathcal{N}_2 = \{P_0, P_1, P_2\} \quad \mathcal{N}_3 = \langle P_0, P_1 \rangle \cup \langle P_0, P_2 \rangle \cup \langle P_1, P_2 \rangle \cup C$$

$$\mathcal{N}_4 = V(w_3) \cup V(w_2) \cup V(g).$$

In particular, \mathcal{N}_4 is the union of two planes (the first one - $V(w_3)$ - contains P_0, P_1 and P_2 and the lines joining these points whereas the second - $V(w_2)$ - is the plane containing the conic C and the line $\langle P_0, P_1 \rangle$) and $V(g)$ (which is a quadric cone with vertex P_2). Notice that, as varieties, we have $\text{Sing}(\mathcal{N}_k) = \mathcal{N}_{k-1}$ for $k = 2, 3, 4$.

3 Strong Lefschetz property in degree 1 for codimension 6

In this section we prove our main result, i.e. the following:

Theorem 3.1 *Let R be a complete intersection SAGA presented by quadrics of codimension 6. Then R satisfies the strong Lefschetz property in degree 1 (SLP_1), i.e. the general element $x \in R^1$ is such that the map $x^4 \cdot : R^1 \rightarrow R^5$ is an isomorphism.*

In order to prove the above theorem we will need some technical results (Propositions 3.4, 3.5 and 3.6) which will be treated separately in Subsection 3.1 as they are interesting on their own because they are valid in a much larger framework than the one of the theorem. Now we will assume them and we will deal with the proof of the main theorem.

Proof (of Theorem 3.1) Let us assume by contradiction that the statement does not hold: the map $x^4 \cdot : R^1 \rightarrow R^5$ is never injective for $x \in R^1$. We are then in the situation described in Construction 1.5 with $\Gamma = \Gamma_{4,1}^{(1,1)} = \{([x], [y]) \in \mathbb{P}(R^1) \times \mathbb{P}(R^1) \mid x^4y = 0\}$: we define Θ, Y and F_y as usual. Let us now focus on the dimensions of Y and of F_y for general $[y] \in Y$.

First of all, let us recall that, by Lemma 1.7, we have

$$1 \leq \dim(Y) \leq 3 \quad 2 \leq \dim(F_y) \leq 4 \quad \dim(Y) + \dim(F_y) = \dim(\Theta) \geq 5. \quad (4)$$

Let us now list in the following table the possible values of the pairs $(\dim(Y), \dim(F_y))$. By using the constraints (4) and the various results proved in Subsection 3.1, we can exclude all the cases: in the table below we specify which result rules out each pair. \square

$\dim(Y)$ VS $\dim(F_y)$	2	3	4
1	(4)	(4)	Prop. 3.4
2	(4)	Prop. 3.6 + Rem. 1.11	Prop. 3.4
3	Prop. 3.5	Prop. 3.5	Prop. 3.4

Since no pair as above is possible for our framework, we get a contradiction and this concludes the proof. \square

We have the following easy but important consequence.

Corollary 3.2 *The jacobian ring of a smooth cubic fourfold satisfies the strong Lefschetz property in degree 1.*

3.1 General preparatory results

In this subsection we consider a complete intersection SAGA presented in degree $d - 1$, i.e. $R = S/I$ where $S = \mathbb{K}[x_0, \dots, x_n]$ and I is generated by a regular sequence of forms of degree $d - 1$. In this case, we recall that R is a SAGA of codimension $n + 1$ with socle in degree $N = (d - 2)(n + 1)$. Moreover, we will assume that R is a SAGA which does not satisfy the strong Lefschetz property in degree 1 at range k , i.e. $SLP_1(k)$, with $2 \leq k \leq N - 2$. Equivalently, the multiplication map $x^k \cdot : R^1 \rightarrow R^{k+1}$ is never injective. Hence, we are in the situation described more generally in Section 1. Under the above assumptions we have

$$\begin{array}{ccc}
 F_y \times [y] & \xrightarrow{\pi_2} & [y] \\
 \searrow & & \downarrow \text{hook} \\
 \Theta & \xrightarrow{\pi_2} & Y \\
 \searrow & & \downarrow \text{hook} \\
 \Gamma_k & \xrightarrow{p_2} & \mathbb{P}(R^1) \\
 \downarrow \pi_1 & & \downarrow p_1 \\
 F_y \hookrightarrow & \xrightarrow{\quad} & \mathbb{P}(R^1)
 \end{array} \tag{5}$$

where we have set $\Gamma_k := \Gamma_{k,1}^{(1,1)} = \{([x], [y]) \in \mathbb{P}(R^1) \times \mathbb{P}(R^1) \mid x^k y = 0\}$. We recall that Θ is the unique irreducible component of Γ_k that dominates $\mathbb{P}(R^1)$ via its first projection π_1 , $Y = \pi_2(\Theta)$ and $F_y = \pi_1(\pi_2^{-1}([y]))$ for $[y] \in Y$.

Remark 3.3 We recall that all complete intersection SAGAs presented in degree $d - 1$ satisfy $SLP_1(1) = WLP_1$ (see [19, Proposition 4.3] and [4, Corollary 4.3]). For this reason in this subsection we do not consider $k = 1$ since in this case it is possible to construct Γ_1 , but p_1 is never dominant (so Θ, Y and F_y cannot be constructed).

Now we prove some results giving restrictions on the dimensions of Y and of the general F_y with $[y] \in Y$. We are ultimately interested into the case where $d = 3$. Nevertheless, we stress that the following proposition (3.4) holds for every $d \geq 3$.

Proposition 3.4 *If we assume $n > \frac{k}{d-2}$, then*

$$\dim(F_y) \leq n - 2.$$

In particular, if $d = 3$, then F_y cannot be an hypersurface.

Proof Recall that $\dim(F_y) \leq n - 1$ by Lemma 1.7 so we have to rule out only the case $\dim(F_y) = n - 1$.

Let us assume, by contradiction, that F_y is an hypersurface. Hence, by denoting with \tilde{F}_y the affine cone over F_y , we have $\dim(\tilde{F}_y) = n$.

Recall that $\Theta := \Theta_k \subseteq \Gamma_k = \{([x], [y]) \mid x^k y = 0\}$ by assumption. We will show that the multiplication map $x^{k-1} \cdot : R^1 \rightarrow R^k$ is never injective so we can define, as we have done for Θ_k , an incidence correspondence Γ_{k-1} with a unique irreducible component Θ_{k-1} which dominates $\mathbb{P}(R^1)$ via its first projection. Moreover, we will have $\Theta_k = \Theta_{k-1}$ so F_y is also the fiber of the second projection from Θ_{k-1} and we can iterate this process.

We claim now that $\Theta \subseteq \Gamma_{k-1}$. If $p = ([x], [y]) \in \Theta$ is general, by using Lemma 1.9 and Proposition 1.10 we can conclude

$$n = \dim(T_{\tilde{F}_y, x}) \leq \dim(K_{x^{k-1}y}^1) \leq \frac{k}{d-2}$$

unless $x^{k-1}y = 0$. Since, by hypothesis, we have that $n > k/(d-2)$, the only possibility is that $x^{k-1}y = 0$. In particular we have shown that $\Theta \subseteq \{([x], [y]) \mid x^{k-1}y = 0\} = \Gamma_{k-1}$ as claimed.

Then, Θ is contained in Θ_{k-1} since it dominates $\mathbb{P}(R^1)$. On the other hand, since $\Gamma_{k-1} \subseteq \Gamma_k$, we have also the other inclusion: $\Theta = \Theta_{k-1}$. In particular, the varieties Y and F_y defined for Θ are the same as the ones defined for Θ_{k-1} . Then, by reasoning as before, we obtain $n = \dim(T_{\tilde{F}_y, x}) \leq \dim(K_{x^{k-2}y}^1)$. If we assume that $x^{k-2}y \neq 0$ for p general in Θ , by Proposition 1.10 we would obtain $n \leq (k-1)/(d-2) \leq k/(d-2)$ which is, as before, incompatible with the hypothesis on n . Then $\Theta \subseteq \Gamma_{k-2}$ and we can iterate this process.

By recursion, we reduce ourselves to the case with $k = 1$. We can then see Θ as a subvariety of Γ_1 which dominates $\mathbb{P}(R^1)$ via its first projection. This implies the failure of the weak Lefschetz Property in degree 1. Then, by Remark 3.3, we get a contradiction: F_y has dimension at most $n - 2$, as claimed. \square

Proposition 3.5 *Assume that $d = 3$. Then, the dimension of Y is at most $n - 3$.*

Proof First of all, let us notice that if $k \leq N - 3 = n - 2$, then by Proposition 1.6, Y is contained in $\mathcal{N}_{k+1} \subseteq \mathcal{N}_{n-1}$, whose dimension is at most $n - 3$ (see Lemma 1.8). Hence, we easily get that in this case $\dim(Y) \leq n - 3$ as claimed.

Let us now consider the remaining case: $k = N - 2 = n - 1$. Recall that $\dim(Y) \leq n - 2$ by Lemma 1.7 so, to prove the proposition, we only have to rule out the case $\dim(Y) = n - 2$. Assume by contradiction that $\dim(Y) = n - 2$. Since $k = n - 1$, proceeding as above, we have that Y is contained in \mathcal{N}_n , which has dimension at most $n - 2$. Hence, we get that Y is a component of \mathcal{N}_n . Then, if $[y] \in Y$ is a general point ($y^{n-1} \neq 0$, since $Y \not\subseteq \mathcal{N}_{n-1}$, for dimension reasons), we can write $T_y(Y) = T_y(\mathcal{N}_n) = \mathbb{P}(K_{y^{n-1}}^1)$ by Lemma 1.8. However, the general $[x] \in F_y$ belongs to $\mathbb{P}(K_{y^{n-1}}^1)$ and then to $T_y(Y)$, contradicting Lemma 1.9. \square

Proposition 3.6 *Assume that $d = 3$. Then, the following conditions are not compatible:*

- (a) *for $y \in Y$ general, $\dim(F_y) = k - 1$;*

(b) $Y \not\subseteq \mathcal{N}_{k-1}$.

Proof Notice that for $k = 1$, condition (a) cannot hold since otherwise we would have $\dim(Y) = n$, which is impossible. Hence we can assume that $k \geq 2$.

Let us observe that by Proposition 1.6 we have that $x^\alpha y^\beta = 0$ for every α, β such that $\alpha + \beta = k + 1$ and $\beta \geq 1$; in particular we have $y^{k+1} = 0$ and $Y \subseteq \mathcal{N}_{k+1}$.

Let us assume by contradiction that both conditions (a) and (b) hold. By (b) we have $y^{k-1} \neq 0$ for $y \in Y$ general so for $([x], [y])$ general in Θ we have $xy^{k-1} \neq 0$. Indeed, otherwise, F_y would be contained in $\mathbb{P}(K_{y^{k-1}}^1)$, whose dimension is at most $k - 2$, which is impossible by assumption. As a consequence, we have that

$$\text{for } ([x], [y]) \in \Theta \text{ general, } x^\alpha y^\beta \neq 0 \text{ for } \alpha + \beta = k \text{ with } \alpha, \beta \geq 1 \tag{6}$$

since, otherwise, by using Proposition 1.6, we would obtain also $xy^{k-1} = 0$.

By property (6) and since $\dim(F_y) = k - 1$ by assumption, we also have

$$T_x(F_y) = \mathbb{P}(K_{x^{k-1}y}^1) = \mathbb{P}(K_{x^{k-2}y^2}^1) \tag{7}$$

by Lemma 1.9.

Let us now claim that

$$\text{for } ([x], [y]) \in \Theta \text{ general, } T_y(Y) \subseteq T_x(F_y). \tag{8}$$

To show this, first of all, recall that $Y \not\subseteq \mathcal{N}_{k-1}$ and $Y \subseteq \mathcal{N}_{k+1}$. Let us now consider two cases:

$$1) Y \not\subseteq \mathcal{N}_k \quad \text{and} \quad 2) Y \subseteq \mathcal{N}_k.$$

Assume that $p = ([x], [y]) \in \Theta$ is general (so that $[y]$ is general in Y and $[x]$ is general in F_y). In the first case, since Y is contained in \mathcal{N}_{k+1} , we have $T_y(Y) \subseteq \mathbb{P}(K_{y^k}^1)$ by Lemma 1.8. Moreover, we have that $\mathbb{P}(K_{y^k}^1) = T_x(F_y)$ since $y^k \neq 0$ and $\dim(F_y) = k - 1$. Analogously, for the second case we have $T_y(Y) \subseteq \mathbb{P}(K_{y^{k-1}}^1) \subseteq \mathbb{P}(K_{xy^{k-1}}^1) = T_x(F_y)$. Here, we have used that $xy^{k-1} \neq 0$ since p is general (by property (6)).

Consider, as in Lemma 1.9, the affine cone \tilde{Y} of Y , the lifting $\tilde{\Theta}$ of $\Theta \subset \mathbb{P}(R^1) \times \mathbb{P}(R^1)$ to $R^1 \times R^1$ and its projection $\tilde{\pi}_2$ on the second factor. By construction, we have that $\tilde{\Theta} \subseteq \{(x, y) \mid x^\alpha y^\beta = 0\} = \tilde{\Gamma}_{\alpha, \beta}$ whenever $\alpha + \beta = k + 1$ and $\beta \geq 1$. As in Lemma 1.9, for $p = (x, y) \in \tilde{\Theta}$ general, the differential map

$$d_p \tilde{\pi}_2 : T_{\tilde{\Theta}, p} \rightarrow T_{\tilde{Y}, y}$$

is surjective.

Let us take any w in $T_{\tilde{Y}, y}$. By Property (8), we have that its class $[w]$ belongs to $T_x(F_y)$. Moreover, by the surjectivity of $d_p \tilde{\pi}_2$, we can take in $T_{\tilde{\Theta}, p}$ an element of the form (v, w) .

The tangent space to $\tilde{\Theta}$ in p is a subspace of the tangent space $T_{\alpha, \beta} = T_{\tilde{\Gamma}_{\alpha, \beta}, p}$ to the locus $\tilde{\Gamma}_{\alpha, \beta}$ in p , so we have $(v, w) \in T_{\alpha, \beta}$. Then, we have

$$0 \equiv (x + tv + t^2(\dots))^\alpha (y + tw + t^2(\dots))^\beta \pmod{t^2}$$

for α, β as above. In particular, by taking (α, β) equal to $(k - 1, 2)$ and $(k, 1)$ we obtain the following relations satisfied by (v, w) :

$$(k - 1)vx^{k-2}y^2 + 2x^{k-1}yw = 0 \quad kvx^{k-1}y + x^k w = 0.$$

Since $[w] \in T_x(F_y)$, by property (7), we have $x^{k-1}yw = 0$. Then, from the first equation we get $v \in K_{x^{k-2}y^2}^1 = K_{x^{k-1}y}^1$ (again by property (7)). Hence the second equation yields $x^k w = 0$. In conclusion, we have proven that $T_y(Y) \subset \mathbb{P}(K_{x^k}^1) = \pi_1^{-1}([x])$. We stress that the last equality holds since $[x]$ is general and then, the whole fiber over $[x]$ with respect to p_1 is contained in Θ so $\mathbb{P}(K_{x^k}^1) = p_1^{-1}([x]) = \pi_1^{-1}([x])$.

This easily brings to a contradiction. Indeed, the above property implies that $\dim(Y) \leq \dim(\pi_1^{-1}([x]))$ for $[x] \in \mathbb{P}(R^1)$ general, and since

$$\dim(\Theta) = \dim(Y) + \dim(F_y) = \dim(\pi_1^{-1}([x])) + n$$

we also get $\dim(Y) \leq \dim(Y) + \dim(F_y) - n$, which is impossible by Lemma 1.7(c). \square

4 A lifting criterion for weak Lefschetz property

It is known that, the *SLP* for a graded algebra is inherited to its quotients by suitable conductor ideals (see, for example, [14, Proposition 3.11]). In this section we prove a sort of converse for *WLP₂* for complete intersection SAGAs presented by quadrics. More precisely, we will give a criterion to reduce the proof of *WLP₂* for a SAGA R as above to a suitable quotient of R , modulo the existence of a non-Lefschetz element. We stress, moreover, that this criterion works for any codimension.

Let us start by setting $S = \mathbb{K}[x_0, \dots, x_n]$ and by proving the following result.

Lemma 4.1 *Assume that $R = S/J$ is a complete intersection SAGA of codimension $n + 1$ presented by quadrics so the socle is in degree $N = n + 1$. Assume that there are $w, z \in R^1 \setminus \{0\}$ such that $zw = 0$. Then $(z) = (0 : w)$ and $\bar{R} = R/(z)$ is a complete intersection SAGA of codimension n presented by quadrics. In particular, $\dim(K_w^s) = \dim(R^{s-1}) - \dim(K_z^{s-1})$.*

Proof By definition, we have $(z) \subseteq (0 : w)$, so we can define an epimorphism of graded \mathbb{K} -algebras

$$\varphi : \bar{R} := R/(z) \rightarrow \tilde{R} := R/(0 : w).$$

By [9, Lemma 2.3], the latter is a SAGA of codimension n and socle in degree $\tilde{N} = n$. Considering \bar{R} , it is clearly an Artinian standard algebra of codimension n . We want to show that \bar{R} is also a complete intersection SAGA presented by quadrics. By hypothesis, we know that $zw \in J$, so we can complete $\{zw\}$ to a regular sequence of the form $\{g_0, \dots, g_{n-1}, zw\}$ spanning J . Notice that, by construction, g_0, \dots, g_{n-1} do not belong to the ideal (z) and the reductions \bar{g}_i of g_i modulo (z) are a regular sequence of quadrics in the polynomial ring $\bar{S} = S/(z)$.

Hence we have

$$\bar{R} = R/(z) = \frac{S/J}{(z)} \simeq \frac{\bar{S}}{(\bar{g}_0, \dots, \bar{g}_{n-1})}$$

so \bar{R} is a complete intersection SAGA presented by quadrics. In particular, it has socle in degree $\tilde{N} = n = \tilde{N}$.

Since φ is an epimorphism and preserve the degrees, the image of a generator $\bar{\sigma}$ of \bar{R}^n is a non-zero multiple of the generator $\tilde{\sigma}$ of \tilde{R}^n . This also implies the injectivity of φ . Indeed, let us take a non-zero element $x \in \bar{R}^i$. There exists $y \in \bar{R}^{n-i}$ such that $xy = \bar{\sigma}$. Hence, we have

$$\lambda \tilde{\sigma} = \varphi(\bar{\sigma}) = \varphi(xy) = \varphi(x)\varphi(y),$$

and so we get that $\varphi(x)$ can not be zero and φ is an isomorphism. In particular, for all s ,

$$R^{s-1} \cdot z = (z)_s = (0 : w)_s = K_w^s$$

then we clearly have

$$\dim(K_w^s) = \dim(R^{s-1}z) = \dim(R^{s-1}) - \dim(K_z^{s-1})$$

as claimed. □

We assume again that $R = S/I$ is a complete intersection SAGA presented by quadrics with codimension $n + 1$ so that it has socle in degree $N = n + 1$. We recall that $[z] \in \mathbb{P}(R^1)$ is a non-Lefschetz element (in degree 1), i.e. $z \notin \mathcal{L}_1$, if and only if there exists $[w] \in \mathbb{P}(R^1)$ such that $zw = 0$ in R or, equivalently $K_z^1 = \langle w \rangle$.

Theorem 4.2 *Let $R = S/I$ be a complete intersection SAGA of codimension $n + 1 \geq 6$ presented by quadrics and assume that z is a non-Lefschetz element for R . If $\bar{R} = R/(z)$ satisfies the WLP_2 , then the same holds for R .*

Proof First of all, by Lemma 4.1, we have that $\bar{R} = R/(z) = R/(0 : w)$ is a complete intersection SAGA presented by quadrics and $K_z^1 = \langle w \rangle$. Let pr_z be the projection $R \rightarrow \bar{R}$.

Assume, by contradiction, that WLP_2 holds for \bar{R} but not for R . In particular, for all $x \in R^1$, the multiplication map $x \cdot : R^2 \rightarrow R^3$ has non trivial kernel, i.e. $K_x^2 \neq \{0\}$. Consider the incidence correspondence

$$\Gamma = \{([x], [v]) \in \mathbb{P}(R^1) \times \mathbb{P}(R^1) \mid xvz = 0\}$$

with its projections p_1 and p_2 on the factors.

We claim that p_1 is surjective. Since Γ is a closed subset, it is enough to show that for $[x] \in \mathbb{P}(R^1)$ general there exists $[v] \in \mathbb{P}(R^1)$ such that $xvz = 0$. Let x be a general element of R^1 . As $K_x^2 \neq 0$ we have that there exists $[q] \in \mathbb{P}(R^2)$ such that $xq = 0$ in R . Then we have also $\text{pr}_z(xq) = \bar{x}\bar{q} = 0$ in \bar{R} . Since $[x]$ is general in $\mathbb{P}(R^1)$, then the same holds for $[\bar{x}] \in \mathbb{P}(\bar{R}^1)$, so we get $\bar{q} = 0$ in \bar{R}^2 , as WLP_2 holds for \bar{R} by assumption. Then, by Lemma 4.1, we have $q \in (0 : w)_2 = (z)_2 = z \cdot R^1$ so there exists $[v] \in \mathbb{P}(R^1)$ such that $0 = xq = xvz$ as claimed.

In analogy with what happens for the construction 1.5, we have that there exists a unique irreducible component Θ of Γ which dominates $\mathbb{P}(R^1)$ via π_1 , where we set π_i to be the restriction of p_i to Θ for $i = 1, 2$. We have that for $[x] \in \mathbb{P}(R^1)$ general

$$\pi_1^{-1}([x]) = p_1^{-1}([x]) = [x] \times \mathbb{P}(K_{xz}^1)$$

so the general fiber of π_1 has dimension at most 1 by Proposition 1.10.

Let us now show that the general fiber of π_1 has dimension 1. Consider $[x] \in \mathbb{P}(R^1)$ general. Firstly, let us observe that $([x], [w])$ belongs to $p_1^{-1}([x]) = \pi_1^{-1}([x])$ since $zw = 0$. As shown above, there exists $[q] \in \mathbb{P}(R^2)$ such that $xq = 0$ and $q = zv$ for suitable $[v] \in \mathbb{P}(R^1)$. Moreover $[v] \neq [w]$ since, otherwise, $[q]$ would be zero, hence $\pi_1^{-1}([x]) = \langle [w], [v] \rangle$ as claimed.

By considering the second projection π_2 , we have that for $[v]$ general in $Y = \pi_2(\Theta)$, the fiber $\pi_2^{-1}([v])$ is such that

$$\pi_2^{-1}([v]) \subseteq p_2^{-1}([v]) = \mathbb{P}(K_{vz}^1) \times [v],$$

which has dimension at most 1 by Proposition 1.10. Since π_1 is dominant, for $([x], [v]) \in \Theta$ general we have

$$n + 1 = \dim(\mathbb{P}(R^1)) + \dim(\pi_1^{-1}([x])) = \dim(\Theta) = \dim(Y) + \dim(\pi_2^{-1}([v])).$$

Since $\dim(Y) \leq n$ and $\dim(\pi_2^{-1}([v])) \leq 1$, for v general, the only possibility is to have $\dim(\pi_2^{-1}([v])) = 1$ and $Y = \mathbb{P}(R^1)$.

We will show now that having $Y = \mathbb{P}(R^1)$ gives a contradiction. First of all, by reasoning as in Proposition 1.6, one can prove that

$$Y \subseteq \{[v] \in \mathbb{P}(R^1) \mid v^2 z = 0\}.$$

Since $Y = \mathbb{P}(R^1)$ and squares of elements of R^1 generates R^2 (as R is standard), we have that $z \cdot R^2 = 0$. This is impossible by Gorenstein duality. \square

In the statement of Theorem 4.2 we require $n + 1 \leq 6$ since for codimension 5 the WLP_2 has already been proved (in [2] or in [4] as consequence of SLP) and in even smaller codimension, it easily follows from WLP_1 that is known to hold. From this, we get the following consequence.

Corollary 4.3 *Let R be a complete intersection SAGA of codimension 6 presented by quadrics (e.g. R is the jacobian ring of a cubic fourfold). If \mathcal{L}_1 is not the whole $\mathbb{P}(R^1)$, R satisfies WLP .*

5 Some results for higher codimensions

In this section, our aim is to obtain some results concerning Lefschetz properties for complete intersection SAGAs presented by quadrics with codimension equal to $n + 1 \geq 4$ by using techniques and results developed in the previous sections.

In particular, our framework will be as follows. Let us consider the polynomial ring $S = \mathbb{K}[x_0 \cdots, x_n]$, an ideal I of S , generated by a regular sequence of quadrics (g_0, \cdots, g_n) and the corresponding complete intersection SAGA $R = S/I$, which has codimension $n + 1$ and socle in degree $N = n + 1$. We recall that we have denoted by $SLP_1(k)$ the strong Lefschetz property in degree 1 at range k : a SAGA R will satisfy the property $SLP_1(k)$ if the multiplication map $x^k \cdot : R^1 \rightarrow R^{k+1}$ is injective for $[x]$ general in $\mathbb{P}(R^1)$. We recall that we have defined the loci $\Gamma_{i,j}^{(a,b)}$ in Section 1. For brevity, we will write Γ_k to mean $\Gamma_{k,1}^{(1,1)}$.

We will prove the following:

Theorem 5.1 *Let $k \in \{2, 3, 4\}$ and assume that R is a complete intersection SAGA of codimension $n + 1$ presented by quadrics. If $n \geq k + 1$ we have that R satisfies $SLP_1(k)$.*

We will show the above theorem by splitting up the proof in 3 cases, which will be treated in the Propositions 5.2, 5.3 and 5.5 respectively.

Proposition 5.2 *Property $SLP_1(2)$ holds for every $n \geq 3$.*

Proof Let us assume by contradiction that the multiplication map $x^2 \cdot : R^1 \rightarrow R^3$ is not injective for $[x] \in \mathbb{P}(R^1)$. Then we can consider the locus $\Gamma_2 = \{([x], [y]) \in \mathbb{P}(R^1) \times \mathbb{P}(R^1) \mid x^2 y = 0\}$, with the corresponding varieties Θ , Y and F_y defined as we have usually done. By Proposition 1.6, we obtain that $Y \subseteq \mathcal{N}_3$. Then, by Lemmas 1.8 and 1.7 we have that $\dim(Y) = 1$: hence F_y must be a hypersurface, which is absurd by Proposition 3.4. \square

The following result has been already proved when $n = 4$ in [4].

Proposition 5.3 *Property $SLP_1(3)$ holds for every $n \geq 4$.*

Proof Let us assume by contradiction that the multiplication map $x^3 \cdot : R^1 \rightarrow R^4$ is not injective for $[x] \in \mathbb{P}(R^1)$. As in the proof of Proposition 5.2, let us construct Γ_3, Θ, Y and F_y . In this case, we have that $Y \subseteq \mathcal{N}_4$ so $\dim(Y) \leq 2$. If $[y] \in Y$ is general, the only possible value for $(\dim(Y), \dim(F_y))$ is $(2, n - 2)$ since F_y can not be a hypersurface by Proposition 3.4. By dimension reasons, $Y \not\subseteq \mathcal{N}_3$, thus we have that the general element $[y]$ of Y is such that $y^3 \neq 0$. Then, by Proposition 1.6, we get F_y must be contained in $\mathbb{P}(K_{y^3}^1)$, whose dimension is at most 2. Then we have proved that $n - 2 = \dim(F_y) \leq 2$ which is impossible for $n \geq 5$. \square

Before showing the analogous result for the $SLP_1(4)$, let us prove a technical result.

Lemma 5.4 *Let R be a SAGA as at the beginning of Section 5 and consider $4 \leq k \leq n - 1$. Assume that R does not satisfy $SLP_1(k)$ so one can consider the varieties Γ_k, Θ, Y and F_y constructed as in Section 1. For $[y] \in Y$ general we have the following properties.*

- (a) *If R satisfies $SLP_1(k - 1)$, then $\dim(F_y) \leq k - 1$;*
- (b) *$(\dim(Y), \dim(F_y)) \neq (k - 1, k - 1)$.*

Proof For (a), let us assume by contradiction that for $[y] \in Y$ general, $\dim(F_y) = h \geq k$. Then, by Lemma 1.9, we have that for $[x] \in F_y$ general

$$T_{[x]}(F_y) = \mathbb{P}(T_{\tilde{F}_{y,x}}) \subseteq \mathbb{P}(K_{x^\alpha y^\beta}^1),$$

where $\alpha + \beta = k$, with $\beta \geq 1$ and \tilde{F}_y is the affine cone over F_y . But since $SLP_1(k - 1)$ holds for R by hypothesis, for $([x], [y]) \in \Theta$ general, we have that $x^{k-1}y \neq 0$, and so

$$h = \dim(F_y) \leq \dim(\mathbb{P}(K_{x^{k-1}y}^1)) \leq k - 1,$$

which is impossible by the assumptions over h .

For (b), let us consider $[y] \in Y$ general and assume by contradiction that $\dim(Y) = \dim(F_y) = k - 1$. By Proposition 1.6 we get that $Y \subseteq \mathcal{N}_{k+1}$, and by Lemma 1.8 we deduce that Y is an irreducible component of \mathcal{N}_{k+1} and we have that $y^k \neq 0$. By reasoning as in the proof of Proposition 5.3 and by Proposition 1.10, we get $F_y = \mathbb{P}(K_{y^k}^1)$. Moreover, since $[y]$ is general in Y , we have $T_{[y]}(Y) = \mathbb{P}(K_{y^k}^1) = F_y$. With these conditions, we can proceed as in the proof of Proposition 3.6 and consider for example the equations

$$0 \equiv (x + tv)^{k-1}(y + tw)^2 \pmod{t^2} \quad 0 \equiv (x + tv)^k(y + tw) \pmod{t^2},$$

where $w \in T_{\tilde{Y},y}$ and $(v, w) \in T_{\tilde{\Theta},(x,y)}$. In this way, we get that $Y \subseteq \pi_1^{-1}([x])$, where, as usual, $\pi_1 : \Theta \subseteq \mathbb{P}(R^1) \times \mathbb{P}(R^1) \rightarrow \mathbb{P}(R^1)$ is the first projection. This leads to a contradiction as shown in Proposition 3.6. \square

We can now show the last case we need to prove Theorem 5.1.

Proposition 5.5 *Property $SLP_1(4)$ holds for every $n \geq 5$.*

Proof First of all, let us notice that the statement has been already proved for $n = 5$ in Section 3: we have to prove $SLP_1(4)$ for $n \geq 6$.

Let us assume that for $x \in R^1$, the multiplication map $x^4 \cdot : R^1 \rightarrow R^5$ is not injective. As usual, we can then consider the incidence correspondence $\Gamma_4 = \{([x], [y]) \in \mathbb{P}(R^1) \times \mathbb{P}(R^1) \mid x^4y = 0\}$ and the corresponding varieties Θ, Y and F_y , for $[y] \in Y$ general.

By Proposition 1.6, we get that $Y \subseteq \mathcal{N}_5$, so $\dim(Y) \leq 3$. By using the bounds of Lemma 1.7(a) and Proposition 3.4, the only possible cases for the values of $(\dim(Y), \dim(F_y))$ are

$$(2, n - 2) \quad (3, n - 2) \quad (3, n - 3).$$

By Proposition 5.3, we know that $SLP_1(3)$ holds for $n \geq 6$. Then, by Lemma 5.4(a), we get that $\dim(F_y)$ is at most 3: the cases $(\dim(Y), \dim(F_y)) = (2, n - 2)$ and $(\dim(Y), \dim(F_y)) = (3, n - 2)$ can not occur for every $n \geq 6$. We also have that $(\dim(Y), \dim(F_y)) \neq (3, n - 3)$ for every $n \geq 7$.

The only case we have still to analyze is the one with $n = 6$ and $\dim(Y) = \dim(F_y) = 3$. By considering Lemma 5.4(b), we can rule out this last possibility too: $SLP_1(4)$ holds for R , for every $n \geq 5$. □

We conclude this section by observing how much can be easily said, by using these methods, for the SLP_1 for complete intersection SAGA of codimension 7 (i.e. $n = 6$) presented by quadrics (e.g. for the Jacobian ring of a smooth cubic fivefold).

Corollary 5.6 *Let R be a complete intersection SAGA of codimension 7 presented by quadrics. We have that $SLP_1(4)$ holds. Moreover, if $SLP_1(5)$ does not hold (i.e. if SLP_1 does not hold), then one can construct the varieties Γ_5, Θ, Y and F_y as usual and we have, for $[y] \in Y$ general, $(\dim(Y), \dim(F_y)) \in \{(2, 4), (3, 3)\}$.*

Proof Property $SLP_1(4)$ holds by Theorem 5.1. Assume that $SLP_1(5)$ does not hold for R . Then, by Lemma 1.7, we have

$$1 \leq \dim(Y) \leq 4 \quad 2 \leq \dim(F_y) \leq 5 \quad \dim(Y) + \dim(F_y) = \dim(\Theta) \geq 6. \quad (9)$$

As in Theorem 3.1, we put in a table the possible values of the pairs $(\dim(Y), \dim(F_y))$ and we specify which result rules out the corresponding case. This concludes the proof.

$\dim(Y)$ VS $\dim(F_y)$	2	3	4	5
1	(9)	(9)	(9)	Prop. 3.4
2	(9)	(9)		Prop. 3.4
3	(9)		Prop. 3.6 + Rem. 1.11	Prop. 3.4
4	Prop. 3.5	Prop. 3.5	Prop. 3.5	Prop. 3.4

This concludes the proof. □

Acknowledgements The authors are thankful to Gian Pietro Pirola for his constant support and his priceless suggestions and to Alberto Alzati and Riccardo Re for the helpful discussions on the topics of the paper. The authors are partially supported by INdAM - GNSAGA and by PRIN “Moduli spaces and Lie theory” and by (MIUR): Dipartimenti di Eccellenza Program (2018-2022) - Dept. of Math. Univ. of Pavia.

Funding Open access funding provided by Universit[Pleaseinsertintopreamble] degli Studi di Pavia within the CRUI-CARE Agreement.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Abdallah, N., Altafi, N., Iarrobino, A., Seceleanu, A., Yaméogo, J.: Lefschetz properties of some codimension three Artinian Gorenstein algebras. (2022)
2. Alzati, A., Re, R.: Complete intersections of quadrics and the weak Lefschetz property. *Collect. Math.* **70**(2), 283–294 (2019)
3. Boij, M., Migliore, J., Miró-Roig, R.M., Nagel, U.: The non-Lefschetz locus. *J. Algebra* **505**, 288–320 (2018). <https://doi.org/10.1016/j.jalgebra.2018.03.006>
4. Bricalli, D., Favale, F., Pirola, G. P.: A theorem of Gordan and Noether via Gorenstein Rings, Preprint available at (2022). [arXiv:2201.07550](https://arxiv.org/abs/2201.07550)
5. Carlson, J., Griffiths, P.A.: Infinitesimal variations of Hodge structure and the global Torelli problem, *Journées de Géométrie Algébrique d'Angers, Juillet 1979, Algebraic Geometry, Angers,; Sijthoff & Noordhoff. Alphen aan den Rijn–Germantown, Md.* **1980**, 51–76 (1979)
6. Carlson, J., Green, M., Griffiths, P.A., Harris, J.: Infinitesimal variations of Hodge structure. I, *Compositio Math.* **50**(2–3), 109–205 (1983)
7. Dimca, A., Gondim, R., Ilardi, G.: Higher order Jacobians, Hessians and Milnor algebras. *Collect. Math.* **71**(3), 407–425 (2020). <https://doi.org/10.1007/s13348-019-00266-1>
8. Dimca, A., Ilardi, G.: Lefschetz properties of Jacobian algebras and singularities of hyperplane sections. [arXiv:2202.02233](https://arxiv.org/abs/2202.02233). (2022)
9. Favale, F.F., Pirola, G.P.: Infinitesimal variation functions for families of smooth varieties. *Appear Milan J. Math.* **3**, 1–20 (2022)
10. Gondim, R., Zappalà, G.: Lefschetz properties for Artinian Gorenstein algebras presented by quadrics. *Proc. Amer. Math. Soc.* **146**(3), 993–1003 (2018). <https://doi.org/10.1090/proc/13822>
11. Guerrieri, L.: Lefschetz properties of Gorenstein graded algebras associated to the Apéry set of a numerical semigroup. *Ark. Mat.* **57**(1), 85–106 (2019). <https://doi.org/10.4310/ARKIV.2019.v57.n1.a5>
12. Griffiths, P.A.: Infinitesimal variations of Hodge structure III. Determinantal varieties and the infinitesimal invariant of normal functions. *Composit. Math.* **50**(2–3), 267–324 (1983)
13. Griffiths, P.A., Harris, J.: Infinitesimal variations of Hodge structure. II. An infinitesimal invariant of Hodge classes. *Compositio Math.* **50**(2–3), 207–265 (1983)
14. Harima, T., Maeno, T., Morita, H., Numata, Y., Wachi, A., Watanabe, J.: The Lefschetz Properties. *Lecture Notes in Mathematics*, pp. 20–80. Springer, Heidelberg (2013)
15. Harima, T., Migliore, J., Nagel, U., Watanabe, J.: The weak and strong Lefschetz properties for Artinian K -algebras. *J. Algebra* **262**(1), 99–126 (2003). [https://doi.org/10.1016/S0021-8693\(03\)00038-3](https://doi.org/10.1016/S0021-8693(03)00038-3)
16. Ilardi, G.: Jacobian ideals, arrangements and the Lefschetz properties. *J. Algebra* **508**, 418–430 (2018). <https://doi.org/10.1016/j.jalgebra.2018.04.029>
17. Mezzetti, E., Miró-Roig, R.M., Ottaviani, G.: Laplace equations and the weak Lefschetz property. *Canad. J. Math.* **65**(3), 634–654 (2013). <https://doi.org/10.4153/CJM-2012-033-x>
18. Migliore, J.C., Miró-Roig, R.M., Nagel, U.: Monomial ideals, almost complete intersections and the weak Lefschetz property. *Trans. Amer. Math. Soc.* **363**(1), 229–257 (2011). <https://doi.org/10.1090/S0002-9947-2010-05127-X>
19. Migliore, J., Nagel, U.: Gorenstein algebras presented by quadrics. *Collect. Math.* **64**(2), 211–233 (2013). <https://doi.org/10.1007/s13348-012-0076-x>
20. Russo, F.: On the geometry of some special projective varieties. *Lecture Notes of the Unione Matematica Italiana*, Springer, Unione Matematica Italiana, Cham, Bologna (2016)