

# The Cauchy Problem for a non Strictly Hyperbolic $3 \times 3$ System of Conservation Laws Arising in Polymer Flooding

Graziano Guerra\*      Wen Shen†

January 19, 2021

## Abstract

We study the Cauchy problem of a  $3 \times 3$  system of conservation laws modeling two-phase flow of polymer flooding in rough porous media with possibly discontinuous permeability function. The system loses strict hyperbolicity in some regions of the domain where the eigenvalues of different families coincide, and BV estimates are not available in general. For a suitable  $2 \times 2$  system, a singular change of variable introduced by Temple [8, 14] could be effective to control the total variation [11]. An extension of this technique can be applied to a  $3 \times 3$  system only under strict hypotheses on the flux functions [4]. In this paper, through an adapted front tracking algorithm we prove the existence of solutions for the Cauchy problem under mild assumptions on the flux function, using a compensated compactness argument.

**Keywords:** Conservation Laws; Discontinuous Flux; Compensated Compactness; Polymer Flooding; Wave Front Tracking; Degenerate Systems

**AMS subject classifications:** 35L65; 35L45; 35L80; 35L40; 35L60

## 1 Introduction

We consider a simple model for polymer flooding in two phase flow in rough media

$$\begin{cases} \partial_t s + \partial_x f(s, c, k) & = 0, \\ \partial_t [cs] + \partial_x [cf(s, c, k)] & = 0, \\ \partial_t k & = 0, \end{cases} \quad (1.1)$$

associated with the initial data

$$(s, c, k)(0, x) = (\bar{s}, \bar{c}, \bar{k})(x), \quad x \in \mathbb{R}. \quad (1.2)$$

Here, the unknown vector is  $(s, c, k)$ , where  $s$  is the saturation of water phase,  $c$  is the fraction of polymer dissolved in water, and  $k$  denotes the permeability of the porous media. We see that  $k$  does not change in time,  $k(t, x) = k(x)$ , and the initial data  $\bar{k}(\cdot)$  might be discontinuous.

---

\*Department of Mathematics and its Applications, University of Milano - Bicocca, Italy, (graziano.guerra@unimib.it).

†Mathematics Department, Pennsylvania State University, USA, (wxs27@psu.edu).

We neglect both the adsorption of polymers in the porous media and the gravitational force, where the solution to the Riemann problem becomes more complex. For such Riemann solvers, see [9, 12] for the effect of the adsorption term, and [11] for the addition of the gravitational force term. In particular, when the adsorption effect is included, the  $c$  family described below would no longer be linearly degenerate, while adding the gravitational force term, the  $s$  waves described below could have negative speed. Both effects would disrupt the carefully designed wave front tracking algorithm we use to prove the main theorem.

The conserved quantities and their fluxes are given by, respectively

$$\mathbf{G}(s, c, k) = \begin{pmatrix} s \\ cs \\ k \end{pmatrix}, \quad \mathbf{F}(s, c, k) = \begin{pmatrix} f(s, c, k) \\ cf(s, c, k) \\ 0 \end{pmatrix}.$$

Denoting the three families as the  $s$ ,  $c$  and  $k$  family, we have the following 3 eigenvalues as functions of the variables  $(\sigma, \gamma, \kappa)$  in the  $(s, c, k)$  space.

$$\lambda_s = \partial_\sigma f(\sigma, \gamma, \kappa), \quad \lambda_c = \frac{f(\sigma, \gamma, \kappa)}{\sigma}, \quad \lambda_k = 0,$$

and the three corresponding right eigenvectors (in the  $(s, c, k)$  space):

$$r_s = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad r_c = \begin{pmatrix} -\partial_\gamma f(\sigma, \gamma, \kappa) \\ \partial_\sigma f(\sigma, \gamma, \kappa) - \frac{f(\sigma, \gamma, \kappa)}{\sigma} \\ 0 \end{pmatrix}, \quad r_k = \begin{pmatrix} -\partial_\kappa f(\sigma, \gamma, \kappa) \\ 0 \\ \partial_\sigma f(\sigma, \gamma, \kappa) \end{pmatrix}.$$

A straight computation shows that both the  $c$  and  $k$  families are linearly degenerate. Furthermore, there exist regions in the domain such that  $\lambda_s = \lambda_c$  or  $\lambda_s = \lambda_c = \lambda_k$ , where the system is parabolic degenerate.

The flux function  $f(\sigma, \gamma, \kappa)$  has the following properties. For any given  $(\gamma, \kappa)$ , the mapping  $\sigma \mapsto f(\sigma, \gamma, \kappa)$  is the well-known S-shaped Buckley-Leverett function [3] with a single inflection point, see Fig. 1. To be specific, we have

$$f(\sigma, \gamma, \kappa) \in [0, 1], \quad \partial_\sigma f(\sigma, \gamma, \kappa) \geq 0, \quad \text{for all } (\sigma, \gamma, \kappa),$$

and, for all  $(\gamma, \kappa)$ ,

$$\begin{aligned} f(0, \gamma, \kappa) &= 0, & f(1, \gamma, \kappa) &= 1, \\ \partial_\sigma f(0, \gamma, \kappa) &= 0, & \partial_\sigma f(1, \gamma, \kappa) &= 0, \\ \partial_{\sigma\sigma} f(0, \gamma, \kappa) &> 0, & \partial_{\sigma\sigma} f(1, \gamma, \kappa) &< 0. \end{aligned} \tag{1.3}$$

Remark that conditions (1.3) guarantee that the eigenvalues and the eigenvectors written above are well defined (can be extended) when  $\sigma = 0$ . For each given  $(\gamma, \kappa)$ , there exists a unique  $\sigma^*(\gamma, \kappa) \in ]0, 1[$  such that

$$\partial_{\sigma\sigma} f(\sigma^*(\gamma, \kappa), \gamma, \kappa) = 0.$$

A detailed analysis of the wave properties for this system is carried out in [13], with the following highlights:

- $k$  waves are the slowest with speed 0. Both  $f$  and  $c$  are continuous across any  $k$  wave;

- $c$  waves travel with non negative speed. Both  $\frac{f}{s}$  and  $k$  are continuous across any  $c$  wave;
- $s$  waves travel with positive speed. Both  $c$  and  $k$  are continuous across any  $s$  wave.

In [13], the global Riemann solver is constructed. Here we give a brief summary. Let  $(s_l, c_l, k_l)$  and  $(s_r, c_r, k_r)$  be the left and right state of a Riemann problem, respectively. In general, the solution of the Riemann problem consists of a  $k$  wave, a  $c$  wave and possibly some  $s$  waves. They can be constructed as follows.

- Let  $(s_m, c_l, k_r)$  denote the right state of the  $k$  wave. The value  $s_m$  is uniquely determined by the condition

$$f(s_m, c_l, k_r) = f(s_l, c_l, k_l).$$

- For the remaining waves, we have  $k \equiv k_r$  throughout. We then solve the Riemann problem for the  $2 \times 2$  sub-system

$$\partial_t s + \partial_x f(s, c, k_r) = 0, \quad \partial_t (cs) + \partial_x (cf(s, c, k_r)) = 0 \quad (1.4)$$

with Riemann data  $(s_m, c_l)$  and  $(s_r, c_r)$  as the left and right states. The solution consists of waves with non-negative speed.

We now give a precise definition of weak solution to the Cauchy problem (1.1)–(1.2) and state the main theorem.

**Definition 1.1.** *The vector-valued function  $(s, c, k) \in \mathbf{L}^\infty([0, +\infty) \times \mathbb{R}, [0, 1]^3)$  is a solution to the Cauchy problem (1.1)–(1.2) if for any  $\phi \in \mathbf{C}_c^1([0, +\infty) \times \mathbb{R}, \mathbb{R})$  the following equalities hold*

$$\begin{aligned} \int_{\Omega} [s \partial_t \phi + f(s, c, k) \partial_x \phi](t, x) dt dx + \int_{\mathbb{R}} \bar{s}(x) \phi(0, x) dx &= 0, \\ \int_{\Omega} [cs \partial_t \phi + cf(s, c, k) \partial_x \phi](t, x) dt dx + \int_{\mathbb{R}} \bar{c}(x) \bar{s}(x) \phi(0, x) dx &= 0, \\ k(t, x) &= \bar{k}(x), \quad \forall (t, x) \in \Omega, \end{aligned}$$

where  $\Omega = ]0, +\infty[ \times \mathbb{R}$ .

**Theorem 1.1.** *If the initial data  $(\bar{s}, \bar{c}, \bar{k})$  satisfy*

$$\bar{s} \in \mathbf{L}^\infty(\mathbb{R}, [0, 1]), \quad \bar{c} \in \mathbf{BV}(\mathbb{R}, [0, 1]), \quad \bar{k} \in \mathbf{BV}(\mathbb{R}, [0, 1]),$$

*then there exists a solution to the Cauchy problem (1.1)–(1.2) in the sense of Definition 1.1.*

We emphasize the fact that  $s=0$  is not excluded in our theorem since we do not make use of Lagrangian coordinates which would have required  $s > 0$ . Indeed, under the hypotheses  $s \geq s^* > 0$ ,  $k(t, x) = \text{const.}$ , system (1.1) is equivalent to its Lagrangian formulation [15]:

$$\begin{cases} \partial_t \left( \frac{1}{s} \right) - \partial_y \left( \frac{f(s, c, k)}{s} \right) &= 0, \\ \partial_t c &= 0, \\ k &= \text{const.}, \end{cases} \quad (1.5)$$

where  $y$  is the Lagrangian coordinate satisfying  $\partial_x y = s$ ,  $\partial_t y = -f(s, c, k)$ . Therefore, under some additional hypotheses, the result in [2], which holds for *scalar* equations since it is based on the maximum principle for Hamilton–Jacobi equation, could be used to prove the existence of a unique vanishing viscosity solution to the first equation in (1.5) and hence a (in some sense) unique solution to system (1.5). However, since we consider the case where  $s$  can become 0, the analysis in [2] cannot be applied. Instead, we need to solve the original non triangular  $2 \times 2$  system in Eulerian coordinates. Furthermore, since we consider rough permeability function  $k$ , the corresponding system in the Lagrangian coordinate is no longer triangular,

$$\begin{cases} \partial_t \left(\frac{1}{s}\right) - \partial_y \left(\frac{f(s, c, k)}{s}\right) & = 0, \\ \partial_t c & = 0, \\ \partial_t \left(\frac{k}{s}\right) - \partial_y \left(\frac{k f(s, c, k)}{s}\right) & = 0. \end{cases}$$

Remark that the Lagrangian coordinates introduced in [2, Section 6] for (1.1) make the system triangular, but still require  $s \geq s^* > 0$  and moreover they mix time and space, therefore the Cauchy problem for (1.1) in Eulerian coordinates is not equivalent to the Cauchy problem in the coordinates introduced in [2, Section 6].

In this paper, the proof for the existence of solution is carried out by showing that wave front tracking approximate solutions are compact by a compensated compactness argument (see for instance [10] for an application of compensated compactness to a  $2 \times 2$  bi-dimensional related model).

The remaining of the paper is organized as follows. In Section 2 the wave front tracking approximate solutions are constructed. In Section 3 we prove the necessary entropy estimates. Finally in Section 4 the compensated compactness argument is carried out to prove Theorem 1.1.

## 2 Front Tracking Approximations

In this section we modify the algorithm constructed in [4] and [11] to adapt it to system (1.1). We define the functions (see Figure 1):

$$g(\sigma, \gamma, \kappa) = \frac{f(\sigma, \gamma, \kappa)}{\sigma}, \quad P(\sigma, \gamma, \kappa) = \int_0^\sigma |\partial_\xi g(\xi, \gamma, \kappa)| d\xi. \quad (2.6)$$

Since at  $\sigma=0$  both  $f$  and its derivative  $\partial_\sigma f$  vanish, we define  $g(0, \gamma, \kappa) = 0$ . Hypotheses (1.3) imply that  $\partial_\sigma g(0, \gamma, \kappa) > 0$  and that the function  $\sigma \mapsto g(\sigma, \gamma, \kappa)$  has one single maximum point somewhere between the single inflexion point of  $f$  and the point  $\sigma = 1$ . The function  $P$  is continuous with respect to its three variables and strictly increasing and invertible with respect to the variable  $\sigma$ . Fixing initial data  $(\bar{s}, \bar{c}, \bar{k})$  satisfying the hypotheses of Theorem 1.1 and fixing an approximation parameter  $\varepsilon > 0$ , we can construct piecewise constant approximate initial data  $(\bar{s}^\varepsilon, \bar{c}^\varepsilon, \bar{k}^\varepsilon)$  with values in  $[0, 1]^3$  such that:

$$\begin{aligned} \|\bar{k}^\varepsilon - \bar{k}\|_{\mathbf{L}^\infty} &< \varepsilon, & \text{Tot. Var. } \bar{k}^\varepsilon &\leq \text{Tot. Var. } \bar{k}, \\ \|\bar{c}^\varepsilon - \bar{c}\|_{\mathbf{L}^\infty} &< \varepsilon, & \text{Tot. Var. } \bar{c}^\varepsilon &\leq \text{Tot. Var. } \bar{c}, \\ \|\bar{s}^\varepsilon - \bar{s}\|_{\mathbf{L}^1\left(\left(-\frac{1}{\varepsilon}, \frac{1}{\varepsilon}\right), \mathbb{R}\right)} &\leq \varepsilon, & & \end{aligned} \quad (2.7)$$




Figure 1: Diagrams of the flux  $\sigma \mapsto f(\sigma, \gamma, \kappa)$  and of the function  $\sigma \mapsto g(\sigma, \gamma, \kappa) = f(\sigma, \gamma, \kappa)/\sigma$  for fixed values of  $\gamma$  and  $\kappa$ .

Let  $\bar{x}_1, \dots, \bar{x}_N$  be the set of points in which  $\bar{k}^\varepsilon$  has jumps such that

$$\bar{k}^\varepsilon(x) = k_0 \chi_{]-\infty, \bar{x}_1]} + \sum_{i=1}^{N-1} k_i \chi_{] \bar{x}_i, \bar{x}_{i+1}]}(x) + k_N \chi_{] \bar{x}_N, +\infty[},$$

and let  $\bar{y}_1, \dots, \bar{y}_M$  be the set of points in which  $\bar{c}^\varepsilon$  has jumps such that

$$\bar{c}^\varepsilon(x) = c_0 \chi_{]-\infty, \bar{y}_1]} + \sum_{j=1}^{M-1} c_j \chi_{] \bar{y}_j, \bar{y}_{j+1}]}(x) + c_M \chi_{] \bar{y}_M, +\infty[}.$$

Without loss of generality, we can suppose that no  $\bar{y}_j$  coincides with any  $\bar{x}_i$ . Define the constant

$$L = \left[ \frac{1}{\varepsilon} \sup_{\gamma, \kappa} \|f(\cdot, \gamma, \kappa)\|_{\mathbf{C}^2} \right] \cdot (N + M),$$

where  $\lceil \alpha \rceil$  denotes the least integer greater than or equal to the real number  $\alpha$ . In the following we denote by  $\wedge$  the logical operator **and**. We consider the following finite sets of possible values for the function  $g$ :

$$\begin{aligned} \mathcal{G}_0^1 &= \left\{ g \mid g = g(\bar{s}^\varepsilon(x), \bar{c}^\varepsilon(x), \bar{k}^\varepsilon(x)), \quad x \in \mathbb{R} \right\}, \\ \mathcal{G}_0^2 &= \left\{ g \mid g = g\left(\frac{\ell}{L}, c_j, k_i\right), \quad i = 0, \dots, N, j = 0, \dots, M, \ell = 0, \dots, L \right\}, \\ \mathcal{G}_0^3 &= \left\{ g \mid g = \max_{0 \leq \sigma \leq 1} g(\sigma, c_j, k_i), \quad i = 0, \dots, N, j = 0, \dots, M \right\}, \\ \mathcal{G}_0^4 &= \left\{ g \mid g = g(\sigma, c_j, k_i) = g(\sigma, c_{j^*}, k_{i^*}) \wedge g_s(\sigma, c_j, k_i) \cdot g_s(\sigma, c_{j^*}, k_{i^*}) < 0, \right. \\ &\quad \left. \text{for some } i, i^* = 0, \dots, N, j, j^* = 0, \dots, M, \sigma \in [0, 1] \right\}, \\ \mathcal{G}_0 &= \mathcal{G}_0^1 \cup \mathcal{G}_0^2 \cup \mathcal{G}_0^3 \cup \mathcal{G}_0^4. \end{aligned}$$

**Remark 2.1.**

- The set  $\mathcal{G}_0^1$  includes all the possible initial values for  $g$ ;




Figure 2: Wave front tracking pattern.  $k$  waves in blue,  $c$  waves in green and  $s$  waves in red

- The set  $\mathcal{G}_0^2$  includes a sufficiently fine grid for  $g$  in order that any  $s$  grid that contains all the counter images of  $\mathcal{G}_0^2$  through  $g(\cdot, c_j, k_i)$ , for any fixed  $j, i$ , is finer than  $\frac{1}{L}$ ;
- The set  $\mathcal{G}_0^3$  includes all the possible maxima of  $g(\cdot, c_j, k_i)$  for any  $j, i$ ;
- The set  $\mathcal{G}_0^4$  includes all the possible values of  $g$  where two graphs of functions of type  $g(\cdot, c_j, k_i)$  intersect with derivatives of different sign. Because of the shape of  $g$ , this set too is finite.

We start the front tracking algorithm from the region  $x < \bar{x}_1$ . For this purpose we define the allowed values for  $s$  in that region:

$$\mathcal{S}_{0,j} = \{\sigma \mid g(\sigma, c_j, k_0) \in \mathcal{G}_0\}.$$

We call  $f^{0,j}(\sigma)$  the linear interpolation of the map  $\sigma \mapsto f(\sigma, c_j, k_0)$  according to the points in  $\mathcal{S}_{0,j}$ . Observe that, since we have included  $\mathcal{G}_0^2$  in  $\mathcal{G}_0$ , the set  $\{\frac{\ell}{L}\}_{\ell=0}^L$  is included in  $\mathcal{S}_{0,j}$  for every  $j$ , hence we have the uniform estimate

$$\begin{cases} |f(\sigma, c_j, k_0) - f^{0,j}(\sigma)| \leq \varepsilon, \\ |\partial_\sigma f(\sigma, c_j, k_0) - \partial_\sigma f^{0,j}(\sigma)| \leq \frac{\varepsilon}{N+M}, \end{cases} \quad \text{for all } \sigma \in [0, 1], j = 0, \dots, M.$$

We solve all the Riemann problems in  $x < \bar{x}_1$  at  $t=0$  in the following way. Let  $\bar{x} \in ]\bar{y}_j, \bar{y}_{j+1}[$  ( $\bar{y}_0 = -\infty$ ) be a jump in  $\bar{s}^\varepsilon$ . Here we take the entropic solution to the Riemann problem

$$\partial_t s + \partial_x f^{0,j}(s) = 0, \quad s(0, x) = \begin{cases} \bar{s}^\varepsilon(\bar{x}-) & \text{for } x < \bar{x}, \\ \bar{s}^\varepsilon(\bar{x}+) & \text{for } x > \bar{x}. \end{cases}$$

Since  $f^{0,j}$  is piecewise linear, the solution to the Riemann problem is piecewise constant and takes values in the set  $\mathcal{S}_{0,j}$  of the kink points of  $f^{0,j}$ , moreover the entropy condition




Figure 3: The graphs of  $G^L$  and  $G^R$  are drawn respectively in blue and red. For each graph, a possible transition level  $\gamma$  (the level at which given  $G^L$  and  $G^R$  intersect) is drawn in green.

in [1, Theorem 4.4] is satisfied. The same Riemann solver is used whenever at  $t > 0$  two  $s$  waves interact in some region  $\Omega_{0,j}$  defined below.

At the points  $\bar{y}_j$ , we solve the Riemann problem according to the minimum jump condition described in [11] (see also [7]) that we briefly outline (see Fig 3). Define

$$s^L = \bar{s}^\varepsilon(\bar{y}_j^-), \quad c^L = \bar{c}^\varepsilon(\bar{y}_j^-) = c_{j-1}, \quad s^R = \bar{s}^\varepsilon(\bar{y}_j^+), \quad c^R = \bar{c}^\varepsilon(\bar{y}_j^+) = c_j,$$

and the two auxiliary monotone functions (the first one non increasing and the second one non decreasing)

$$G^L(\sigma) = \begin{cases} \max \{g(\varsigma, c^L, k_0) \mid \varsigma \in [\sigma, s^L]\}, & \text{for } \sigma \leq s^L, \\ \min \{g(\varsigma, c^L, k_0) \mid \varsigma \in [s^L, \sigma]\}, & \text{for } \sigma \geq s^L, \end{cases}$$

$$G^R(\sigma) = \begin{cases} \min \{g(\varsigma, c^R, k_0) \mid \varsigma \in [\sigma, s^R]\}, & \text{for } \sigma \leq s^R, \\ \max \{g(\varsigma, c^R, k_0) \mid \varsigma \in [s^R, \sigma]\}, & \text{for } \sigma \geq s^R. \end{cases}$$

Call  $\gamma$  the unique level at which  $G^L$  and  $G^R$  intersect. Because of the hypotheses on  $g$ ,  $\gamma$  is equal to either  $g(s^L, c^L, k_0)$ ,  $g(s^R, c^R, k_0)$ , a maximum of either  $g(\cdot, c^L, k_0)$  or  $g(\cdot, c^R, k_0)$ , or a point in which these two function intersect with derivatives having opposite sign. In any case  $\gamma \in \mathcal{G}_0$  holds. Define the closed intervals

$$I^L = [G^L]^{-1}(\{\gamma\}), \quad I^R = [G^R]^{-1}(\{\gamma\}).$$

Finally call  $s^-$  and  $s^+$  respectively the unique projections of  $s^L$  and  $s^R$  on the closed strictly convex sets  $I^L$  and  $I^R$ . It is not difficult to show that

$$\begin{aligned}\gamma &= G^L(s^-) = g(s^-, c^L, k_0), & s^L, s^- &\in \mathcal{S}_{0,j-1}, \\ \gamma &= G^R(s^+) = g(s^+, c^R, k_0), & s^R, s^+ &\in \mathcal{S}_{0,j}.\end{aligned}$$

Take any wave, with left and right states  $s_l, s_r \in \mathcal{S}_{0,j-1}$ , of the entropic solution to the Riemann problem

$$\partial_t s + \partial_x f^{0,j-1}(s) = 0, \quad s(0, x) = \begin{cases} s^L & \text{for } x < \bar{y}_j, \\ s^- & \text{for } x > \bar{y}_j, \end{cases} \quad (2.8)$$

and suppose  $s^L < s^-$ . Then,  $s^L \leq s_l < s_r \leq s^-$  and, because of the entropy condition [1, Theorem 4.4], its speed satisfies ( $f^{0,j-1}$  coincides with  $f(\cdot, c^L, k_0)$  on  $\mathcal{S}_{0,j-1}$ ):

$$\begin{aligned}\lambda_s &= \frac{f(s_r, c^L, k_0) - f(s_l, c^L, k_0)}{s_r - s_l} \leq \frac{f(s^-, c^L, k_0) - f(s_l, c^L, k_0)}{s^- - s_l} \\ &= g(s^-, c^L, k_0) + s_l \frac{g(s^-, c^L, k_0) - g(s_l, c^L, k_0)}{s^- - s_l} \\ &= \lambda_c + s_l \frac{G^L(s^-) - g(s_l, c^L, k_0)}{s^- - s_l} \\ &\leq \lambda_c,\end{aligned}$$

with  $\lambda_c = \gamma = g(s^-, c^L, k_0) = g(s^+, c^R, k_0)$  and where we used the definition of  $G^L$ . If instead  $s^- < s^L$ , then  $s^- \leq s_r < s_l \leq s^L$  and, as in the previous computation, we have

$$\lambda_s \leq \lambda_c + s_l \frac{G^L(s^-) - g(s_l, c^L, k_0)}{s^- - s_l} \leq \lambda_c,$$

Therefore, in any case, the solution to the Riemann problem (2.8) can be patched with a  $c$  wave that travel with speed  $\lambda_c$  and connect the left state  $(s^-, c^L, k_0)$  to the right state  $(s^+, c^R, k_0)$ . Similar computations can be done at the right of the  $c$  wave so that the complete solution includes a  $c$  wave travelling with speed  $\lambda_c$ , possibly together some entropic  $s$  waves to its left (solutions to  $\partial_t s + \partial_x f^{0,j-1}(s) = 0$ ) and some entropic  $s$  waves to its right (solutions to  $\partial_t s + \partial_x f^{0,j}(s) = 0$ ). We also use this Riemann solver whenever, for  $t > 0$ , a  $c$  wave interact with one or more  $s$  waves.

We point out the following properties of this Riemann solver that will be needed in the proof of the main theorem.

- The  $c$  wave satisfies Rankine-Hugoniot

$$\begin{cases} f(s^+, c^R, k_0) - f(s^-, c^L, k_0) = \lambda_c (s^+ - s^-), \\ c^R f(s^+, c^R, k_0) - c^L f(s^-, c^L, k_0) = \lambda_c (c^R s^+ - c^L s^-). \end{cases}$$

- The  $c$  wave is an “admissible” path as defined in [11] and satisfies the following entropy condition:

– if  $s^- < s^+$  there exists  $s^* \in [s^-, s^+]$  such that

$$\begin{cases} g(\sigma, c^L, k_0) \geq \lambda_c & \text{for all } \sigma \in [s^-, s^*], \\ g(\sigma, c^R, k_0) \geq \lambda_c & \text{for all } \sigma \in [s^*, s^+]. \end{cases} \quad (2.9)$$



– if  $s^+ < s^-$  there exists  $s^* \in [s^+, s^-]$  such that

$$\begin{cases} g(\sigma, c^R, k_0) \leq \lambda_c & \text{for all } \sigma \in [s^+, s^*], \\ g(\sigma, c^L, k_0) \leq \lambda_c & \text{for all } \sigma \in [s^*, s^-]. \end{cases} \quad (2.10)$$

Let  $y_1(t), \dots, y_M(t)$  denote all the  $c$  wave fronts at the time  $t$ . Their initial positions are the discontinuity points of  $c^\varepsilon$ . We will show that they do not interact between each other and keep the same number and order as time goes on.

We define the open regions

$$\Omega_{0,j} = \{(t, x) \in [0, +\infty) \times \mathbb{R} \mid x < \bar{x}_1 \wedge y_j(t) < x < y_{j+1}(t)\},$$

and the flux

$$F^\varepsilon(t, x, \sigma) = f^{0,j}(\sigma), \text{ for } (t, x) \in \Omega_{0,j}.$$

The wave front tracking approximation  $s^\varepsilon$  so constructed is an exact weak entropic solution to

$$\begin{cases} \partial_t s^\varepsilon + \partial_x [F^\varepsilon(t, x, s^\varepsilon)] = 0, \\ \partial_t (c^\varepsilon s^\varepsilon) + \partial_x [c^\varepsilon F^\varepsilon(t, x, s^\varepsilon)]_x = 0, \end{cases}$$

in the region  $x < \bar{x}_1$ .

Since the  $c$  family is linearly degenerate,  $c$  waves will not interact with each other. Indeed, given two consecutive  $c$  waves located respectively in  $y_j(t)$  and  $y_{j+1}(t)$ , the first conservation law in the previous system implies

$$\begin{aligned} \frac{d}{dt} \int_{y_j(t)}^{y_{j+1}(t)} s^\varepsilon(t, x) dx &= \dot{y}_{j+1}(t) s^\varepsilon(t, y_{j+1}(t)-) - \dot{y}_j(t) s^\varepsilon(t, y_j(t)+) \\ &\quad - f^{0,j}(s^\varepsilon(t, y_{j+1}(t)-)) + f^{0,j}(s^\varepsilon(t, y_j(t)+)) = 0 \end{aligned}$$

since  $\dot{y} = \lambda_c = \frac{f}{\sigma}$ . Hence  $c$  waves cannot interact with each other (if  $s^\varepsilon = 0$  between two  $c$  waves, then they both must travel with zero speed and therefore even in this case they cannot interact).

Since any interaction with the  $k$  wave located at  $\bar{x}_1$  cannot give rise to waves entering the region  $x < \bar{x}_1$ , following [11], the wave front tracking algorithm can be carried out for all times in that region. Observe that for a fixed  $\varepsilon$  the total variation of the singular variable  $P$  introduced in (2.6) is bounded. Since the grid  $\mathcal{S}_{0,j}$  contains all the possible maximum points of  $g(\cdot, c_j, k_0)$ , in the regions  $\Omega_{0,j}$ ,  $P(\sigma, c_j, k_0) = \int_0^\sigma \left| \partial_\xi \frac{f^{0,j}(\xi)}{\xi} \right| d\xi$  for any  $\sigma \in \mathcal{S}_{0,j}$ . The variable  $P$  is well behaved in the interplay between the two resonant waves  $s$  and  $c$ . Unfortunately, this behavior is disrupted by the third family of waves, the  $k$  waves, except in the case where very strong hypothesis are assumed on the flux as in [4]. In fact, in [4] it is assumed that the point of maximum of  $g$  does not change with  $k$  (actually in [4] our  $k$  waves correspond to discontinuities in time because of Lagrangian coordinates). Since such assumptions are not realistic for our model, we are not able to prove a bound on the total variation of  $P$  uniformly in  $\varepsilon$ . Instead, we resolve this difficulty by applying a compensated compactness argument.

Up to now, all the values of  $(s^\varepsilon, c^\varepsilon, k^\varepsilon)$  are determined for  $x < \bar{x}_1$ . Since  $k^\varepsilon$  is constant in time, its value at the right of  $\bar{x}_1$  is known. The jump conditions  $\Delta f = \Delta c = 0$  determine all the values of  $(s^\varepsilon, c^\varepsilon, k^\varepsilon)$  to the right of  $\bar{x}_1$ . These values could introduce new values for the function  $g$  that must be added to the grid, i.e.,

$$\mathcal{G}_1 = \mathcal{G}_0 \cup \{g(s^\varepsilon(t, \bar{x}_1+), c^\varepsilon(t, \bar{x}_1+), k^\varepsilon(\bar{x}_1+)) \mid t \geq 0\}.$$

This gives the new allowed values for  $s$  for  $\bar{x}_1 < x < \bar{x}_2$ :

$$\mathcal{S}_{1,j} = \{\sigma \mid g(\sigma, c_j, k_1) \in \mathcal{G}_1\}.$$

Using these values we now build the corresponding approximations  $f^{1,j}(\sigma)$  of the flux as the linear interpolation of  $f(\sigma, c_j, k_1)$  according to the points in  $\mathcal{S}_{1,j}$ . Then we solve, as before, all the Riemann problems at  $t=0$ ,  $\bar{x}_1 < x < \bar{x}_2$  and  $t \geq 0$ ,  $x = \bar{x}_1$ . As before  $c$  waves cannot interact with each other so, by induction, we can carry out the wave front tracking algorithm on the semi plane  $t \geq 0$ .

We define the open regions  $(\bar{x}_0 = y_0(t) = -\infty, \bar{x}_{N+1} = y_{M+1}(t) = +\infty)$

$$\Omega_{i,j} = \{(t, x) \in [0, +\infty) \times \mathbb{R} \mid \bar{x}_i < x < \bar{x}_{i+1} \wedge y_j(t) < x < y_{j+1}(t)\}.$$

The wave front tracking approximations so obtained are weak entropic solutions to

$$\begin{cases} \partial_t s^\varepsilon + \partial_x [F^\varepsilon(t, x, s^\varepsilon)] = 0, \\ \partial_t (c^\varepsilon s^\varepsilon) + \partial_x [c^\varepsilon F^\varepsilon(t, x, s^\varepsilon)] = 0, \\ \partial_t k^\varepsilon = 0, \\ (s^\varepsilon, c^\varepsilon, k^\varepsilon)(0, x) = (\bar{s}^\varepsilon, \bar{c}^\varepsilon, \bar{k}^\varepsilon)(x), \end{cases} \quad (2.11)$$

where the flux  $F^\varepsilon$  is defined by

$$F^\varepsilon(t, x, \sigma) = f^{i,j}(\sigma), \text{ for } (t, x) \in \Omega_{i,j}.$$

For all  $(t, x) \in [0, +\infty[ \times \mathbb{R}$  and  $\sigma \in [0, 1]$ , the flux satisfies the estimates

$$\begin{cases} |F^\varepsilon(t, x, \sigma) - f(\sigma, c^\varepsilon(t, x), k^\varepsilon(x))| \leq \varepsilon, \\ |\partial_\sigma F^\varepsilon(t, x, \sigma) - \partial_\sigma f(\sigma, c^\varepsilon(t, x), k^\varepsilon(x))| \leq \frac{\varepsilon}{N+M}. \end{cases} \quad (2.12)$$

We remark that, in any region  $\Omega_{i,j}$ ,  $s^\varepsilon$  is an entropic solution to the scalar conservation law

$$\partial_t s^\varepsilon + \partial_x [f^{i,j}(s^\varepsilon)] = 0.$$

### 3 Entropy estimates

Given a smooth (not necessarily convex) entropy function  $\eta(\sigma)$  with  $\eta(0) = 0$ , we define the corresponding entropy flux  $q^\varepsilon$  (relative to the approximate flux  $F^\varepsilon$ ) as

$$q^\varepsilon(t, x, \sigma) = \int_0^\sigma \eta'(\zeta) \partial_\zeta F^\varepsilon(t, x, \zeta) d\zeta. \quad (3.13)$$

**Theorem 3.1.** *For a fixed **convex** entropy  $\eta$ , the positive part of the measure*

$$\mu_\varepsilon = \partial_t [\eta(s^\varepsilon)] + \partial_x [q^\varepsilon(t, x, s^\varepsilon)] \quad (3.14)$$

*is uniformly (with respect to the approximation parameter  $\varepsilon$ ) locally bounded. More precisely, for any compact set  $K \subset ]0, +\infty[ \times \mathbb{R}$  there exists a constant  $C_K$  such that*

$$\mu_\varepsilon^+(K) \leq C_K.$$

*Here the constant  $C_K$  may depend on  $\eta$ ,  $f$  and on the total variation of the initial data  $\bar{c}$  and  $\bar{k}$ , but it does not depend on the approximation parameter  $\varepsilon$ .*

*Proof.* Fixing a non negative test function  $\phi \in C_c^\infty([0, +\infty[ \times \mathbb{R})$ , we compute

$$\begin{aligned} \langle \partial_t [\eta(s^\varepsilon)] + \partial_x [q^\varepsilon(t, x, s^\varepsilon)], \phi \rangle &= - \int \eta(s^\varepsilon) \partial_t \phi + q^\varepsilon(t, x, s^\varepsilon) \partial_x \phi \, dt dx \\ &= \int_0^{+\infty} \sum_{\ell=1}^{\mathcal{N}(t)} \left[ \Delta q_\ell^\varepsilon(t) - \Delta \eta_\ell(t) \dot{\xi}_\ell(t) \right] \phi(t, \xi_\ell(t)) \, dt. \end{aligned} \quad (3.15)$$

Here  $\xi_1, \dots, \xi_{\mathcal{N}(t)}$  are the locations of the discontinuities in  $(s^\varepsilon, c^\varepsilon, k^\varepsilon)$ , and the notation  $\Delta$  denotes the jumps:

$$\begin{cases} \Delta \eta_\ell(t) = \eta(s^\varepsilon(t, \xi_\ell(t) +)) - \eta(s^\varepsilon(t, \xi_\ell(t) -)), \\ \Delta q_\ell^\varepsilon(t) = q^\varepsilon(t, \xi_\ell(t) +, s^\varepsilon(t, \xi_\ell(t) +)) - q^\varepsilon(t, \xi_\ell(t) -, s^\varepsilon(t, \xi_\ell(t) -)). \end{cases}$$

We study separately the three different kinds of waves and denote with the superscripts “−” and “+” the values computed respectively to the left and the right of the discontinuities. We omit the superscript for the values that do not change across the discontinuities.

**s waves:** Both  $c$  and  $k$  are constant while the jump in  $s$  satisfies Rankine-Hugoniot and is entropic according to the approximate flux. If  $(t, \xi_\ell(t)) \in \Omega_{i,j}$ , then

$$\dot{\xi}_\ell(s^+ - s^-) = f^{i,j}(s^+) - f^{i,j}(s^-) = f^+ - f^-.$$

Hence, applying the definition of  $q^\varepsilon$  and integrating by parts, we compute

$$\begin{aligned} \Delta q_\ell^\varepsilon - \Delta \eta_\ell \dot{\xi}_\ell &= \int_{s^-}^{s^+} \eta'(\varsigma) \left[ \partial_\varsigma f^{i,j}(\varsigma) - \dot{\xi}_\ell \right] d\varsigma \\ &= \left[ \eta'(\varsigma) \left( f^{i,j}(\varsigma) - f^- - \dot{\xi}_\ell(\varsigma - s^-) \right) \right]_{s^-}^{s^+} \\ &\quad - \int_{s^-}^{s^+} \eta''(\varsigma) \left[ f^{i,j}(\varsigma) - f^- - \dot{\xi}_\ell(\varsigma - s^-) \right] d\varsigma \\ &\leq 0. \end{aligned}$$

Since  $\eta'' \geq 0$  and the  $s$  wave in  $\xi_\ell$  is an entropic wave for the flux  $f^{i,j}$ , therefore for all  $\varsigma \in [\min\{s^-, s^+\}, \max\{s^-, s^+\}]$  one has

$$\text{sign}(s^+ - s^-) \left[ f^{i,j}(\varsigma) - f^- - \frac{f^+ - f^-}{s^+ - s^-} (\varsigma - s^-) \right] \geq 0.$$

**c waves:** Both  $k$  and  $g = \frac{f}{s}$  are constants and the speed  $\dot{\xi}_\ell$  of the wave equals  $g(s^-, c^-, k) = g(s^+, c^+, k)$ , where  $\xi_\ell$  is the boundary between the regions  $\Omega_{i,j}$  and  $\Omega_{i,j+1}$ . Denoting by  $C$  a generic constant that depends only on  $\eta$  and  $f$ , the uniform esti-

mates (2.12) lead to

$$\begin{aligned}
\Delta q_{\ell}^{\varepsilon} - \Delta \eta_{\ell} \dot{\xi}_{\ell} &= \int_0^{s^+} \eta'(\varsigma) \partial_{\varsigma} f^{i,j+1}(\varsigma) d\varsigma - \int_0^{s^-} \eta'(\varsigma) \partial_{\varsigma} f^{i,j}(\varsigma) d\varsigma - \dot{\xi}_{\ell} (\eta(s^+) - \eta(s^-)) \\
&\leq C \frac{\varepsilon}{N+M} + \int_0^{s^+} \eta'(\varsigma) \partial_{\varsigma} f(\varsigma, c^+, k) d\varsigma - \int_0^{s^-} \eta'(\varsigma) \partial_{\varsigma} f(\varsigma, c^-, k) d\varsigma \\
&\quad - \int_0^{s^+} \eta'(\varsigma) \dot{\xi}_{\ell} d\varsigma + \int_0^{s^-} \eta'(\varsigma) \dot{\xi}_{\ell} d\varsigma \\
&\leq C \frac{\varepsilon}{N+M} + \int_0^{s^+} \eta'(\varsigma) [\partial_{\varsigma} f(\varsigma, c^+, k) - \dot{\xi}_{\ell}] d\varsigma - \int_0^{s^-} \eta'(\varsigma) [\partial_{\varsigma} f(\varsigma, c^-, k) - \dot{\xi}_{\ell}] d\varsigma \\
&\leq C \frac{\varepsilon}{N+M} - \int_0^{s^+} \eta''(\varsigma) [f(\varsigma, c^+, k) - \dot{\xi}_{\ell} \varsigma] d\varsigma + \int_0^{s^-} \eta''(\varsigma) [f(\varsigma, c^-, k) - \dot{\xi}_{\ell} \varsigma] d\varsigma.
\end{aligned}$$

Here we have integrated by parts and used the relations

$$f(0, c^{\pm}, k) = 0, \quad f(s^{\pm}, c^{\pm}, k) = s^{\pm} g(s^{\pm}, c^{\pm}, k) = s^{\pm} \dot{\xi}_{\ell}.$$

Suppose  $s^- \leq s^+$ , the other case being symmetric. Because of the entropy condition on  $c$  waves (2.9) there exists  $s^* \in [s^-, s^+]$  such that

$$\begin{cases} g(\varsigma, c^-, k) \geq \dot{\xi}_{\ell} & \text{for all } \varsigma \in [s^-, s^*], \\ g(\varsigma, c^+, k) \geq \dot{\xi}_{\ell} & \text{for all } \varsigma \in [s^*, s^+]. \end{cases}$$

The estimates (2.12) further lead to

$$\begin{aligned}
\Delta q_{\ell}^{\varepsilon} - \Delta \eta_{\ell} \dot{\xi}_{\ell} &\leq - \int_{s^*}^{s^+} \eta''(\varsigma) [f(\varsigma, c^+, k) - \dot{\xi}_{\ell} \varsigma] d\varsigma + \int_{s^*}^{s^-} \eta''(\varsigma) [f(\varsigma, c^-, k) - \dot{\xi}_{\ell} \varsigma] d\varsigma \\
&\quad + C \left( \frac{\varepsilon}{N+M} + |c^+ - c^-| \right) \\
&= - \int_{s^*}^{s^+} \eta''(\varsigma) \varsigma [g(\varsigma, c^+, k) - \dot{\xi}_{\ell}] d\varsigma + \int_{s^*}^{s^-} \eta''(\varsigma) \varsigma [g(\varsigma, c^-, k) - \dot{\xi}_{\ell}] d\varsigma \\
&\quad + C \left( \frac{\varepsilon}{N+M} + |c^+ - c^-| \right) \\
&\leq C \left( \frac{\varepsilon}{N+M} + |\Delta c_{\ell}| \right).
\end{aligned}$$

**$k$  waves:** For a  $k$  wave, both  $c$  and  $f$  are constant and  $\dot{\xi}_{\ell} = 0$ , where  $\xi_{\ell}$  is the boundary

between two regions  $\Omega_{i,j}$  and  $\Omega_{i+1,j}$ . We have

$$\begin{aligned}
\Delta q_\ell^\varepsilon - \Delta \eta_\ell \dot{\xi}_\ell &= \int_0^{s^+} \eta'(\varsigma) \partial_\varsigma f^{i+1,j}(\varsigma) d\varsigma - \int_0^{s^-} \eta'(\varsigma) \partial_\varsigma f^{i,j}(\varsigma) d\varsigma \\
&\leq C \frac{\varepsilon}{N+M} + \int_0^{s^+} \eta'(\varsigma) \partial_\varsigma f(\varsigma, c, k^+) d\varsigma - \int_0^{s^-} \eta'(\varsigma) \partial_\varsigma f(\varsigma, c, k^-) d\varsigma \\
&\leq C \left( \frac{\varepsilon}{N+M} + |k^+ - k^-| \right) + \int_{s^-}^{s^+} \eta'(\varsigma) \partial_\varsigma f(\varsigma, c, k^-) d\varsigma \\
&\leq C \left( \frac{\varepsilon}{N+M} + |\Delta k_\ell| \right) + \|\eta'\|_\infty \text{sign}(s^+ - s^-) \int_{s^-}^{s^+} \partial_\varsigma f(\varsigma, c, k^-) d\varsigma \\
&= C \left( \frac{\varepsilon}{N+M} + |\Delta k_\ell| + |f(s^+, c, k^-) - f(s^-, c, k^-)| \right) \\
&= C \left( \frac{\varepsilon}{N+M} + |\Delta k_\ell| + |f(s^+, c, k^-) - f(s^+, c, k^+)| \right) \\
&= C \left( \frac{\varepsilon}{N+M} + |\Delta k_\ell| \right)
\end{aligned}$$

where we used the fact that  $\partial_\varsigma f(\varsigma, c, k^-) \geq 0$  and that  $f(s^-, c, k^-) = f(s^+, c, k^+)$ .

Finally, if the compact support of  $\phi$  is contained in  $]0, T[ \times \mathbb{R}$ , equality (3.15) and the previous analysis on the three types of waves lead to

$$\begin{aligned}
\langle \partial_t [\eta(s^\varepsilon)] + \partial_x [q^\varepsilon(t, x, s^\varepsilon)], \phi \rangle &\leq CT \left( \frac{N\varepsilon + M\varepsilon}{N+M} + \text{Tot. Var. } \{\bar{c}\} + \text{Tot. Var. } \{\bar{k}\} \right) \|\phi\|_\infty \\
&\leq CT (1 + \text{Tot. Var. } \{\bar{c}\} + \text{Tot. Var. } \{\bar{k}\}) \|\phi\|_\infty
\end{aligned}$$

for any  $\varepsilon \in ]0, 1[$ , proving the theorem.  $\square$

**Theorem 3.2.** *For any smooth entropy  $\eta$  (even non convex) and decreasing sequence  $\varepsilon_j \rightarrow 0$  there exists a compact set  $\mathcal{K} \subset H_{loc}^{-1}(\Omega)$ , independent of  $j$ , such that*

$$\mu_{\varepsilon_j} = \partial_t [\eta(s^{\varepsilon_j})] + \partial_x [q^{\varepsilon_j}(t, x, s^{\varepsilon_j})] \in \mathcal{K}.$$

*Proof.* We apply standard arguments in compensated compactness theory [6]. Integrating the measure  $\mu_\varepsilon$  over a rectangle (with  $t_1 > 0$ )  $R = [t_1, t_2] \times [-L, L]$  we obtain

$$\begin{aligned}
\mu_\varepsilon(R) &= \int_{t_1}^{t_2} q^\varepsilon(t, L+, s^\varepsilon(t, L+)) - q^\varepsilon(t, -L-, s^\varepsilon(t, -L-)) dt \\
&\quad + \int_{-L}^L \eta(s^\varepsilon(t_2+, x)) - \eta(s^\varepsilon(t_1-, x)) dx.
\end{aligned}$$

Since  $s^\varepsilon$  is uniformly bounded, there exists a constant  $\bar{C}_R$  such that  $|\mu_\varepsilon(R)| \leq \bar{C}_R$  for any  $\varepsilon \in ]0, 1[$ . If  $\eta$  is convex, we can apply Theorem 3.1 to estimate the total variation of  $\mu_\varepsilon$  uniformly with respect to  $\varepsilon$ :

$$|\mu_\varepsilon|(R) = \mu_\varepsilon^+(R) + \mu_\varepsilon^-(R) = 2\mu_\varepsilon^+(R) - \mu_\varepsilon(R) \leq 2C_R + \bar{C}_R.$$

If  $\eta$  is not convex, then we take a strictly convex entropy  $\eta^*$  (for instance  $\eta^*(\sigma) = \sigma^2$ ) and define  $\tilde{\eta} = \eta + H\eta^*$ . The entropy  $\tilde{\eta}$  is convex for a sufficiently big constant  $H$ . We

denote by  $\mu_\varepsilon$ ,  $\mu_\varepsilon^*$  and  $\tilde{\mu}_\varepsilon$  the measures corresponding to the entropies  $\eta$ ,  $\eta^*$  and  $\tilde{\eta}$ . Since the definition of the entropy flux (3.13) is linear with respect to the associated entropy, the measures satisfy  $\tilde{\mu}_\varepsilon = \mu_\varepsilon + H\mu_\varepsilon^*$ . Hence the inequality

$$|\mu_\varepsilon|(R) \leq |\tilde{\mu}_\varepsilon|(R) + H|\mu_\varepsilon^*|(R)$$

holds. This means that  $|\mu_\varepsilon|(R)$  is bounded uniformly with respect to  $\varepsilon$  since both  $\tilde{\mu}_\varepsilon$  and  $\mu_\varepsilon^*$  are associated with convex entropies. Since the measure  $\mu_\varepsilon = \partial_t[\eta(s^\varepsilon)] + \partial_x[q^\varepsilon(t, x, s^\varepsilon)]$  restricted to  $R$  lies both in a bounded set of the space of measures  $\mathcal{M}(R)$  and in a bounded set of  $W^{-1, \infty}(R)$ , [5, Lemma 17.2.2] allows us to conclude the proof of the theorem.  $\square$

## 4 Strong Convergence

The following result is a step towards the proof of Theorem 1.1.

**Theorem 4.1.** *There exists a sequence  $\varepsilon_j \rightarrow 0$  such that  $(s^{\varepsilon_j}, c^{\varepsilon_j}, k^{\varepsilon_j}) \rightarrow (\tilde{s}, \tilde{c}, \tilde{k})$  in  $L^1_{loc}(\Omega)$ .*

*Proof.* We suitably modify the proof of [2, Theorem 4.2], omitting some computations already written there. The proof takes several steps.

1. Observe that by construction we have

$$\text{Tot. Var. } \{c^\varepsilon(t, \cdot)\} = \text{Tot. Var. } \{\tilde{c}^\varepsilon\} \leq \text{Tot. Var. } \{\tilde{c}\}$$

and the wave speeds are uniformly bounded. Hence Helly's theorem implies that there exist a sequence  $c^{\varepsilon_j} \rightarrow \tilde{c}$  in  $L^1_{loc}(\Omega)$ . Since  $k^\varepsilon$  is constant in time, we have  $k^{\varepsilon_j} \rightarrow \tilde{k} = \bar{k}$  in  $L^1_{loc}(\Omega)$  as well. In the following we always take subsequences of this sequence and we will drop the index  $j$  to simplify notations. We define the limit flux

$$F(t, x, \sigma) = f\left(\sigma, \tilde{c}(t, x), \tilde{k}(x)\right), \quad \text{for all } (t, x) \in \Omega, \text{ and } \sigma \in [0, 1]$$

and for any entropy  $\eta$  we define the limit entropy flux

$$q(t, x, \sigma) = \int_0^\sigma \eta'(\varsigma) \partial_\varsigma F(t, x, \varsigma) d\varsigma.$$

The estimate (uniform in  $\sigma \in [0, 1]$ )

$$\begin{aligned} |q(t, x, \sigma) - q^\varepsilon(t, x, \sigma)| &\leq \int_0^1 |\eta'(\varsigma)| \left( \left| \partial_\varsigma f\left(\varsigma, \tilde{c}(t, x), \tilde{k}(x)\right) - \partial_\varsigma f\left(\varsigma, c^\varepsilon(t, x), k^\varepsilon(x)\right) \right| \right. \\ &\quad \left. + \left| \partial_\varsigma f\left(\varsigma, c^\varepsilon(t, x), k^\varepsilon(x)\right) - \partial_\varsigma F^\varepsilon(t, x, \varsigma) \right| \right) d\varsigma \\ &\leq C \left( |\tilde{c}(t, x) - c^\varepsilon(t, x)| + |\tilde{k}(x) - k^\varepsilon(x)| + \varepsilon \right) \rightarrow 0 \quad \text{in } L^1_{loc}(\Omega) \end{aligned}$$

implies that

$$\partial_x [q(t, x, s^\varepsilon) - q^\varepsilon(t, x, s^\varepsilon)] \rightarrow 0, \quad \text{in } H^{-1}_{loc}(\Omega).$$

Together with Theorem 3.2, it implies that the sequence

$$\partial_t [\eta(s^\varepsilon)] + \partial_x [q(t, x, s^\varepsilon)] = \partial_t [\eta(s^\varepsilon)] + \partial_x [q^\varepsilon(t, x, s^\varepsilon)] + \partial_x [q(t, x, s^\varepsilon) - q^\varepsilon(t, x, s^\varepsilon)]$$

belongs to a compact set in  $H^{-1}_{loc}(\Omega)$ .

2. For any  $(t, x) \in \Omega$  and  $v, w \in [0, 1]$  we define

$$I(t, x, v, w) \doteq (v-w) \int_w^v [\partial_\sigma F(t, x, \sigma)]^2 d\sigma - [F(t, x, v) - F(t, x, w)]^2. \quad (4.16)$$

The following properties hold.

- (i)  $(v, w) \mapsto I(t, x, v, w)$  is continuous with  $I(t, x, v, v) = 0$  for any  $v \in [0, 1]$ .
- (ii)  $I(t, x, v, w) > 0$  for any  $v, w \in [0, 1]$  with  $v \neq w$ .

Indeed, (i) is trivial, while (ii) follows from Jensen's inequality and the fact that  $\sigma \mapsto f(\sigma, \gamma, \kappa)$  and hence  $\sigma \mapsto F(t, x, \sigma)$  have a unique inflection point. Indeed suppose  $w < v$ , we observe that  $\sigma \mapsto \partial_\sigma F(t, x, \sigma)$  is not constant over the interval  $\omega \in [w, v]$ , and we compute

$$\begin{aligned} I(t, x, v, w) &= (v-w) \int_w^v [\partial_\sigma F(t, x, \sigma)]^2 d\sigma - (v-w)^2 \left[ \frac{1}{v-w} \int_w^v \partial_\sigma F(t, x, \sigma) d\sigma \right]^2 \\ &> (v-w) \int_w^v [\partial_\sigma F(t, x, \sigma)]^2 d\sigma - (v-w)^2 \frac{1}{v-w} \int_w^v [\partial_\sigma F(t, x, \sigma)]^2 d\sigma \\ &= 0. \end{aligned}$$

3. Fixing  $(\tau, y) \in \Omega$  and we consider the following entropies and corresponding limit fluxes

$$\begin{aligned} \eta(\sigma) &= \sigma, & q(t, x, \sigma) &= F(t, x, \sigma), \\ \eta_{(\tau, y)}(\sigma) &= F(\tau, y, \sigma), & q_{(\tau, y)}(t, x, \sigma) &= \int_0^\sigma \partial_\varsigma F(\tau, y, \varsigma) \partial_\varsigma F(t, x, \varsigma) d\varsigma. \end{aligned}$$

The same computations as the ones used to obtain [2, (4.16)] prove that there exists a constant  $C_2 \geq 0$  such that

$$\begin{aligned} (v-w) [q_{(\tau, y)}(t, x, v) - q_{(\tau, y)}(t, x, w)] \\ \geq I(t, x, v, w) + [F(t, x, v) - F(t, x, w)]^2 - C_2 \sup_{\sigma \in [0, 1]} |F(\tau, y, \sigma) - F(t, x, \sigma)|. \end{aligned} \quad (4.17)$$

4. By possibly taking subsequences, we can achieve the following weak\* convergences in  $L^\infty(\Omega)$ :

$$\begin{cases} s^\varepsilon(t, x) \xrightarrow{*} \tilde{s}(t, x), \\ F(t, x, s^\varepsilon(t, x)) \xrightarrow{*} \tilde{F}(t, x), \\ I(t, x, s^\varepsilon(t, x), \tilde{s}(t, x)) \xrightarrow{*} \tilde{I}(t, x). \end{cases} \quad (4.18)$$

Taking further subsequences (which this time may depend on  $(\tau, y)$ ) we can achieve these further weak\* convergences in  $L^\infty(\Omega)$

$$F(\tau, y, s^\varepsilon(t, x)) \xrightarrow{*} \tilde{F}_{(\tau, y)}(t, x), \quad q_{(\tau, y)}(t, x, s^\varepsilon(t, x)) \xrightarrow{*} \tilde{q}_{(\tau, y)}(t, x). \quad (4.19)$$

Notice that the weak limits  $\tilde{s}$ ,  $\tilde{f}$ ,  $\tilde{I}$  in (4.18) do not depend on the values  $(\tau, y)$ . Step 1 implies

$$\partial_t [s^\varepsilon(t, x)] + \partial_x [F(t, x, s^\varepsilon(t, x))], \quad \partial_t [F(\tau, y, s^\varepsilon(t, x))] + \partial_x [q_{(\tau, y)}(t, x, s^\varepsilon(t, x))] \in \mathcal{K},$$

where  $\mathcal{K}$  is a compact set (independent of the subsequence index) in  $H_{loc}^{-1}(\Omega)$ . By an application of the *div-curl lemma*, see for example Theorem 16.2.1 in [5], one obtains

$$\begin{aligned} s^\varepsilon(t, x)q_{(\tau, y)}(t, x, s^\varepsilon(t, x)) - F(t, x, s^\varepsilon(t, x))F(\tau, y, s^\varepsilon(t, x)) \\ \xrightarrow{*} \tilde{s}(t, x)\tilde{q}_{(\tau, y)}(t, x) - \tilde{F}(t, x)\tilde{F}_{(\tau, y)}(t, x). \end{aligned} \quad (4.20)$$

Following the proof of [2, Theorem 4.2] we set  $v = s^\varepsilon(t, x)$  and  $w = \tilde{s}(t, x)$  in (4.17) and take the weak\* limit as  $\varepsilon \rightarrow 0$  to obtain

$$\begin{aligned} \tilde{I}(t, x) - \left[ \tilde{s}(t, x)\tilde{q}_{(\tau, y)}(t, x) - \tilde{F}(t, x)\tilde{F}_{(\tau, y)}(t, x) \right] + \tilde{s}(t, x)\tilde{q}_{(\tau, y)}(t, x) \\ - 2\tilde{F}(t, x)F(t, x, \tilde{s}(t, x)) + F(t, x, \tilde{s}(t, x))^2 \leq C_3 \sup_{\sigma \in [0, 1]} |F(\tau, y, \sigma) - F(t, x, \sigma)|. \end{aligned}$$

This can be written as

$$\begin{aligned} \tilde{I}(t, x) + \left[ \tilde{F}(t, x) - F(t, x, \tilde{s}(t, x)) \right]^2 \leq C_3 \sup_{\sigma \in [0, 1]} |F(\tau, y, \sigma) - F(t, x, \sigma)| \\ + \left| \tilde{F}(t, x) \right| \left| \tilde{F}_{(\tau, y)}(t, x) - \tilde{F}(t, x) \right|, \end{aligned}$$

which holds for any fixed  $(\tau, y) \in \Omega$  and a.e.  $(t, x) \in \Omega$ . Taking the weak\* limit in

$$\begin{aligned} - \sup_{\sigma \in [0, 1]} |F(\tau, y, \sigma) - F(t, x, \sigma)| \leq F(\tau, y, s^\varepsilon(t, x)) - F(t, x, s^\varepsilon(t, x)) \\ \leq \sup_{\sigma \in [0, 1]} |F(\tau, y, \sigma) - F(t, x, \sigma)|, \end{aligned}$$

we obtain

$$\begin{aligned} - \sup_{\sigma \in [0, 1]} |F(\tau, y, \sigma) - F(t, x, \sigma)| \leq \tilde{F}_{(\tau, y)}(t, x) - \tilde{F}(t, x) \\ \leq \sup_{\sigma \in [0, 1]} |F(\tau, y, \sigma) - F(t, x, \sigma)|. \end{aligned}$$

Hence for any fixed  $(\tau, y) \in \Omega$ , we have for a.e.  $(t, x) \in \Omega$

$$\tilde{I}(t, x) + \left[ \tilde{F}(t, x) - F(t, x, \tilde{s}(t, x)) \right]^2 \leq C_4 \sup_{\sigma \in [0, 1]} |F(\tau, y, \sigma) - F(t, x, \sigma)|. \quad (4.21)$$

**4.** We call  $E_1$  the set of Lebesgue points of the left hand side of (4.21). Moreover, for each  $\sigma \in [0, 1]$  let  $E_\sigma$  be the set of Lebesgue points of the map  $(t, x) \mapsto F(t, x, \sigma)$ . Defining

$$E \doteq E_1 \cap \left( \bigcap_{q \in \mathbb{Q} \cap [0, 1]} E_q \right),$$

we observe that its complement  $\Omega \setminus E$  has zero measure. Take any  $(\tau, y) \in E$  and fix  $\epsilon > 0$ . Let  $\mathcal{F}_\epsilon \subset \mathbb{Q} \cap [0, 1]$  be a finite set such that  $\inf_{q \in \mathcal{F}_\epsilon} |q - \sigma| < \epsilon$  for every  $\sigma \in [0, 1]$ . Then we have

$$\sup_{\sigma \in [0, 1]} |F(\tau, y, \sigma) - F(t, x, \sigma)| \leq \max_{q \in \mathcal{F}_\epsilon} |F(\tau, y, q) - F(t, x, q)| + 2L\epsilon$$



$$\leq \sum_{q \in \mathcal{F}_\epsilon} |F(\tau, y, q) - F(t, x, q)| + 2L\epsilon, \quad (4.22)$$

where  $L$  is a uniform Lipschitz constant for  $\varsigma \mapsto F(t, x, \varsigma)$ . Let  $B_\delta(\tau, y)$  be the disc in  $\Omega$  centered in  $(\tau, y)$  with radius  $\delta > 0$  whose area is  $\pi\delta^2$ . Integrating (4.21) and using (4.22) we obtain

$$\begin{aligned} & \frac{1}{\pi\delta^2} \int_{B_\delta(\tau, y)} \left( \tilde{I}(t, x) + [\tilde{F}(t, x) - F(t, x, \tilde{s}(t, x))]^2 \right) dt dx \\ & \leq \frac{C_4}{\pi\delta^2} \sum_{q \in \mathcal{F}_\epsilon} \int_{B_\delta(\tau, y)} |F(\tau, y, q) - F(t, x, q)| dt dx + 2C_4L\epsilon. \end{aligned}$$

Since  $(\tau, y)$  is a Lebesgue point for the map  $(t, x) \mapsto F(t, x, q)$ , for all  $q \in \mathcal{F}_\epsilon$ , letting  $\delta \rightarrow 0$  we obtain

$$\tilde{I}(\tau, y) + [\tilde{F}(\tau, y) - F(\tau, y, \tilde{s}(\tau, y))]^2 \leq C_4L\epsilon.$$

Since  $\epsilon > 0$  is arbitrary, this implies

$$\tilde{I}(\tau, y) + [\tilde{F}(\tau, y) - F(\tau, y, \tilde{s}(\tau, y))]^2 \leq 0 \quad \text{for every } (\tau, y) \in E.$$

Hence  $\tilde{I}(t, x) \leq 0$  a.e. in  $\Omega$ . Since, by Step 2,  $I(t, x, s^\epsilon(t, x), \tilde{s}(t, x)) \geq 0$ , its weak\* limit  $\tilde{I}(t, x)$  must be greater or equal to zero almost everywhere. Therefore we get

$$\tilde{I}(t, x) = 0, \quad \text{and} \quad \tilde{F}(t, x) = F(t, x, \tilde{s}(t, x)), \quad \text{a.e. in } \Omega.$$

Since  $I(t, x, s^\epsilon(t, x), \tilde{s}(t, x)) \geq 0$  converges weakly\* to zero, we conclude that it converges strongly in  $L^1_{loc}(\Omega)$ . We can thus take a subsequence such that  $I(t, x, s^\epsilon(t, x), \tilde{s}(t, x)) \rightarrow 0$  a.e. in  $\Omega$ . Finally, property (ii) proved in Step 2 implies  $s^\epsilon(t, x) \rightarrow \tilde{s}(t, x)$  a.e. in  $\Omega$ , completing the proof.  $\square$

*Proof of Theorem 1.1.* By Theorem 4.1 we know that there exists a subsequence of wave front tracking approximate solutions constructed in Section 2 ( $s^\epsilon, c^\epsilon, k^\epsilon$ ) which converges strongly in  $L^1_{loc}(\Omega)$  to a limit  $(\tilde{s}, \tilde{c}, \tilde{k})$ . Clearly  $\tilde{k}_t = 0$ . Let  $\phi$  be a test function with compact support in  $[0, +\infty[ \times \mathbb{R}$ . By construction (see Section 2) the approximate solutions satisfy

$$\begin{aligned} & \int_{\Omega} [s^\epsilon \phi_t + F^\epsilon(t, x, s^\epsilon) \phi_x](t, x) dt dx + \int_{\mathbb{R}} \bar{s}^\epsilon(x) \phi(0, x) dx = 0, \\ & \int_{\Omega} [c^\epsilon s^\epsilon \phi_t + c^\epsilon F^\epsilon(t, x, s^\epsilon) \phi_x](t, x) dt dx + \int_{\mathbb{R}} \bar{c}^\epsilon(x) \bar{s}^\epsilon(x) \phi(0, x) dx = 0, \\ & k^\epsilon(t, x) = \bar{k}^\epsilon(x), \quad \forall (t, x) \in \Omega. \end{aligned}$$

The uniform estimate (2.12) and the strong convergence of approximate solutions allows us to pass to the limit and to conclude that the limit  $(\tilde{s}, \tilde{c}, \tilde{k})$  satisfies Definition 1.1.  $\square$

**Acknowledgment:** The present work was supported by the PRIN 2015 project *Hyperbolic Systems of Conservation Laws and Fluid Dynamics: Analysis and Applications* and by GNAMPA 2019 project *Equazioni alle derivate parziali di tipo iperbolico o non locale ed applicazioni*. The authors would like to thank the anonymous referee for carefully reading the manuscript and providing many useful suggestions.

## References

- [1] Alberto Bressan, *Hyperbolic systems of conservation laws*, Oxford Lecture Series in Mathematics and its Applications, vol. 20, Oxford University Press, Oxford, 2000. The one-dimensional Cauchy problem. MR1816648 ↑[2](#), [2](#)
- [2] Alberto Bressan, Graziano Guerra, and Wen Shen, *Vanishing viscosity solutions for conservation laws with regulated flux*, J. Differential Equations **266** (2019), no. 1, 312–351. MR3870565 ↑[1](#), [4](#), [4](#), [4](#)
- [3] S.E. Buckley and M. Leverett, *Mechanism of fluid displacement in sands*, Transactions of the AIME **146** (1942), 107–116. ↑[1](#)
- [4] Giuseppe Maria Coclite and Nils Henrik Risebro, *Conservation laws with time dependent discontinuous coefficients*, SIAM J. Math. Anal. **36** (2005), no. 4, 1293–1309. MR2139451 ↑[\(document\)](#), [2](#), [2](#)
- [5] Constantine M. Dafermos, *Hyperbolic conservation laws in continuum physics*, Fourth, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 325, Springer-Verlag, Berlin, 2016. MR3468916 ↑[3](#), [4](#)
- [6] R. J. DiPerna, *Convergence of approximate solutions to conservation laws*, Arch. Rational Mech. Anal. **82** (1983), no. 1, 27–70. MR684413 ↑[3](#)
- [7] Tore Gimse and Nils Henrik Risebro, *Riemann problems with a discontinuous flux function*, Third International Conference on Hyperbolic Problems, Vol. I, II (Uppsala, 1990), 1991, pp. 488–502. MR1109304 ↑[2](#)
- [8] Eli L. Isaacson and J. Blake Temple, *Analysis of a singular hyperbolic system of conservation laws*, J. Differential Equations **65** (1986), no. 2, 250–268. MR861520 ↑[\(document\)](#)
- [9] Thormod Johansen and Ragnar Winther, *The solution of the Riemann problem for a hyperbolic system of conservation laws modeling polymer flooding*, SIAM J. Math. Anal. **19** (1988), no. 3, 541–566. MR937469 ↑[1](#)
- [10] Kenneth H. Karlsen, Michel Rascle, and Eitan Tadmor, *On the existence and compactness of a two-dimensional resonant system of conservation laws*, Commun. Math. Sci. **5** (2007), no. 2, 253–265. MR2334842 ↑[1](#)
- [11] Wen Shen, *On the Cauchy problems for polymer flooding with gravitation*, J. Differential Equations **261** (2016), no. 1, 627–653. MR3487271 ↑[\(document\)](#), [1](#), [2](#), [2](#), [2](#), [2](#)
- [12] ———, *On the uniqueness of vanishing viscosity solutions for Riemann problems for polymer flooding*, NoDEA Nonlinear Differential Equations Appl. **24** (2017), no. 4, Art. 37, 25. MR3663611 ↑[1](#)
- [13] ———, *Global Riemann solvers for several  $3 \times 3$  systems of conservation laws with degeneracies*, Math. Models Methods Appl. Sci. **28** (2018), no. 8, 1599–1626. MR3827966 ↑[1](#)
- [14] Blake Temple, *Global solution of the Cauchy problem for a class of  $2 \times 2$  nonstrictly hyperbolic conservation laws*, Adv. in Appl. Math. **3** (1982), no. 3, 335–375. MR673246 ↑[\(document\)](#)
- [15] David H. Wagner, *Equivalence of the Euler and Lagrangian equations of gas dynamics for weak solutions*, J. Differential Equations **68** (1987), no. 1, 118–136. MR885816 (88i:35100) ↑[1](#)